

8 Literaturverzeichnis

- Absolom, D. R., Zingg, W., Neumann, A. W. (1987):
Protein adsorption to polymer particles: role of surface properties.
J. Biomed. Mater. Res. 21, 161-171
- Aiello, L. P., Avery, R., Arrigg, R. G., Keyt, B. A., Jampel, H. D., Shah, S. T., Pasquale, L. R., Thieme, H., Iwamoto, M. A., Park, J. E. (1994):
Vascular endothelial growth factor in ocular fluid of patients with diabetic retinopathy and other retinal disorders.
N. Engl. J. Med. 331, 1480-1487
- Allemann, E., Gravel, P., Leroux, J.-C., Balant, L., Gurny, R. (1997):
Kinetics of blood component adsorption on poly(d,l-lactic acid) nanoparticles: evidence of complement C3 component involvement.
J. Biomed. Mater. Res. 37, 229-234
- Ali-Osman, F., Sriram, R., Raikar, A. (1991):
Inhibition of repair of DNA-interstrand cross links in human malignant astrocytoma cells: mediation by glutathione depletion.
AACR-Proc. 32, A 2509
- Altenschöpfer, P., Spruß, T., Bernhardt, G., Buschauer, A (1998):
Comparison of subcutaneously and orthotopically implanted glioblastomas.
Poster, Institut für Pharmazie der Universität Regensburg
- Alyautdin, R. N., Gothier, D., Petrov, V. E., Kharkevich, D. A., Kreuter, J. (1995):
Analgesic activity of the hexapeptide dalargin adsorbed on the surface of polysorbate 80-coated poly(butylcyanoacrylate) nanoparticles.
Eur. J. Pharm. Biopharm. 41, 44-48
- Alyautdin, R. N., Petrov, V. E., Langer, K., Berthold, A., Kharkevich, D. A., Kreuter, J. (1997):
Delivery of loperamide across the blood-brain barrier with polysorbate 80-coated polybutylcyanoacrylate nanoparticles.
J. Pharm. Res. 14, 325-328
- Ammoury, N., Fessi, H., Devissaguet, J.-P., Dubrasquet, M., Benita, S. (1991):
Jejunal absorption, pharmacological activity, and pharmacokinetic evaluation of indomethacin-loaded poly(d,l-lactide) and poly(isobutyl-cyanoacrylate) nanoparticles in rats.
Pharm. Res. 8, 101-105
- Anderson, N. L., Anderson, N. G. (1991):
A two-dimensional gel database of human plasma proteins.
Electrophoresis 12, 883-906
- Andreesen, R., Osterlaz, J., Bodemann, H. H., Bross, K. J., Costabel, U., Löhr, G. W. (1984):
Expression of transferrin receptors and intracellular ferritin during terminal differentiation of human monocytes.
Blut 49, 195-202
- Antman, K. (1996):
Hochdosis-Therapie und Knochenmark-Transplantation:
In Spektrum der Wissenschaft: Spezial 5: Krebsmedizin.65
- Armstrong, T. I., Davies, M. C. Illum, L. (1997):
Human serum albumin as a probe for protein adsorption to nanoparticles: relevance to biodistribution.
J. Drug Targ. 4, 6, 389-398
- Asahara, T., Bauters, C., Zheng, L. P., Takeshita, S., Bunting, S., Ferrara, N., Symes, J. S., Isner, J. M. (1995):
Synergistic effect of vascular endothelial growth factor and basic fibroblast growth factor on angiogenesis in vivo.
Circulation 92, 365-371

Astier, A., Doat, B., Ferrer, M.-J., Benoit, G., Fleury, J., Rolland, A., Leverage, R. (1988):
Enhancement of Adriamycin antitumor activity by its binding with an intracellular sustained-release form,
polymethacrylate nanospheres, in U-937 cells.
Cancer Res. 48, 1835-1841

Bates, S. R., Coughlin, B. A., Mazzone, T., Borensztajn, J., Getz, G. S. (1987):
Apoprotein E mediates the interaction of B-VLDL with macrophages.
J. Lipid Res. 28, 787-797

Bartsch, H. (1996):
DNA adducts in human carcinogenesis: etiological relevance and structure-activity relationship.
Mutat. Res. 385, 67-79

Bastert, G., Costa, S. D. (1996):
Spezielle Diagnostik und Therapie: Therapie des Mammakarzinoms.
In : Zeller, W. J., zur Hausen, H.: Onkologie: Grundlagen-Diagnostik-Therapie-Entwicklungen, ecomed
Verlagsgesellschaft AG&CoKG.1-34

Bazile, D.V., Ropert, C., Huve, P., Verrecchia, T., Marlard, M., Frydman, A., Veillard, M., Spenlehauer, G.
(1992):
Body distribution of fully biodegradable (¹⁴C)-poly(lactid acid) albumin after parenteral administration to rats.
Biomaterials 13, 15, 1093-1102

Beck, P. H. (1993):
Entwicklung und Charakterisierung von Nanopartikelformulierungen als Trägersysteme für Cytostatika.
Frankfurt am Main, Fachbereich Biochemie, Pharmazie und Lebensmittelchemie der J. W. Goethe- Universität,
Diss.

Beck, P., Kreuter, J., Reszka, R., Fichtner, I. (1993):
Influence of polybutylcyanoacrylate nanoparticles and liposomes on the efficacy and toxicity of the anticancer
drug mitoxantrone in murine tumor models.
J. Microencapsul. 10, 101-114

Becker, N., Wahrendorf, J. (1991):
Eine Bewertung aus epidemiologischer Sicht.
Dtsch. Ärztebl. 88, 3636-3640

Becker, G., Krone, A., Koulis, D., Lindner, A., Hofmann, E., Roggendorf, W., Bogdahn, U. (1994):
Reliability of transcranial colour-coded real-time sonography in assessment of brain tumours: correlation of
ultrasound, computed tomography and biopsy findings.
Neuroradiology 1994, 585-590

Bentele, V., Berg, U. E., Kreuter, J. (1983):
Molecular weights of poly(methylmethacrylate) nanoparticles.
Int. J. Pharm. 13, 109-113

Berg, U. E. (1979):
Immunstimulation durch hochdisperse Polymersuspension.
ETH Nr. 6481, Zürich, Diss., 31-49

Berndt, A. (1998):
mdl. Mitteilung

Berschneider, F. (1991):
In Wiesner, E., Ribbeck, R. (eds.): Wörterbuch der Veterinärmedizin, 797, 3. Aufl., Gustav Fischer Verlag Jena,
Stuttgart

Bertino, J., R. (1997):
Encyclopedia of cancer.
Volume I-III, Academic Press, San Diego-London-Boston-New York-Sydney-Tokyo-Toronto

Bigner, D. D., Bigner, S. H., Ponten, J., Westermark, B., Mahaley, M. S., Ruoslahti, E., Herschmann, H., Eng, L. F., Wikstrand, C. J. (1981):
Heterogeneity of genotypic and phenotypic characteristics of fifteen permanent cell lines derived from human gliomas.
J. Neuropathol. Exp. Neurol. 40, 201-229

Bitran, J. (1990):
High-dose chemotherapy with autologous hematopoietic stem cell support in the treatment of women with breast cancer: novel therapeutic approach or passing fad?
Oncology Journal Club 2, 4, 15-16

Black, P. M. (1991):
Brain tumors.
N. Engl. J. Med. 324, 1471-1476

Bleehen, N. M., Ford, J. M. (1993):
Radiotherapy, hyperthermia and photodynamic therapy for central nervous system tumors.
Curr. Opin. Oncol. 5, 458-463

Blunk, T., Hochstrasser, D.F., Sanchez, J., Müller, B.W., Müller, R.H. (1993):
Colloidal carriers for intravenous drug targeting: plasma protein adsorption patterns on surface-modified latex particles evaluated by two-dimensional polyacrylamide gel electrophoresis.
Electrophoresis 14, 1382-1387

Blunk, T. (1994):
Plasmaproteinadsorption auf kolloidalen Arzneistoffträgern.
Kiel, Mathematisch-Naturwissenschaftliche Fakultät der Christian-Albrechts-Universität, Diss.

Bodner, W. F. (1994):
Cancer genetics.
Br. Med. Bull. 50, 517-526

Bogdahn, U., Jachimczak, P. (1996):
Spezielle Diagnostik und Therapie: Maligne Gliome.
In : Zeller, W. J., zur Hausen, H.: Onkologie: Grundlagen-Diagnostik-Therapie-Entwicklungen, ecomed Verlagsgesellschaft AG&CoKG.1-22

Bonadonna, G., Gianni, A. M. (1990):
High-dose chemotherapy and autologous bone marrow transplant for adjuvant treatment of poor-risk breast cancer.
Oncology Journal Club 2, 4, 3-6

Borchard, G. (1993):
Untersuchungen von Interaktionen kolloidalen Arzneistoffträger (Nanopartikel) mit Bestandteilen des retikuloendothelialen Systems bei Ratten: Charakterisierung der die Körperverteilung beeinflussenden Faktoren.
Frankfurt am Main, Fachbereich Biochemie, Pharmazie und Lebensmittelchemie der J. W. Goethe- Universität, Diss.

Borchard, G., Kreuter, J. (1993):
Interaction of serum components with poly(methylmethacrylate) nanoparticles and the resulting body distribution after intravenous injection in rats.
J. Drug Target. 1, 15-19

Borchard, G., Kreuter, J. (1996):
The role of serum complement on the organ distribution of intravenously administered poly(methyl methacrylate) nanoparticles: effects of pre-coating with plasma and with serum complement.
Pharm. Res. 13, 1055-1058

Brasseur, F., Couvreur, P., Kante, B., Deckers-Passau, L., Roland, M., Deckers, C., Speiser, P. (1980):
Actinomycin D adsorbed on polymethylmethacrylate nanoparticles: an increased efficiency against an experimental tumor.
Eur. J. Cancer 16, 1441-1445

- Brown, J. M., Giaccia, A. J. (1998):
The unique physiology of solid tumors: opportunities (and problems) for cancer therapy.
Cancer Res. 58, 1408-1416
- Bucke, W. E., Leitzke, S., Diederichs, J. E., Borner, K., Hahn, H., Ehlers, S., Müller, R. H. (1997):
Surface-modified amikacin-liposomes: organ distribution and interaction with plasma proteins.
J. Drug Target. 5, 99-108
- Buckton, G. (1995):
Interfacial phenomena in drug delivery and targeting.
Harwood Academic Publishers, Chur, 237-281
- Cairns, J. (1975):
Das Krebsproblem.
In: Schirrmacher, V.: Krebs-Tumoren, Zellen, Gene, 3. Aufl, Spektrum der Wissenschaft / Verständliche Forschung (1989), 10-21
- Chiannilkulchai, N., Driouich, Z., Benoit, J. P., Poradi, A. L., Couvreur, P. (1989):
Doxorubicin-loaded nanoparticles: increased efficiency against murine hepatic metastases.
Sel. Cancer Ther. 5, 1-11
- Chiannilkulchai, N., Ammoury, N., Caillou, B., Devissaguet, J. Ph., Couvreur, P. (1990):
Hepatic tissue distribution of doxorubicin-loaded nanoparticles after i.v. administration in reticulosarcoma M5076 metastasis-bearing mice.
Cancer Chemother. Pharmacol. 26, 122-126
- Couvreur, P., Kante, B., Roland, M., Guiot, P., Bauduin, P., Speiser, P. (1979):
Polycyanoacrylate nanocapsules as potential lysosomotropic carriers: preparation, morphological and sorptive properties.
J. Pharm. Pharmacol. 31, 331-332
- Couvreur, P., Kante, B., Grislain, L., Roland, M., Speiser, P. (1982):
Toxicity of polyalkylcyanoacrylate nanoparticles II: Doxorubicin-loaded nanoparticles.
J. Pharm. Sci. 71, 790-792
- Couvreur, P., Aubry, J. (1983):
Monoclonal antibodies for the targeting of drugs: application to nanoparticles.
In: Breimer, D. D.: Topics in Pharmaceutical Sciences.
305-316, Elsevier Science Publishers B. V.
- Couvreur, P., Lenaerts, V., Leyh, D., Roland, M., Speiser, P. (1984):
Design of biodegradable polyalkylcyanoacrylate nanoparticles as a drug carrier.
In: Davis, S. S., Illum, L., Mc Vie, J. G., Tomlinson, E. (Eds.): Microspheres and drug therapy, Elsevier Science Publishers B. V., Amsterdam, 103-115
- Couvreur, P., Grislain, L., Lenaerts, V. (1986):
Biodegradable polymeric nanoparticles as a drug carrier for antitumor agents.
In: Guiot, P., Couvreur, P. (Eds.): Polymeric nanoparticles and microspheres, CRC Press, Boca Raton
- Couvreur, P., Roblot-Treupel, L., Poupon, M., Brasseur, F., Puisieux, F. (1990):
Nanoparticles as microcarriers for anticancer drugs.
Ad. Drug Delivery Rev. 5, 209-230
- Couvreur, P., Fattal, E., Andremont, A. (1991):
Liposomes and nanoparticles in the treatment of intracellular bacterial infections.
Pharm. Res. 8, 1079-1086
- Crommelin, D. J. A., Schreier, H. (1994):
Liposomes.
In Kreuter, J., Dekker, M. (eds.): Colloidal drug delivery systems., 73-190, Inc., New York,

- Culver, K. W., Rigolet, M., Raffel, C. (1997):
 Gene therapy strategies for brain cancer.
 In Bertino, J., R. (1997):
 Encyclopedia of cancer, 704-717, Volume II, Academic Press, San Diego-London-Boston-New York-Sydney-Tokyo-Toronto
- Cuvier, C., Roblot-Treupel, L., Millot, J. M., Lizard, G., Chevillard, S., Mainfait, M., Couvreur, P., Poupon, M. F. (1992):
 Doxorubicin-loaded nanospheres bypass tumor cell multidrug resistance.
Biochem. Pharmacol. 44, 509-517
- Davis, S. S., Illum, L. (1986):
 Review: colloidal carriers and drug targeting.
Acta Pharm. Technol. 32, 4-9
- Davis, S. S., Illum, L. (1988):
 Polymeric microspheres as drug carriers.
Biomaterials 9, 111-115
- De Dycker, R. P. (1992):
 Regional drug delivery in cancer treatment- an overview.
Eur. J. Hosp. Pharm. 2, 22-25
- De Keyser, J.-L., Poupaert, J. H., Dumont, P. (1991):
 Poly(dimethyl methylenemalonate) nanoparticles as a potential drug carrier: preparation, distribution and elimination after intravenous and peroral administration to mice.
J. Pharm. Sci. 80, 67-70
- De Vries, C., Escobedo, J. A., Ueno, H., Houck, K. A., Ferrara, N., Williams, L. T. (1992):
 The fms-like tyrosine kinase , a receptor for vascular endothelial growth factor.
Science 255, 989-991
- Dewey, W. C., Hopwood, L. E., Sapareto, S. A., Gerweck, L. E. (1977):
 Cellular responses to combinations of hyperthermia and radiation.
Radiology 123, 463-474
- Dewhirst, M. W., Tso, C. Y., Oliver, R., Gustafson, C. S., Secomb, T. W., Gross, J. F. (1989):
 Morphologic and hemodynamic comparison of tumor and healing normal tissue microvasculature.
Int. J. Radiat. Oncol. Biol. Phys. 17, 91-99
- Dickman, C. A., Rekate, H. L., Bird, C. R., Drayer, B. R., Medina, M. (1989):
 Unenhanced and gadolinium-DTPA-enhanced MR-imaging in postoperative evaluation in pediatric brain tumors.
J. Neurosurg. 71, 49-53
- Doll, R., Peto, R. (1981):
 The causes of cancer: quantitatives estimates of avoidable risk in the United States today.
J. Natl. Cancer Inst. 68, 1191-1303
- Douglas, S. J., Davies, S. S., Illum, L. (1986):
 Biodistribution of poly (butyl 2-cyanoacrylate) nanoparticles in rabbits.
Int. J. Pharm. 34, 145-152
- Dunn, S. E., Coombes, A. G. A., Garnett, M. C., Davis, S. S., Davies, M. C., Illum, L. (1997):
 In vitro cell interaction and in vivo biodistribution of poly (lactide-co-glycolide) nanospheres surface modified by poloxamer and poloxamine copolymers.
J. Control. Rel. 44, 65-76
- Egea, M. A., Gamisans, F., Valero, J., Garcia, M. E., Garcia, M. L. (1994):
 Entrapment of cisplatin into biodegradable polyalkylcyanoacrylate nanoparticles.
Il Farmaco 49, 211-217

Ehrlich, P. (1906):
Collected studies on immunology.
Vol.11, Wiley, New York, 442-447

Endrich, B., Reinhold, H. S., Gross, J. F., Intaglietta, M. (1979):
Tissue perfusion inhomogeneity during early tumor growth in rats.
J. Natl. Cancer Inst. 62, 387-395

Ferrara, N., Henzel, W. J. (1989):
Pituitary follicular cells secrete a novel heparin-binding growth factor specific for vascular endothelial cells.
Biochem. Biophys. Res. Commun. 161, 851-859

Ferrara, N. (1996):
Vascular endothelial growth factor.
Europ. J. Cancer 32A, 2413-2422

Fichtner, I., Kniest, A., Arndt, D. (1992):
Measurment of carbon clearance on mice as toxicity parameter for liposomal preparations.
In Vivo 6, 113-118

Fidler, I. J., Hart, I. R. (1982):
Biological diversity in metastatic neoplasms: origins and implications.
Science 217, 998-1003

Fiedler, H. P. (1989):
Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzende Gebiete.
3.Aufl., Editio Cantor Verlag

Folkman, J. (1971):
Tumorangiogenesis: therapeutic implications.
N. Engl. J. Med. 285, 1182-1186

Folkman, J., Klagsbrun, M. (1987):
Angiogenetic factors
Science 235, 442-447

Folkmann, J., Shing, Y. (1992):
Angiogenesis.
J. Biol. Chem. 267, 10931-10934

Fransen, A., De Tribolet, N. (1993):
Surgery of supratentorial tumors.
Curr. Opin. Oncol. 5, 450-457

Garner, A. (1994):
Vascular disease.
In: Garner, A.: Pathobiology of ocular disease. A dynamic approach, 2.Aufl., Marcel Dekker, New York,
(1994), 1625-1710

Gaspar, M. M., Blanco, D., Cruz, M. E. M., Alonso, M. J. (1997):
Formulation of L-asparaginase-loaded poly(lactide-co-glycolide) nanoparticles: influence of polymer properties
on enzyme loading, activity and in vitro release:
J. Control. Rel. 52, 53-62

Geran, R. I., Greenberg, N. H., Macdonald, M. M. (1967):
Protocols for screening chemical agents and natural products against animal tumors and other biological systems.
Cancer Chemother. Rep 3, 1-103

Giles, G. G., Thursfield, V., Staples, M et al. (1993):
Incidence and survival from childhood cancers in Victoria 1970-1979 and 1980-1989.
Anticancer Council of Victoria, Melbourne

Gipps, E. M., Arshady, R., Kreuter, J., Groscurth, P., Speiser, P. P. (1986):
Distribution of polyhexylcyanoacrylate nanoparticles in nude mice bearing human osteosarcoma.
J. Pharm. Sci. 75, 256-258

Gipps, E. M., Groscurth, P., Kreuter, J., Speiser, P. (1988):
Distribution of polyhexylcyanoacrylate nanoparticles in nude mice over extended times and after repeated injection.
J. Pharm. Sci. 77, 208-209

Glick, R. P., Lichtor, T., Kim, T. S., Ilangoan, S., Cohen, E. P. (1995):
Fibroblasts genetically engineered to secrete cytokines suppress tumor growth and induce antitumor immunity to a murine glioma in vivo.
Neurosurgery 36, 548-555

Golaz, O., Hughes, G. T., Frutiger, S., Paquet, N., Bairoch, A., Pasquali, C., Sanchez, J.-C., Tissot, J.-D., Appel, R. D., Walzer, C., Balant, L., Hochstrasser, D. F. (1993):
Plasma and red blood cell protein maps: Update 1993.
Electrophoresis 14, 1223-1231

Goldman, C. K., Kim, J., Wong, W. L., King, V., Brock, T., Gillespie, G. Y. (1993):
Epidermal growth factor stimulates vascular endothelial growth factor production by human malignant glioma cells: a model of glioblastoma multiforme pathophysiology.
Mol. Biol. Cell 4, 121-133

Goldring, S. R., Schiller, A. L., Roelke, M., Rourke, C. M., O'Neil, D. A., Harris, W. H. (1983):
The synovial-like membrane at the bone-cement interface in loose total hip replacements and its proposed role in bone lysis.
J. Bone Jt. Surg. 65-A, 575-584

Gombotz, W. R., Guanghui, W., Horbett, T. A., Hoffman, A. S. (1992):
Protein adsorption to and elution from polyether surfaces.
In: Harris, J. M. (Ed.): Poly (ethylene glycol) chemistry: biotechnical and biomedical applications, Plenum Press, New York, 247-261

Gray, L. H., Conger, A. D., Ebert, M., Hornsey, S., Scott, O. C. (1953):
Concentration of oxygen dissolved in tissues at the time of irradiation as a factor in radiotherapy.
Br. J. Radiol. 26, 638-648

Gref, R., Minamitake, Y., Peracchia, M. T., Trubetskoy, V., Langer, R. (1994):
Biodegradable long-circulating-polymeric nanospheres.
Science 263, 1600-1603

Gregoriadis, G. (1977):
Review: targeting of drugs.
Nature 265, 407-411

Grislain, L., Couvreur, P., Lenaerts, V., Roland, M., Deprez-Decampeneere, D., Speiser, P. (1983):
Pharmacokinetics and distribution of a biodegradable drug-carrier.
Int. J. Pharm. 15, 335-345

Gudewicz, P. W., Molnar, J., Lai, M. Z., Beezhold, D. W., Siefring, G. E., Ofedo, R. B., Lorand, L. (1980):
Fibronectin-mediated uptake of gelatin-coated latex particles by peritoneal macrophages.
J. Cell. Biol. 87, 427-433

Hahn, G. M. (1982):
Hyperthermia and cancer.
Plenum Press, New York, 1982, 7-54

Hanahan, D., Folkman, J. (1996):
Patterns and emerging mechanisms of the angiogenic switch during tumorigenesis.
Cell 86, 353-364

Harper, G. R., Davies, M. C., Davis, S. S., Tadros, Th. F., Taylor, D. C., Irving, M. P., Waters, J. A. (1991):
Steric stabilisation of microspheres with grafted polyoxyethylene oxide reduces phagocytosis by rat Kupffer
cells in vitro.
Biomaterials 12, 695-700

Harris, J. R., Lippman, M. E., Veronesi, U., Willett, W. (1992):
Breast cancer (first of three parts).
N. Engl. J. Med. 327/5, 319-328

Hertig, A. T. (1935):
Angiogenesis in the early human chorion and in the primary placenta of the macaque monkey.
Contrib. Embryol. 25, 37-81

Hochstrasser, D. F., Patchornik, A., Merril, C. R. (1988):
Development of polyacrylamide gels that improve the separation of proteins and their detection by silver
staining.
Anal. Biochem. 173, 412-423

Hsu, J. J., Juliano, R. L. (1982):
Interactions of liposomes with the reticuloendothelial system. II. Nonspecific and receptor mediated uptake of
liposomes by mouse peritoneal macrophages.
Biochem. Biophys. Acta 720, 411-419

IARC (1987):
Monographs on the evaluation of carcinogenic risk of chemicals to humans: overall evaluations of
carcinogenicity: an updating of IARC Monographs Volumes 1-42.
IARC Scientific Publications (Suppl. 7), Lyon

Ibrahim, A., Couvreur, P., Roland, M., Speiser, P. (1983):
New magnetic drug carrier.
J. Pharm. Pharmacol. 35, 59-61

Illum, L., Davis, S. S., Wilson, C. G., Thomas, N. W., Frier, M., Hardy, J. G. (1982):
Blood clearance and organ deposition of intravenously administered colloidal particles. The effects of particle
size, nature and shape.
Int. Pharm. Sci. 72, 1086-1089

Illum, L., Davis, S. S. (1983):
Effect of the nonionic surfactant poloxamer 338 on the fate and deposition of polystyrene microspheres
following intravenous administration.
J. Pharm. Sci. 72, 1086-1089

Illum, L., Jones, P. D. E., Kreuter, J., Baldwin, R. W., Davies, S. S. (1983):
Adsorption of monoclonal antibodies to polyhexylcyanoacrylate nanoparticles and subsequent immunospecific
binding to tumor cells in vitro.
Int. J. Pharm. 17, 65-76

Illum, L., Jones, P. D. E., Baldwin, R. W., Davies, S. S. (1984):
Tissue distribution of poly (hexyl 2-cyanoacrylate) nanoparticles coated with monoclonal antibodies in mice
bearing human tumor xenografts.
J. Pharmacol. Exp. Ther. 230, 733-736

Illum, L., Davis, S. S. (1984):
The organ uptake of intravenously administered colloidal particles can be altered using a non-ionic surfactant
(poloxamer 338)
FEBS Lett. 167, 79-82

Illum, L., Thomas, N. W., Davies, S. S. (1986):
Effect of a selected suppression of the reticuloendothelial system on the distribution of model carrier particles.
J. Pharm. Sci. 75, 1, 16-22

Illum, L., Davis, S. S. (1987):
Targeting of colloidal particles to the bone marrow.
Life Sci. 40, 1553-1560

Illum, L., Jacobsen, L. O., Müller, R. H., Mak, E., Davis, S. S. (1987):
Surface characteristics and the interaction of colloidal particles with mouse peritoneal macrophages.
Biomaterials 8, 113-117

Illum, L., Wright, J., Davies, S. S. (1989):
Targeting of microspheres to the sites of inflammation.
Int. J. Pharm. 52, 221-224

Innerarity, T. L., Arnold, K. S., Weisgraber, K. H., Mahley, R. (1986):
Apolipoprotein E is the determinant that mediates the receptor uptake of B-very low density lipoproteins by
mouse macrophages.
Arteriosclerosis 6, 114-122

Jain, R. K. (1997):
Delivery of molecular and cellular medicine to solid tumors.
Microcirculation 4, 1, 1-23

Jeon, S. I., Lee, J. H., Andrade, J. D., De Gennes, P. G. (1991):
Protein-surface interactions in the presence of polyethylene oxide.
Colloid. Interface 142, 149-158

Johnson, S. A., Thomas, N. W., Warren, N. W., Wilson, C. G., Fry, J. (1986):
Uptake of latex microparticle by rat hepatocytes in tissue culture.
J. Pharm. Pharmacol., Suppl. 101

Juliano, R. L. (1988):
Factors affecting the clearance kinetics and tissue distribution of liposomes, microspheres and emulsions.
Adv. Drug Deliv. Rev. 2, 31-54

Jungblut, P. (1995):
Hochauflösende zweidimensionale Elektrophorese.
Biospektrum 5, 69-72

Kabanov, A. V., Chekhonin, V. P., Alakhov, V. Y., Batrakova, E. V., Lebedev, A. S., Melik-Nubarov, N. S., Arzhakov, S. A., Levashov, A. V., Morozov, G. V., Severin, E. S. (1989):
The neuroleptic activity of haloperidol increases after its solubilization in surfactant micelles. Micelles as
microcontainer for drug targeting.
FEBS Lett. 258, 343-345

Kalenikova, E. I., Dmitrieva, O. F., Korobov, N. N., Zhukova, S. V., Tischenko, V. A. (1988):
Pharmacokinetics of dalargin.
Vopr. Med. Khim. 34, 75-83

Kanke, M., Simmons, G. H., Weiss, D. L., Bivins, B. A., DeLuca, P. P. (1980):
Clearance of ¹⁴¹Ce-labeled microspheres from blood and distribution in specific organs following intravenous
and intraarterial administration in beagle dogs.
J. Pharm. Sci. 69, 755-762

Kante, B., Couvreur, P., Dubois-Krack, G., De Meester, C., Guiot, P., Roland, M., Mercier, M., Speiser, P. (1982):
Toxicity of polyalkylcyanoacrylate nanoparticles I: free nanoparticles.
J. Pharm. Sci. 71, 786-790

Karino, A., Hayashi, H., Yamada, K., Ozawa, Y. (1987):
Effect of particle size and emulsifiers on the blood clearance and deposition of injected emulsions.
J. Pharm. Sci. 76, 273-276

Kattan, J., Droz, J.-P., Couvreur, P., Marino, J.-P., Boutan-Laroze, A., Rougier, P., Brault, P., Vranckx, H., Grognet, J.-M., Morge, X., Sancho-Garnier, H. (1992):
Phase I clinical trial and pharmacokinetic evaluation of doxorubicin carried by polyisohexylcyanoacrylate nanoparticles.
Invest. New Drugs 10, 191-199

Kaufmann, R., Proebstle, T., Sterry, W. (1996):
Spezielle Diagnostik und Therapie: Malignes Melanom.
In : Zeller, W. J., zur Hausen, H.: Onkologie: Grundlagen-Diagnostik-Therapie-Entwicklungen, ecomed Verlagsgesellschaft AG&CoKG.1-12

Ke, L. D., Chen, X. S., Steck, P. A., Yung, W. K. A. (1995):
Molecular modulation of vascular endothelial growth factor (VEGF) expression in glioma cells by a ribozyme.
11th International conference on brain tumor research and therapy, November 1995

Kim,K.J., Li, B., Winer, J., Armanini, M., Gillett, N., Phillips, H. S., Ferrara, N. (1993):
Inhibition of vascular endothelial growth factor-induced by targeted suppresses tumour growth in vivo.
Nature 362, 841-844

Kinoshita, T. (1991):
Biology of complement: the overture.
Immunol. Today 12, 291-295

Kissel, T., Roser, M. (1991):
Influence of chemical surface-modifications on the phagocytic properties of albumin nanoparticles.
Proceed. Intern. Symp. Control. Rel. Bioact. Mater. 18, 275-276

Kleihues, P., Soylemezoglu, F., Schäuble, B., Scheithauer, B. W., Burger, P. C. (1995):
Histopathology, classification and grading of gliomas.
Glia 15, 211-221

Knudson, A. G. (1971):
Statistical study of retinoblastoma.
Proc. Natl. Acad. Sci. USA 68, 820-823

Knudson, A., G. (1995):
Mutations and cancer: a personal odyssey
Adv. Cancer Res. 67, 1-23

Kost, J., Shefer, S., Yashpe, M. (1991):
Controlled release from swellable crosslinked starch particles for tumor chemoembolism.
Proceed. Intern. Symp. Control. Rel. Bioact. Mater. 18, 217-218

Kreuter, J., Speiser, P. P. (1976):
In vitro studies of poly(methylmethacrylate) adjuvants.
J. Pharm. Sci. 65, 1624-1627

Kreuter, J., Zehnder, H. J. (1978):
The use of ^{60}Co -irradiation for the production of vaccines.
Radiat. Effects 35, 161-166

Kreuter, J., Taeuber, U., Illi, V. (1979):
Distribution and elimination of poly(methyl-2- ^{14}C -methacrylate) nanoparticle radioactivity after injection in rats and mice.
J. Pharm. Sci. 68, 1443-1447

Kreuter, J. (1981):
Evaluation of nanoparticles as drug delivery systems.
Habilitationsschrift, ETH Zürich

- Kreuter, J., Liehl, E. (1981):
Long-term studies of microencapsulated and adsorbed influenza vaccine nanoparticles.
J. Pharm. Sci. 70, 367-371
- Kreuter, J. (1983):
Evaluation of nanoparticles as drug delivery systems II: comparison of the body distribution of nanoparticles with the body distribution of microspheres (diameter > 1μm), liposomes, and emulsions.
Pharm. Acta Helv. 58, 242-250
- Kreuter, J. (1983a):
Evaluation of nanoparticles as drug delivery systems I: preparation methods.
Pharm. Acta Helv. 58, 196-209
- Kreuter, J., Hartmann, H. R. (1983b):
Comparative study on the cytostatic effects and the tissue distribution of 5-fluorouracil in a free form and bound to polybutylcyanoacrylate nanoparticles in sarcoma 180-bearing mice.
Oncology 40, 363-366
- Kreuter, J., Nefzger, M., Liehl, E., Czok, R., Voges, R. (1983c):
Distribution and elimination of poly(methyl methacrylate) nanoparticles after subcutaneous administration to rats.
J. Pharma. Sci. 72, 10, 1146-1149
- Kreuter, J. (1983d):
Evaluation of nanoparticles as drug delivery systems III: materials, stability, toxicity, possibilities of targeting and use.
Pharm. Acta Helv. 58, 242-250
- Kreuter, J. (1985):
Factors influencing the body distribution of polyacrylic nanoparticles.
In Buri, P., Gumma, A. (eds.): *Drug targeting*, Elsevier Science Publishers B. V., Amsterdam,
- Kreuter, J. (1994):
Drug targeting with nanoparticles.
Eur. J. Drug Metab. Pharmacokinet. 3, 253-256
- Kreuter, J. (1994a):
Nanoparticles.
In: Kreuter, J. (Ed.): *Colloidal drug delivery systems*, Marcel Dekker, Inc., New York, 219-342
- Kreuter, J., Alyautdin, R. N., Kharkevich, D. A., Ivanov, A. A. (1995):
Passage of peptides through the blood-brain barrier with colloidal polymer particles (nanoparticles).
Brain Res. 674, 171-174
- Kreuter, J. (1997):
mdl. Mitteilung
- Kreuter, J., Alyaudin, R. N. (1998):
mdl. Mitteilung
- Labarre, D., Montdargent, B., Carreno, M.-J., Maillet, F. (1993):
Strategy for in vitro evaluation of the interactions between biomaterials and complement systems.
J. Appl. Biomater. 4, 231-240
- Lee, J. H., Kopecek, J., Andrade, J. D. (1989):
Protein-resistant surfaces prepared by PEO-containing block copolymer surfactants.
J. Biomed. Mater. Res. 23, 351-168
- Lee, J., Martic, P. A., Tan, J. S. (1989a):
Protein adsorption on pluronic co-polymer coated polystyrene particles.
J. Colloid. Interface Sci. 131, 252-266

Lenaerts, V., Couvreur, P., Christiaens-Leyh, D., Joiris, E., Rolland, M., Rollman, B., Speiser, P. (1984):
Degradation of poly(isobutylcyanoacrylate) nanoparticles.
Biomaterials 5, 65-68

Le Ray, A. M., Vert, M., Benoit, J. P. (1991):
Synthesis of radiolabeled poly(alpha-hydroxy acid). Preparation and in vivo studies of ^{14}C - poly(d,l-lactide-co-glycolide) nanoparticles.
Proceed. Intern. Symp. Control. Rel. Bioact. Mater. 18, 461-462

Leroux, J.-C., Gravel, P., Balant, L., Volet, B., Anner, B. M., Allemand, E., Doelker, E., Gurny, R. (1994):
Internalization of poly(d,l-lactic acid) nanoparticles by isolated human leukocytes and analysis of plasma proteins adsorbed onto the particles.
J. Biomed. Mater. Res. 28, 471-481

Leroux, J.-C., De Jaeghere, F., Anner, B., Doelker, E., Gurny, R. (1995):
An investigation on the role of plasma and serum opsonins on the internalization of biodegradable poly(d,l-lactic acid) nanoparticles by human monocytes.
Life Sci. 57, 7, 695-703

Lescure, F., Zimmer, C., Roy, D., Teulon, J. M., Couvreur, P. (1991):
Synthesis and evaluation of a new biodegradable monomer.
Proceed. Intern. Symp. Control. Rel. Bioact. Mater. 18, 325-326

Leu, D., Manthey, B., Kreuter, J., Speiser, P., DeLuca, P. P. (1984):
Distribution and elimination of coated polymethyl($2-^{14}\text{C}$)methacrylate nanoparticles after intravenous injection in rats.
J. Pharm. Sci. 73, 1433-1437

Leung, D. W., Cachianes, G., Kuang, W.-J., Goeddel, D.V., Ferrara, N. (1989):
Vascular endothelial growth factor is a secreted angiogenic mitogen.
Science 246, 1306-1312

Levine, A. J. (1995):
The genetic origins of neoplasia.
JAMA 273,592

Lherm, C., Müller, R. H., Puisieux, F., Couvreur, P. (1992):
Alkylcyanoacrylate drug carrier. II Cytotoxicity of cyanoacrylate nanoparticles with different alkyl chain length.
Int. J. Pharm. 84, 13-22

Lück, M. (1997):
Plasmaproteinadsorption als möglicher Schlüsselfaktor für eine kontrollierte Arzneistoffapplikation mit partikulären Trägern.
Fachbereich Pharmazie der FU-Berlin, Diss.

Maaßen, S., Fattal, E., Müller, R. H., Couvreur, P. (1993):
Cell cultures for the assessment of toxicity und uptake of polymeric particulate drug carriers.
S. T. P. Pharma Sciences 3, 1, 11-22

Malter, M., Süss, R. (1989):
Krebs im Blickpunkt.
1. Aufl., Hüthig Verlaggemeinschaft Decker & Müller, Heidelberg

Marty, J. J. (1977):
The preparation, purification and properties of nanoparticles.
Thesis, Victorian College of Pharmacy, Parksville, Australia

Mehta, K., Juliano, R. L., Lopez-Berestein, G. (1984):
Stimulation of macrophage protease secretion via liposomal delivery of muramyl dipeptide derivate to intracellular sites.
Immunology 51, 517-527

Millauer, B., Wizigmann-Voos, S., Schnurch, H. et al. (1993):
High affinity binding and developmental expression suggest flk-1 as a major regulator of vasculogenesis and angiogenesis.
Cell 72, 835-846

Millauer, B., Shawver, L. K., Plate, K. H., Risau, W., Ullrich, A. (1994):
Glioblastoma growth inhibited in vivo by a dominant-negative FLK-1 mutant.
Nature 367, 576-578

Moghimi, S. M., Patel, H. M. (1988):
Tissue specific opsonins for phagocytic cells and their different affinity for cholesterol-rich liposomes.
FEBS Lett. 233, 143-147

Moghimi, S. M., Porter, C. J. H., Muir, I. S., Illum, L., Davis, S. S. (1991):
Non-phagocytic uptake of intravenously injected microspheres in rat spleen: influence of particle size and hydrophilic coating.
Res. Commun. 177, 861-866

Moghimi, S. M., Davis, S. S. (1994):
Innovations in avoiding particle clearance from blood by Kupffer cells: some cause for reflection.
Crit. Rev. Ther. Drug Carrier Syst. 11, 31-59

Moghimi, S. M., Patel, H. M. (1996):
Altered tissue-specific opsonic activities and opsono-recognition of liposomes in tumour-bearing rats.
Biochim. Biophys. Acta 1285, 56-64

Mokuno, H., Yamada, N., Shimano, H., Ishibashi, S., Mori, N., Takahashi, K., Oka, T., Yoon, T. H., Takaku, F. (1991):
The enhanced cellular uptake of very-low-density lipoprotein enriched in apolipoprotein E.
Biochim. Biophys. Acta 1082, 63-70

Morimoto, Y., Akimoto, M., Sugibayashi, K., Nadai, T., Kato, Y. (1980):
Drug-carrier property of albumin microspheres in chemotherapy. IV. Antitumor effect of single-shot administration of microsphere-entrapped 5-fluorouracil on Ehrlich ascites or solid tumor in mice.
Chem. Pharm. Bull. 28, 3087-3092

Müller, R. H., Heinemann, S. (1989):
Surface modelling of microparticles as parenteral systems with high tissue affinity.
In: Gurny, R., Junginger, H. E. (Eds.): Bioadhesion - possibilities and future trends, WVG, Stuttgart, 202-213

Müller, R. H., Herbst, S. (1990):
Nanopartikel - Gewebespezifische Arzneiform für Cytostatika.
DAZ 130, 1509-1511

Müller, R. H. (1991):
Colloidal carriers for controlled drug delivery and targeting.
Wissenschaftliche Verlagsgesellschaft, Stuttgart.

Müller, H., Aigner, K. R. (1992):
Isolierte Beckenperfusion (IPP) bei inoperablen Beckentumoren.
Chemotherapy J. 1, 11-16

Müller, R. H., Wallis, K. H. (1993):
Surface modification of i.v. injectable biodegradable nanoparticles with poloxamer polymers and poloxamine 908.
Int. J. Pharm. 89, 25-31

Müller, R. H., Maaßen, S., Weyhers, H., Mehnert, W. (1996):
Phagocytic uptake and cytotoxicity of solid lipid nanoparticles (SLN) sterically stabilized with poloxamine 908 and Poloxamer 407.
J. Drug Target. 4, 3, 161-170

Müller, R. H. (1996):
Zetapotential und Partikelladung in der Laborpraxis.
Wissenschaftliche Verlagsgesellschaft mbH Stuttgart, Bd. 37

Mutschler, E. (1991):
Arzneimittelwirkungen- Chemotherapie maligner Tumoren.
6.Aufl., Wissenschaftliche Verlagsgesellschaft Stuttgart.

Newell, D. R., Hendriks, H. R. (1998):
Standard operating procedure: detailed protocol for preliminary toxicological and pharmacokinetic studies with compounds derived from the NCI in vitro drug evaluation programme.
EORTC-SPG, Standard operating procedure

Nicklas, W. (1995):
Hinweise zur Blutentnahme bei kleinen Versuchstieren.
Tierärztliche Vereinigung für Tierschutz e.V., Iltisstieg 5, 22159 Hamburg

Nicolson, G. L. (1979):
Krebsmetastasen.
In: Schirrmacher, V.: Krebs- Tumoren, Zellen, Gene, 3.Aufl., Spektrum der Wissenschaft / Verständliche Forschung (1989), 148-159

Norde, W. (1986):
Adsorption of proteins from solution at the solid-liquid interface.
Adv. Coll. Interf. Sci. 25, 267-340

Norde, W., MacRitchie, F., Nowicka, G., Lyklema, J. (1986):
Protein adsorption at solid-liquid interfaces: reversibility and conformation aspects.
J. Coll. Interf. Sci. 112, 447-456

Norman, M. E., Williams, P., Illum, L. (1993):
Influence of block copolymers on the adsorption of plasma proteins to microspheres.
Biomaterials 14, 3, 193-202

O'Farrell, P. H. (1975):
High resolution two-dimensional electrophoresis of proteins.
J. Biol. Chem. 250, 4007-4021

Old, L. J. (1996):
Immuntherapie gegen Krebs.
In Spektrum der Wissenschaft: Spezial 5: Krebsmedizin.77-84

O'Mullane, J.E., Artursson, P., Tomlinson, E. (1987):
Biopharmaceutics of microparticle drug carriers.
Ann. N. Y. Acad. Sci. 507,100-140

O'Mullane, J.E., Petrak, K., Hutchinson, L.E.F., Tomlinson, E. (1990):
The effect of adsorbed coats of Poloxamers 237 and 338 on the in vitro aggregation and in vivo distribution of polystyrene latex (PSL) particles.
Int. J. Pharm. 63, 177-180

Oppenheim, R. C. (1981):
Solid colloidal drug delivery systems: nanoparticles.
Int. J. Pharm. 8, 217-234

Oppenheim, R. C., Gipps, E. M., Forbes, J. F., Whitehead, R. H. (1984):
Development and testing of proteinaceous nanoparticles containing cytotoxins.
In: Davis, S. S., Illum, L., McVie, J. G., Tomlinson, E. (Eds.), Elsevier, Amsterdam

- Papatheofanis, F. J., Barmada, R. (1991):
 Polymorphonuclear leukocyte degranulation with exposure to polymethylmethacrylate nanoparticles.
J. Biomed. Mater. Res. 25, 761-771
- Pardridge, W. M. (1996):
 Physiologic-based strategies for protein drug delivery to the brain.
J. Control. Rel. 39, 281-286
- Patel, K. R., Li, M. P., Baldeschwiler, J. D. (1983):
 Suppression of liver uptake of liposomes by dextran sulfate 500.
Proc. Natl. Acad. Sci. USA 80, 6518-6522
- Patel, H. M. (1992):
 Serum opsonins and liposomes: their interaction and opsonophagocytosis.
Crit. Rev. Ther. Drug Carr. Syst. 9, 39-90
- Paul, M., Durand, R., Boulard, Y., Fusai, T., Fernandez, C., Rivollet, D., Deniau, M., Astier, A. (1998):
 Physicochemical characteristics of Pentamidine-loaded polymethacrylate nanoparticles: implication in the intracellular drug release in *Leishmania major* infected mice.
J. Drug Targeting 5, 6, 481-490
- Pepper, M. S., Ferrara, N., Orci, L., Montesano, R. (1992):
 Potent synergism between vascular endothelial growth factor and basic fibroblast growth factor in the induction of angiogenesis in vitro.
Res. Comm. 189, 824-831
- Pilkington, G. J. (1994):
 Tumour cell migration in the central nervous system.
Brain. Pathol. 4, 157-166
- Plate, K. H., Breier, G., Weich, H. A., Risau, W. (1992):
 Vascular endothelial growth factor is a potential tumour angiogenesis factor in human gliomas in vivo.
Nature 359, 845-848
- Plate, K. H., Breier, G., Millauer, B., Ullrich, A., Risau, W. (1993):
 Up-regulation of vascular endothelial growth factor and its cognate receptors in a rat glioma model of tumor angiogenesis.
Cancer Res. 53, 5822-5827
- Ploüet, J., Schilling, J., Gospodarowicz, D. (1989):
 Isolation and characterisation of a newly identified endothelial cell mitogen produced by AtT20 cells.
EMBO J. 8, 3801-3807
- Pötgens, A.J. G., Lubsen, N. H., van Altena, M. C., Schoenmakers, J. G. G., Ruiter, D. J., de Waal, R. M. W. (1995):
 Vascular permeability factor expression influences tumor angiogenesis in human melanoma lines xenografted to nude mice.
Amer. J. Pathol. 146, 197-209
- Porter, C. J. H., Moghimi, S. M., Illum, L., Davis, S. S. (1992):
 The polyoxyethylene/polyoxypropylene block co-polymer poloxamer-407 selectively redirects intravenously injected microspheres to sinusoidal endothelial cells of rabbit bone marrow.
FEBS Lett. 305, 62-66
- Poste, G., Kirsh, R., Koestler, T. (1984):
 The challenge of liposome targeting in vivo.
 In Gregoriadis, G. (ed): *Liposome technology Vol. III: targeted drug delivery and biological interactions*. CRC Press., Florida, 1984, 1-28
- Reinhart, T., Bauer, K. H. (1995):
 Untersuchungen zum Hämolyse- und Solubilisationsverhalten einiger nichtionischer polymerer Tensidklassen.
Pharmazie 50, 403-407

Reszka, R., Beck, P., Fichtner, I., Hentschel, M., Richter, J., Kreuter, J. (1997):
Body distribution of free, liposomal and nanoparticle- associated mitoxantrone in B16-Melanoma-bearing mice.
J. Pharmacol. Exp. Ther. 280, 232-237

Risau, W. (1996):
What, if anything, is an angiogenic factor?
Cancer Metastasis Rev. 15, 149-151

Rollot, J. M., Couvreur, P., Roblot-Treupel, L., Puisieux, F. (1986):
Physicochemical and morphological characterization of polyisobutyl cyanoacrylate nanoparticles.
J. Pharm. Sci. 75, 361-364

Romeis, B. (1989):
Mikroskopische Technik.
Urbahn und Schwarzenbeck, 17 Auflage, München, Wien, Baltimore

Roth, J. A., Mukhopadhyay, T., Zhang, W.-W., Fujiwara, T., Georges, R. (1995):
Gene replacement strategies for the prevention and therapy of cancer.
In Sobol, R. E., Scanlon, K. J. (eds): *The internet book of gene therapy*. Appleton & Lange 1995, 283-296

Rudt, S. (1992):
Untersuchungen zur Phagocytose und zur Toxizität von Arzneistoffträger-Systemen zur parenteralen Applikation.
Ph. D.Thesis, Univeristät Kiel

Sachs, L. (1986):
Leukämie: Umkehr der Entartung.
In: Schirrmacher,V.: *Krebs-Tumoren, Zellen, Gene*, 3. Aufl, Spektrum der Wissenschaft / Verständliche Forschung (1989), 124-132

Sampson, J. H., Archer, G. E., Kurpad, S. N., Castleberry, K. A., Fuchs, H. E., Gilboa, E., Dranoff, G., Bigner, D. D. (1995):
Cytokine producing tumor cells demonstrate reduced intracerebral tumorigenicity and reduce the growth of intracerebral tumors when as subcutaneous vaccinations.
11th International Conference on Brain tumor research and therapy, November, 1995

Schäfer, V., v. Briesen, H., Andreesen, R., Steffan, A.-M., Royer, C., Tröster, S., Kreuter, J., Rübsamen-Waigmann, H. (1992):
Phagocytosis of nanoparticles by human immunodeficiency virus (HIV)-infected macrophages: a possibility for antiviral drug targeting.
Pharm. Res. 9, 541-546

Scherneck, S., Kölble, K. (1998):
Mechanismen der Entstehung genetisch bedingter Krebsformen.
In: Ganter, D., Ruckpaul, K.: *Handbuch der Molekularen Medizin, Band II: Tumorerkrankungen*, Springer, Berlin-Heidelberg-New York

Schapel, A. (1998):
mdl. Mitteilung

Schröder, U., Sabel, B. A. (1995):
Nanoparticles, a drug carrier system to pass the blood-brain barrier, permit central analgesic effects of i.v. dalargin injections.
Brain Res. 710, 121-124

Senger, D.R., Galli, S. J., Dvorak, A. M., Perruzzi, C.A., Harvey, V. S., Dvorak, H. F. (1983)
Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid.
Science 219, 983-985

Sezaki, H., Hashida, M. (1984):
Macromolecule-drug conjugates in targeted cancer chemotherapy.
CRC critical reviews in therapeutic drug carrier systems 1, 1-38

- Shah-Yukich, A. A., Nelson, A. C. (1988):
 Characterization of solid tumor microvasculature: a three-dimensional analysis using the polymer casting technique.
Lab. Invest. 58, 236-244
- Sharma, D., Chelvi, T., Kaur, J., Chakravorty, K., De, T., Maitra, A., Ralhan, R. (1996):
 Novel Taxol® formulation: polyvinylpyrrolidone nanoparticle-encapsulated Taxol® for drug delivery in cancer therapy.
Oncol. Res. 8, 7/8, 281-286
- Sherman, L. A., Lee, J. (1977):
 Specific binding of soluble fibrin to macrophages.
J. Exp. Med. 145, 76-85
- Shibuya, M., Yamaguchi, S., Yamane A. et al. (1990):
 Nucleotide sequence and expression of a novel human receptor-type tyrosine kinase (flt) closely related to the fms family.
Oncogene 8, 519-527
- Shweiki, D., Itin, A., Soffer, D., Keshet, E. (1992):
 Vascular endothelial growth factor induced by hypoxia may mediate hypoxia-initiated angiogenesis.
Nature 359, 843-845
- Smith-McCune, K. S., Weidner, N. (1994):
 Demonstration and characterisation of the angiogenetic properties of cervical dysplasia.
Cancer Res. 54, 804-808
- Sobol, R. E., Scanlon, K. J. (1995):
 Cancer therapeutics.
 In Sobol, R. E., Scanlon, K. J. (eds): *The internet book of gene therapy*. Appleton & Lange 1995, 283-296
- Sobol, R. E., Shawler, D., Dorigo, O., Gold, D., Royson, I., Fakhrai, H. (1995):
 Immunogene therapy of cancer.
 In Sobol, R. E., Scanlon, K. J. (eds): *The internet book of gene therapy*. Appleton & Lange 1995, 175-180
- Souhami, R. L., Bradfield, J. W. B. (1974):
 The recovery of hepatic phagocytosis after blockade of Kupffer cells.
J. Reticuloendothel. Soc. 16, 75-86
- Spennlehauer, G., Vert, M., Benoit, J.-P., Chabot, F., Veillard, M. (1988):
 Biodegradable cisplatin microspheres prepared by the solvent evaporation method: morphology and release characteristics.
J. Control. Rel. 7, 217-229
- Stan, A. C., Nemati, M. N., Pietsch, T., Walter, G. F., Dietz, H. (1995):
 In vivo inhibition of angiogenesis and growth of the human U-87 malignant glia tumor by treatment with an antibody against basic fibroblast growth factor.
J. Neurosurg. 82, 1044-1052
- Stolnik, S., Davies, M. C., Illum, L., Davis, S. S., Boustta, M., Vert, M. (1994):
 The preparation of sub-200nm biodegradable colloidal particles from poly (β -malic acid-co-benzyl malate) copolymers and their surface modification with Poloxamer and Poloxamine surfactants.
J. Control. Rel. 30, 57-67
- Storm, G., Belliot, S. O., Daemen, T., Lasic, D. D. (1995):
 Surface modification of nanoparticles to oppose uptake by the mononuclear phagocyte system.
Adv. Drug Deliv. Rev. 17, 31-48
- Stricker, H. (1987):
 Physikalische Pharmazie.
 3. Aufl., Wissenschaftliche Verlagsgesellschaft mbH Stuttgart.

Strawn, L. M., McMahon, G., App, H., Schreck, R., Kuchler, W. R., Longhi, M. P., Hui, T. H., Tang, C., Levitzki, A., Gazit, A., Chen, I., Keri, G., Orfi, L., Risau, W., Flamme, I., Ullrich, A., Hirth, K. P., Shawver, L. K. (1996):
Flk-1 as a target for tumor growth inhibition.
Cancer Res. 56, 3540-3545

Stünzi, H., Weiss, E. (1990):
Allgemeine Pathologie für Tierärzte und Studierende der Veterinärmedizin, 8. Aufl., Paul Parey Verlag, Berlin-Hamburg, 317-376

Sugibayashi, K., Morimoto, Y., Nadai, T. (1979):
Drug-carrier property of albumin microspheres in chemotherapy. II. Preparation and tissue distribution in mice of microsphere-entrapped 5-fluorouracil.
Chem. Pharm. Bull. 27, 204-209

Takahashi, Y., Kitadai, Y., Bucana, C. D., Cleary, K. R., Ellis, L. M. (1995):
Expression of vascular endothelial growth factor and its receptor, KDR, correlates with vascularity, metastasis and proliferation of human colon cancer.
Cancer Res. 55, 3964-3968

Thews, G., Mutschler, E., Vaupel, P. (1989):
Anatomie, Physiologie, Phatophysiologie des Menschen.
3.Aufl., Wissenschaftliche Verlagsgesellschaft mbH Stuttgart

Toi, M., Hoshina, S., Takayanagi, T., Tominaga, T. (1994):
Association of vascular endothelial growth factor expression with tumor angiogenesis and with early relapse in primary breast cancer.
Jpn. J. Cancer Res. 85, 1045-1049

Tooyama, I., Kawamata, T., Akiyama, H., Kimura, H., Moestrup, S. K., Gliemann, J., Matsuo, A., McGeer, P. L. (1995):
Subcellular localization of the low density lipoprotein receptor-related protein (α -2-Macroglobuline receptor) in human brain.
Brain Res. 691, 235-238

Tröster, S. D., Müller, U., Kreuter, J. (1990):
Modification of the body distribution of poly (methyl methacrylate) nanoparticles in rats by coating with surfactants.
Int. J. Pharm. 61, 85-100

Tröster, S. (1990):
Veränderung der Köperverteilung kolloidalen Arzneistoffträger, speziell von Nanopartikeln, nach intravenöser Applikation.
Frankfurt am Main, Fachbereich Biochemie, Pharmazie und Lebensmittelchemie der J. W. Goethe- Universität, Diss.

Tröster, S. D., Wallis, K. H., Müller, R. H., Kreuter, J. (1992):
Correlation of the surface hydrophobicity of ^{14}C -poly(methyl methacrylate) nanoparticles to their body distribution.
J. Control. Rel. 20, 247-260

Tsai, J.-C., Goldman, C. K., Gillespie, G. E. (1995):
Vascular endothelial growth factor in human glioma cell lines: induced secretion by EGF, PDGF-BB and bFGF.
J. Neurosurg. 82, 864-873

Tuder, R. M., Flook, B. E., Voelkel, N. F. (1995):
Increased gene expression of VEGF and VEGF receptor KDR/Flk and Flt in lungs exposed to acute or chronic hypoxia.
J. Clin. Invest. 95, 1798-1807

United Kingdom Co-ordinating Committee on Cancer Research (UKCCCR) (1998):
Guidelines for the welfare of animals in experimental neoplasia (2. Edition).
Brit. J. Cancer 77, 1-10

Van Berkel, Th. J. C. (1993):
Drug targeting: application of endogenous carriers for site-specific delivery of drugs
J. Control. Rel. 24, 145-155

van Oss, C. J. (1978):
Phagocytosis as a surface phenomenon.
Ann. Rev. Microbiol. 32, 19-39

Van Leuven, F., Marynen, P., Sottrup-Jensen, Cassiman, J.-J., Van Den Berghe, H. (1986):
The receptor-binding domain of human α 2-macroglobulin. Isolation after limited proteolysis with a bacterial proteinase.
J. Biol. Chem. 261, 11369-11373

Varma, R. K., Kaushal, R., Junnarkar, A. Y., Thomas, G. P., Naidu, M. U. R., Singh, P. P., Tripathi, R. M., Shridhar, D. R. (1985):
Polysorbate 80: a pharmacological study.
Drug Res. 35, 804-808

Vogel, W., Bomford, A., Young, S., Williams, R. (1987):
Heterogeneous distribution of transferrin receptors on parenchymal and nonparenchymal liver cells.
Biochemical and morphological evidence, Blood 69, 264-270

Vogelstein, B., Kinzler, K. W. (1993):
The multistep nature of cancer.
TIG 9,138-141

Weidner, N., Folkman, J., Pozza, F. (1992):
Tumor angiogenesis: a new significant and independent prognostic indicator in early-stage breast carcinoma.
J. Natl. Cancer Inst. 84, 1875-1888

Weinberg, R. A. (1984):
Molekulare Grundlagen von Krebs.
In: Schirrmacher,V.: Krebs-Tumoren, Zellen, Gene, 3. Aufl, Spektrum der Wissenschaft / Verständliche
Forschung (1989),66-77

Weinberg, R. B., Patton, C. S. (1990):
Binding of human apolipoprotein A-IV to human hepatocellular plasma membranes.
Biochim. Biophys. Acta 1044, 255-261

Weiss, E., Korbe, E. (1990):
Geschwülste.
In: Stünzi, H.,Weiss, E.,: Allgemeine Pathologie für Tierärzte und Studierende der Veterinärmedizin, 317-376, 8.
Aufl., Paul Parey, Berlin-Hamburg

Wesemeyer, H. (1994):
Mikrokalorimetrische und chromatographische Untersuchungen zur Oberflächenmodifikation von partikulären
Arzneistoffträgern durch Tenside.
Christian-Albrechts-Universität, Kiel, Diss.

Westermark, B., Ponten, J., Hugosson, R. (1973):
Determinants for the establishment of permanent tissue culture lines from human gliomas.
Acta. Pathol. Microbiol. Scand. A. 81, 791-805

Widder, K. J., Senyei, A. E., Ranney, D. F. (1979):
Magnetically responsive microspheres and other carriers for the biophysical targeting of antitumor agents.
Advances in Pharmacology and Chemotherapy 16, 213-271

- Widder, K. J., Senyei, A. E., Ranney, D. F. (1980):
 In vitro release of biologically active adriamycin by magnetically responsive albumin microspheres.
Cancer Research 40, 3512-3517
- Widder, K. J., Morris, R. M., Poore, G. A., Howard, D. P., Senyei, A. E. (1983):
 Selective Targeting of magnetic albumin microspheres containing low-dose doxorubicin: total remission in Yoshida sarcoma-bearing rats.
Eur. J. Cancer Clin. Oncol. 19, 1, 135-139
- Widmaier, R., Wildner, G. P., Papsdorf, G., Graffi, A. (1974):
 Über eine neue, in vitro unbegrenzt wachsende Zelllinie, MaTu, von Mamma-Tumorzellen des Menschen.
Archiv für Geschwulstforschung 44, 1-10
- Wilkens, D. J., Myers, P. A. (1966):
 Studies on the relationship between the electrophoretic properties of colloids and their blood clearance and organ distribution in the rat.
Brit. J. Exp. Path. 47, 568-576
- Wilkowski, J. A. (1990):
 The inherited character of cancer- an historical survey.
Cancer Cells 2, 229-257
- Yoshioka, T., Hashida, M., Muranishi, S., Sezaki, H. (1981):
 Specific delivery of mitomycin C to liver, spleen and lung: nano- and micropartical carriers of gelatin.
Int. J. Pharm. 81, 131-141
- Zagzag, D. (1995):
 Angiogenic growth factors in neural embryogenesis and neoplasia.
Amer. J. Pathol. 146, 293-309
- Zheng, G., Bachinsky, D. R., Stamenkovic, I., Stickland, D. H., Brown, D., Andres, G., McCluskey, R. T. (1994):
 Organ distribution in rats of two members of the low-density lipoprotein receptor gene family, gp 330 and LRP/alpha 2MR, and the receptor-associated protein (RAP).
J. Histochem. Cytochem. 42, 531-542
- Zhu, J. (1997):
 Experimentelle Ansätze zur molekularen Therapie maligner Hirntumore mittels Gen- und Antisense-Oligonukleotid-Transfer:
 Berlin, Mathematisch Naturwissenschaftliche Fakultät I am Institut für Biologie der Humboldt-Universität zu Berlin, Diss.
- Zimmer, A., Mutschler, E., Lambrecht, G., Mayer, D., Kreuter, J. (1994):
 Pharmacokinetic and pharmacodynamic aspects of an ophthalmic Pilocarpine nanoparticle-delivery-system.
Pharm. Res. 11, 10, 1435-1442
- Zlokovic, B. V., Martel, C. L., Matsubara, E., McComb, J. G., Zheng, G., McCluskey, R. T., Frangione, B., Ghiso, J. (1996):
 Glycoprotein 330/megalin: probable role in receptor-mediated transport of apolipoproteins J alone and in a complex with Alzheimer disease amyloid beta at the blood-brain and blood-cerebrospinal fluid barriers.
J. Proc. Natl. Acad. Sci., USA, 93, 4229-4234
- Zolle, I., Hosain, F., Rhodes, B. A., Wagner, H. N. (1970):
 Abstract: human serum albumin millimicrospheres for studies of the reticuloendothelial system.
J. Nucl. Med. 11, 379
- zur Hausen, H. (1990):
 Geleitwort.
 In Becker, N., Wahrendorf, J.: *Krebsatlas der Bundesrepublik Deutschland 1981-1990*, 3. Aufl., Springer,