

Molecular basis of PI3-kinase γ regulation

Molekulare Grundlagen der PI3-Kinase- γ -Regulation

Dissertation zur Erlangung des akademischen Grades des
Doktors der Naturwissenschaften (Dr. rer. nat.)

eingereicht im Fachbereich Biologie, Chemie, Pharmazie
der Freien Universität Berlin

vorgelegt von

Philipp Voigt

Diplom-Biochemiker aus Berlin

Juni 2007

1. Gutachter: Prof. Dr. M. Schaefer, Charité – Universitätsmedizin Berlin

2. Gutachter: Prof. Dr. G. Multhaup, Freie Universität Berlin

Disputation am 17. Juli 2007

Table Of Contents

1	Introduction	1
1.1	The family of phosphoinositide 3-kinases	1
1.1.1	Class II and III PI3Ks	1
1.1.2	Class I PI3Ks	3
Activation of class IA PI3Ks	4	
Activation of class IB PI3K	6	
1.2	Effector systems of class I PI3Ks	7
1.3	Physiological roles of PI3K γ	13
1.3.1	Leukocyte systems	14
1.3.2	Heart and vascular system	16
1.3.3	Other tissues	17
1.4	Structure and interaction partners of PI3K γ	17
1.4.1	Structure of p110 γ	18
1.4.2	Properties of the p101 regulatory subunit	18
1.4.3	Ras as an activator of p110 γ	20
1.4.4	Interaction with GRK2 and PDE3B in heart	20
2	Aims	23
3	Materials and methods	25
3.1	Culture and transfection of mammalian cells	25
3.2	Molecular biology	26
3.2.1	Generation of expression plasmids	27
p101 deletion mutants	28	
Cloning of p87 ^{PIKAP}	29	
Vectors for baculovirus generation	30	
Miscellaneous constructs	32	
3.2.2	Generation of plasmids for RNAi	34
3.2.3	Multiplex and semi-quantitative RT-PCR	35
3.2.4	Northern blot analysis	37
3.2.5	<i>In silico</i> methods	38

Table Of Contents

3.3	Immunochemical assays	38
3.3.1	Gel electrophoresis and immunoblot	38
3.3.2	Immunoprecipitation	40
3.4	Biochemical assays	41
3.4.1	PDE assay	41
3.4.2	β -Hexosaminidase assay	42
3.4.3	Peptide SPOT overlay assay	43
3.5	Protein purification	44
3.5.1	Culture and infection of Sf21 cells	45
3.5.2	Purification of PI3K γ subunits and G $\beta\gamma$ by affinity chromatography	45
	His-p110 γ	45
	His-p87 ^{PIKAP}	46
	His-p110 γ /His-p87 ^{PIKAP} dimer	47
	His-TEV-p110 γ /p87 ^{PIKAP} dimer	47
	Purification of G $\beta\gamma$ complexes	48
3.6	Imaging techniques	49
3.6.1	Confocal microscopy	49
3.6.2	Imaging of intracellular calcium concentrations	50
3.6.3	FRET	50
4	Results	53
4.1	Assignment of functional domains in p101	53
4.1.1	Establishing a sketchy functional map	53
4.1.2	Fine-mapping of the p110 γ binding domain	56
4.1.3	Fine-mapping of the G $\beta\gamma$ binding domain	59
4.1.4	Interference with PI3K γ signaling	60
4.1.5	Conservation of the functional domains in orthologues of p101	62
4.1.6	Data base mining for novel p110 γ - and G $\beta\gamma$ -binding proteins	65
4.2	p87 ^{PIKAP} – a novel PI3K γ regulatory subunit	68
4.2.1	Cloning and basal characterization of p87 ^{PIKAP}	68
4.2.2	Interaction of p87 ^{PIKAP} with p110 γ	69
4.2.3	Interaction of p87 ^{PIKAP} with G $\beta\gamma$	74
4.2.4	Reconstitution of PI3K γ signaling	75
4.2.5	Expression pattern of p87 ^{PIKAP}	77
4.2.6	Interaction of p87 ^{PIKAP} with PDE3B	80
4.2.7	Analysis of p87 ^{PIKAP} function by RNA interference	82
4.3	Structural studies on the interaction of p101 and p87 ^{PIKAP} with p110 γ and G $\beta\gamma$.	88
4.3.1	Purification of proteins for peptide SPOT analysis	89

4.3.2	Interaction of regulatory subunits with p110 γ	90
4.3.3	Interaction of regulatory subunits with G $\beta\gamma$	96
4.3.4	Purification of heterodimeric PI3K γ for crystallization	98
5	Discussion	103
5.1	The domain structure of p101	104
5.2	A novel PI3K γ regulatory subunit	108
5.2.1	Genealogy of p101 and p87 ^{PIKAP}	108
5.2.2	General features and expression pattern of p87 ^{PIKAP}	110
5.2.3	Interaction of p87 ^{PIKAP} with PDE3B	113
5.2.4	Role of p87 ^{PIKAP} in mast cell signaling	115
5.3	Structural studies on interactions involved in activation of PI3K γ	116
5.4	Potential implications for the development of isotype-specific PI3K γ inhibitors	119
6	Summary	125
7	Zusammenfassung	127
References		129
List of own publications		159
Curriculum vitae		163
Acknowledgments		165

Abbreviations

aa	amino acid(s)
AMP	adenosine 5'-monophosphate
APS	ammoniumpersulfate
β -ME	β -mercaptoethanol
BSA	bovine serum albumin
Btk	Bruton's tyrosine kinase
bp	base pair(s)
β AR	β -adrenergic receptor
β ARK	β -adrenergic receptor kinase
[Ca ²⁺] _i	intracellular concentration of calcium ions
cAMP	adenosine 3',5'-cyclic monophosphate
<i>C. elegans</i>	<i>Caenorhabditis elegans</i>
CD	cluster of differentiation
CFP	cyan fluorescent protein, color variant of GFP
cLSM	confocal laser scanning microscope
CV	column volume(s)
DC	dendritic cell
DMEM	Dulbecco's modified Eagle medium
DNA-PK	DNA-dependent protein kinase
DNP	dinitrophenol
DTT	dithiothreitol
<i>E. coli</i>	<i>Escherichia coli</i>
EDTA	ethylenediaminetetraacetic acid
EGF	epidermal growth factor
ERK	extracellular signal-regulated kinase
EST	expressed sequence tag
FBS	foetal bovine serum
fMLP	formyl-methionyl-leucyl-phenylalanine

FRET	fluorescence resonance energy transfer
GAP	GTPase-activating protein
GAPDH	glyceraldehyde 3-phosphate dehydrogenase
GEF	guanine nucleotide exchange factor
GFP	green fluorescent protein
GPCR	G protein-coupled receptor
GRK	G protein-coupled receptor kinase
Grp1	general receptor for phosphoinositides 1
GSK-3	glycogen synthase kinase 3
GTPase	GTP hydrolase
HEK	human embryonic kidney
HEPES	2-[4-(2-hydroxyethyl)-1-piperazinyl]-ethanesulfonic acid
HM	hydrophobic motif
HRP	horseradish peroxidase
IP	immunoprecipitation
MAPK	mitogen-activated protein kinase
MEM	minimal essential medium
mTOR	mammalian target of rapamycin
NLS	nuclear localization sequence(s)
NRTK	nonreceptor tyrosine kinase
PAGE	polyacrylamide gel electrophoresis
p87 ^{PIKAP}	PI3K γ adapter protein of 87 kDa
PCR	polymerase chain reaction
PDK-1	3-phosphoinositide-dependent kinase 1
PDE	phosphodiesterase
PH domain	pleckstrin homology domain
PI	phosphatidylinositol
PI(3)P	phosphatidylinositol 3-phosphate
PI(3,4)P ₂	phosphatidylinositol 3,4-bisphosphate
PI(4)P	phosphatidylinositol 4-phosphate
PI(4,5)P ₂	phosphatidylinositol 4,5-bisphosphate
PIP ₃	phosphatidylinositol 3,4,5-trisphosphate
PI3K	phosphoinositide 3-kinase
PKA	protein kinase A

PKC	protein kinase C
PLC	phospholipase C
PTEN	phosphatase and tensin homolog deleted on chromosome 10
PX domain	Phox domain
raptor	regulatory-associated protein of mTOR
RBD	Ras binding domain
rictor	rapamycin-insensitive companion of mTOR
RNAi	RNA interference
ROS	reactive oxygen species
rpm	revolutions per minute
RT	reverse transcription <i>or</i> reverse transcriptase
RTK	receptor tyrosine kinase
S6K	p70 ribosomal S6 kinase
SDS	sodium dodecyl sulfate
S.E.	standard error
Sf	<i>Spodoptera frugiperda</i>
SHIP	SH2-containing inositol 5-phosphatase
shRNA	short hairpin RNA
siRNA	short interfering RNA
SMART	simple modular architecture research tool
TEMED	N,N,N',N'-tetramethylethylenediamide
TEV	tobacco etch virus
Tris	Tris(hydroxymethyl)-aminomethan
TRPV	transient receptor potential subfamily V
TSC	tuberous sclerosis complex
VSMC	vascular smooth muscle cell(s)
YFP	yellow fluorescent protein, color variant of GFP

The amino acid three-letter code is used when referring to *e.g.* phosphorylation sites, whereas peptide sequences are given in the one-letter code.

