

8. Literaturverzeichnis:

Amaral, D.G., and Witter, M.P. (1995) Hippocampal formation. In G. Paxinos (ed.): The Rat Nervous System. San Diego, Acad. Press Inc., pp. 443-493.

Angata, K., Long, JM., Bukalo, O., Lee, W., Dityatev, A., Wynshaw-Boris, A., Schachner, M., Fukuda, M., Marth, JD. (2004). Sialyltransferase ST8Sia-II assembles a subset of polysialic acid that directs hippocampal axonal targeting and promotes fear behavior. *J. Biol. Chem.* 279(31):32603-13.

Angevine, JB Jr. (1965). Time of neuron origin in the hippocampal region. An autoradiographic study in the mouse. *Exp. Neurol.* 2:1-70.

Bagri, A Cheng, H.J., Yaron, A., Pleasure, S.J., and Tessier-Lavigne, M. (2003). Stereotyped pruning of long hippocampal axon branches triggered by retraction inducers of the semaphorin family. *Cell* 113: 285-299.

Baldwin, TJ., Fazeli, MS., Doherty, P., Walsh, FS. (1996). Elucidation of the molecular actions of NCAM and structurally related cell adhesion molecules. *J. Cell Biochem.* 61(4):502-13.

Barthels, D., Santoni M.J., Wille W., Ruppert C., Chaix J.C., Hirsch, M.R., Fontecilla-Camps, J.C., and Gordis C. (1987). Isolation and nucleotide sequence of mouse NCAM cDNA that codes for a Mr 79,000 polypeptide without a membrane-spanning region. *EMBO J.* 6(4):907-14. Erratum in: *EMBO J.* 6(10):3202.

Blackstad, T.W. (1958). On the termination of some afferents to the hippocampus and fascia dentata: and experimental study in the rat. *Acta Anat.* 35: 202-214.

Blackstad, T.-W., K. Fuxe, and T. Hokfelt. (1967). Noradrenaline nerve terminals in the hippocampal region of the rat and the guinea pig. *Z. Zellforsch. Mikrosk. Anat.* 78:463-473.

Bolz, J., Novak, N., Götz, M., and Bonhoeffer, T. (1990). Formation of target-specific neuronal projections in organotypic slice cultures from rat visual cortex. *Nature* 346: 359-363.

Büttner, B., Kannicht, C., Schmidt, C., Loster, K., Reutter, W., Lee, HY., Nohring, S., Horstkorte, R. (2002). Biochemical engineering of cell surface sialic acids stimulates axonal growth. *J. Neurosci.* 22(20):8869-75.

Brummendorf, T., Rathjen, FG. (1995). Cell adhesion molecules 1: immunoglobulin superfamily. *Protein Profile.* 2(9):963-1108.

Catalano, SM., Messersmith, EK., Goodman, CS., Shatz, CJ., and Chédotal, A. (1998). Many major CNS axon projections develop normally in the absence of semaphorin 3. *Mol. Cell. Neurosci.* 11:173-182.

Caviness, VS. (1973). Time of neuron origin in the hippocampus and dentate gyrus of normal and reeler mutant mice: an autoradiographic analysis. *J. Comp. Neurol.* 151:113-120.

Caviness, VS. and Rakic, P. (1978). Mechanisms of cortical development: a view from mutations in mice. *Annu. Rev. Neurosci.* 12:97-326.

Cecchi, C. and Boncinelli, E. (2000). Emx homeogenes and mouse brain development. *Trends. Neurosci.* 23:347-352.

Ceranik, K., Deng, J., Heimrich, B., Lübke, J., Zhao, S., Förster, E., and Frotscher, M. (1999). Hippocampal Cajal-retzius cells project to the entorhinal cortex: retrograde tracing and intracellular labeling studies. *Eur. J. Neurosci.* 11: 4278-4290.

Chen, H., He, Z., Bagri, A., and Tessier-Lavigne, M. (1998). Semaphorin-neuropilin interactions underline sympathetic axon responses to class III semaphorins. *Neuron* 21:1283-1290.

Cheng, HJ., Bagri, A., Yaron A., Stein, E., Pleasure, SJ., and Tessier-Lavigne, M. (2001). Plexin-A3 mediates semaphorin signaling and regulates the development of hippocampal axonal projections. *Neuron* 32:249-263.

Chédotal, A., Del Rio, JA., Riuz, M., He, Z., Borrell, V., del Castro, F., Ezan, F., Goodman, CS., Tessier-Lavigne, M., Sotelo, C., Soriano, E. (1998). Semaphorins 3 and 4 repell hippocampal axons via two distinct receptors. *Development* 125:4313-4323.

Clark, GD., Mizuguchi, M., Antalffy, B., Barnes, J., Armstrong, D. (1997). Predominant localization of the LIS family of gene products to Cajal-Retzius cells and ventricular neuroepithelium in the developing human cortex. *J. Neuropathol. Exp. Neurol.* 56(9):1044-52.

Clark, PHG and Oppenheim, RW. (1995). Neuron death in vertebrate development: in vivo methods. *Methods Cell Biol.* 46: 277-321.

Colmarino, SA., and Tessier-Lavigne, M. (1995). The axonal chemoattractant Netrin1 is also a chemorepellent for trochlear motor axons. *Cell* 81:621-629.

Cowan, W.-M., B.-B. Stanfield, and K. Kishi. (1980). The development of the dentate gyrus. *Curr. Top. Dev. Biol.* 15 Pt 1103-57:57.

Cremer, H., Lange, R., Christoph, A., Plomann, M., Vopper, G., Roes, J., Brown, R., Baldwin, S., Kraemer, P., Scheffs, S., et al. (1994).

Inactivation of the N-CAM gene in mice results in size reduction of the olfactory bulb and deficits in spatial learning. *Nature* 367(6462):455-9.

Cunningham, BA., Hemperly, J., Murray, BA., Prediger, EA., Brackenbury, R., Edelman, GM. (1987). Neural cell adhesion molecule: structure, immunoglobulin-like domains, cell surface modulation, and alternative RNA splicing. *Science* 15;236(4803):799-806.

Dalton, D., Chadwick, R., and W. McGinnis. (1989). Expression and embryonic function of empty spiracles: a *Drosophila* homeobox gene with two patterning functions on the anterior-posterior axis of the embryo. *Genes Dev.* 3:1940-1956.

Deller, T., Drakew A, and Frotscher, M. (1999). Different primary target cells are important for fiber lamination in the fascia dentata: a lesson from reeler mutant mice. *Exp. Neurol.* 156:239-253.

Del Rio, J.A., Heimrich, B., Borrell, V., Förster, E., Drakew, A., Alcantara, S., Nakajima, K., Miyata, T., Ogawa, M., Mikoshiba, K., Derer, P., Frotscher, M., and Soriano, E. (1997) A role for Cajal-Retzius cells and reelin in the development of hippocampal connections. *Nature* 385: 70-74.

Drakew, A., Frotscher, M., and Heimrich, B. (1999). Blockade of neuronal activity alters spine maturation of dentate granule cells but not their dendritic arborization. *Neuroscience* 94: 767-774.

D'Eustachio, P., Owens, GC., Edelman, GM., Cunningham, BA. (1985). Chromosomal location of the gene encoding the neural cell adhesion molecule (N-CAM) in the mouse. *Proc. Natl. Acad. Sci. USA.* 82(22):7631-5.

Finne J., Finne U., Deagostini-Bazin, H., and Goridis, C. (1983). Occurrence of alpha 2-8 linked polysialosyl units in a neural cell adhesion molecule. *Biochem. Biophys. Res. Commun.* 29;112(2):482-7.

Förster, E., Kaltschmidt, C., Deng, J., Cremer, H., Deller, T., and Frotscher, M. (1998). Lamina-specific cell adhesion on living slices of hippocampus. *Development* 125: 3399-3410.
Förster, E., Zhao, S., Frotscher, M. (2006). Laminating the hippocampus. *Nat. Rev. Neurosci.* 7(4):259-67.

Frosch, M., Gorgen, I., Boulnois, GJ., Timmis KN., Bitter-Suermann, D. (1985). NZB mouse system for production of monoclonal antibodies to weak bacterial antigens: isolation of an IgG antibody to the polysaccharide capsules of Escherichia coli K1 and group B meningococci. *Proc. Natl. Acad. Sci. USA.* 82(4):1194-8.

Frotscher, M., and Heimrich, B. (1993). Formation of layer-specific fiber projections to the hippocampus in vitro. *Proc. Natl. Acad. Sci. USA* 90: 10400-10403.

Frotscher, M., Zafirov, S., and Heimrich, B. (1995). Development of identified neuronal types and of specific synaptic connections in slice cultures of rat hippocampus. *Prog. Neurobiol.* 45: 143-164.

Frotscher, M., Seress, L., and Heimrich, B. (2001). Early generated Cajal-Retzius cells have different functions in cortical development, p. 43-49. In: M. Thorndyke, P. Beesley, J. Miyan, and C. Bannister (eds.), *Brain Stem Cells*.

Fujisawa, H., Kitsukawa, T., Kawakami, A., Takagi, S., Shimizu, M., and Hirata T. (1997). Roles of a neuronal cell-surface molecule, neuropilin, in nerve fiber fasciculation and guidance. *Cell Tissue Res.* 290(2):465-70.

Gähwiler, B.H. (1984). Slice cultures of cerebellar, hippocampal and hypothalamic tissue. *Experientia* 40: 235-243.

Gähwiler, B.H. (1988) Organotypic cultures of neural tissue. *TINS* 11: 484-489.

Gähwiler, B.H., and Brown, D.A. (1985) Functional innervation of cultured hippocampal neurons by cholinergic afferents from co-cultured septal explants. *Nature* 313: 577-579.

Gähwiler, B. H., Capogna, M., Debanne, D., McKinney, R.A., and Thompson, S.M. (1997). Organotypic slice cultures: a technique has come of age. *TINS* 20: 471-477.

Glüer, S., Schelp, C., Gerady-Schahn, R., and von Schweinitz, D. (1998). Polysialylated neural cell adhesion molecule as a marker for differential diagnosis in pediatric tumors. *J Pediatr Surg.* 33(10):1516-20.

Glumm, R., Klöting A., and Heimrich, B. (2002). Development of axonal projections in cocultures of the hippocampal formation visualized with β -Aktin-gfp transgenic slices. *Neuroembryology* :1 17-22.

Gulisano, M., Broccoli, V., Pardini, C., and Boncinelli, E. (1996). Emx1 and Emx2 show different patterns of expression during proliferation and differentiation of the developing cerebral cortex in the mouse. *Eur. J. Neurosci.* 8:1037-1050.

Gundersen, HJ., Jensen, EB. (1987). The efficiency of systematic sampling in stereology and its prediction. *J. Microsc.* 147(Pt 3):229-63.

Heimrich, B., and Frotscher, M. (1991). Differentiation of dentate granule cells in slice cultures of rat hippocampus: a Golgi/electron microscopic study. *Brain Res.* 538: 263-268.

Heimrich, B., and Frotscher, M. (1994). Slice cultures as a tool to study neuronal development and the formation of specific connections. *Neuroscience Protocols* 94-030-05-01-09.

Heimrich, B., Vogt, J., Simbürger, E., Skutella, T. and Glumm, R. (2002). Axon guidance and the formation of specific connections in the hippocampus. *Neuroembryology* 1: 154-160.

Hildebrandt, H., Becker, C., Murau, M., Gerady-Schahn, R., and Rahmann, H. (1998) Heterogeneous expression of the polysialyltransferases ST8Sia II and ST8Sia IV during postnatal rat brain development. *J. Neurochem.* 71(6):2339-48.

Horstkorte, R., Lee, HY, Lucka, L., Danker, K., Mantey, L., Reutter, W. (2001). Biochemical engineering of the side chain of sialic acids increases the biological stability of the highly sialylated cell adhesion molecule CEACAM1. *Biochem Biophys Res Commun.* 283(1):31-5.

Horstkorte, R., Muhlenhoff, M., Reutter, W., Nohring, S., Zimmermann-Kordmann, M-. Gerady-Schahn, R. (2004). Selective inhibition of polysialyltransferase ST8SiaII by unnatural sialic acids. *Exp. Cell Res.* 298(1):268-74.

Hynes, RO., Lander, AD. (1992). Contact and adhesive specificities in the associations, migrations, and targeting of cells and axons. *Cell.* 24;68(2):303-22.

Kean, L. (1991). Topographical orientation in microsomal vesicles of the N-acetylglucosaminyltransferases which catalyze the biosynthesis of N-acetylglucosaminylpyrophosphoryldolichol and N-acetylglucosaminyl-N-acetylglucosaminylpyrophosphoryldolichol. *J. Biol. Chem.* 266(2):942-6.

Kennedy, TE., Serafini, T., de la Torre, J., Tessier-Lavigne, M. (1994). Netrins are diffusible chemotropic factors for commissural axons in the embryonic spinal cord. *Cell* 78:425-435.

Keppler, OT., Stehling, P., Herrmann, M., Kayser, H., Grunow, D., Reutter, W., Pawlita, M. (1995). Biosynthetic modulation of sialic acid-dependent virus-receptor interactions of two primate polyoma viruses. *J. Biol. Chem.* 270(3):1308-14.

Keppler, OT., Horstkorte, R., Pawlita, M., Schmidt, C., Reutter, W. (2001). Biochemical engineering of the N-acyl side chain of sialic acid: biological implications. *Glycobiology*. 11(2):11R-18R.

Kern, WF., Spier, CM., Hanneman, EH., Miller, TP., Matzner, M., Grogan, TM. (1992). Neural cell adhesion molecule-positive peripheral T-cell lymphoma: a rare variant with a propensity for unusual sites of involvement. *Blood*. 79(9):2432-7.

Kidd, T., Bland, KS., and Goodman, CS. (1999). Slit is the midline repellent for the Robo receptor in *Drosophila*. *Cell*;96:785-794

Lanier, ER., Brown, RM., Kraig, E. (1991). Binding of thymic factors to the conserved decanucleotide promoter element of the T-cell receptor V beta gene is developmentally regulated and is absent in SCID mice. *Proc. Natl. Acad. Sci. USA*. 88(18):8131-5.

Laurberg, S. and Hjorth Simonsen, A. (1977). Growing central axons deprived of normal target neurones by neonatal X-ray irradiation still terminate in a precisely laminated fashion. *Nature* 269:158-160.

Li, D., Field, PM., Starega, U., Li, Y., and Raisman, G. (1993). Entorhinal axons project to dentate gyrus in organotypic slice co-cultures. *Neuroscience* 52: 799-813.

Lipp, H.P., Schwegler, H., Heimrich, B., and Driscol, P. (1988). Infrapyramidal mossy fibers and two-way avoidance learning: developmental modification of hippocampal circuitry and adult behavior of rats and mice. *J. Neurosci.* 8(6):1905-21.

Mahal, LK., Charter, NW., Angata, K., Fukuda, M., Koshland, DE. Jr., Bertozzi, CR. (2001). A small-molecule modulator of poly-alpha 2,8-sialic acid expression on cultured neurons and tumor cells. *Science* 294(5541):380-1.

Mallamaci, A., R. Iannone, P. Briata, L. Pintonello, S. Mercurio, E. Boncinelli, and G. Corte. (1998). EMX2 protein in the developing mouse brain and olfactory area. *Mech. Dev.* 77:165-172.

Mallamaci, A., Muzio, L., Chan, CH., Parnavelas, J., and Boncinelli, E. (2000 a). Area identity shifts in the early cerebral cortex of Emx2^{-/-} mutant mice. *Nat. Neurosci.* 3:679-686.

Mallamaci, A., Mercurio, S., Muzio, L., Cecchi, C., Pardini, CL., Gruss, P., and Boncinelli, E. (2000 b). The lack of Emx2 causes impairment of Reelin signaling and defects of neuronal migration in the developing cerebral cortex. *J. Neurosci.* 20:1109-1118.

Marin Padilla, M. (1978). Dual origin of the mammalian neocortex and evolution of the cortical plate. *Anat. Embryol. (Berl)* 152:109-126.

Meyer, G., und Gofinet, AM. (1998). Prenatal development of reelin-immunoreactive neurons in the human neocortex. *J. Comp. Neurol.* 397(1):29-40.

Michalides, R., Kwa, B., Springall, D., van Zandwijk, N., Koopman, J., Mooi, W. (1994). NCAM and lung cancer. *Int. J. Cancer Suppl.* 8:34-7.

Münster, AK., Weinhold, B., Gotza, B., Mühlenhoff, M., Frosch, M., Gerady-Schahn, R. (2002). Nuclear localization signal of murine CMP-Neu5Ac synthetase includes residues required for both nuclear targeting and enzymatic activity. *J. Biol. Chem.* 277(22):19688-96.

Muller, D., Buchs, PA., and Stoppini, L. (1993). Time course of synaptic development in hippocampal organotypic cultures. *Devl. Brain Res.* 71: 93-100.

Muller, D., Stoppini, L., Wang, C., and Kiss, JZ. (1994). A role for polysialysated neural cell adhesion molecule in lesion-induced sprouting in hippocampal organotypic cultures. *Neuroscience* 61: 441-445.

Nakamura, F., Kalb, RG., and Strittmatter, SM. (2000). Molecular basis of semaphorin-mediated axon guidance. *J. Neurobiol.* 44:219-229.

Naumann, T., Peterson, GM., Frotscher, M. (1992). Fine structure of rat septohippocampal neurons: II. A time course analysis following axotomy. *J. Comp. Neurol.* 325(2):219-42.

Nguyen Ba-Charvet, KT., Brose, K., Marillat, V., Kidd, T., Goodman, CS., Tessier-Lavigne, M., Sotelo, C., and Chédotal, A. (1999). Slit2-mediated chemorepulsion and collapse of developing forebrain axons. *Neuron* 22: 463-473.

Nguyen, C., Mattei, MG, Mattei, JF., Santoni, MJ., Goridis, C., Jordan, BR. (1986). Localization of the human NCAM gene to band q23 of chromosome 11: the third gene coding for a cell interaction molecule mapped to the distal portion of the long arm of chromosome 11. *J. Cell. Biol.* 102(3):711-5.

Nguyen, LB., Ricciardi, TN., Malouf, AT. (1996). Reinnervation of stratum lucidum by hippocampal mossy fibers is developmentally regulated. *Brain Res. Dev. Brain. Res.* 95(2):184-93.

Obst, K., and Wahle, P. (1995). Areal differences of NPY mRNA-expressing neurons are established in the late postnatal rat visual cortex *in vivo*, but not in organotypic cultures. *Eur. J. Neurosci.* 7: 2139-2158.

Özdinler, PH., and Erzurumlu, RS. (2002). Slit2, a branching-arborization factor for sensory axons in the mammalian CNS. *J. Neurosci.* 22:4540-4549.

Pellegrini, M., Mansouri, A., Simeone, A., Boncinelli, E., and Gruss, P. (1996). Dentate gyrus formation requires Emx2. *Development* 122:3893-3898.

Perl, AK., Dahl, U., Wilgenbus, P., Cremer, H., Semb, H., Christofori, G. (1999). Reduced expression of neural cell adhesion molecule induces metastatic dissemination of pancreatic beta tumor cells. *Nat Med.* 5(3):286-91.

Pollerberg, GE., Nolte, C., Schachner, M. (1990). Accumulation of N-CAM 180 at Contact Sites Between Neuroblastoma Cells and Latex Beads Coated with Extracellular Matrix Molecules. *Eur. J. Neurosci.* 2(10):879-887.

Pon, RA., Lussier, M., Yang, QL., Jennings, HJ. (1997). N-Propionylated group B meningococcal polysaccharide mimics a unique bactericidal capsular epitope in group B *Neisseria meningitidis*. *J. Exp. Med.* 185(11):1929-38.

Pozas, E., Pascual, M., Nguyen Ba-Charvet, KT., Guijarro, P., Sotelo, C., Chédotal, A., Del Rio, JA., and Soriano, E. (2001). Age dependent effects of secreted semaphorines 3A, 3F, 3E on developing hippocampal axons: in vitro effects and phenotype of semaphorin 3A (-/-) mice. *Mol. Cell. Neurosci.* 18:26-43.

Raisman, G., Cowan, WM., and Powell, TP. (1966). An experimental analysis of the efferent projection of the hippocampus. *Brain* 89:83-108.

Rutishauser, U., Thiery, JP., Brackenbury, R., Sela, BA., Edelman, GM. (1976). Mechanisms of adhesion among cells from neural tissues of the chick embryo. *Proc Natl Acad Sci U S A.* 73(2):577-81.

Rutishauser, U., Landmesser, L. (1996). Polysialic acid in the vertebrate nervous system: a promoter of plasticity in cell-cell interactions. *Trends Neurosci.* 19(10):422-7.

Sanes, JR., and Yamagata. (1999). M. Formation of laminar-specific synaptic connections. *Curr. Opin. Neurobiol.* 9:79-87.

Savaskan, NE., Alvarez-Bolado, G., Glumm, R., Nitsch, R., Skutella, T. and Heimrich, B. (2002). Impaired postnatal development of hippocampal neurons and axon projections in the Emx2-/- mutants. *J. Neurochem.* 83: 1-12.

Schwab, MH., Bartholomae, A., Heimrich, B., Feldmeyer, D., Druffel, AS., Goebbels, S., Naya, FJ., Zhao, S., Frotscher, M., Tsai, MJ., and Nave, KA. (2000). Neuronal basic helix-loop-helix proteins (NEX and BETA2/Neuro D) regulate terminal granule cell differentiation in the hippocampus. *J. Neurosci.* 20:3714-3724.

Schwegler, H., Heimrich, B., Keller, F., Renner, P., and Crusio, WE. (1988). Strain-specific development of the mossy fiber system in organotypic cultures of the mouse hippocampus. *Neurosci. Lett.* 87:7-10.

Seki, T., Arai, Y. (1993). Highly polysialylated neural cell adhesion molecule (NCAM-H) is expressed by newly generated granule cells in the dentate gyrus of the adult rat. *J. Neurosci.* 13(6):2351-8.

Seki, T., Rutishauser, U. (1998). Removal of polysialic acid-neural cell adhesion molecule induces aberrant mossy fiber innervation and ectopic synaptogenesis in the hippocampus. *J. Neurosci.* 18(10):3757-66.

Semaphorin Nomenclature Committee (1999). Unified nomenclature for the semaphorins/collapsins. *Semaphorin Nomenclature Committee. Cell* 97:551-552.

Serafini, T., Kennedy, TE., Galko, MJ., Mirzyan, C., Jessel, TM., and Tessier-Lavigne, M. (1994). The Netrins define a family of outgrowth promoting proteins homologous to *C elegans* UNC-6. *Cell* 78:409-424.

Simeone, A., Gulisano, M., Acampora, D., Stornaiuolo, A., Rambaldi, M., and Boncinelli, E. (1992). Two vertebrate homeobox genes related to the *Drosophila* empty spiracles gene are expressed in the embryonic cerebral cortex. *EMBO J.* 11:2541-2550.

Skaliora, I., Singer, W., Betz, H., and Püschel, AW. (1998). Differential patterns of semaphorin expression in the developing rat brain. *Eur. J. Neurosci.* 10:1215-1229.

Skutella, T. and Nitsch, R. (2001). New molecules for hippocampal development. *Trends Neurosci.* 24:107-113.

Stanfield, BB., and Cowan, WM. (1979). The development of the hippocampus and dentate gyrus in normal and reeler mice. *J. Comp. Neurol.* 185:423-459.

Stein, E., Savaskan, NE., Ninnemann, O., Nitsch, R., Zhou, R., and Skutella, T. (1999). A role for the Eph-ligand Ephrin-A3 in entorhino-hippocampal axon targeting. *J. Neurosci.* 19: 8885-8893.

Steup, A., Ninnemann, O., Savaskan, N.E., Nitsch, R., Püschel, A.W., and Skutella, T. (1999). Semaphorin D acts as a repulsive factor for entorhinal and hippocampal neurons. *Eur. J. Neurosci.* 11: 729-734.

Steup, A., Lohrum, M., Hamscho, H., Savaskan, NE., Ninneman, O., Nitsch, R., Fujisawa, H., Püschel, AW., and Skutella, T. (2000). Sema3C and Netrin-1 differentially affect axon growth in the hippocampal formation. *Mol. Cell. Neurosci.* 15:141-155.

Stevens, A. and Jacobs, JR. (2002). Integrins regulate responsiveness to slit repellent signals. *J. Neurosci.* 22:4448-4455.

Stoppini, L., Buchs, PA., Muller, D. (1991) A simple method for organotypic cultures of nervous tissue. *J. Neurosci. Meth.* 37: 173-182.

Stoppini, L., Buchs, PA., and Muller, D. (1993). Lesion-induced sprouting and synapse formation in hippocampal organotypic cultures. *Neuroscience* 57: 985-994.

Supèr, H., Martinez, A., Del Rio, JA., and Soriano, E. (1998). Involvement of distinct pioneer neurons in the formation of layer-specific connections in the hippocampus. *J. Neurosci.* 18: 4616-4626.

Szele, FG., Dowling, JJ., Gonzales, C., Theveniau, M., Rougon, G., and Chesselet, MF. (1994). Pattern of expression of highly polysialylated neural cell adhesion molecule in the developing and adult rat striatum. *Neuroscience* 60(1):133-44.

Tamagnone, L., Artigiani, S., Chen, H., He, Z., Ming, GI., Song, H., Chédotal, A., Winberg, ML., Goodman, CS., Poo, M., Tessier-Lavigne, M., and Comoglio, PM. (1999). Plexins are a large family of receptors for transmembrane secreted and GPE-anchored semaphorins in vertebrates. *Cell* 99:71-80.

Tanaka, F., Otake, Y., Nakagawa, T. Kawano, Y., Miyahara, R., Li, M., Yanagihara, K., Inui, K., Oyanagi, H., Yamada, T., Nakayama, J., Fujimoto, I., Ikenaka K., and Wada, H. (2001). Prognostic significance of polysialic acid expression in resected non-small cell lung cancer. *Cancer Res.* 61(4):1666-70.

Tessier-Lavigne, M., and Goodman. CS. (1996). The molecular biology of axon guidance. *Science* 247:1123-133.

Their JP., Brackenbury R., Ruithauser, U., and Edelman GM. (1977). Adhesion among neural cells of the chick embryo. II. Purification and characterization of a cell adhesion molecule from neural retina. *J. Biol. Chem.* 10;252(19):6841-5.

Tole, S., Goudreau, G., Assimacopoulos, S., and Grove, EA. (2000). Emx2 is required for growth of the hippocampus but not for hippocampal field specification. *J. Neurosci.* 20:2618-2625.

Toni, N., Buchs, PA., Nikonenko, I., Bron, CR., and Muller, D. (1999). LTP promotes formation of multiple spine synapses between a single axon terminal and dendrite. *Nature* 402: 421-425.

Varela, FG., Maturana, HR., Uribe, R. (1974). Autopoiesis: the organization of living systems, its characterization and a model. *Curr. Mod. Biol.* 5(4):187-96.

Woodhams, PL. and Terashima, T. (1999). Laminar boundaries persist in the hippocampal dentate molecular layer of the mutant Shaking Rat Kawasaki despite aberrant granule cell migration. *J. Comp. Neurol.* 409:57-70.

Yoshida, M., Suda, Y., Matsuo, I., Miyamoto, N., Takeda, N., Kuratani, S., and Aizawa, S. (1997). Emx1 and Emx2 functions in development of dorsal telencephalon. *Development* 124:101-111.

Zafirov, S., Heimrich, B., and Frotscher, M., (1994). Dendritic development of dentate granule cells in the absence of the specific extrinsic afferents. *J. Comp. Neurol.* 345: 472-480.

Zimmer, J., Gähwiler, BH. (1984). Cellular and connective organization of slice cultures of the rat hippocampus and fascia dentata. *J. Comp. Neurol.* 228(3):432-46.

Zuber, C., Lackie, PM., Catterall, WA. and Roth, J. (1992). Polysialic acid is associated with sodium channels and the neural cell adhesion molecule N-CAM in adult rat brain. *J. Biol. Chem.* 15;267(14):9965-71.