

7. Literaturverzeichnis

- Abel, T., Nguyen, P.V., Barad, M., Deuel, T.A., Kandel, E.R. und Bourtchouladze, R. (1997). Genetic demonstration of a role for PKA in the late phase of LTP and in hippocampus-based long-term memory. *Cell* 88, 615-626.
- Abraham, W.C. und Bear, M.F. (1996). Metaplasticity: the plasticity of synaptic plasticity. *Trends Neurosci.* 19, 126-130.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. und Lipman, D.J. (1990). Basic local alignment search tool. *J. Mol. Biol.* 215, 403-410.
- Amaral, D.G. (1993). Emerging principles of intrinsic hippocampal organization. *Curr. Opin. Neurobiol.* 3, 225-229.
- Amaral, D.G. und Witter, M.P. (1989). The three-dimensional organization of the hippocampal formation: a review of anatomical data. *Neuroscience* 31, 571-591.
- Atkins, C.M., Selcher, J.C., Petraitis, J.J., Trzaskos, J.M. und Sweatt, J.D. (1998). The MAPK cascade is required for mammalian associative learning. *Nat. Neurosci.* 1, 602-609.
- Auernhammer, C.J., Bousquet, C., and Melmed, S. (1999). Autoregulation of pituitary corticotroph SOCS-3 expression: characterization of the murine SOCS-3 promoter. *Proc. Natl. Acad. Sci. U. S. A* 96, 6964-6969.
- Bagni, C., Mannucci, L., Dotti, C.G. und Amaldi, F. (2000). Chemical stimulation of synaptosomes modulates alpha -Ca²⁺/calmodulin-dependent protein kinase II mRNA association to polysomes. *J. Neurosci.* 20, RC76.
- Banbury Conference (1997). Mutant mice and neuroscience: recommendations concerning genetic background. *Banbury Conference on genetic background in mice. Neuron* 19, 755-759.
- Bayer, K.U., De Koninck, P., Leonard, A.S., Hell, J.W. und Schulman, H. (2001). Interaction with the NMDA receptor locks CaMKII in an active conformation. *Nature* 411, 801-805.
- Beer, J., Mielke, K., Zipp, M., Zimmermann, M. und Herdegen, T. (1998). Expression of c-jun, junB, c-fos, fra-1 and fra-2 mRNA in the rat brain following seizure activity and axotomy. *Brain Res.* 794, 255-266.
- Ben Ari, Y. und Cossart, R. (2000). Kainate, a double agent that generates seizures: two decades of progress. *Trends Neurosci.* 23, 580-587.

- Benson,D.L. (1997). Dendritic compartmentation of NMDA receptor mRNA in cultured hippocampal neurons. *Neuroreport* 8, 823-828.
- Berman,D.E., Hazvi,S., Rosenblum,K., Seger,R. und Dudai,Y. (1998). Specific and differential activation of mitogen-activated protein kinase cascades by unfamiliar taste in the insular cortex of the behaving rat. *J. Neurosci.* 18, 10037-10044.
- Bick-Sander,A. (2002). Integration eines auf dem Cre/loxP-System basierenden *targeting* Vektors am genomischen Arg3.1-Lokus mittels homologer Rekombination. Dissertation, Universität Hamburg.
- Bienenstock,E.L., Cooper,L.N. und Munro,P.W. (1982). Theory for the development of neuron selectivity: orientation specificity and binocular interaction in visual cortex. *J. Neurosci.* 2, 32-48.
- Bliss,T.V. und Collingridge,G.L. (1993). A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 361, 31-39.
- Bliss,T.V. und Lømo,T. (1973). Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J. Physiol* 232, 331-356.
- Blum,S., Moore,A.N., Adams,F. und Dash,P.K. (1999). A mitogen-activated protein kinase cascade in the CA1/CA2 subfield of the dorsal hippocampus is essential for long-term spatial memory. *J. Neurosci.* 19, 3535-3544.
- Bockers,T.M., Mameza,M.G., Kreutz,M.R., Bockmann,J., Weise,C., Buck,F., Richter,D., Gundelfinger,E.D. und Kreienkamp,H.J. (2001). Synaptic scaffolding proteins in rat brain. Ankyrin repeats of the multidomain Shank protein family interact with the cytoskeletal protein alpha-fodrin. *J. Biol. Chem.* 276, 40104-40112.
- Bradford,M.M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248-254.
- Brakeman,P.R., Lanahan,A.A., O'Brien,R., Roche,K., Barnes,C.A., Huganir,R.L. und Worley,P.F. (1997). Homer: a protein that selectively binds metabotropic glutamate receptors. *Nature* 386, 284-288.
- Brenman,J.E., Chao,D.S., Gee,S.H., McGee,A.W., Craven,S.E., Santillano,D.R., Wu,Z., Huang,F., Xia,H., Peters,M.F., Froehner,S.C. und Brecht,D.S. (1996). Interaction of nitric oxide synthase with the postsynaptic density protein PSD-95 and alpha1-syntrophin mediated by PDZ domains. *Cell* 84, 757-767.
- Burgin,K.E., Waxham,M.N., Rickling,S., Westgate,S.A., Mobley,W.C. und Kelly,P.T. (1990). In situ hybridization histochemistry of Ca²⁺/calmodulin-dependent protein kinase in developing rat brain. *J. Neurosci.* 10, 1788-1798.

- Buzsáki,G. (1989). Two-stage model of memory trace formation: a role for “noisy” brain states. *Neuroscience* 31, 551-570.
- Casanova,E., Fehsenfeld,S., Mantamadiotis,T., Lemberger,T., Greiner,E., Stewart,A.F. und Schutz,G. (2001). A CamKIIalpha iCre BAC allows brain-specific gene inactivation. *Genesis*. 31, 37-42.
- Chen,L., Chetkovich,D.M., Petralia,R.S., Sweeney,N.T., Kawasaki,Y., Wenthold,R.J., Brecht,D.S. und Nicoll,R.A. (2000). Stargazing regulates synaptic targeting of AMPA receptors by two distinct mechanisms. *Nature* 408, 936-943.
- Chiu,R., Boyle,W.J., Meek,J., Smeal,T., Hunter,T. und Karin,M. (1988). The c-Fos protein interacts with c-Jun/AP-1 to stimulate transcription of AP-1 responsive genes. *Cell* 54, 541-552.
- Crusio,W.E. (1996). Gene-targeting studies: new methods, old problems. *Trends in Neurosciences* 19, 186-187.
- Dammermann,B. (1999). Herstellung und Charakterisierung einer arg3.1-Knockout-Maus. Dissertation, Universität Hamburg.
- De Deyn,P.P., Marescau,B. und MacDonald,R.L. (1990). Epilepsy and the GABA-hypothesis a brief review and some examples. *Acta Neurol. Belg.* 90, 65-81.
- Deadwyler,S.A., Dunwiddie,T. und Lynch,G. (1987). A critical level of protein synthesis is required for long-term potentiation. *Synapse* 1, 90-95.
- Debiec,J., LeDoux,J.E. und Nader,K. (2002). Cellular and systems reconsolidation in the hippocampus. *Neuron* 36, 527-538.
- Deisseroth,K., Bito,H. und Tsien,R.W. (1996). Signaling from synapse to nucleus: postsynaptic CREB phosphorylation during multiple forms of hippocampal synaptic plasticity. *Neuron* 16, 89-101.
- Deisseroth,K., Mermelstein,P.G., Xia,H. und Tsien,R.W. (2003). Signaling from synapse to nucleus: the logic behind the mechanisms. *Curr. Opin. Neurobiol.* 13, 354-365.
- Donai,H., Sugiura,H., Ara,D., Yoshimura,Y., Yamagata,K., and Yamauchi,T. (2003). Interaction of Arc with CaM kinase II and stimulation of neurite extension by Arc in neuroblastoma cells expressing CaM kinase II. *Neurosci. Res.* 47, 399-408.
- Dragunow,M., Abraham,W.C., Goulding,M., Mason,S.E., Robertson,H.A. und Faull,R.L. (1989). Long-term potentiation and the induction of c-fos mRNA and proteins in the dentate gyrus of unanesthetized rats. *Neurosci. Lett.* 101, 274-280.
- Dudai,Y. (2002). Molecular bases of long-term memories: a question of persistence. *Curr. Opin. Neurobiol.* 12, 211-216.

- Ehrlich,I. und Malinow,R. (2004). Postsynaptic Density 95 controls AMPA Receptor Incorporation during Long-Term Potentiation and Experience-Driven Synaptic Plasticity. *J. Neurosci.* 24, 916-927.
- Eichenbaum,H. (2001). The hippocampus and declarative memory: cognitive mechanisms and neural codes. *Behav. Brain Res.* 127, 199-207.
- Engert,F. und Bonhoeffer,T. (1999). Dendritic spine changes associated with hippocampal long-term synaptic plasticity. *Nature* 399, 66-70.
- English,J.D. und Sweatt,J.D. (1997). A requirement for the mitogen-activated protein kinase cascade in hippocampal long term potentiation. *J. Biol. Chem.* 272, 19103-19106.
- Fanselow,M.S. und LeDoux,J.E. (1999). Why we think plasticity underlying Pavlovian fear conditioning occurs in the basolateral amygdala. *Neuron* 23, 229-232.
- Fleischmann,A., Hvalby,O., Jensen,V., Strekalova,T., Zacher,C., Layer,L.E., Kvello,A., Reschke,M., Spanagel,R., Sprengel,R., Wagner,E.F. und Gass,P. (2003). Impaired long-term memory and NR2A-type NMDA receptor-dependent synaptic plasticity in mice lacking c-Fos in the CNS. *J. Neurosci.* 23, 9116-9122.
- Freund,T.F. und Buzsáki,G. (1996). Interneurons of the hippocampus. *Hippocampus* 6, 347-470.
- Frey,U., Frey,S., Schollmeier,F. und Krug,M. (1996). Influence of actinomycin D, a RNA synthesis inhibitor, on long-term potentiation in rat hippocampal neurons in vivo and in vitro. *J. Physiol* 490, 703-711.
- Frey,U., Huang,Y.Y. und Kandel,E.R. (1993). Effects of cAMP simulate a late stage of LTP in hippocampal CA1 neurons. *Science* 260, 1661-1664.
- Frey,U., Krug,M., Reymann,K.G. und Matthies,H. (1988). Anisomycin, an inhibitor of protein synthesis, blocks late phases of LTP phenomena in the hippocampal CA1 region in vitro. *Brain Res.* 452, 57-65.
- Gallagher,M., Burwell,R. und Burchinal,M. (1993). Severity of spatial learning impairment in aging: development of a learning index for performance in the Morris water maze. *Behav. Neurosci.* 107, 618-626.
- Garner,C.C., Tucker,R.P. und Matus,A. (1988). Selective localization of messenger RNA for cytoskeletal protein MAP2 in dendrites. *Nature* 336, 674-677.
- Gille,H. und Downward,J. (1999). Multiple ras effector pathways contribute to G(1) cell cycle progression. *J. Biol. Chem.* 274, 22033-22040.
- Goelet,P., Castellucci,V.F., Schacher,S. und Kandel,E.R. (1986). The long and the short of long-term memory - a molecular framework. *Nature* 322, 419-422.

- Grant,S.G., O'Dell,T.J., Karl,K.A., Stein,P.L., Soriano,P. und Kandel,E.R. (1992). Impaired long-term potentiation, spatial learning, and hippocampal development in fyn mutant mice. *Science* 258, 1903-1910.
- Greenberg,M.E., Ziff,E.B. und Greene,L.A. (1986). Stimulation of neuronal acetylcholine receptors induces rapid gene transcription. *Science* 234, 80-83.
- Grewal,S.S., Fass,D.M., Yao,H., Ellig,C.L., Goodman,R.H. und Stork,P.J. (2000). Calcium and cAMP signals differentially regulate cAMP-responsive element-binding protein function via a Rap1-extracellular signal-regulated kinase pathway. *J. Biol. Chem.* 275, 34433-34441.
- Guzowski,J.F., Lyford,G.L., Stevenson,G.D., Houston,F.P., McGaugh,J.L., Worley,P.F. und Barnes,C.A. (2000). Inhibition of activity-dependent arc protein expression in the rat hippocampus impairs the maintenance of long-term potentiation and the consolidation of long-term memory. *J. Neurosci.* 20, 3993-4001.
- Guzowski,J.F., McNaughton,B.L., Barnes,C.A. und Worley,P.F. (1999). Environment-specific expression of the immediate-early gene Arc in hippocampal neuronal ensembles. *Nat. Neurosci.* 2, 1120-1124.
- Guzowski,J.F., Setlow,B., Wagner,E.K. und McGaugh,J.L. (2001). Experience-dependent gene expression in the rat hippocampus after spatial learning: a comparison of the immediate-early genes Arc, c-fos, and zif268. *J. Neurosci.* 21, 5089-5098.
- Haan,S., Ferguson,P., Sommer,U., Hiremath,M., McVicar,D.W., Heinrich,P.C., Johnston,J.A. und Cacalano,N.A. (2003). Tyrosine phosphorylation disrupts elongin interaction and accelerates SOCS3 degradation. *J. Biol. Chem.* 278, 31972-31979.
- Halazonetis,T.D., Georgopoulos,K., Greenberg,M.E. und Leder,P. (1988). c-Jun dimerizes with itself and with c-Fos, forming complexes of different DNA binding affinities. *Cell* 55, 917-924.
- Hazzalin,C.A., Le Panse,R., Cano,E. und Mahadevan,L.C. (1998). Anisomycin selectively desensitizes signalling components involved in stress kinase activation and fos and jun induction. *Mol. Cell Biol.* 18, 1844-1854.
- Hebb,D.O. (1949). *The Organization of Behaviour: a Neurophysiological Theory*. New York: Wiley.
- Hollmann,M. und Heinemann,S. (1994). Cloned glutamate receptors. *Annu. Rev. Neurosci.* 17, 31-108.
- Huang,Y.Y., Li,X.C. und Kandel,E.R. (1994). cAMP contributes to mossy fiber LTP by initiating both a covalently mediated early phase and macromolecular synthesis-dependent late phase. *Cell* 79, 69-79.

- Husi,H., Ward,M.A., Choudhary,J.S., Blackstock,W.P. und Grant,S.G. (2000). Proteomic analysis of NMDA receptor-adhesion protein signaling complexes. *Nat. Neurosci.* 3, 661-669.
- Husi,H. und Grant,S.G.N. (2001). Proteomics of the nervous system. *Trends in Neurosciences* 24, 259-266.
- Hyman,S.E. und Malenka,R.C. (2001). Addiction and the brain: the neurobiology of compulsion and its persistence. *Nat. Rev. Neurosci.* 2, 695-703.
- Impey,S., Mark,M., Villacres,E.C., Poser,S., Chavkin,C. und Storm,D.R. (1996). Induction of CRE-mediated gene expression by stimuli that generate long-lasting LTP in area CA1 of the hippocampus. *Neuron* 16, 973-982.
- Inoue,H., Nojima,H. und Okayama,H. (1990). High efficiency transformation of *Escherichia coli* with plasmids. *Gene* 96, 23-28.
- Irie,M., Hata,Y., Takeuchi,M., Ichtchenko,K., Toyoda,A., Hirao,K., Takai,Y., Rosahl,T.W. und Sudhof,T.C. (1997). Binding of neuroligins to PSD-95. *Science* 277, 1511-1515.
- Johnston,M.V. (2004). Clinical disorders of brain plasticity. *Brain Dev.* 26, 73-80.
- Kang,H. und Schuman,E.M. (1996). A requirement for local protein synthesis in neurotrophin-induced hippocampal synaptic plasticity. *Science* 273, 1402-1406.
- Kang,H., Sun,L.D., Atkins,C.M., Soderling,T.R., Wilson,M.A. und Tonegawa,S. (2001). An important role of neural activity-dependent CaMKIV signaling in the consolidation of long-term memory. *Cell* 106, 771-783.
- Karin,M., Liu,Z. und Zandi,E. (1997). AP-1 function and regulation. *Curr. Opin. Cell Biol.* 9, 240-246.
- Kauselmann,G., Weiler,M., Wulff,P., Jessberger,S., Konietzko,U., Scafidi,J., Staubli,U., Bereiter-Hahn,J., Strebhardt,K. und Kuhl,D. (1999). The polo-like protein kinases Fnk and Snk associate with a Ca(2+)- and integrin-binding protein and are regulated dynamically with synaptic plasticity. *EMBO J.* 18, 5528-5539.
- Kelly,M.P. und Deadwyler,S.A. (2002). Acquisition of a novel behavior induces higher levels of Arc mRNA than does overtrained performance. *Neuroscience* 110, 617-626.
- Kennedy,M.B. (2000). Signal-processing machines at the postsynaptic density. *Science* 290, 750-754.
- Kim,E., Cho,K.O., Rothschild,A. und Sheng,M. (1996). Heteromultimerization and NMDA receptor-clustering activity of Chapsyn-110, a member of the PSD-95 family of proteins. *Neuron* 17, 103-113.

- Kim,E., Naisbitt,S., Hsueh,Y.P., Rao,A., Rothschild,A., Craig,A.M. und Sheng,M. (1997). GKAP, a novel synaptic protein that interacts with the guanylate kinase-like domain of the PSD-95/SAP90 family of channel clustering molecules. *J. Cell Biol.* *136*, 669-678.
- Kim,E., Niethammer,M., Rothschild,A., Jan,Y.N. und Sheng,M. (1995). Clustering of Shaker-type K⁺ channels by interaction with a family of membrane-associated guanylate kinases. *Nature* *378*, 85-88.
- Kim,J.H., Liao,D., Lau,L.F. und Huganir,R.L. (1998). SynGAP: a synaptic RasGAP that associates with the PSD-95/SAP90 protein family. *Neuron* *20*, 683-691.
- Kim,J.J., Rison,R.A. und Fanselow,M.S. (1993). Effects of amygdala, hippocampus, and periaqueductal gray lesions on short- and long-term contextual fear. *Behav. Neurosci.* *107*, 1093-1098.
- Kistner,A., Gossen,M., Zimmermann,F., Jerecic,J., Ullmer,C., Lubbert,H. und Bujard,H. (1996). Doxycycline-mediated quantitative and tissue-specific control of gene expression in transgenic mice. *Proc. Natl. Acad. Sci. U. S. A* *93*, 10933-10938.
- Klebaur,J.E., Ostrander,M.M., Norton,C.S., Watson,S.J., Akil,H. und Robinson,T.E. (2002). The ability of amphetamine to evoke arc (Arg 3.1) mRNA expression in the caudate, nucleus accumbens and neocortex is modulated by environmental context. *Brain Res.* *930*, 30-36.
- Komiyama,N.H., Watabe,A.M., Carlisle,H.J., Porter,K., Charlesworth,P., Monti,J., Strathdee,D.J., O'Carroll,C.M., Martin,S.J., Morris,R.G., O'Dell,T.J. und Grant,S.G. (2002). SynGAP regulates ERK/MAPK signaling, synaptic plasticity, and learning in the complex with postsynaptic density 95 and NMDA receptor. *J. Neurosci.* *22*, 9721-9732.
- Konietzko,U., Kauselmann,G., Scafidi,J., Staubli,U., Mikkers,H., Berns,A., Schweizer,M., Waltereit,R. und Kuhl,D. (1999). Pim kinase expression is induced by LTP stimulation and required for the consolidation of enduring LTP. *EMBO J.* *18*, 3359-3369.
- Kornau,H.C., Schenker,L.T., Kennedy,M.B. und Seeburg,P.H. (1995). Domain interaction between NMDA receptor subunits and the postsynaptic density protein PSD-95. *Science* *269*, 1737-1740.
- Kornau,H.C., Seeburg,P.H., and Kennedy,M.B. (1997). Interaction of ion channels and receptors with PDZ domain proteins. *Curr. Opin. Neurobiol.* *7*, 368-373.
- Krebs,D.L. und Hilton,D.J. (2001). SOCS proteins: negative regulators of cytokine signaling. *Stem Cells* *19*, 378-387.
- Krug,M., Lossner,B. und Ott,T. (1984). Anisomycin blocks the late phase of long-term potentiation in the dentate gyrus of freely moving rats. *Brain Res. Bull.* *13*, 39-42.

- Kuhl, D. Learning about activity dependent genes. In: Advances in synaptic plasticity (Baudry M, Davis JL, Thompson RF, eds). Boston: MIT, 1-31. 2000.
- Kuhl,D. und Skehel,P. (1998). Dendritic localization of mRNAs. *Curr. Opin. Neurobiol.* 8, 600-606.
- Lamprecht,R. und LeDoux,J. (2004). Structural plasticity and memory. *Nat. Rev. Neurosci.* 5, 45-54.
- Lanahan,A. und Worley,P. (1998). Immediate-Early Genes and Synaptic Function. *Neurobiol. Learn. Mem.* 70, 37-43.
- Lashley,K.S. (1929). *Brain Mechanisms and Intelligence: a Quantitative Study of Injuries to the Brain.* Chicago: Chicago University Press.
- Link,W., Konietzko,U., Kauselmann,G., Krug,M., Schwanke,B., Frey,U. und Kuhl,D. (1995). Somatodendritic expression of an immediate early gene is regulated by synaptic activity. *Proc. Natl. Acad. Sci. U. S. A* 92, 5734-5738.
- Lipp,H.P. und Wolfer,D.P. (1998). Genetically modified mice and cognition. *Curr. Opin. Neurobiol.* 8, 272-280.
- Lisman,J. (2003). Long-term potentiation: outstanding questions and attempted synthesis. *Philos. Trans. R. Soc. Lond B Biol. Sci.* 358, 829-842.
- Lisman,J., Schulman,H. und Cline,H. (2002). The molecular basis of CaMKII function in synaptic and behavioural memory. *Nat. Rev. Neurosci.* 3, 175-190.
- Liu,D., Bei,D., Parmar,H. und Matus,A. (2000). Activity-regulated, cytoskeleton-associated protein (Arc) is essential for visceral endoderm organization during early embryogenesis. *Mech. Dev.* 92, 207-215.
- Liu,E., Cote,J.F. und Vuori,K. (2003). Negative regulation of FAK signaling by SOCS proteins. *EMBO J.* 22, 5036-5046.
- Logue,S.F., Paylor,R. und Wehner,J.M. (1997). Hippocampal lesions cause learning deficits in inbred mice in the Morris water maze and conditioned-fear task. *Behav. Neurosci.* 111, 104-113.
- Lyford,G.L., Yamagata,K., Kaufmann,W.E., Barnes,C.A., Sanders,L.K., Copeland,N.G., Gilbert,D.J., Jenkins,N.A., Lanahan,A.A. und Worley,P.F. (1995). Arc, a growth factor and activity-regulated gene, encodes a novel cytoskeleton-associated protein that is enriched in neuronal dendrites. *Neuron* 14, 433-445.
- Lynch,G., Larson,J., Kelso,S., Barrionuevo,G., and Schottler,F. (1983). Intracellular injections of EGTA block induction of hippocampal long-term potentiation. *Nature* 305, 719-721.
- Malenka,R.C., Lancaster,B. und Zucker,R.S. (1992). Temporal limits on the rise in postsynaptic calcium required for the induction of long-term potentiation. *Neuron* 9, 121-128.

- Malinow, R. und Malenka, R.C. (2002). AMPA receptor trafficking and synaptic plasticity. *Annu. Rev. Neurosci.* 25, 103-126.
- Malinow, R., Schulman, H. und Tsien, R.W. (1989). Inhibition of postsynaptic PKC or CaMKII blocks induction but not expression of LTP. *Science* 245, 862-866.
- Martin, K.C., Casadio, A., Zhu, H., Yaping, E., Rose, J.C., Chen, M., Bailey, C.H., and Kandel, E.R. (1997). Synapse-specific, long-term facilitation of aplysia sensory to motor synapses: a function for local protein synthesis in memory storage. *Cell* 91, 927-938.
- Martin, S.J., Grimwood, P.D. und Morris, R.G. (2000). Synaptic plasticity and memory: an evaluation of the hypothesis. *Annu. Rev. Neurosci.* 23:649-711., 649-711.
- Matus, A. (2000). Actin-based plasticity in dendritic spines. *Science* 290, 754-758.
- Mauch, D.H., Nagler, K., Schumacher, S., Goritz, C., Muller, E.C., Otto, A. und Pfrieder, F.W. (2001). CNS synaptogenesis promoted by glia-derived cholesterol. *Science* 294, 1354-1357.
- McHugh, T.J., Blum, K.I., Tsien, J.Z., Tonegawa, S. und Wilson, M.A. (1996). Impaired hippocampal representation of space in CA1-specific NMDAR1 knockout mice. *Cell* 87, 1339-1349.
- McNamara, T.P. und Shelton, A.L. (2003). Cognitive maps and the hippocampus. *Trends Cogn Sci.* 7, 333-335.
- Migaud, M., Charlesworth, P., Dempster, M., Webster, L.C., Watabe, A.M., Makhinson, M., He, Y., Ramsay, M.F., Morris, R.G., Morrison, J.H., O'Dell, T.J. und Grant, S.G. (1998). Enhanced long-term potentiation and impaired learning in mice with mutant postsynaptic density-95 protein. *Nature* 396, 433-439.
- Miller, S., Yasuda, M., Coats, J.K., Jones, Y., Martone, M.E. und Mayford, M. (2002). Disruption of dendritic translation of CaMKIIalpha impairs stabilization of synaptic plasticity and memory consolidation. *Neuron* 36, 507-519.
- Milner, B., Squire, L.R. und Kandel, E.R. (1998). Cognitive neuroscience and the study of memory. *Neuron* 20, 445-468.
- Minichiello, L., Korte, M., Wolfer, D., Kuhn, R., Unsicker, K., Cestari, V., Rossi-Arnaud, C., Lipp, H.P., Bonhoeffer, T. und Klein, R. (1999). Essential role for TrkB receptors in hippocampus-mediated learning. *Neuron* 24, 401-414.
- Montag-Sallaz, M., Welzl, H., Kuhl, D., Montag, D. und Schachner, M. (1999). Novelty-induced increased expression of immediate-early genes c-fos and arg 3.1 in the mouse brain. *J. Neurobiol.* 38, 234-246.
- Morgan, J.I., Cohen, D.R., Hempstead, J.L. und Curran, T. (1987). Mapping patterns of c-fos expression in the central nervous system after seizure. *Science* 237, 192-197.

- Morris,R. (1984). Developments of a water-maze procedure for studying spatial learning in the rat. *J. Neurosci. Methods* *11*, 47-60.
- Morris,R.G., Anderson,E., Lynch,G.S. und Baudry,M. (1986). Selective impairment of learning and blockade of long-term potentiation by an N-methyl-D-aspartate receptor antagonist, AP5. *Nature* *319*, 774-776.
- Morris,R.G., Garrud,P., Rawlins,J.N. und O'Keefe,J. (1982). Place navigation impaired in rats with hippocampal lesions. *Nature* *297*, 681-683.
- Muller,R., Tremblay,J.M., Adamson,E.D. und Verma,I.M. (1983). Tissue and cell type-specific expression of two human c-onc genes. *Nature* *304*, 454-456.
- Nader,K. (2003). Memory traces unbound. *Trends in Neurosciences* *26*, 65-72.
- Naisbitt,S., Kim,E., Tu,J.C., Xiao,B., Sala,C., Valtschanoff,J., Weinberg,R.J., Worley,P.F. und Sheng,M. (1999). Shank, a novel family of postsynaptic density proteins that binds to the NMDA receptor/PSD-95/GKAP complex and cortactin. *Neuron* *23*, 569-582.
- Nedivi,E., Hevroni,D., Naot,D., Israeli,D. und Citri,Y. (1993). Numerous candidate plasticity-related genes revealed by differential cDNA cloning. *Nature* *363*, 718-722.
- Nguyen,P.V., Abel,T., Kandel,E.R. und Bourtchouladze,R. (2000). Strain-dependent differences in LTP and hippocampus-dependent memory in inbred mice. *Learn. Mem.* *7*, 170-179.
- Nguyen,P.V. und Kandel,E.R. (1996). A macromolecular synthesis-dependent late phase of long-term potentiation requiring cAMP in the medial perforant pathway of rat hippocampal slices. *J. Neurosci.* *16*, 3189-3198.
- Nowak,L., Bregestovski,P., Ascher,P., Herbet,A. und Prochiantz,A. (1984). Magnesium gates glutamate-activated channels in mouse central neurones. *Nature* *307*, 462-465.
- Okazaki,Y., Furuno,M., Kasukawa,T., Adachi,J., Bono,H., Kondo,S., Nikaido,I., et al. (2002). Analysis of the mouse transcriptome based on functional annotation of 60,770 full-length cDNAs. *Nature* *420*, 563-573.
- O'Keefe,J. (1976). Place units in the hippocampus of the freely moving rat. *Exp. Neurol.* *51*, 78-109.
- O'Keefe,J. und Nadel,L. (1978). *The hippocampus as a cognitive map*. Clarendon Press.
- Osten,P., Valsamis,L., Harris,A. und Sacktor,T.C. (1996). Protein synthesis-dependent formation of protein kinase Mzeta in long-term potentiation. *J. Neurosci.* *16*, 2444-2451.
- Otani,S. und Abraham,W.C. (1989). Inhibition of protein synthesis in the dentate gyrus, but not the entorhinal cortex, blocks maintenance of long-term potentiation in rats. *Neurosci. Lett.* *106*, 175-180.

- Penfield,W. und Milner,B. (1958). Memory deficit produced by bilateral lesions in the hippocampal zone. *AMA. Arch. Neurol. Psychiatry* 79, 475-497.
- Pinaud,R., Penner,M.R., Robertson,H.A. und Currie,R.W. (2001). Upregulation of the immediate early gene arc in the brains of rats exposed to environmental enrichment: implications for molecular plasticity. *Brain Res. Mol. Brain Res.* 91, 50-56.
- Pinaud,R., Tremere,L.A., Penner,M.R., Hess,F.F., Barnes,S., Robertson,H.A. und Currie,R.W. (2002). Plasticity-driven gene expression in the rat retina. *Brain Res. Mol. Brain Res.* 98, 93-101.
- Qian,Z., Gilbert,M.E., Colicos,M.A., Kandel,E.R. und Kuhl,D. (1993). Tissue-plasminogen activator is induced as an immediate-early gene during seizure, kindling and long-term potentiation. *Nature* 361, 453-457.
- Rial-Verde,E.M., Worley,P., Malinow,R. und Cline,H.T. The immediate-early gene Arc regulates AMPA-receptor trafficking. Abstract Viewer and Itinerary Planner. Washington, DC: Society for Neuroscience, 2003. Online.
- Robinson,R. (2004). RNAi Therapeutics: How Likely, How Soon? *PLoS Biology* 2, e28.
- Romanski,L.M., Clugnet,M.C., Bordi,F. und LeDoux,J.E. (1993). Somatosensory and auditory convergence in the lateral nucleus of the amygdala. *Behav. Neurosci.* 107, 444-450.
- Rosell,D.R., Akama,K.T., Nacher,J. und McEwen,B.S. (2003). Differential expression of suppressors of cytokine signaling-1, -2, and -3 in the rat hippocampus after seizure: implications for neuromodulation by gp130 cytokines. *Neuroscience* 122, 349-358.
- Rosenblum,K., Berman,D.E., Hazvi,S., Lamprecht,R. und Dudai,Y. (1997). NMDA receptor and the tyrosine phosphorylation of its 2B subunit in taste learning in the rat insular cortex. *J. Neurosci.* 17, 5129-5135.
- Saffen,D.W., Cole,A.J., Worley,P.F., Christy,B.A., Ryder,K. und Baraban,J.M. (1988). Convulsant-induced increase in transcription factor messenger RNAs in rat brain. *Proc. Natl. Acad. Sci. U. S. A* 85, 7795-7799.
- Sakimura,K., Kutsuwada,T., Ito,I., Manabe,T., Takayama,C., Kushiya,E., Yagi,T., Aizawa,S., Inoue,Y. und Sugiyama,H. (1995). Reduced hippocampal LTP and spatial learning in mice lacking NMDA receptor epsilon 1 subunit. *Nature* 373, 151-155.
- Sambrook,J. und Russel, D.W. (2001). *Molecular Cloning. A laboratory manual.* Cold Spring Harbor Press, New York.
- Sanger,F., Nicklen,S. und Coulson,A.R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. U. S. A* 74, 5463-5467.

- Sassone-Corsi,P. (1995). Transcription factors responsive to cAMP. *Annu. Rev. Cell Dev. Biol.* *11*, 355-377.
- Schnell,E., Sizemore,M., Karimzadegan,S., Chen,L., Brecht,D.S. und Nicoll,R.A. (2002). Direct interactions between PSD-95 and stargazin control synaptic AMPA receptor number. *Proc. Natl. Acad. Sci. U. S. A* *99*, 13902-13907.
- Scoville,W.B. und Milner,B. (1957). Loss of recent memory after bilateral hippocampal lesions. *J. Neurochem.* *20*, 11-21.
- Setlow,B. und McGaugh,J.L. (1999). Involvement of the posteroventral caudate-putamen in memory consolidation in the Morris water maze. *Neurobiol. Learn. Mem.* *71*, 240-247.
- Sheng,M. (2001). Molecular organization of the postsynaptic specialization. *Proc. Natl. Acad. Sci. U. S. A* *98*, 7058-7061.
- Sheng,M. und Greenberg,M.E. (1990). The regulation and function of c-fos and other immediate early genes in the nervous system. *Neuron* *4*, 477-485.
- Sheng,M. und Kim,E. (2000). The Shank family of scaffold proteins. *J. Cell Sci.* *113*, 1851-1856.
- Sheng,M. und Kim,M.J. (2002). Postsynaptic signaling and plasticity mechanisms. *Science* *298*, 776-780.
- Shimizu,E., Tang,Y.P., Rampon,C. und Tsien,J.Z. (2000). NMDA receptor-dependent synaptic reinforcement as a crucial process for memory consolidation. *Science* *290*, 1170-1174.
- Siekevitz,P. (1985). The postsynaptic density: a possible role in long-lasting effects in the central nervous system. *Proc. Natl. Acad. Sci. U. S. A* *82*, 3494-3498.
- Silva,A.J. (2003). Molecular and cellular cognitive studies of the role of synaptic plasticity in memory. *J. Neurobiol.* *54*, 224-237.
- Silva,A.J., Paylor,R., Wehner,J.M. und Tonegawa,S. (1992). Impaired spatial learning in alpha-calmodulin-calmodulin kinase II mutant mice. *Science* *257*, 206-211.
- Soderling,T.R. und Derkach,V.A. (2000). Postsynaptic protein phosphorylation and LTP. *Trends Neurosci.* *23*, 75-80.
- Soriano,S.F., Hernanz-Falcon,P., Rodriguez-Frade,J.M., De Ana,A.M., Garzon,R., Carvalho-Pinto,C., Vila-Coro,A.J., Zaballos,A., Balomenos,D., Martinez,A. und Mellado,M. (2002). Functional inactivation of CXCR4-mediated responses through SOCS3 up-regulation. *J. Exp. Med.* *196*, 311-321.
- Sprengel,R., Suchanek,B., Amico,C., Brusa,R., Burnashev,N., Rozov,A., Hvalby,O., Jensen,V., Paulsen,O., Andersen,P., Kim,J.J., Thompson,R.F., Sun,W., Webster,L.C., Grant,S.G., Eilers,J.,

- Konnerth,A., Li,J., McNamara,J.O. und Seeburg,P.H. (1998). Importance of the intracellular domain of NR2 subunits for NMDA receptor function in vivo. *Cell* 92, 279-289.
- Squire,L.R., Knowlton,B. und Musen,G. (1993). The Structure and Organization of Memory. *Annual Review of Psychology* 44, 453-495.
- Stanton,P.K. und Sarvey,J.M. (1984). Blockade of long-term potentiation in rat hippocampal CA1 region by inhibitors of protein synthesis. *J. Neurosci.* 4, 3080-3088.
- Steward,O. und Levy,W.B. (1982). Preferential localization of polyribosomes under the base of dendritic spines in granule cells of the dentate gyrus. *J. Neurosci.* 2, 284-291.
- Steward,O. und Schuman,E.M. (2003). Compartmentalized synthesis and degradation of proteins in neurons. *Neuron* 40, 347-359.
- Steward,O., Wallace,C.S., Lyford,G.L. und Worley,P.F. (1998). Synaptic activation causes the mRNA for the IEG Arc to localize selectively near activated postsynaptic sites on dendrites. *Neuron* 21, 741-751.
- Steward,O. and Worley,P.F. (2001). A cellular mechanism for targeting newly synthesized mRNAs to synaptic sites on dendrites. *Proc. Natl. Acad. Sci. U. S. A* 98, 7062-7068.
- Sweatt,J.D. (2001). The neuronal MAP kinase cascade: a biochemical signal integration system subserving synaptic plasticity and memory. *J. Neurochem.* 76, 1-10.
- Tan,A., Moratalla,R., Lyford,G.L., Worley,P. und Graybiel,A.M. (2000). The activity-regulated cytoskeletal-associated protein arc is expressed in different striosome-matrix patterns following exposure to amphetamine and cocaine. *J. Neurochem.* 74, 2074-2078.
- Tang,Y.P., Shimizu,E., Dube,G.R., Rampon,C., Kerchner,G.A., Zhuo,M., Liu,G. und Tsien,J.Z. (1999). Genetic enhancement of learning and memory in mice. *Nature* 401, 63-69.
- Thomas,G.M. und Huganir,R.L. (2004). MAPK cascade signalling and synaptic plasticity. *Nat. Rev. Neurosci.* 5, 173-183.
- Tischmeyer,W. und Grimm,R. (1999). Activation of immediate early genes and memory formation. *Cell Mol. Life Sci.* 55, 564-574.
- Tonegawa,S., Li,Y., Erzurumlu,R.S., Jhaveri,S., Chen,C., Goda,Y., Paylor,R., Silva,A.J., Kim,J.J. und Wehner,J.M. (1995). The gene knockout technology for the analysis of learning and memory, and neural development. *Prog. Brain Res.* 105, 3-14.
- Toni,N., Buchs,P.A., Nikonenko,I., Bron,C.R. und Muller,D. (1999). LTP promotes formation of multiple spine synapses between a single axon terminal and a dendrite. *Nature* 402, 421-425.

- Toni, N., Buchs, P.A., Nikonenko, I., Povilaitite, P., Parisi, L. und Muller, D. (2001). Remodeling of synaptic membranes after induction of long-term potentiation. *J. Neurosci.* *21*, 6245-6251.
- Tsien, J.Z., Chen, D.F., Gerber, D., Tom, C., Mercer, E.H., Anderson, D.J., Mayford, M., Kandel, E.R. und Tonegawa, S. (1996a). Subregion- and cell type-restricted gene knockout in mouse brain. *Cell* *87*, 1317-1326.
- Tsien, J.Z., Huerta, P.T. und Tonegawa, S. (1996b). The essential role of hippocampal CA1 NMDA receptor-dependent synaptic plasticity in spatial memory. *Cell* *87*, 1327-1338.
- Tu, J.C., Xiao, B., Naisbitt, S., Yuan, J.P., Petralia, R.S., Brakeman, P., Doan, A., Aakalu, V.K., Lanahan, A.A., Sheng, M. und Worley, P.F. (1999). Coupling of mGluR/Homer and PSD-95 complexes by the Shank family of postsynaptic density proteins. *Neuron* *23*, 583-592.
- Turnley, A.M. und Bartlett, P.F. (2000). Cytokines that signal through the leukemia inhibitory factor receptor-beta complex in the nervous system. *J. Neurochem.* *74*, 889-899.
- Wallace, C.S., Lyford, G.L., Worley, P.F. und Steward, O. (1998). Differential intracellular sorting of immediate early gene mRNAs depends on signals in the mRNA sequence. *J. Neurosci.* *18*, 26-35.
- Waltereit, R., Dammermann, B., Wulff, P., Scafidi, J., Staubli, U., Kauselmann, G., Bundman, M. und Kuhl, D. (2001). Arg3.1/Arc mRNA induction by Ca²⁺ and cAMP requires protein kinase A and mitogen-activated protein kinase/extracellular regulated kinase activation. *J. Neurosci.* *21*, 5484-5493.
- Wang, J. und Campbell, I.L. (2002). Cytokine signaling in the brain: putting a SOCS in it? *J. Neurosci. Res.* *67*, 423-427.
- Warren, S.T. und Ashley, C.T., Jr. (1995). Triplet repeat expansion mutations: the example of fragile X syndrome. *Annu. Rev. Neurosci.* *18*, 77-99.
- Watabe, A.M., Zaki, P.A. und O'Dell, T.J. (2000). Coactivation of beta-adrenergic and cholinergic receptors enhances the induction of long-term potentiation and synergistically activates mitogen-activated protein kinase in the hippocampal CA1 region. *J. Neurosci.* *20*, 5924-5931.
- Weiler, I.J., Irwin, S.A., Klintsova, A.Y., Spencer, C.M., Brazelton, A.D., Miyashiro, K., Comery, T.A., Patel, B., Eberwine, J. und Greenough, W.T. (1997). Fragile X mental retardation protein is translated near synapses in response to neurotransmitter activation. *Proc. Natl. Acad. Sci. U. S. A* *94*, 5395-5400.
- Welzl, H., D'Adamo, P. und Lipp, H.P. (2001). Conditioned taste aversion as a learning and memory paradigm. *Behav. Brain Res.* *125*, 205-213.

- Winder,D.G., Martin,K.C., Muzzio,I.A., Rohrer,D., Chruscinski,A., Kobilka,B. und Kandel,E.R. (1999). ERK plays a regulatory role in induction of LTP by theta frequency stimulation and its modulation by beta-adrenergic receptors. *Neuron* 24, 715-726.
- Wolfer,D.P., Crusio,W.E. und Lipp,H.P. (2002). Knockout mice: simple solutions to the problems of genetic background and flanking genes. *Trends Neurosci.* 25, 336-340.
- Wolfer,D.P. und Lipp,H.P. Meta-analysis of strategy choices by 85 mutant mouse lines in a standardized place navigation task identifies behavioral flexibility as performance limiting factor. Abstract Viewer and Itinerary Planner. Washington, DC: Society for Neuroscience, 2003. Online
- Wolfer,D.P., Stagljjar-Bozicevic,M., Errington,M.L. und Lipp,H.P. (1998). Spatial Memory and Learning in Transgenic Mice: Fact or Artifact? *News Physiol Sci.* 13, 118-123.
- Worley,P.F., Bhat,R.V., Baraban,J.M., Erickson,C.A., McNaughton,B.L. und Barnes,C.A. (1993). Thresholds for synaptic activation of transcription factors in hippocampus: correlation with long-term enhancement. *J. Neurosci.* 13, 4776-4786.
- Wormald,S. und Hilton,D.J. (2004). Inhibitors of cytokine signal transduction. *J. Biol. Chem.* 279, 821-824.
- Xia,Z., Dudek,H., Miranti,C.K. und Greenberg,M.E. (1996). Calcium influx via the NMDA receptor induces immediate early gene transcription by a MAP kinase/ERK-dependent mechanism. *J. Neurosci.* 16, 5425-5436.
- Xiao,B., Tu,J.C., Petralia,R.S., Yuan,J.P., Doan,A., Breder,C.D., Ruggiero,A., Lanahan,A.A., Wenthold,R.J. und Worley,P.F. (1998). Homer regulates the association of group 1 metabotropic glutamate receptors with multivalent complexes of homer-related, synaptic proteins. *Neuron* 21, 707-716.
- Yamagata,K., Andreasson,K.I., Kaufmann,W.E., Barnes,C.A. und Worley,P.F. (1993). Expression of a mitogen-inducible cyclooxygenase in brain neurons: regulation by synaptic activity and glucocorticoids. *Neuron* 11, 371-386.
- Yin,Y., Edelman,G.M. und Vanderklish,P.W. (2002). The brain-derived neurotrophic factor enhances synthesis of Arc in synaptoneurosome. *Proc. Natl. Acad. Sci. U. S. A* 99, 2368-2373.
- Zalfa,F., Giorgi,M., Primerano,B., Moro,A., Di Penta,A., Reis,S., Oostra,B. und Bagni,C. (2003). The fragile X syndrome protein FMRP associates with BC1 RNA and regulates the translation of specific mRNAs at synapses. *Cell* 112, 317-327.
- Zamanillo,D., Sprengel,R., Hvalby,O., Jensen,V., Burnashev,N., Rozov,A., Kaiser,K.M., Koster,H.J., Borchardt,T., Worley,P., Lubke,J., Frotscher,M., Kelly,P.H., Sommer,B., Andersen,P.,

Seeburg,P.H. und Sakmann,B. (1999). Importance of AMPA receptors for hippocampal synaptic plasticity but not for spatial learning. *Science* 284, 1805-1811.

Zan,Y., Haag,J.D., Chen,K.S., Shepel,L.A., Wigington,D., Wang,Y.R., Hu,R., Lopez-Guajardo,C.C., Brose,H.L., Porter,K.I., Leonard,R.A., Hitt,A.A., Schommer,S.L., Elegbede,A.F. und Gould,M.N. (2003). Production of knockout rats using ENU mutagenesis and a yeast-based screening assay. *Nat. Biotechnol.* 21, 645-651.

Abkürzungsverzeichnis

'	Minuten
+/+	Wildtyp
-/-	Knockout
3'UTR	3'-untranslatierte Region
5'UTR	5'-untranslatierte Region
AMPA	α -Amino-3-Hydroxy-5-Methyl-4-Isoxazol-Propionat
Arc	<i>activity-regulated, cytoskeleton-associated protein</i> (=Arg3.1)
Arg3.1	<i>activity-regulated gene, 3.1 base pairs in length</i> (=Arc)
BDNF	<i>brain derived neurotrophic factor</i>
Ca ⁺⁺	Kalziumion
CA1-3 Region	Cornu ammonis Region 1-3
CaMKII	Ca ²⁺ /Calmodulin abhängige Proteinkinase
Cis	<i>cytokine-inducible SH2-containing protein</i>
Co-IP	Co-Immunopräzipitation
CS	<i>conditioned stimulus</i> (zu konditionierender Stimulus)
CTA	<i>conditioned taste aversion</i>
DNA	<i>Deoxyribonucleic Acid</i> (Desoxyribonukleinsäure)
dNTP	Desoxyribonukleotidtriphosphat
E-LTP	frühe (<i>early</i>) Phase der LTP
FEM	Forschungseinrichtung für experimentelle Medizin der Charité Berlin
GAPDH	Glycerinaldehyd-3-Phosphat Dehydrogenase
GSF	Gesellschaft für Strahlenforschung, heute: Forschungszentrum für Umwelt und Gesundheit
h	Stunden
i.p.	intraperitoneal
JAB	<i>Janus kinase-binding protein</i> (=SOCS)
JAK	Janus Kinase
Kb	Kilobasen
ko	<i>knockout</i>
loxP	<i>Locus of Crossover</i> in P1 (Erkennungssequenz der Cre-Rekombinase)
LTP	Langzeit-Potenzierung
L-LTP	späte (<i>late</i>) Phase der LTP
m	Meter
MAP2	<i>microtubule-associated protein2</i>
mM	milliMolar
mRNA	<i>messenger RNA</i>

Na ⁺	Natriumion
NIH	<i>National Institute of Health</i>
NMDA	N-Methyl-D-Aspartat
NMDAR	NMDA-Rezeptor-Kanal
NR1	NMDAR Untereinheit 1
NRC	<i>NMDA-receptor-complex</i>
MPI	Max-Planck-Institut
ORF	<i>open reading frame</i>
PBS	Phosphat gepufferte Salzlösung
PIAS	<i>protein inhibitor of activated STAT</i>
PKA	Proteinkinase A
PKC	Proteinkinase C
(rt)PCR	<i>(reverse transcriptase) polymerase chain reaction</i>
PSD	Postsynaptische Dichte
RNA	<i>Ribonucleic acid</i> (Ribonukleinsäure)
RT	Raumtemperatur
Rpm	Umdrehungen pro Minute (<i>rounds per minute</i>)
sec	Sekunden
SD	<i>standard deviation</i> , Standardabweichung
SDS	<i>Sodium</i> (Natrium) Dodecylsulfat
SOCS	<i>suppressor of cytokine signalling</i>
SSI	<i>STAT-induced STAT inhibitor</i> (=SOCS)
STAT	<i>signaling transducer and activator of transcription</i>
t-PA	<i>tissue-plasminogen activator</i>
Tris	Tris-(hydroxymethyl)-aminomethan
U	<i>Unit</i> , Enzymeinheit
UKE	Universitätskrankenhaus Eppendorf, Hamburg
UR	unkonditionierte Reaktion
US	unkonditionierter Stimulus
Vol	Volumen
vs.	versus
WT	Wildtyp
ZMNH	Zentrum für Molekulare Neurobiologie Hamburg
ZNS	zentrales Nervensystem

Danksagung

An erster Stelle möchte ich mich bei Herrn Prof. Dr. Dietmar Kuhl bedanken, der es mir ermöglicht hat, mich mit diesem faszinierenden Thema zu beschäftigen. Bei der Betreuung der Arbeit hat er mir seine ständige Diskussionsbereitschaft angeboten und mich mit wissenschaftlichem und freundschaftlichem Rat unterstützt. Darüber hinaus hat er es mir ermöglicht, zahlreiche Mitglieder der nationalen und internationalen Forschungsgemeinschaft kennenzulernen und mich mit diesen auszutauschen.

Herrn Prof. Dr. Randolf Menzel danke ich für sein anhaltendes Interesse an meiner Arbeit, die langjährige Begleitung durch mein Studium sowie die Übernahme des Gutachtens dieser Dissertation.

Allen Mitarbeitern der Arbeitsgruppe Kuhl in Hamburg und Berlin möchte ich ganz herzlich für ihre große Hilfsbereitschaft und die angenehme Arbeitsatmosphäre danken. Dieser Dank geht an Anastasia, Anita, André, Arne, Claudia M., Claudia S., Derk, Fang, Franziska, Kathrin, Micky, Ria, Stefan, Tina, Ulla, Xiasong und ganz speziell Nina, die von einer Kollegin zu einer guten Freundin geworden ist.

Ich danke den Mitarbeitern des Instituts für Anatomie der Universität Zürich unter der Leitung von Prof. Dr. Hans-Peter Lipp und Dr. David Wolfer für die Einarbeitung in die Verhaltensexperimente und ihre Gastfreundlichkeit.

Mange tak til jhk for den elektroniske støtte.

Mein ganz besonderer Dank gilt meinen Eltern, Carine, meiner Schwester Kerstin und Michael für ihr Verständnis und ihre unendliche Unterstützung.

Lebenslauf

Niels Plath

Geboren am 12. Juni 1973 in Oberhausen

- 1979 – 1983 Falkenstein Grundschule in Oberhausen
- 1983 – 1986 Bertha-von-Suttner Gymnasium in Oberhausen
- 1986 – 1992 Louise-Schroeder-Gymnasium in München
- 1993 – 1994 Zivildienst an der DJH in München
- 1994 – 2000 Biologie Studium an der FU Berlin
- 1996 Vordiplom
- 1997 – 1998 *Visiting and Exchange Student* an der *University of Sussex* in Brighton, England
- 2000 Diplom in Biologie an der FU Berlin
- Juni 2000 Beginn der Dissertation am Zentrum für Molekulare Neurobiologie Hamburg (ZMNH) unter der Anleitung von Prof. Dr. D. Kuhl
- August 2002 Umzug an die FU Berlin und Weiterführung der Dissertation unter der Anleitung von Prof. Dr. D. Kuhl

Publikationen

- Hermey G., Plath N., Hubner C.A., Kuhl D., Schaller H.C., and Hermans-Borgmeyer I. (2004). The three sorCS genes are differentially expressed and regulated by synaptic activity. *J. Neurochem.* 88, 1470-1476.
- Malun D., Giurfa M., Galizia C.G., Plath N., Brandt R., Gerber B., and Eisermann B. (2002). Hydroxyurea-induced partial mushroom body ablation does not affect acquisition and retention of olfactory differential conditioning in honeybees. *J. Neurobiol.* 53, 343-360.
- Malun D., Plath N., Giurfa M., Moseleit A.D., and Müller U. (2002). Hydroxyurea-induced partial mushroom body ablation in the honeybee *Apis mellifera*: volumetric analysis and quantitative protein determination. *J. Neurobiol.* 50, 31-44.

Tagungsbeiträge

- Plath N., Malun D., and Müller U. (2000). Chemical mushroom body ablation in the honeybee: Effects on various olfactory learning tasks and on protein amounts. *Proceedings of the Forum of European Neuroscience (FENS) 2000.* Eur. J. Neurosci. 12, Suppl. 11, p.92.
- Plath N., Ohana O., Dammermann B., Welzl H., Waltereit R., Bick-Sander A., Wolfer, D.P., Therstappen E., Husi H., Errington M.L., Blanquet V., Wurst W., Bliss T.V.P., Grant S.G., Lipp H.P., Bösl M., and Kuhl D. Arg3.1 is associated with the NMDA receptor complex and is required for memory formation. *Soc. Neurosci. 33rd Annual Meeting, Abstract Viewer and Itinerary Planner.* Washington, DC: Society for Neuroscience, 2003. Online.