

## 8. Literaturverzeichnis

Ammann RW. Idiopathic “juvenile” chronic pancreatitis. Dtsch Med Wochenschr 1976;101:1789-1794.

Ammann RW, Bühler H, Münch R, et al. Differences of the natural history of idiopathic (nonalcoholic) and alcoholic chronic pancreatitis. A comparative long-term study of 287 patients. Pancreas 1987;2:368-377.

Ammann RW. Chronic pancreatitis in the elderly. Gastroenterol Clin North Am 1990;19:905-914.

Audrézet MP, Chen JM, Le Maréchal C, et al. Determination of the relative contribution of three genes—the cystic fibrosis transmembrane conductance regulator gene, the cationic trypsinogen gene, and the pancreatic secretory trypsin inhibitor gene—to the etiology of idiopathic chronic pancreatitis. Eur J Hum Genet 2002;10:100-106.

Bernardino AL, Guarita DR, Mott CB, et al. *CFTR*, *PRSS1* and *SPINK1* mutations in the development of pancreatitis in Brazilian patients. JOP 2003;4:169-177.

Bhatia E, Choudhuri G, Sikora SS, et al. Tropical calcific pancreatitis: strong association with *SPINK1* trypsin inhibitor mutations. Gastroenterology 2002;123:1020-1025.

Bombieri C, Benetazzo MG, Saccomani A, et al. Complete mutational screening of the *CFTR* gene in 120 patients with pulmonary disease. Hum Genet 1998;103:718-722.

Braxel C, Versieck J, Lemey G, et al. Alpha<sub>1</sub>-antitrypsin in pancreatitis. Digestion 1982;23:93-96.

Casanova ML, Bravo A, Ramirez A, et al. Exocrine pancreatic disorders in transgenic mice expressing human keratin 8. J Clin Invest 1999;103:1587-1595.

Castellani C, Bonizzato A, Rolfini R, et al. Increased prevalence of mutations of the cystic fibrosis gene in idiopathic chronic and recurrent pancreatitis (letter). Am J Gastroenterol 1999;94:1993-1995.

Castellani C, Gomez Lira M, Frulloni L, et al. Analysis of the entire coding region of the cystic fibrosis transmembrane regulator gene in idiopathic pancreatitis. Hum Mutat 2001;18:166.

Cavestro GM, Frulloni L, Nouvenne A, et al. Association of keratin 8 gene mutation with chronic pancreatitis. Dig Liver Dis 2003;35:416-420.

Chandak GR, Idris MM, Reddy DN, et al. Mutations in the pancreatic secretory trypsin inhibitor gene (PST1/SPINK1) rather than the cationic trypsinogen gene (PRSS1) are significantly associated with tropical calcific pancreatitis. J Med Genet 2002;39:347-351.

Chiari H. Über Selbstverdauung des menschlichen Pankreas. Zeitschrift für Heilkunde 1896;17:69-96.

Chillon M, Casals T, Mercier B, et al. Mutations in the cystic fibrosis gene in patients with congenital absence of the vas deferens. N Engl J Med 1995;332:1475-1480.

Chu CS, Trapnell BC, Murtagh JJ Jr, et al. Variable deletion of exon 9 coding sequences in cystic fibrosis transmembrane conductance regulator gene mRNA transcripts in normal bronchial epithelium. EMBO J 1991;10:1355-1363.

Chu CS, Trapnell BC, Curristin S, et al. Genetic basis of variable exon 9 skipping in cystic fibrosis transmembrane conductance regulator mRNA. Nat Genet 1993;3:151-156.

Claustres M, Guittard C, Bozon D, et al. Spectrum of *CFTR* mutations in cystic fibrosis and in congenital absence of the vas deferens in France. Hum Mutat 2000;16:143-156.

Cohn JA, Friedman KJ, Noone PG, et al. Relation between mutations of the cystic fibrosis gene and idiopathic pancreatitis. *New Engl J Med* 1998;339:653-658.

Cohn JA, Neoptolemos JP, Feng J, et al. Increased risk of idiopathic chronic pancreatitis in cystic fibrosis carriers. *Hum Mutat* 2005;26:303-307.

Colomb E, Guy O, Deprez P, et al. The two human trypsinogens: Catalytic properties of the corresponding trypsins. *Biochem Biophys Acta* 1978;525:186-193.

Colomb E, Figarella C. Comparative studies on the mechanism of activation of the two human trypsinogens. *Biochem Biophys Acta* 1979;571:343-351.

Comfort MW, Steinberg AG. Pedigree of a family with hereditary chronic relapsing pancreatitis. *Gastroenterology* 1952;21:54-63.

Cuppens H, Marynen P, De Boeck C, et al. Detection of 98.5% of the mutations in 200 Belgian cystic fibrosis alleles by reverse dot-blot and sequencing of the complete coding region and exon/intron junctions of the *CFTR* gene. *Genomics* 1993;18:693-697.

Cuppens H, Lin W, Jaspers M, et al. Polyvariant mutant cystic fibrosis transmembrane conductance regulator genes. The polymorphic (Tg)m locus explains the partial penetrance of the T5 polymorphism as a disease mutation. *J Clin Invest* 1998;15:487-496.

DiMagno EP, Layer P, Clain JE. Chronic pancreatitis. In *The Pancreas: Biology, Pathobiology, and Disease* (eds Go VLW et al.) 665-706 (Raven, New York, 1993).

Dörk T, Dworniczak B, Aulehla-Scholz C, et al. Distinct spectrum of *CFTR* gene mutations in congenital absence of vas deferens. *Hum Genet* 1997;100:365-377.

Drenth JP, te Morsche R, Jansen JB. Mutations in serine protease inhibitor Kazal type 1 are strongly associated with chronic pancreatitis. Gut 2002;50:687-692.

Dumur V, Gervais R, Rigot JM, et al. Abnormal distribution of CF deltaF508 allele in azoospermic men with congenital aplasia of epididymis and vas deferens. Lancet 1990;336:512.

Durbec JP, Sarles H. Multicenter survey on the etiology of pancreatic diseases. Relationship between the relative risk of developing chronic pancreatitis and alcohol, protein, and lipid consumption. Digestion 1978;18:337-350.

Durno C, Corey M, Zielinski J, et al. Genotype and phenotype correlations in patients with cystic fibrosis and pancreatitis. Gastroenterology 2002;123:1857-1864.

Férec C, Raguénès O, Salomon R, et al. Mutations in the cationic trypsinogen gene and evidence for genetic heterogeneity in hereditary pancreatitis. J Med Genet 1999;36:228-232.

Freedman SD, Blanco P, Shea JC, et al. Mechanisms to explain pancreatic dysfunction in cystic fibrosis. Gastroenterol Clin North Am 2000;84:657-664.

Gomez Lira M, Benetazzo MG, Marzari MG, et al. High frequency of cystic fibrosis transmembrane regulator mutation L997F in patients with recurrent idiopathic pancreatitis and in newborns with hypertrypsinemia. Am J Hum Genet 2000;66:2013-2014.

Gomez Lira M, Bonamini D, Castellani C, et al. Mutations in the *SPINK1* gene in idiopathic pancreatitis italiano patients. Eur J Hum Genet 2003;11:543-546.

Gordon D, Heath SC, Liu X, et al. A transmission/disequilibrium test that allows for genotyping errors in the analysis of single-nucleotide polymorphism data. Am J Hum Genet 2001; 69:371-380.

Gordon D, Haynes C, Johnnidis C, et al. A transmission disequilibrium test for general pedigrees that is robust to the presence of random genotyping errors and any number of untyped parents. Eur J Hum Genet 2004;12:752-761.

Gorry MC, Gabbaizedeh D, Furey W, et al. Mutations in the cationic trypsinogen gene are associated with recurrent acute and chronic pancreatitis. Gastroenterology 1997;113:1063-1068.

Groman JD, Meyer ME, Wilmott RW, et al. Variant cystic fibrosis phenotypes in the absence of *CFTR* mutations. N Engl J Med 2002;347:401-407.

Guy O, Lombardo D, Bartelt DC, et al. Two human trypsinogens: purification, molecular properties and N-terminal sequences. Biochemistry 1978;17:1669-1675.

Haber PS, Wilson JS, McGarity BH, et al.  $\alpha_1$  antitrypsin phenotypes and alcoholic pancreatitis. Gut 1991;32:945-948.

Hassan Z, Mohan V, Ali L, et al. SPINK1 is a susceptibility gene for fibrocalculus pancreatic diabetes in subjects from the Indian subcontinent. Am J Hum Genet 2002;71:964-968.

Johnson MP, Haupt LM, Griffiths LR. Locked nucleic acid (LNA) single nucleotide polymorphism (SNP) genotype analysis and validation using real-time PCR. Nucleic Acids Res 2004;32:e55.

Kazal LA, Spicer DS, Brahinsky RA. Isolation of a crystalline trypsin inhibitor-anticoagulant protein from the pancreas. J Am Chem Soc 1948;70:304-340.

Kerem BS, Rommens JM, Buchanan JA, et al. Identification of the cystic fibrosis gene: genetic analysis. Science 1989;245:1073-1080.

Kiesewetter S, Macek M Jr, Davis C, et al. A mutation in CFTR produces different phenotypes depending on chromosomal background. *Nat Genet* 1993;5:274-278.

Király O, Guan L, Szepessy E, et al. Expression of human cationic trypsinogen with an authentic N terminus using intein-mediated splicing in aminopeptidase P deficient Escherichia coli. *Protein Expr Purif* 2006;48:104-111.

Király O, Boulling A, Witt H, et al. Signal peptide variants that impair secretion of pancreatic secretory trypsin inhibitor (SPINK1) cause autosomal dominant hereditary pancreatitis. *Hum Mutat*. 2007;28:469-476.

Koch C, Cuppens H, Rainisio M, et al. European Epidemiologic Registry of Cystic Fibrosis (ERCF): comparison of major disease manifestations between patients with different classes of mutations. *Pediatr Pulmonol* 2001;31:1-12.

Koshkin AA, Singh SK, Nielsen P et al. LNA (Locked Nucleic Acids): synthesis of the Adenine, Cytosine, Guanine, 5-Methylcytosine, Thymine and Uracil bicyclonucleoside monomers, oligomerisation, and unprecedented nucleic acid recognition. *Tetrahedron* 1998;54:3607-3630.

Ku, Nam-On et al. The cytoskeleton of digestive epithelia in health and disease. *Am J Physiol Gastrointest Liver Physiol* 1999;277:G1108-G1137.

Lankisch PG, Koop H, Winckler K, et al.  $\alpha_1$ -antitrypsin in pancreatic diseases. *Digestion* 1978;18:138-140.

Larsson C. Natural history and life expectancy in severe alpha-1-antitrypsin deficiency, Pi Z. *Acta Med Scand* 1978;204:345-351.

Laskowski M, Wu FC. Temporary inhibition of trypsin. *J Biol Chem* 1953;204:797-805.

Layer P, Yamamoto H, Kalthoff L, et al. The different courses of early- and late-onset idiopathic and alcoholic chronic pancreatitis. *Gastroenterology* 1994;107:1481-1487.

Le Bodic L, Bignon JD, Raguénès O, et al. The hereditary pancreatitis gene maps to long arm of chromosome 7. *Hum Mol Genet* 1996;5:549-554.

Lester LA, Kraut J, Lloyd-Still J, et al. Delta F508 genotype does not predict disease severity in an ethnically diverse cystic fibrosis population. *Pediatrics* 1994;93:114-118.

Marchbank T, Freeman TC, Playford RJ. Human pancreatic secretory trypsin inhibitor. *Digestion* 1998;59:167-174.

Mihas AA, Hirschowitz BI. Alpha<sub>1</sub>-antitrypsin and chronic pancreatitis. *Lancet* 1976;ii:1032-1033.

Mote VL, Anderson RL. An investigation of the effect of misclassification on the properties of Chi-2-tests in the analysis of categorical data. *Biometrika* 1965;52:95-109.

Niel F, Legendre M, Bienvenu T, et al. A new large *CFTR* rearrangement illustrates the importance of searching for complex alleles. *Hum Mutat* 2006;27:716-717.

Nemoda Z, Sahin-Tóth M. Chymotrypsin C (caldecrin) stimulates autoactivation of human cationic trypsinogen. *J Biol Chem* 2006;281:11879-11886.

Noone PG, Zhou Z, Silverman LM, et al. Cystic fibrosis gene mutations and pancreatitis risk: relation to epithelial ion transport and trypsin inhibitor gene mutations. *Gastroenterology* 2001;121:1310-1319.

Novis BH, Young GO, Bank S, Marks IN. Chronic pancreatitis and alpha-1-antitrypsin. Lancet 1975;ii:748-749.

Obika S, Nanbu D, Hari Y, et al. Stability and structural features of the duplexes containing nucleoside analogues with a fixed N-type conformation, 2'-O,4'-C-methyleneribonucleosides. Tetrahedron Lett 1998;39:5401-5404.

Orita M, Iwahana H, Kanazawa H, et al. Detection of polymorphisms of human DNA by gel electrophoresis as single-strand conformation polymorphism. Proc Natl Acad Sci 1989;86:2766-2770.

Pagani F, Buratti E, Stuani C, et al. Splicing factors induce cystic fibrosis transmembrane regulator exon 9 skipping through nonevolutionary conserved intronic elements. J Biol Chem 2000;275:21041-21047.

Pandya A, Blanton SH, Landa B, et al. Linkage studies in a large kindred with hereditary pancreatitis confirms mapping of the gene to a 16-cM region on 7q. Genomics 1996;38:227-230.

Perlmutter DH. Alpha-1-antitrypsin deficiency. Semin Liver Dis 1998;18:217-225.

Perrault J. Hereditary pancreatitis. Gastroenterol Clin North Am 1994;23:743-752.

Petersen M, Bondensgaard K, Wengel J et al. Locked nucleic acid (LNA) recognition of RNA: NMR solution structures of LNA:RNA hybrids. J Am Chem Soc 2002;124:5974-5982.

Pfützer RH, Barmada NM, Brunskill APJ, et al. SPINK1/PSTI polymorphisms act as disease modifiers in familial and idiopathic chronic pancreatitis. Gastroenterology 2000;119:615-623.

Ren CL. Mutations of the Cystic fibrosis gene and pancreatitis. *N Engl J Med* 1999;340:238-239.

Riesner D, Steger G, Wiese U, et al. Temperature gradient gel electrophoresis of nucleic acids: analysis of conformational transitions, sequence variations and protein nucleic acid interactions. *Electrophoresis* 1989;10:377-389.

Riordan JR, Rommens JM, Kerem B, et al. Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. *Science* 1989;245:1066-1073.

Riordan JR. The cystic fibrosis transmembrane conductance regulator. *Annu Rev Physiol* 1993;55:609-630.

Rohlf EM, Zhou Z, Sugarman EA, et al. The I148T CFTR allele occurs on multiple haplotypes: a complex allele is associated with cystic fibrosis. *Genet Med* 2002;4:319-323.

Rosendahl J, Bödeker H, Mössner J, et al. Hereditary chronic pancreatitis. *Orphanet J Rare Dis* 2007;2:1.

Rosenstein BJ, Cutting GR. The diagnosis of cystic fibrosis: a consensus statement. Cystic Fibrosis Foundation Consensus Panel. *J Pediatr* 1998;132:589-595.

Sanger F, Nicklen S, Coulson AR. DNA sequencing with chain terminating inhibitors. *Proc Natl Acad Sci* 1977;74:5463-5467.

Sahin-Tóth M, Gráf L, Tóth M. Trypsinogen stabilization by mutation arg117-to-his: a unifying pathomechanism for hereditary pancreatitis? *Biochem Biophys Res Commun* 1999;264:505-508.

Sahin-Tóth M, Tóth M. Gain-of-function mutations associated with hereditary pancreatitis enhance autoactivation of human cationic trypsinogen. *Biochem Biophys Res Commun* 2000;278:286-289.

Scheele GA, Bartelt D, Bieger W. Characterization of human exocrine pancreatic proteins by two-dimensional isoelectric focusing/SDS gel electrophoresis. *Gastroenterology* 1981;80:461-473.

Schneider A, Suman A, Rossi L, et al. SPINK1/PST1 mutations are associated with tropical pancreatitis and type II diabetes mellitus in Bangladesh. *Gastroenterology* 2002;123:1026-1030.

Secknus R, Mössner J. Incidenz- und Prävalenzveränderungen der akuten und chronischen Pankreatitis in Deutschland. *Chirurg* 2000;71:249-252.

Sharer N, Schwartz M, Malone G, et al. Mutations of the cystic fibrosis gene in patients with chronic pancreatitis. *New Engl J Med* 1998;339:645-652.

Sheppard DN, Welsh MJ. Structure and function of the CFTR chloride channel. *Physiol Rev* 1999;79(Suppl):S23-45.

Shibata T, Ogawa M, Matsuda K, et al. Purification and characterisation of pancreatic secretory trypsin inhibitor in human gastric mucosa. *Clin Chim Acta* 1986;159:27-36.

Shibata T, Ogawa M, Takata N. Distribution of pancreatic secretory trypsin inhibitor in various human tissues and its inactivation in the gastric mucosa. *Res Commun Pathol Pharmacol* 1987;55:243-248.

Shwachman H, Lebenthal E, Khaw W. Recurrent acute pancreatitis in patients with cystic fibrosis with normal pancreatic enzymes. *Pediatrics* 1975;55:86-94.

Singer MV, Gyr K, Sarles H. Report of the Second International Symposium on the Classification of Pancreatitis in Marseille, France, March 28-30, 1984. *Gastroenterology* 1985;89:683-685.

Singh SK, Nielsen P, Koshkin AA et al. LNA (Locked Nucleic Acids): synthesis and high-affinity nucleic acid recognition. *Chem Commun* 1998;4:455-456.

Sobczynska-Tomaszewska A, Bak D, Oralewska B, et al. Analysis of CFTR, SPINK1, PRSS1 and AAT mutations in children with acute or chronic pancreatitis. *J Pediatr Gastroenterol Nutr.* 2006;43:299-306.

Steer ML, Meldolesi J. The cell biology of experimental pancreatitis. *N Engl J Med* 1987;316:144-150.

Stern RC. The diagnosis of cystic fibrosis. *N Engl J Med* 1997;336:487-491.

Strong TV, Wilkinson DJ, Mansoura MK, et al. Expression of an abundant alternatively spliced form of the cystic fibrosis transmembrane conductance regulator (*CFTR*) gene is not associated with a cAMP-activated chloride conductance. *Hum Mol Genet* 1993;2:225-230.

Teich N, Mössner J, Keim V. Mutations of the cationic trypsinogen in hereditary pancreatitis. *Hum Mutat* 1998;12:39-43.

Teich N, Ockenga J, Hoffmeister A, et al. Chronic pancreatitis associated with an activation peptide mutation that facilitates trypsin activation. *Gastroenterology* 2000;119:461-465.

Teich N, Walther K, Bödeker H, et al. Relevance of variants in serum antiproteinases for the course of chronic pancreatitis. *Scand J Gastroenterol* 2002;37:360-365.

The Cystic Fibrosis Genotype-Phenotype Consortium. Correlation between genotype and phenotype in patients with cystic fibrosis. *N Engl J Med* 1993;329:1308-1313.

The Cystic Fibrosis Genetic Analysis Consortium. Population variation of common cystic fibrosis mutations. *Hum Mutat* 1994;4:167-177.

Treiber M, Schulz HU, Landt O, et al. Keratin 8 sequence variants in patients with pancreatitis and pancreatic cancer. *J Mol Med* 2006;84(12):1015-1022.

Ugozzoli LA, Latorra D, Puckett R, et al. Real-time genotyping with oligonucleotide probes containing locked nucleic acids. *Anal Biochem* 2004;324:143-152.

Várallyay É, Pál G, Patthy A, et al. Two mutations in the rat trypsin confer resistance against autolysis. *Biochem Biophys Res Commun* 1998;243:56-60.

Wei L, Vankeerberghen A, Jaspers M, et al. Suppressive interactions between mutations located in the two nucleotide binding domains of CFTR. *FEBS Lett* 2000;473:149-153.

Weiss FU, Simon P, Bogdanova N, et al. Complete cystic fibrosis transmembrane conductance regulator gene sequencing in patients with idiopathic chronic pancreatitis and controls. *Gut* 2005;54:1456-1460.

Whitcomb DC, Preston RA, Aston CE, et al. A gene for hereditary pancreatitis maps to chromosome 7q35. *Gastroenterology* 1996;110:1975-1980.

Whitcomb DC, Gorry MC, Preston RA, et al. Hereditary pancreatitis is caused by a mutation in the cationic trypsinogen gene. *Nat Genet* 1996;14:141-145.

Witt H, Luck W, Becker M. A signal peptide cleavage site mutation in the cationic trypsinogen gene is strongly associated with chronic pancreatitis. *Gastroenterology* 1999;117:7-10.

Witt H, Luck W, Hennies HC, et al. Mutations in the gene encoding the serine protease inhibitor, Kazal type 1 are associated with chronic pancreatitis. *Nat Genet* 2000;25:213-216.

Witt H, Luck W, Becker M, et al. Mutation in the SPINK1 trypsin inhibitor gene, alcohol use, and chronic pancreatitis. *JAMA* 2001;285:2716-7.

Witt H, Kage A, Luck W, et al. Alpha-1-antitrypsin genotypes in patients with chronic pancreatitis. *Scand J Gastroenterol* 2002;37:356-359.

Witt H. Chronic pancreatitis and cystic fibrosis. *Gut* 2003;52 Suppl 2:ii31-41.

Witt H, Sahin-Tóth M, Landt O, et al. A degradation-sensitive anionic trypsinogen (PRSS2) variant protects against chronic pancreatitis. *Nat Genet* 2006;38:668-673.

You Y, Moreira BG, Behlke M, et al. Design of LNA probes that improve mismatch discrimination. *Nucleic Acids Res* 2006;34(8):e60.

Zielenski J, Roznahel R, Bozon D, et al. Genomic DNA sequence of the cystic fibrosis transmembrane conductance regulator (CFTR) gene. *Genomics* 1991;10:214-228.

Zielenski J, Patrizio P, Corey M, et al. *CFTR* gene variant for patients with congenital absence of vas deferens. *Am J Hum Genet* 1995;57:958-960.