

#### 4. Literatur

- Abramowicz MJ, Duprez L, Parma J, Vassart G, Heinrichs C (1997): Familial congenital hypothyroidism due to inactivating mutation of the thyrotropin receptor causing profound hypoplasia of the thyroid gland. *J Clin Invest*; **99**:3018-3024.
- Abramowicz MJ, Tarkovnik HM, Varela V, Cochaux P, Krawiec L, Pisarev MA, Propato FVE, Juvenal G, Chester HA, Vassart G (1992): Identification of a mutation in the coding sequence of the human thyroid peroxidase gene causing congenital goiter. *J Clin Invest*; **90**:1200-1204.
- Acharya S, Karnik SS (1996). Modulation of GDP release from transducin by the conserved Glu134-Arg135 sequence in rhodopsin. *J Biol Chem*; **271**:25406-11.
- Alberti L, Proverbio MC, Costagliola S, Romoli R, Boldrighini B, Vigone MC, Weber G, Chiumello G, Beck-Peccoz P, Persani L (2002) Germline mutations of TSH receptor gene as cause of nonautoimmune subclinical hypothyroidism. *J Clin Endocrinol Metab*; **87**:2549-55.
- Ambrugger P, Stoeva I, Biebermann H, Torresani T, Leitner C, Grüters A. (2001) Novel mutations of the thyroid peroxidase gene in patients with permanent congenital hypothyroidism. *Eur J Endocrinol*; **145**:19-24.
- Bai M, Quinn S, Trivedi S, Kifor O, Pearce SH, Pollak MR, Krapcho K, Hebert SC, Brown EM (1996). Expression and characterization of inactivating and activating mutations in the human Ca<sup>2+</sup>-sensing receptor. *J Biol Chem*; **271**:19537-45.
- Benkirane M, Jin DY, Chun RF, Koup RA, Jeang KT (1997). Mechanism of transdominant inhibition of CCR5-mediated HIV-1 infection by ccr5delta32. *J Biol Chem*; **272**:30603-6.
- Biebermann H, Schöneberg T, Hess C, Germak J, Gudermann T, Grüters A. (2001) The first activating TSH receptor mutation in transmembrane domain 1 identified in a family with nonautoimmune hyperthyroidism. *J Clin Endocrinol Metab*; **86**:4429-33.
- Biebermann H, Schöneberg T, Krude H, Gudermann T, Grüters A. (2000): Constitutively activating TSH-receptor mutations as a molecular cause of non-autoimmune hyperthyroidism in childhood. *Langenbecks Arch Surg*; **385**:390-2.
- Biebermann H, Schöneberg T, Krude H, Schultz G, Gudermann T, Grüters A (1997). Mutations of the human thyrotropin receptor gene causing thyroid hypoplasia and persistent congenital hypothyroidism. *J Clin Endocrinol Metab*; **82**:3471-80.
- Bikker H, Vulsma T, Baas F, de Vijlder JJM (1995): Identification of five novel inactivating mutations in the human thyroid peroxidase gene by denaturing gradient gel electrophoresis. *Hum Mutations*; **6**:9-16.
- Bouvier M (2001). Oligomerization of G-protein-coupled transmitter receptors. *Nat Rev Neurosci* , **2**:274-86.
- Brumm H, Pfeufer A, Biebermann H, Schnabel D, Deiss D, Grüters A. (2002): Congenital central hypothyroidism due to homozygous thyrotropin beta 313 Delta T mutation is caused by a Founder effect. *J Clin Endocrinol Metab*; **87**:4811-6.
- Challis BG, Pritchard LE, Creemers JW, Delplanque J, Keogh JM, Luan J, Wareham NJ, Yeo GS, Bhattacharyya S, Froguel P, White A, Farooqi IS, O'Rahilly S (2002). A missense mutation disrupting a dibasic prohormone processing site in pro-opiomelanocortin (POMC) increases susceptibility to early-onset obesity through a novel molecular mechanism. *Hum Mol Genet*; **11**:1997-2004.
- Cook JV, Eidne KA (1997). An intramolecular disulfide bond between conserved extracellular cysteines in the gonadotropin-releasing hormone receptor is essential for binding and activation. *Endocrinology*; **138**: 2800-6.
- Dacou-Voutetakis C, Feltquate DM, Drakopoulou M, Kourides IA, Dracopoli NC. (1990) Familial hypothyroidism caused by a nonsense mutation in the thyroid-stimulating hormone beta-subunit gene. *Am J Hum Genet*; **46**:988-93.
- De Lean A, Stadel JM, Lefkowitz RJ (1980). A ternary complex model explains the agonist-specific binding properties of the adenylate cyclase-coupled beta-adrenergic receptor. *J Biol Chem*; **255**:7108-17.
- De Roux N, Polak M, Couet J, Leger J, Czernichow P, Milgrom E, Misrahi M. (1996): A neomutation of the thyroid-stimulating hormone receptor in a severe neonatal hyperthyroidism. *J Clin Endocrinol Metab*; **81**: 2023-2026.
- Doeker BM, Pfaffle RW, Pohlenz J, Andler W. (1998) Congenital central hypothyroidism due to a homozygous mutation in the thyrotropin beta-subunit gene follows an autosomal recessive inheritance. *J Clin Endocrinol Metab*; **83**:1762-5.

- Duprez L, Parma J, Van Sande J, Allgeier A, Leclere J, Schwartz C, Delisle MJ, Decoulx M, Orgiazzi J, Dumont J, Vassart G (1994): Germline mutations in the thyrotropin receptor gene cause non-autoimmune autosomal dominant hyperthyroidism. *Nat Genet*; **7**:396-401.
- Farooqi IS, Keogh JM, Yeo GS, Lank EJ, Cheetham T, O'Rahilly S (2003). Clinical spectrum of obesity and mutations in the melanocortin 4 receptor gene. *N Engl J Med*. **20**; **348**:1085-95.
- Fernandez LM, Puett D. (1996b): Lys583 in the third extracellular loop of the lutropin/choriogonadotropin receptor is critical for signaling. *J Biol Chem*; **271**:925-930.
- Fotiadis D, Liang Y, Filipek S, Saperstein DA, Engel A, Palczewski K (2003): Atomic-force microscopy: Rhodopsin dimers in native disc membranes. *Nature*; **421**:127-8.
- Führer D, Wonerow P, Willgerodt H, Paschke (1997) RIdentification of a new thyrotropin receptor germline mutation (Leu629Phe) in a family with neonatal onset of autosomal dominant nonautoimmune hyperthyroidism. *J Clin Endocrinol Metab.*; **82**: 4234-8.
- Gagné N, Parma J, Deal C, Vassart G, Van Vliet G (1998): Apparent congenital athyreosis contrasting with normal plasma levels and associated with inactivating mutations in the thyrotropin receptor gene: are athyreosis and ectopic thyroid distinct entities? *J Clin Endocrinol Metab*; **83**:1771-1775.
- Gantz I, Miwa H, Konda Y, Shimoto Y, Tashiro T, Watson SJ, DelValle J, Yamada T (1993): Molecular cloning, expression, and gene localization of a fourth melanocortin receptor. *J Biol Chem*; **268**:15174-9.
- George SR, O'Dowd BF, Lee SP (2002): G-protein-coupled receptor oligomerization and its potential for drug discovery. *Nat Rev Drug Discov*; **1**:808-20.
- Grosse R, Schöneberg T, Schultz G, Gudermann T (1997). Inhibition of gonadotropin-releasing hormone receptor signaling by expression of a splice variant of the human receptor. *Mol Endocrinol*; **11**:1305-18.
- Grüters A (1992): Congenital hypothyroidism. *Ped Annals*; **21**:15-21.
- Grüters A, Schöneberg T, Biebermann H, Krude H, Krohn HP, Dralle H, Gudermann T (1998): Severe congenital hyperthyroidism caused by a germline neo-mutation in the extracellular portion of the thyrotropin receptor. *J Clin Endocrinol*; **83**:1431-1436.
- Hayashizaki Y, Hiraoka Y, Endo Y, Miyai K, Matsubara K. (1989) Thyroid-stimulating hormone (TSH) deficiency caused by a single base substitution in the CAGYC region of the beta-subunit. *EMBO J*; **8**:2291-6.
- Hinney A, Becker I, Heibult O, Nottebom K, Schmidt A, Ziegler A, Mayer H, Siegfried W, Blum WF, Remschmidt H, Hebebrand J (1998). Systematic mutation screening of the pro-opiomelanocortin gene: identification of several genetic variants including three different insertions, one nonsense and two missense point mutations in probands of different weight extremes. *J Clin Endocrinol Metab*; **83**:3737-41.
- Hinney A, Schmidt A, Nottebom K, Heibult O, Becker I, Ziegler A, Gerber G, Sina M, Gorg T, Mayer H, Siegfried W, Fichter M, Remschmidt H, Hebebrand J (1999). Several mutations in the melanocortin-4 receptor gene including a nonsense and a frameshift mutation associated with dominantly inherited obesity in humans. *J Clin Endocrinol Metab*; **84**:1483-6.
- Ho G, MacKenzie RG (1999). Functional characterization of mutations in melanocortin-4 receptor associated with human obesity. *J Biol Chem*; **274**:35816-22.
- Huszar D, Lynch CA, Fairchild-Huntress V, Dunmore JH, Fang Q, Berkemeier LR, Gu W, Kesterson RA, Boston BA, Cone RD, Smith FJ, Campfield LA, Burn P, Lee F (1997). Targeted disruption of the melanocortin-4 receptor results in obesity in mice. *Cell*; **88**:131-41.
- Kendall SK, Samuelson LC, Saunders TL, Wood RI, Camper SA (1995): Target disruption of the pituitary glycoprotein hormone  $\alpha$ -subunit produces hypogonadal and hypothyroid mice. *Genes & Development*; **9**:2007-2019.
- Kopp P, Van Sande J, Parma J, Duprez L, Gerber H, Joss E, Jameson JL, Dumont JE, Vassart G. (1995): Brief report: congenital hyperthyroidism caused by a mutation in the thyrotropin-receptor gene. *N Engl J Med*; **332**: 150-154.
- Krude H, Biebermann H, Gopel W, Grüters A (1996): The gene for the thyrotropin receptor (TSHR) as a candidate gene for congenital hypothyroidism with thyroid dysgenesis. *Exp Clin Endocrinol Diabetes*; **104 Suppl 4**:117-120.
- Krude H, Grüters A (2000). Implications of proopiomelanocortin (POMC) mutations in humans: the POMC deficiency syndrome. *Trends Endocrinol Metab.*; **11**:15-22.
- Laugwitz K-L, Allgeier A, Offermanns S, Spicher K, Van Sande J, Dumont JE, Schultz G (1996): The human thyrotropin receptor: A heptahelical receptor capable of stimulating members of all four G protein families. *Proc Natl Acad Sci USA*; **93**:116-120.

- Libert F, Lefort A, Gerard C, Parmentier M, Perret J, Ludgate M, Dumont JE, Vassart G (1989): Cloning, sequencing and expression of the human thyrotropin (TSH) receptor: evidence for binding of autoantibodies. *Biochem Biophys Res Commun*; **165**:1250-1255.
- Liu G, Duranteau L, Carel JC, Monroe J, Doyle DA, Shenker A (1999) Leydig-cell tumors caused by an activating mutation of the gene encoding the luteinizing hormone receptor. *N Engl J Med*; **341**:1731-6.
- Maggio R, Vogel Z, Wess J (1993). Coexpression studies with mutant muscarinic/adrenergic receptors provide evidence for intermolecular "cross-talk" between G-protein-linked receptors. *Proc Natl Acad Sci USA*.; **90**:3103-7.
- Matthews CH, Borgato S, Beck-Peccoz P, Adams M, Tone Y, Gambino G, Casagrande S, Tedeschini G, Benedetti A, Chatterjee VK.(1993) Primary amenorrhoea and infertility due to a mutation in the beta-subunit of follicle-stimulating hormone. *Nat Genet*; **5**:83-6.
- Medeiros-Neto G, Herodotou DT, Rajan S, Kommareddi S, de Lacerda L, Sandrini R, Boguszewski MC, Hollenberg AN, Radovick S, Wondisford FE. (1996) A circulating, biologically inactive thyrotropin caused by a mutation in the beta subunit gene. *J Clin Invest*; **97**:1250-6.
- Miraglia del Giudice E, Cirillo G, Santoro N, D'Urso L, Carbone MT, Di Toro R, Perrone L. (2001) Molecular screening of the proopiomelanocortin (POMC ) gene in Italian obese children: report of three new mutations. *Int J Obes Relat Metab Disord*; **25**:61-7.
- Nagayama Y, Kaufman KD, Seto P, Rapoport B. (1989): Molecular cloning, sequence and functional expression of the cDNA for the human thyrotropin receptor. *Biochem Biophys Res Commun*; **165**:1184-1190.
- Palczewski K, Kumasaka T, Hori T, Behnke CA, Motoshima H, Fox BA, Le Trong I, Teller DC, Okada T, Stenkamp RE, Yamamoto M, Miyano M (2000) Crystal structure of rhodopsin: A G protein-coupled receptor. *Science*; **289**:739-45.
- Parma J, Duprez L, Van Sande J, Cochaux P, Gervy C, Mockel J, Dumont J, Vassart G (1993): Somatic mutations in the thyrotropin receptor gene cause hyperfunctioning thyroid adenomas. *Nature*; **365**:649-651.
- Pritchard LE, Turnbull AV, White A. (2002) Pro-opiomelanocortin processing in the hypothalamus: impact on melanocortin signalling and obesity. *J Endocrinol*; **172**:411-21.
- Rodien P, Bremont C, Sanson ML, Parma J, Van Sande J, Costagliola S, Luton JP, Vassart G, Duprez L (1998) Familial gestational hyperthyroidism caused by a mutant thyrotropin receptor hypersensitive to human chorionic gonadotropin. *N Engl J Med*.; **339**: 1823-6.
- Russo D, Wong MG, Costante G, Chiefari E, Treseler PA, Arturi F, Filetti S, Clark OH (1999). A Val 677 activating mutation of the thyrotropin receptor in a Hurthle cell thyroid carcinoma associated with thyrotoxicosis. *Thyroid*; **9**:13-7.
- Schaffer JV, Bologna JL (2001). The melanocortin-1 receptor: red hair and beyond. *Arch Dermatol*.; **137**:1477-85.
- Schertler GF (1998): Structure of rhodopsin. *Eye*.; **12**:504-10.
- Schöneberg T, Schultz G, Gudermann T (1999). Structural basis of G protein-coupled receptor function. *Mol Cell Endocrinol* ; **151**:181-93.
- Schöneberg T, Yun J, Wenkert D, Wess J (1996). Functional rescue of mutant V2 vasopressin receptors causing nephrogenic diabetes insipidus by a co-expressed receptor polypeptide. *EMBO J*; **15**:1283-91.
- Schulz A, Bruns K, Henklein P, Krause G, Schubert M, Gudermann T, Wray V, Schultz G, Schöneberg T (2000a). Requirement of specific intrahelical interactions for stabilizing the inactive conformation of glycoprotein hormone receptors. *J Biol Chem*.; **275**:37860-9.
- Schulz A, Grosse R, Schultz G, Gudermann T, Schöneberg T (2000): Structural implication for receptor oligomerization from functional reconstitution studies of mutant V2 vasopressin receptors. *J Biol Chem*.; **275**:2381-9.
- Schulz A, Schöneberg T, Paschke R, Schultz G, Gudermann T (1999): Role of the third intracellular loop for the activation of gonadotropin receptors. *Mol Endocrinol*; **13**:181-90.
- Schwab KO, Gerlich M, Broecker M, Sohlmann P, Derwahl M, Lohse MJ (1997) Constitutively active germline mutation of the thyrotropin receptor gene as a cause of congenital hyperthyroidism. *J Pediatr*.; **131**: 899-904.
- Sealfon S C , Chi L , Ebersole B J , Rodic V , Zhang D , Ballesteros J A , Weinstein H (1995) Related contribution of specific helix 2 and 7 residues to conformational activation of the serotonin 5-HT<sub>2A</sub> receptor. *J Biol Chem*, **270**:16683-16688.
- Shenker A, Laue L, Kosugi S, Merendino JJ Jr, Minegishi T, Cutler GB Jr. (1993) A constitutively activating mutation of the luteinizing hormone receptor in familial male precocious puberty. *Nature*; **365**:652-4.

- Sina M, Hinney A, Ziegler A, Neupert T, Mayer H, Siegfried W, Blum WF, Remschmidt H, Hebebrand J (1999). Phenotypes in three pedigrees with autosomal dominant obesity caused by haploinsufficiency mutations in the melanocortin-4 receptor gene. *Am J Hum Genet*; **65**:
- Smits G, Olatunbosun O, Delbaere A, Pierson R, Vassart G, Costagliola S (2003): Ovarian hyperstimulation syndrome due to a mutation in the follicle-stimulating hormone receptor. *N Engl J Med*; **349**:760-6.
- Stein SA, Oates EL, Hall CR, Grumbles RM, Fernandez LM, Taylor NA, Puett D, Jin S (1994): Identification of a point mutation in the thyrotropin receptor of the *hyt/hyt* mouse. *Mol Endocrinol*. **8**:129-138.
- Sunthornthepvarakul T, Gottschalk ME, Hayashi Y, Refetoff S (1995): Resistance to TSH caused by mutations in the thyrotropin-receptor gene. *N Engl J Med*. **332**:155-160.
- Tarnow P, Schöneberg T, Krude H, Grüters A, Biebermann H (2003). Mutationally induced disulfide bond formation within the third extracellular loop causes melanocortin 4 receptor inactivation in patients with obesity. *J Biol Chem*; **278**:48666-48673.
- Tatsumi K, Miyai K, Notomi T, Kaibe K, Amino N, Mizuno Y, Kohno H (1992) Cretinism with combined hormone deficiency caused by a mutation in the PIT1 gene. *Nat Genet*; **1**:56-8.
- Vaisse C, Clement K, Durand E, Hercberg S, Guy-Grand B, Froguel P. (2000): Melanocortin-4 receptor mutations are a frequent and heterogeneous cause of morbid obesity. *J Clin Invest*.; **106**:253-62.
- Vaisse C, Clement K, Guy-Grand B, Froguel P (1998). A frameshift mutation in human MC4R is associated with a dominant form of obesity. *Nat Genet*; **20**:113-4.
- Valverde P, Healy E, Sikkink S, Haldane F, Thody AJ, Carothers A, Jackson IJ, Rees JL (1996) The Asp84Glu variant of the melanocortin 1 receptor (MC1R) is associated with melanoma. *Hum Mol Genet*, **5**:1663-1666.
- Van Sande J, Parma J, Tonacchera M, Swillens S, Dumont J, Vassart G. (1995): Somatic and germline mutations of the TSH receptor gene in thyroid diseases. *J Clin Endocrinol Metab*.; **80**:2577-85.
- Vassart G, Dumont JE (1992): The thyrotropin receptor and the regulation of thyrocyte function and growth. *Endocr Rev*; **13**:596-611.
- Vuissoz JM, Deladoey J, Buyukgebiz A, Cemeroglu P, Gex G, Gallati S, Mullis PE. (2001) New autosomal recessive mutation of the TSH-beta subunit gene causing central isolated hypothyroidism. *J Clin Endocrinol Metab*; **86**:4468-71.
- Wang CD, Buck MA, Fraser CM (1991): Site-directed mutagenesis of alpha 2A-adrenergic receptors: identification of amino acids involved in ligand binding and receptor activation by agonists. *Mol. Pharmacol.*; **40**:168-179.
- Wang Z, Hipkin RW, Ascoli M (1996): Progressive cytoplasmic tail truncations of the lutropin-choriogonadotropin receptor prevent agonist- or phorbol ester-induced phosphorylation, impair agonist- or phorbol ester-induced desensitization, and enhance agonist-induced receptor down-regulation. *Mol Endocrinol*; **10**:748-59.
- Weiss J, Axelrod L, Whitcomb RW, Harris PE, Crowley WF, Jameson JL (1992). Hypogonadism caused by a single amino acid substitution in the beta subunit of luteinizing hormone. *N Engl J Med*; **326**:179-83.
- White JH, Wise A, Main MJ, Green A, Fraser NJ, Disney GH, Barnes AA, Emson P, Foord SM, Marshall FH (1998). Heterodimerization is required for the formation of a functional GABA(B) receptor. *Nature*; **396**:679-82.
- Wonerow P, Schöneberg T, Schultz G, Gudermann T, Paschke R (1998): Deletion in the third intracellular loop of the thyrotropin receptor. A new mechanism for constitutive activation. *J Biol Chem*; **273**:7900-7905.
- Wu W, Cogan JD, Pfaffle RW, Dasen JS, Frisch H, O'Connell SM, Flynn SE, Brown MR, Mullis PE, Parks JS, Phillips JA 3rd, Rosenfeld MG (1998). Mutations in PROP1 cause familial combined pituitary hormone deficiency. *Nat Genet*; **18**:147-9.
- Xie J, Pannain S, Pohlenz J, Weiss RE, Moltz K, Morlot M, Asteria C, Persani L, Beck-Peccoz P, Parma J, Vassart G, Refetoff S (1997): Resistance to thyrotropin (TSH) in three families is not associated with mutations in the TSH receptor or TSH. *J Clin Endocrinol Metab*; **82**:3933-3940.
- Yaswen L, Diehl N, Brennan MB, Hochgeschwender U (1999). Obesity in the mouse model of pro-opiomelanocortin deficiency responds to peripheral melanocortin. *Nat Med*; **5**:1066-70.
- Yeo GS, Farooqi IS, Aminian S, Halsall DJ, Stanhope RG, O'Rahilly S (1998). A frameshift mutation in MC4R associated with dominantly inherited human obesity. *Nat Gene*; **20**:111-2.