

**Extracellular osmolality and solute composition participate in the  
expressional regulation of Aquaporin-2 in renal inner medullary  
collecting duct cells**

**Evidence for an involvement of the TonE/TonEBP pathway**

Dissertation to obtain the academic degree

Doctor rerum naturalium (Dr. rer. nat.)

submitted to the Department of Biology

of Freie Universität Berlin, Germany

by

Robert Storm

born in Berlin, Germany

April, 2004

1. Expert: Prof. Dr. Walter Rosenthal

2. Expert: Prof. Dr. Horst Kreß

Disputation on the 12<sup>th</sup> of July, 2004

## Acknowledgements:

I would like to thank Prof. Dr. W. Rosenthal for the possibility to perform this study in his group at the Forschungsinstitut für Molekulare Pharmakologie (FMP) in Berlin.

Moreover, I would like to thank Dr. K. Maric and Prof. Dr. W. Rosenthal for their support and for helpful discussions and practical advice concerning this work.

I would also like to thank Prof. Dr. H. Kreß of the Freie Universität Berlin biology department for his cooperativeness and support during my studies and with this work.

Many thanks to all the colleagues at the FMP who helped with practical advice and inspiring discussions, in particular Dr. E. Klussmann, Dr. E. Klauschenz, A. Geelhaar and all members of the Klussmann lab.

In addition, I would like to thank my parents and friends for their support.

## Table of contents

	page
<b>1. Introduction</b>	1
1.1 Structure and function of the mammalian kidney	1
1.2 Aquaporins: structure and function	4
1.3 Molecular mechanisms regulating the expression and trafficking of AQP2 in renal principal cells: description of the classical AVP-triggered pathway	5
1.4 States with impaired water homeostasis due to altered AQP2 expression	7
1.5 Introduction of the primary cultured inner medullary collecting duct (IMCD) cell model used in this work	9
1.6 The aim of this study	10
<b>2. Materials &amp; Experimental procedures</b>	11
<b>2.1 Materials</b>	11
2.1.1 Chemicals, reagents, antibodies, cells, and animals	11
2.1.2 Apparatus and software	14
<b>2.2 Experimental procedures</b>	16
2.2.1 Preparation of primary cultures of IMCD cells from rat inner medullae	16
2.2.2 Coating of glass and plastic ware with collagen type IV for cell culture	16
2.2.3 Media used for cell culture	17
2.2.4 Protein analysis	19
2.2.4.1 Determination of protein concentrations by the method of Bradford	19
2.2.4.2 Protein preparations	20
2.2.4.3 Size separation of proteins by SDS-PAGE	20
2.2.4.4 Semi-dry Western blotting	21
2.2.4.5 Immunocytochemistry	22
2.2.5 Nucleic acid analysis	23
2.2.5.1 Isolation of total RNA from IMCD cells	23
2.2.5.2 Horizontal agarose gel electrophoresis of DNA	23
2.2.5.3 Recovery of DNA fragments from agarose gels	24
2.2.5.4 Gel electrophoresis of RNA and Northern blot analysis	24
2.2.5.5 Radioactive random prime labelling of cDNA fragments	26
2.2.5.6 Purification of [ $\alpha$ 32-P]dCTP-labeled cDNA fragments using Sephadex G50 columns	26
2.2.5.7 Sequence searches and alignments	27
2.2.5.8 Generation of GFP-TonEBP and dominant negative GFP-TonEBP	27

2.2.6	Transfection of competent bacteria	28
2.2.7	Immunofluorescence	29
<b>3.</b>	<b>Results</b>	<b>31</b>
3.1	Elevated osmolality is required for the DBcAMP-elicited expression of robust AQP2 protein levels in IMCD cells	31
3.1.1	AQP2 protein levels are regulated by extracellular osmolality and solute composition	32
3.1.2	The effects of osmolality and solute composition on AQP2 protein levels are due to altered AQP2 mRNA expression	34
3.1.3	Altered osmolality has no effect on the cytomegalovirus-governed expression of AQP2 in MDCK1 cells stably transfected with AQP2	35
3.1.4	Evaluation of an alternative to the quantification of AQP2 protein levels in membrane preparations: reducing animal consumption	37
3.2	Osmolality and solute composition have no effect on the hormone-stimulated trafficking of AQP2 in IMCD cells	40
3.3	The role of DBcAMP in the expression of AQP2 in IMCD cells	41
3.3.1	Ser-133 phosphorylation of CREB is stimulated by DBcAMP	42
3.3.2	The DBcAMP-induced phosphorylation of CREB is not affected by changes in extracellular osmolality and solute composition	43
3.4	Hypertonic challenge, not simple elevation of osmolality by the membrane permeating solute urea, promotes AQP2 expression	44
3.4.1	The expression of AQP2 correlates with extracellular sodium concentrations	46
3.4.2	The effect of hypertonic challenge on AQP2 expression is enhanced by urea	47
3.5	The kinetics of AQP2 regulation in response to hypo- and hypertonic challenge	48
3.6	The kinetics of AQP2 regulation in response to DBcAMP withdrawal and addition	50
3.7	The human, rat, and mouse AQP2 promoter regions contain a conserved tonicity responsive element (TonE)	51
3.7.1	The expression of the transcription factor TonEBP is regulated by osmolality and solute composition in IMCD cells	53
3.7.2	TonEBP activity (nuclear localization) corresponds with AQP2 expression in response to hyper- or hypotonic challenge	55

3.8	Investigations of the effect of compounds reported to interfere with the TonE/TonEBP pathway on the expression of AQP2	58
3.8.1	Proteasome inhibition abrogates up-regulation of AQP2 expression in response to hypertonic challenge	58
3.8.2	Rottlerin impairs up-regulation of AQP2 expression in response to hypertonic challenge	60
3.8.3	MEK / p38 inhibition reduces the detrimental effect of hypotonic challenge on AQP2 expression	62
3.8.4	Influence of proteasome inhibition, rottlerin-treatment, and inhibition of MEK1/p38 on cell morphology, cellular localization and expression of AQP2, TonEBP and the actin cytoskeleton in IMCD cells	67
4.	<b>Discussion</b>	74
5.	<b>Abstract (English/German)</b>	91
6.	<b>References</b>	96
7.	<b>Appendix</b>	110
7.1	Publications	110
7.2	Abbreviations	111