

1. Literaturverzeichnis

Abelseth, M. (1986):

Clinical Values and Procedures
The Merck Veterinary Manual.

Adal, Y., Smit, M.F., Osicka, T.M., Comper, W.D. (1995):

Albumin interaction with the glomerular capillary wall in vitro.
Kidney Int 47(4): 1031-1038.

Ajayi, A.A., Newaz, M., Hercule, H., Saleh, M., Bode, C.O., Oyekan, A.O. (2003):

Endothelin-like action of Pausinystalia yohimbe aqueous extract on vascular and renal regional hemodynamics in Sprague Dawley rats.

Methods Find Exp Clin Pharmacol 25(10): 817-822.

Alcorn, D., Emslie, K.R., Ross, B.D., Ryan, G.B., Tange, J.D. (1981):

Selective distal nephron damage during isolated kidney perfusion.
Kidney Int 19(5): 638-647.

Ametani, M.S., Southard, J.H., Belzer, F.O. (1990):

Importance of glutathione and adenosine in cold storage of the kidney.
Transplant Proc 22(2): 469-471.

Arnaud, F.G., Khirabadi, B.S., Fahy, G.M. (2000):

Normothermic blood perfusion of isolated rabbit kidneys. II. In vitro evaluation of renal function followed by orthotopic transplantation.

Asaio J 46(6): 707-718.

Arnould, T., Michiels, C., Remacle, J. (1993):

Increased PMN adherence on endothelial cells after hypoxia: involvement of PAF, CD18/CD11b, and ICAM-1.

Am J Physiol 264(5 Pt 1): C1102-1110.

Baeyer, H.v., Haerberle, D.A., van Liew, J.B., Hare, D. (1980):

Glomerular tubular balance of renal D-glucose transport during hyperglycemia: clearance and micropuncture studies on its characterisation at saturated transport conditions.

Pflugers Arch 384(1): 39-47.

Baeyer, H.v., Stahl, K., Häusler, M., Meissler, M., Unger, V., Frank, J., Grosse-Siestrup, C., Kaczmarczyk, G., Affeld, K., Flaig, H.-J., Steinbach, B. (1997):

Eine neue Methode zur Ex-vivo-Vollblut-Perfusion isolierter Warmblüterorgane, dargestellt an der Niere von Schweinen.

Biomedizinische Technik 42(3): 61-68.

Barber, W.H., Laskow, D.A., Deierhoi, M.H., Poplawski, S.C., Diethelm, A.G. (1991):

Comparison of simple hypothermic storage, pulsatile perfusion with Belzer's gluconate-albumin solution, and pulsatile perfusion with UW solution for renal allograft preservation.

Transplant Proc 23(5): 2394-2395.

Beck, F., Thurau, K., Straunthaler, G. (1992):

Pathophysiology and pathobiochemistry of acute renal failure., in "The Kidney: Physiology and Pathophysiology".

Seldin, DW and Giebisch, G.

New York, Raven Press Ltd.: 3157-3179.

Bedell, S.E., Bush, B.T. (1985):

Erythrocyte sedimentation rate. From folklore to facts.

Am J Med 78(6 Pt 1): 1001-1009.

Behring. (2004):

Albumin Produktinformation.

URL: <http://www.hsaexcipient.com/ZLBBPRODUCT/Excipient/English/home.jsp>.

Belzer, F.O., Ashby, B.S., Downes, G.L. (1968):

Lactic acid dehydrogenase as an index of future function of cadaver kidneys during isolated perfusion.
Surg Forum 19: 205-206.

Bennett, M., Horton, S., Thuys, C., Augustin, S., Rosenberg, M., Brizard, C. (2004):

Pump-induced haemolysis: a comparison of short-term ventricular assist devices.
Perfusion 19(2): 107-111.

Berliner, R.W., Levinsky, N.G., Davidson, D.G., Eden, M. (1958):

Dilution and concentration of the urine and the action of antidiuretic hormone.
Am J Med 24(5): 730-744.

Bessems, M., Doorschodt, B.M., van Vliet, A.K., van Gulik, T.M. (2004):

Preservation of rat livers by cold storage: a comparison between the University of Wisconsin solution and Hypothermosol.
Ann Transplant 9(2): 35-37.

Bessems, M., t Hart, N.A., Tolba, R., Doorschodt, B.M., Leuvenink, H.G., Ploeg, R.J., Minor, T., van Gulik, T.M. (2006):

The isolated perfused rat liver: standardization of a time-honoured model.
Lab Anim 40(3): 236-246.

Bigaud, M., Gfeller, P., Deveze, S., Vogt, G., Evenou, J.P., Bruns, C., Zerwes, H.G. (1998):

Transplantation-induced ischemia/reperfusion injury in the rat heart.
Transplant Proc 30(5): 2311-2313.

Bilzer, M. (1998):

"Hormonelle und antioxidative Prävention von Ischämie-Reperfusionsschäden der Leber durch atriales natriuretisches Peptid und Glutathion"
Lengerich, Pabst Science Publishers.

Bilzer, M., Witthaut, R., Paumgartner, G., Gerbes, A.L. (1994):

Prevention of ischemia/reperfusion injury in the rat liver by atrial natriuretic peptide.
Gastroenterology 106(1): 143-151.

Blauhut, B. (1986):

"Albumin and the Systemic Circulation"
Freiburg, Karger Verlag für Medizin und Naturwissenschaften GmbH.

BMVEL, B.f.V., Ernährung und Landwirtschaft. (2005):

Tierschutzbericht 2005.

URL: www.verbraucherministerium.de.

Boehnert, M.U., Hilbig, H., Armbruster, F.P. (2005):

Relaxin as an additional protective substance in preserving and reperfusion solution for liver transplantation, shown in a model of isolated perfused rat liver.
Ann N Y Acad Sci 1041: 434-440.

Boldt, J. (2000):

The good, the bad, and the ugly: should we completely banish human albumin from our intensive care units?
Anesth Analg 91(4): 887-895.

Boldt, J. (2003):

New light on intravascular volume replacement regimens: what did we learn from the past three years?
Anesth Analg 97(6): 1595-1604.

Boldt, J., Suttner, S. (2005):

Plasma substitutes.

Minerva Anesthesiol 71(12): 741-758.

Bonventre, J.V., Weinberg, J.M. (1992):

Kidney preservation ex vivo for transplantation.

Annu Rev Med 43: 523-553.

Bowman, R.H., Maack, T. (1974):

Effect of albumin concentration and ADH on H₂O and electrolyte transport in perfused rat kidney.

Am J Physiol 226(2): 426-430.

Brasile, L., Green, E., Haisch, C. (1997):

Ex vivo resuscitation of kidneys following postmortem warm ischemia.

Transplant Proc 29(8): 3518-3519.

Brasile, L., Green, E., Haisch, C. (1997):

Oxygen consumption in warm-preserved renal allografts.

Transplant Proc 29(1-2): 1322-1323.

Brasile, L., Green, E., Haisch, C. (1997):

Warm ex vivo perfusion prevents reperfusion injury in warm ischemically damaged kidneys.

Transplant Proc 29(8): 3422-3423.

Breimer, M.E., Svalander, C.T., Haraldsson, B., Bjorck, S. (1996):

Physiological and histological characterisation of a pig kidney in vitro perfusion model for xenotransplantation studies.

Scand J Urol Nephrol 30(3): 213-221.

Brezis, M., Rosen, S., Silva, P., Epstein, F.H. (1984):

Selective vulnerability of the medullary thick ascending limb to anoxia in the isolated perfused rat kidney.

J Clin Invest 73(1): 182-190.

Brezis, M., Rosen, S., Silva, P., Epstein, F.H. (1984):

Renal ischemia: a new perspective.

Kidney Int 26(4): 375-383.

Canelo, R., Hakim, N.S., Ringe, B. (2003):

Experience with histidine tryptophan ketoglutarate versus University Wisconsin preservation solutions in transplantation.

Int Surg 88(3): 145-151.

Cassutto, B.H., Gfeller (2003):

Use of intravenous lidocaine to prevent reperfusion injury and subsequent multiple organ dysfunction syndrome.

J Vet Emerg Crit Care 13 (3): 137 - 148.

Chamorro, V., Moreno, J.M., Wangenstein, R., Sainz, J., Rodriguez-Gomez, I., Osuna, A., Vargas, F. (2004):

Effects of deoxycorticosterone on renal vascular reactivity and flow-pressure curve in spontaneously hypertensive rats.

J Physiol Pharmacol 55(1 Pt 1): 17-26.

Chan, L., Chittinandana, A., Shapiro, J.I., Shanley, P.F., Schrier, R.W. (1994):

Effect of an endothelin-receptor antagonist on ischemic acute renal failure.

Am J Physiol 266(1 Pt 2): F135-138.

Chien, C., Lee PH., Chen, C., Ma, M., Lai, M., Hsu, S. (2001):

De novo demonstration and co-localization of free-radical production and apoptosis formation in rat kidney subjected to ischemia/reperfusion.

J Am Soc Nephrol 12 (5): 973 - 982.

Collins, G.M., Wicomb, W.N. (1992):

New organ preservation solutions.

Kidney Int Suppl 38: S197-202.

Compagnon, P., Lindell, S., Ametani, M.S., Gilligan, B., Wang, H.B., D'Alessandro, A.M., Southard, J.H., Mangino, M.J. (2005):

Ischemic preconditioning and liver tolerance to warm or cold ischemia: experimental studies in large animals.

Transplantation 79(10): 1393-1400.

Comper, W.D., Glasgow, E.F. (1995):

Charge selectivity in kidney ultrafiltration.

Kidney Int 47(5): 1242-1251.

Cowley, A.W., Jr. (1997):

Role of the renal medulla in volume and arterial pressure regulation.

Am J Physiol 273(1 Pt 2): R1-15.

Cuypers, Y., Vandenreyt, I., Bipat, R., Toelsie, J., Van Damme, B., Steels, P. (2000):

The functional state of the isolated rabbit kidney perfused with autologous blood.

Pflugers Arch 440(4): 634-642.

Datta, P.R., Nelson, M.J. (1968):

p,p'-DDT detoxication by isolated perfused rat liver and kidney.

IMS Ind Med Surg 37(7): 521.

Daugharty, T.M., Brenner, B.M. (1975):

Reversible hemodynamic defect in glomerular filtration rate after ischemic injury.

Am J Physiol 228(5): 1436-1439.

Daugharty, T.M., Ueki, I.F., Mercer, P.F., Brenner, B.M. (1974):

Dynamics of glomerular ultrafiltration in the rat. V. Response to ischemic injury.

J Clin Invest 53(1): 105-116.

Davidson, I.J. (2006):

Renal impact of fluid management with colloids: a comparative review.

Eur J Anaesthesiol 23(9): 721-738.

De Mello, G., Maack, T. (1976):

Nephron function of the isolated perfused rat kidney.

Am J Physiol 231(6): 1699-1707.

Deen, W.M., Bohrer, M.P., Robertson, C.R., Brenner, B.M. (1977):

Determinants of the transglomerular passage of macromolecules.

Fed Proc 36(12): 2614-2618.

Deetjen, P., Boylan, J., Kramer, K. (1976):

Tubulärer Transport

Niere und Wasserhaushalt - Physiologie des Menschen. Gauer, O, Kramer, K and Jung, R. München, Urban und Schwarzenberg: 15-37.

Dittrich, S., Schuth, A., von Baeyer, H., Grosse-Siestrup, C., Lange, P.E., Kaczmarczyk, G. (1998):

Effect of blood viscosity on the function of isolated perfused porcine kidney after cold preservation.

Zentralbl Chir 123(7): 809-813.

Dittrich, S., Schuth, A., Aurich, H., vonLoeper, J., Grosse-Siestrup, C., Lange, P.E. (2000):

Haemodilution improves organ function during normothermic cardiopulmonary bypass: investigations in isolated perfused pig kidneys.

Perfusion 15(3): 225-229.

- Dittrich, S., Lippek, F., Gratopp, A., Grosse-Siestrup, C., Lange, P.E., Buhrer, C. (2002):**
Intravascular granulocyte aggregates caused by the selectin-binding carbohydrate fucoidin in pig kidneys.
Clin Exp Pharmacol Physiol 29(10): 909-914.
- Eckhart, J., Forst, H., Burchardi, H. (2004):**
Akutes Nierenversagen, in "Intensivmedizin-Kopendium und Repetitorium zur interdisziplinären Weiter- und Fortbildung".
München, ecomed: 24-37.
- Edelstein, C.L., Ling, H., Schrier, R.W. (1997):**
The nature of renal cell injury.
Kidney Int 51(5): 1341-1351.
- Eder, H. (1987):**
Blut und Lymphe, in "Lehrbuch der Veterinärphysiologie".
Scheunert A. and Trautmann, A.
Berlin; Hamburg, Parey Verlag
160-207.
- Eppel, G.A., Osicka, T.M., Pratt, L.M., Jablonski, P., Howden, B., Glasgow, E.F., Comper, W.D. (2001):**
The return of glomerular filtered albumin to the rat renal vein--the albumin retrieval pathway.
Ren Fail 23(3-4): 347-363.
- Erkasap, N., Ates, E., Erkasap, S., Kaygisiz, Z. (2002):**
Lidocaine-containing Euro-Collins solution prevents renal injury in the isolated perfused canine kidney exposed to prolonged cold ischemia.
Physiol Res 51(5): 493-499.
- Erkasap, S., Ates, E. (2000):**
L-Arginine-enriched preservation solution decreases ischaemia/reperfusion injury in canine kidneys after long-term cold storage.
Nephrol Dial Transplant 15(8): 1224-1227.
- Faber, M.D., Narins, R.G. (1992):**
Insidious renal failure.
Hosp Pract (Off Ed) 27(5): 159-164, 173-176.
- Fehrenberg, C., von Baeyer, H., Unger, V., Schmitt, R., Haider, W., Quarcoo, D., Groneberg, D.A., Grosse-Siestrup, C. (2004):**
Protective effects of B2 preservation solution in comparison to a standard solution (histidine-tryptophan-ketoglutarate/Bretschneider) in a model of isolated autologous hemoperfused porcine kidney.
Nephron Physiol 96(2): p52-58.
- Feola, M., Simoni, J., Tran, R., Canizaro, P.C. (1990):**
Nephrotoxicity of hemoglobin solutions.
Biomater Artif Cells Artif Organs 18(2): 233-249.
- Frodin, L., Engberg, A., Aallskog, O., Wolgast, M. (1975):**
Renal transplantation in the rat. IV. Cortical pressure gradients measured in vivo and during isolated perfusion with reference to renal function after transplantation.
Scand J Clin Lab Invest 35(5): 463-472.
- Fromm, M., Hierholzer, K. (2000):**
Niere, in "Physiologie der Haustiere".
Engelhardt, Wv, Breves, G.
Stuttgart, Enke Verlag: 254 - 294.

- Ghinea, N., Fixman, A., Alexandru, D., Popov, D., Hasu, M., Ghitescu, L., Eskenasy, M., Simionescu, M., Simionescu, N. (1988):**
Identification of albumin-binding proteins in capillary endothelial cells.
J Cell Biol 107(1): 231-239.
- Ghitescu, L., Fixman, A., Simionescu, M., Simionescu, N. (1986):**
Specific binding sites for albumin restricted to plasmalemmal vesicles of continuous capillary endothelium: receptor-mediated transcytosis.
J Cell Biol 102(4): 1304-1311.
- Giebisch, G. (1979):**
Renal tubular control of potassium transport.
Klin Wochenschr 57(19): 1001-1008.
- Glazier, W.B., Chaudry, I.H., Siegel, N.J., Kashgarian, M., Lytton, B., Baue, A.E. (1978):**
Enhanced recovery from severe ischemic renal injury with adenosine triphosphate-magnesium chloride: administration after the insult.
Surg Forum 29: 82-84.
- Gotoh, S., Ogihara, T., Ohde, H., Nakamaru, M., Masuo, K., Saeki, S., Kumahara, Y. (1982):**
Release of prostaglandin I₂ from hog kidney by propranolol.
Prostaglandins Leukot Med 8(6): 537-544.
- Grosse-Siestrup, C., Unger, V., Fehrenberg, C., v Baeyer, H., Fischer, A., Schaper, F., Groneberg, D.A. (2002):**
A model of isolated autologously hemoperfused porcine slaughterhouse kidneys.
Nephron 92(2): 414-421.
- Grosse-Siestrup, C., Unger, V., Meissler, M., Nagel, S., Wussow, A., Peiser, C., Fischer, A., Schmitt, R., Groneberg, D.A. (2003):**
Hemoperfused isolated porcine slaughterhouse kidneys as a valid model for pharmacological studies.
J Pharm Sci 92(6): 1147-1154.
- Guarrera, J.V., Polyak, M.M., O'Mar Arrington, B., Boykin, J., Brown, T., Jean-Jacques, M.A., Kapur, S., Stubenbord, W.T., Kinkhabwala, M. (2004):**
Pushing the envelope in renal preservation; improved results with novel perfusate modifications for pulsatile machine perfusion of cadaver kidneys.
Transplant Proc 36(5): 1257-1260.
- Gyrd-Hansen, N., Lund, F., Hansen, R.I., Fernandes, A., Kemp, E., Lokkegaard, H., Rasmussen, F. (1969):**
Renal clearances in unilaterally nephrectomized pigs and in pigs with an autotransplanted kidney.
Acta Med Scand 186(5): 383-391.
- Harper, S., Hosgood, S., Kay, M., Nicholson, M. (2006):**
Leucocyte depletion improves renal function during reperfusion using an experimental isolated haemoperfused organ preservation system.
Br J Surg 93(5): 623-629.
- Hartmann, H. (1994):**
Pathophysiologie der Organsysteme - Funktionsstörungen der Nieren und ableitenden Harnwege, in "Klinische Pathologie der Haustiere".
Hartmann H. and Meyer, H.
Jena, Stuttgart, Gustav Fischer Verlag. 1: 412 - 430.
- Hartmann, H., Ungemach, F.R. (1996):**
Pharmakologie des Wasser- und Elektrolythaushaltes, in "Lehrbuch der Pharmakologie und Toxikologie für die Veterinärmedizin".
Frey, HH and Löscher, W.
Stuttgart, Enke Verlag. 2: 170-189.

- Hauet, T., Baumert, H., Amor, I.B., Goujon, J.M., Gibelin, H. (2000):**
Protection of autotransplanted pig kidneys from ischemia-reperfusion injury by polyethylene glycol.
Transplantation 70(11): 1569-1575.
- Hauet, T., Faure, J.P., Baumert, H., Bardou, A., Gibelin, H., Beguinot, S. (1998):**
Influence of different colloids on hemodynamic and renal functions: comparative study in an isolated perfused pig kidney model.
Transplant Proc 30(6): 2796-2797.
- Hellige, G., Spiekermann, P.G. (2000):**
Herzinsuffizienz und Nierenfunktion.
Herz-Kreislauf Transparenz 2: 1-28.
- Henderson, L.W., Colton, C., Leber, H., Quelhorst, E. (1978):**
Hemofiltration.
Trans Am Soc Artif Intern Organs 24: 788-789.
- Hendriks, J.M., Grootenboers, M.J., Schramel, F.M., van Boven, W.J., Stockman, B., Seldenrijk, C.A., ten Broecke, P., Knibbe, C.A., Slee, P., De Bruijn, E., Vlaeminck, R., Heeren, J., Vermorken, J.B., van Putte, B., Romijn, S., Van Marck, E., Van Schil, P.E. (2004):**
Isolated lung perfusion with melphalan for resectable lung metastases: a phase I clinical trial.
Ann Thorac Surg 78(6): 1919-1927.
- Heyman, S.N., Rosen, S., Brezis, M. (1997):**
The renal medulla: life at the edge of anoxia.
Blood Purif 15(4-6): 232-242.
- Hochel, J., Lehmann, D., Fehrenberg, C., Unger, V., Groneberg, D.A., Grosse-Siestrup, C. (2003):**
Effects of different perfusates on functional parameters of isolated perfused dog kidneys.
Nephrol Dial Transplant 18(9): 1748-1754.
- Hrabalova, M., Bachleda, P., Lubuska, L., Kojecky, Z., Zadrazil, J., Krejci, K., Al-Jabry, S. (2003):**
Effect of various protective solutions on function after kidney transplantation.
Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub 147(2): 197-202.
- Huxley, V.H., Curry, F.E. (1985):**
Albumin modulation of capillary permeability: test of an adsorption mechanism.
Am J Physiol 248(2 Pt 2): H264-273.
- Hvistendahl, J.J., Pedersen, T.S., Hvistendahl, G.M., Djurhuus, J.C., Frokiaer, J. (2001):**
Reduced renal vascular resistance in response to verapamil during graded ureter obstruction in pigs.
Urol Res 29(5): 350-358.
- Imai, M., Kokko, J.P. (1972):**
Effect of peritubular protein concentration on reabsorption of sodium and water in isolated perfused proximal tubules.
J Clin Invest 51(2): 314-325.
- Imaizumi, T.A., Stafforini, D.M., Yamada, Y., McIntyre, T.M., Prescott, S.M., Zimmerman, G.A. (1995):**
Platelet-activating factor: a mediator for clinicians.
J Intern Med 238(1): 5-20.
- Inman, S.R., Davis, N.A., Olson, K.M., Lukaszek, V.A., McKinley, M.R., Seminerio, J.L. (2003):**
Rapamycin preserves renal function compared with cyclosporine A after ischemia/reperfusion injury.
Urology 62(4): 750-754.
- Jablonski, P., Howden, B.O., Rae, D.A., Birrell, C.S., Marshall, V.C., Tange, J. (1983):**
An experimental model for assessment of renal recovery from warm ischemia.
Transplantation 35(3): 198-204.

Literaturverzeichnis

- Jacob, M., Bruegger, D., Rehm, M., Welsch, U., Conzen, P., Becker, B.F. (2006):**
Contrasting effects of colloid and crystalloid resuscitation fluids on cardiac vascular permeability.
Anesthesiology 104(6): 1223-1231.
- Joannidis, M. (2003):**
Das akute Nierenversagen.
URL: http://www.intensiv-innsbruck.at/education/nierenversagen_joannidis.htm.
- Joannidis, M., P.G., M. (2005):**
Epidemiology and natural history of acute renal failure in the ICU.
Crit Care Clin 21(2): 239 - 249.
- Johnston, P.A., Bernard, D.B., Perrin, N.S., Levinsky, N.G. (1981):**
Prostaglandins mediate the vasodilatory effect of mannitol in the hypoperfused rat kidney.
J Clin Invest 68(1): 127-133.
- Jorres, A. (2002):**
Acute renal failure. Extracorporeal treatment strategies.
Minerva Med 93(5): 329-324.
- Jorres, A.J. (2002):**
Acute renal failure: pathogenesis, diagnosis and conservative treatment.
Minerva Med 93(2): 85-93.
- Kirby, R. (1989):**
Acute renal failure as a complication in the critically ill animal.
Vet Clin North Am Small Anim Pract 19(6): 1189-1208.
- Klausner, J.M., Paterson, I.S., Goldman, G., Kobzik, L., Rodzen, C. (1989):**
Postischemic renal injury is mediated by neutrophils and leukotrienes.
Am J Physiol 256(5 Pt 2): F794-802.
- Knight, A.J., O'Leary, E.A., Nicholson, M.L. (2001):**
Cold ischaemia further increases intrarenal resistance when non-heart-beating kidneys are pulsatile machine perfused.
Transplant Proc 33(1-2): 893-894.
- Kohrmann, K.U., Back, W., Bensemman, J., Florian, J., Weber, A., Kahmann, F., Rassweiler, J., Alken, P. (1994):**
The isolated perfused kidney of the pig: new model to evaluate shock wave-induced lesions.
J Endourol 8(2): 105-110.
- Kraft, H, Schillinger, D. (1989)**
Normwerttabellen in "Klinische Labormethoden der Veterinärmedizin bei Haussäugetieren."
Stuttgart, Thieme Verlagsgruppe, Enke-Verlag
- Lang, F., Fromm, M. (2000):**
Funktionen der Niere, in "Physiologie des Menschen".
Schmidt, R, Thews, G and Lang, F.
Berlin, Springer Verlag. 28: 737-776.
- Lauriat, S., Linas, S.L. (1998):**
The role of neutrophils in acute renal failure.
Semin Nephrol 18(5): 498-504.
- Lazar, N., Dallos, G., Nemes, B., Nemeth, T., Sotonyi, P., Kobori, L. (1997):**
Experimental investigation of preservation injury in animal kidneys after reperfusion with Euro-Collins.
Acta Chir Hung 36(1-4): 192-194.

Lehninger, A.L., Nelson, D.L., Cox, M.M. (1998):

Enzyme, in "Prinzipien der Biochemie".

Tscheche, H.

Heidelberg, Spektrum Akademischer Verlag. 2: 227-279.

Leverett, L.B., Hellums, J.D., Alfrey, C.P., Lynch, E.C. (1972):

Red blood cell damage by shear stress.

Biophys J 12(3): 257-273.

Lieberthal, W., Stephens, G.W., Wolf, E.F., Rennke, H.G., Vasilevsky, M.L., Valeri, C.R., Levinsky, N.G. (1987):

Effect of erythrocytes on the function and morphology of the isolated perfused rat kidney.

Ren Physiol 10(1): 14-24.

Linas, S.L., Shanley, P.F., Whittenburg, D., Berger, E., Repine, J.E. (1988):

Neutrophils accentuate ischemia-reperfusion injury in isolated perfused rat kidneys.

Am J Physiol 255(4 Pt 2): F728-735.

List, W.F. (2001):

Albumin, der erste Blutersatz.

Anesthesiol Intensivmed Notfallmed Schmerzther 36, Suppl 2: 76 - 78.

Little, J.R., Cohen, J.J. (1974):

Effect of albumin concentration on function of isolated perfused rat kidney.

Am J Physiol 226(3): 512-517.

Lohfert, H. (1973):

Untersuchungen am zellfrei perfundierten isolierten Nierenpräparat der Ratte unter besonderer Berücksichtigung des Einflusses der Natriumkonzentration im Perfusionsmedium.

Dissertationsschrift, Freie Universität Berlin.

Lozano, F.S., Lopez-Novoa, J.M., Rodriguez, J.M., Barros, M.B., Garcia-Criado, F.J., Nicolas, J.L., Parreno, A., Revilla, J., Gomez-Alonso, A. (2005):

Exogenous nitric oxide modulates the systemic inflammatory response and improves kidney function after risk-situation abdominal aortic surgery.

J Vasc Surg 42(1): 129-139.

Margarson, M.P., Soni, N.C. (2002):

Effects of albumin supplementation on microvascular permeability in septic patients.

J Appl Physiol 92(5): 2139-2145.

Mason, J., Welsch, J., Takabatake, T. (1983):

Disparity between surface and deep nephron function early after renal ischemia.

Kidney Int 24(1): 27-36.

Mason, J., Welsch, J., Torhorst, J. (1987):

The contribution of vascular obstruction to the functional defect that follows renal ischemia.

Kidney Int 31(1): 65-71.

Messmer, K., Rosenthal, L. (1966):

Isolated perfusion of dog's kidneys in situ before and during mannitol diuresis.

Z Gesamte Exp Med 141(3): 281-300.

Nakamoto, M., Shapiro, J.I., Mills, S.D., Schrier, R.W., Chan, L. (1988):

Improvement of renal preservation by verapamil with 24-hour cold perfusion in the isolated rat kidney.

Transplantation 45(2): 313-315.

Navar, L.G. (1978):

Renal autoregulation: perspectives from whole kidney and single nephron studies.

Am J Physiol 234(5): F357-370.

- Neew-Galuschka, E., Stahl, K., Fehrenberg, C., Kaczmarczyk, G. (1998):**
Pathophysiology of the isolated blood-perfused porcine kidney - better understanding of the Renin-Angiotensin-System (Abstract)
Seminar on isolated perfused organs.
Hamburg, Society for Laboratory Animal Science (GV-SOLAS).
- Nielsen, T., Maaske, C. (1966):**
Some comparative aspects of porcine renal function, in "Swine in biomedical research".
MacClellan, R and Bustad, L.
Seattle, Frayn Publ. Comp. : 529-536.
- Niemi, T.T., Suojaranta-Ylinen, R.T., Kukkonen, S.I., Kuitunen, A.H. (2006):**
Gelatin and hydroxyethyl starch, but not albumin, impair hemostasis after cardiac surgery.
Anesth Analg 102(4): 998-1006.
- Nui, A., Katsuramaki, T., Kikuchi, H., Kukita, K., Kimura, H., Meguro, M., Nagayama, M., Isobe, M., Hirata, K. (2006):**
The functional integrity of a normothermic perfusion system using artificial blood in pig liver.
J Surg Res 131(2): 189-198.
- Ohde, H., Ogihara, T., Nakamaru, M., Higaki, J., Gotoh, S., Masuo, K., Ohtsuka, A., Saeki, S., Kumahara, Y. (1982):**
Effect of prostacyclin infusion on active and inactive renin release in the isolated perfused kidney.
Life Sci 31(26): 3031-3035.
- Paquet, K.J. (1965 a):**
Experimental studies on the function of an isolated kidney in pulsatory and continuous perfusion. 2.
Bruns Beitr Klin Chir 211(4): 463-475.
- Paquet, K.J. (1965 b):**
On methodology and hemodynamics of extracorporeal perfusion and on the performance of the isolated kidney.
Bruns Beitr Klin Chir 211(3): 358-374.
- Pardridge, W.M., Buciak, J.L., Kang, Y.S., Boado, R.J. (1993):**
Protamine-mediated transport of albumin into brain and other organs of the rat. Binding and endocytosis of protamine-albumin complex by microvascular endothelium.
J Clin Invest 92(5): 2224-2229.
- Pedersen, L.M., Terslev, L., PG, S.L., Stokholm, K.H. (2000):**
Urinary albumin excretion and transcapillary escape rate of albumin in malignancies.
Med Oncol 17(2): 117-122.
- Peters, T., Jr. (1985):**
Serum Albumin.
Adv Protein Chem 37: 161-245.
- Rieger, C. (1989):**
Histometrische Untersuchungen an den Nieren gesunder Schweine unterschiedlichen Alters.
Dissertationsschrift, Tierärztliche Hochschule Hannover.
- Riera, M., Torras, J., Herrero, I., Valles, J., Paubert-Braquet, M., Cruzado, J.M., Alsina, J., Grinyo, J.M. (1997):**
Neutrophils accentuate renal cold ischemia-reperfusion injury. Dose-dependent protective effect of a platelet-activating factor receptor antagonist.
J Pharmacol Exp Ther 280(2): 786-794.
- Robinson, J. (1971):**
Control of water content of non-metabolizing kidney slices by sodium chloride and polyethylene glycol (PEG 6000).
J Physiol 213(1): 227-234.

Rosenzweig, L.J., Kanwar, Y.S. (1982):

Removal of sulfated (heparan sulfate) or nonsulfated (hyaluronic acid) glycosaminoglycans results in increased permeability of the glomerular basement membrane to ¹²⁵I-bovine serum albumin.
Lab Invest 47(2): 177-184.

Rozich, J.D., Paul, R.V. (1989):

Acute renal failure precipitated by elevated colloid osmotic pressure.
Am J Med 87(3): 359-360.

Saltzstein, E.C., Villarica, J.D., Bortin, M.M. (1966):

Perfusion of an isolated renal allograft.
Am J Surg 111(6): 877-880.

Santos-Neto, M.S., Carrithers, S.L., Carvalho, A.F., Monteiro, H.S., Greenberg, R.N., Forte, L.R., Fonteles, M.C. (2003):

Guanylin and its lysine-containing analogue in the isolated perfused rat kidney: interaction with chymotrypsin inhibitor.
Pharmacol Toxicol 92(3): 114-120.

Schauer, R. (2000):

Ischemic preconditioning prevents microcirculatory failure following extended of the rat liver.
Hepatology 32: 498(A).

Schauer, R.J., Gerbes, A.L., Vonier, D., op den Winkel, M., Fraunberger, P., Bilzer, M. (2003):

Induction of cellular resistance against Kupffer cell-derived oxidant stress: a novel concept of hepatoprotection by ischemic preconditioning.
Hepatology 37(2): 286-295.

Schmidl, M. (1981)

Laboruntersuchungen für die Diagnose und Verlaufskontrolle in der Veterinärmedizin.
Mannheim, Boehringer Mannheim GmbH

Schnermann, J., Valtin, H., Thurau, K., Nagel, W., Horster, M., Fischbach, H., Wahl, M., Liebau, G. (1969):

Micropuncture studies on the influence of antidiuretic hormone on tubular fluid reabsorption in rats with hereditary diabetes insipidus.
Pflugers Arch 306(2): 103-118.

Schrör, K., Darius, H. (2004):

Therapie mit Antikoagulantien, Plättchenfunktionshemmern und Thrombolytika, in "Pharmakotherapie: klinische Pharmakologie".
Lemmer, B and Brune, K.
Berlin, Springer Verlag: 115-133.

Schumer M., Colombel M.C., Sawczuk I.S., Gobe G., Connor J., O'Toole K.M., Olsson C.A., Wise G.J., R., B. (1992):

Morphologic, biochemical, and molecular evidence of apoptosis during the reperfusion phase after brief periods of renal ischemia.
Am J Pathol 140: 831 - 838.

Schurek, H.J., Alt, J.M. (1981):

Effect of albumin on the function of perfused rat kidney.
Am J Physiol 240(6): F569-576.

Schurek, H.J., Brecht, J.P., Lohfert, H., Hierholzer, K. (1975):

The basic requirements for the function of the isolated cell free perfused rat kidney.
Pflugers Arch 354(4): 349-365.

Schwartz, G.J., Al-Awqati, Q. (1985):

Carbon dioxide causes exocytosis of vesicles containing H⁺ pumps in isolated perfused proximal and collecting tubules.
J Clin Invest 75(5): 1638-1644.

Semb, B.K., Williams, M., Hume, D.H. (1968):

The effect of allogeneic lymphocytes on the isolated perfused kidney.
Transplantation 6(9): 977-985.

Siegel, N.J., Glazier, W.B., Chaudry, I.H., Gaudio, K.M., Lytton, B., Baue, A.E., Kashgarian, M. (1980):

Enhanced recovery from acute renal failure by the postischemic infusion of adenine nucleotides and magnesium chloride in rats.
Kidney Int 17(3): 338-349.

Sinert, R., Peacock, P. (2005):

Acute renal failure.

URL: www.emedicine.com/emerg/topic500.htm.

Sone, M., Ohno, A., Albrecht, G.J., Thureau, K., Beck, F.X. (1995):

Restoration of urine concentrating ability and accumulation of medullary osmolytes after chronic diuresis.
Am J Physiol 269(4 Pt 2): F480-490.

Sonnenberg, H., Deetjen, P. (1964):

Methods for Perfusing Single Nephron Segments.

Pflügers Arch Gesamte Physiol Menschen Tiere 278: 669-674.

Southard, J.H., den Butter, B., Marsh, D.C., Lindell, S., Belzer, F.O. (1991):

The role of oxygen free radicals in organ preservation.
Klin Wochenschr 69(21-23): 1073-1076.

Spitzer, A., Windhager, E.E. (1970):

Effect of peritubular oncotic pressure changes on proximal tubular fluid reabsorption.
Am J Physiol 218(4): 1188-1193.

Sümpelmann, R., Zander, R. (2001):

Gelatin protects erythrocytes in vivo and in vitro against mechanical stress.
Anesthesiol Intensivmed Notfallmed Schmerzther 36 Suppl 1: S62-68.

Sümpelmann, R., Schurholz, T., Marx, G., Zander, R. (2000):

Protective effects of plasma replacement fluids on erythrocytes exposed to mechanical stress.
Anaesthesia 55(10): 976-979.

Swindle, M.M. (1992):

Preface, in "Swine as Models in Biomedical Research".

Swindle, MM.

Ames IA, Iowa State University Press. 1: 9-10.

Swindle, M.M., Smith, A.C., Laber-Laird, K., Dungan, L. (1994):

Swine in Biomedical Research: Management and Models.

ILAR News 36(1): 1 - 5.

Tack, B.F., Dean, J., Eilat, D., Lorenz, P.E., Schechter, A.N. (1980):

Tritium labeling of proteins to high specific radioactivity by reduction methylation.
J Biol Chem 255(18): 8842-8847.

Thadhani, R., Pascual, M., Bonventre, J.V. (1996):

Acute renal failure.

N Engl J Med 334(22): 1448-1460.

Thiessen, H. (1976):

Untersuchungen an den Nieren von Haus- und Wildschweinen.

Zeitschr Tierzücht Züchtgsbiol 93: 178 - 216.

Thun, R., Schwartz-Porsche, D. (1994):

Regulation der Mineralocorticoidsekretion - Das Renin-Angiotensin-Aldosteron-System, in "Veterinärmedizinische Endokrinologie".

Döcke, F.

Jena, Gustav-Fischer-Verlag: 327 - 328.

Thurau, K., Schnermann, J. (1998):

The Na concentration of the macula densa cells as a factor regulating glomerular filtration rate (micropuncture studies). 1965.

J Am Soc Nephrol 9(5): 925-934.

Treska, V., Molacek, J., Kobr, J., Racek, J., Trefil, L., Hes, O. (2006):

Ischemic training and immunosuppressive agents reduce the intensity of ischemic reperfusion injury after kidney transplantation.

Exp Clin Transplant 4(1): 439-444.

Tullis, J.L. (1977):

Albumin. 1. Background and Use.

Jama 237(4): 355-360

Ullrich, K., Drenckhahn, F.O., Jaraus, K.H. (1956):

Untersuchungen zum Problem der Harnkonzentrierung und -verdünnung.

Pflügers Arch 261(1): 62-77.

Ullrich, K.J., Frömter, E., Murer, H. (1979):

Prinzipien des epithelialen Transportes in Niere und Darm.

J Mol Med 57(19): 977-991.

Unger, J.K., Horn, N.A., Kashefi, A., Blumberg, A., Klosterhalfen, B., Rossaint, R. (2001):

The influence of hypoalbuminemia on maximal flow rates and transmembrane pressure during plasmapheresis--an in vitro study.

Blood Purif 19(4): 408-416.

Unger, J.K., Haltern, C., Dohmen, B., Gressner, A., Grosse-Siestrup, C., Groneberg, D.A., Rossaint, R. (2005):

Albumin and hydroxyethyl starch 130 kDa/0.4 improve filter clearance and haemocompatibility in haemo- and plasmafiltration--an in vitro study.

Nephrol Dial Transplant 20(9): 1922-1931.

Unger, V., Grosse-Siestrup, C., Groneberg, D.A. (2006):

Evaluation of renal functional parameters in different settings of isolated organ hemoperfusions.

Physiol Meas 27(11): 1167-1175.

Unger, V., Grosse-Siestrup, C., Fehrenberg, C., Fischer, A., Meissler, M., Groneberg, D.A. (2007):

Reference values and physiological characterization of a specific isolated pig kidney perfusion model.

J Occup Med Toxicol 2(1): in press.

van den Eijnden, M.M., Leuvenink, H.G., Ottens, P.J., t Hart, N.A., van Oeveren, W., Morariu, A.M., van Goor, H., Ploeg, R.J. (2003):

Effect of brain death and non-heart-beating kidney donation on renal function and injury: an assessment in the isolated perfused rat kidney.

Exp Clin Transplant 1(2): 85-95.

van der Hoeven, T., Kilic, A., Vogel, P., Loudovici, D.A., Grosse-Siestrup, C., Affeld, K. (1998):

Vergleichende Untersuchungen der Hämolyse von Blutpumpen verschiedener Bauart

Biomed Tech (Berl). Band 43, Ergänzungsband 1: 332 - 334.

Van der Linden, P., Ickx, B.E. (2006):

The effects of colloid solutions on hemostasis.

Can J Anaesth 53(6 Suppl): S30-39.

van Etten, B., Eggermont, A.M., van Tiel, S.T., Ambagtsheer, G., de Wilt, J.H., ten Hagen, T.L. (2005):

Gene therapy in in vivo isolated perfusion models.

Curr Gene Ther 5(2): 195-202.

Wagner, S.M., Nogueira, A.C., Paul, M., Heydeck, D., Klug, S., Christ, B. (2003):

The isolated normothermic hemoperfused porcine forelimb as a test system for transdermal absorption studies.

J Artif Organs 6(3): 183-191.

Waldmann, K.-H., Wendt, M. (1991):

Kreatininclearance als Grundlage klinischer Nierenfunktionsbestimmung beim Schwein.

Tierärztl Prax 19: 373-380.

Weiss, E. (1999):

Harnorgane - Niere, in "Grundriß der speziellen pathologischen Anatomie der Haustiere".

Dahme, E and Weiss, E.

München, Gießen, Enke Verlag. 5: 243-271.

Wendt, M., Waldmann, K.-H., Bickhardt, K. (1990):

Vergleichende Untersuchung der Inulin- und Kreatinin-Clearance beim Schwein.

J Vet Med A37: 752-759.

Willard, M.D., Refsal, K., Thacker, E. (1987):

Evaluation of plasma aldosterone concentrations before and after ACTH administration in clinically normal dogs and in dogs with various diseases.

Am J Vet Res 48(12): 1713-1718.

Yamauchi, J., Schramm, R., Richter, S., Vollmar, B., Menger, M.D., Minor, T. (2003):

Improvement of microvascular graft equilibration and preservation in non-heart-beating donors by warm preflush with streptokinase.

Transplantation 75(4): 449-453.

Yoshida, H., Okuno, H., Kamoto, T., Habuchi, T., Toda, Y., Hasegawa, S., Nakamura, T., Wada, H., Ogawa, O., Yamamoto, S. (2002):

Comparison of the effectiveness of ET-Kyoto with Euro-Collins and University of Wisconsin solutions in cold renal storage.

Transplantation 74(9): 1231-1236.

Zager, R.A., Fuerstenberg S.M., Baehr P.H., Myerson D., B., T.-S. (1994):

An evaluation of antioxidant effects on recovery from postischemic acute renal failure.

J Am Soc Nephrol 4: 1588 - 1597.