

## 5 Literaturverzeichnis

- Acland, G.M., Aguirre, G.D., Ray, J., Zhang, Q., Aleman, T.S., Cideciyan, A.V., Pearce-Kelling, S.E., Anand, V., Zeng, Y., Maguire, A.M., Jacobson, S.G., Hauswirth, W.W. and Bennett, J. (2001) Gene therapy restores vision in a canine model of childhood blindness. *Nat Genet* 28(1), 92-5.
- Antoni, B.A., Rabson, A.B., Miller, I.L., Trempe, J.P., Chejanovsky, N. and Carter, B.J. (1991) Adeno-associated virus Rep protein inhibits human immunodeficiency virus type 1 production in human cells. *J Virol* 65(1), 396-404.
- Atchison, R.W., Casto, B.C. and Hammon, W.M. (1965) Adenovirus-Associated Defective Virus Particles. *Science* 149, 754-6.
- Balague, C., Kalla, M. and Zhang, W.W. (1997) Adeno-associated virus Rep78 protein and terminal repeats enhance integration of DNA sequences into the cellular genome. *J Virol* 71(4), 3299-306.
- Bantel-Schaal, U. and zur Hausen, H. (1988) Adeno-associated viruses inhibit SV40 DNA amplification and replication of herpes simplex virus in SV40-transformed hamster cells. *Virology* 164(1), 64-74.
- Bartlett, J.S., Wilcher, R. and Samulski, R.J. (2000) Infectious entry pathway of adeno-associated virus and adeno-associated virus vectors. *J Virol* 74(6), 2777-85.
- Batchu, R.B., Kotin, R.M. and Hermonat, P.L. (1994) The regulatory rep protein of adeno-associated virus binds to sequences within the c-H-ras promoter. *Cancer Lett* 86(1), 23-31.
- Batterson, W. and Roizman, B. (1983) Characterization of the herpes simplex virion-associated factor responsible for the induction of alpha genes. *J Virol* 46(2), 371-7.
- Beaton, A., Palumbo, P. and Berns, K.I. (1989) Expression from the adeno-associated virus p5 and p19 promoters is negatively regulated in trans by the rep protein. *J Virol* 63(10), 4450-4.
- Beck, S.E., Jones, L.A., Chesnut, K., Walsh, S.M., Reynolds, T.C., Carter, B.J., Askin, F.B., Flotte, T.R. and Guggino, W.B. (1999) Repeated delivery of adeno-associated virus vectors to the rabbit airway. *J Virol* 73(11), 9446-55.
- Berns, K.I. and Linden, R.M. (1995) The cryptic life style of adeno-associated virus. *Bioessays* 17(3), 237-45.
- Booth, M.J., Mistry, A., Li, X., Thrasher, A. and Coffin, R.S. (2004) Transfection-free and scalable recombinant AAV vector production using HSV/AAV hybrids. *Gene Ther* 11(10), 829-37.
- Brister, J.R. and Muzyczka, N. (1999) Rep-mediated nicking of the adeno-associated virus origin requires two biochemical activities, DNA helicase activity and transesterification. *J Virol* 73(11), 9325-36.
- Brister, J.R. and Muzyczka, N. (2000) Mechanism of Rep-mediated adeno-associated virus origin nicking. *J Virol* 74(17), 7762-71.
- Brockstedt, D.G., Podskakoff, G.M., Fong, L., Kurtzman, G., Mueller-Ruchholtz, W. and Engleman, E.G. (1999) Induction of immunity to antigens expressed by recombinant adeno-associated virus depends on the route of administration. *Clin Immunol* 92(1), 67-75.
- Brown, C.R., Nakamura, M.S., Mosca, J.D., Hayward, G.S., Straus, S.E. and Perera, L.P. (1995) Herpes simplex virus trans-regulatory protein ICP27 stabilizes and binds to 3' ends of labile mRNA. *J Virol* 69(11), 7187-95.
- Brune, W., Messerle, M. and Koszinowski, U.H. (2000) Forward with BACs: new tools for herpesvirus genomics. *Trends Genet* 16(6), 254-9.
- Buller, R.M., Janik, J.E., Sebring, E.D. and Rose, J.A. (1981) Herpes simplex virus types 1 and 2 completely help adenovirus-associated virus replication. *J Virol* 40(1), 241-7.
- Buller, R.M. and Rose, J.A. (1978) Characterization of adenovirus-associated virus-induced polypeptides in KB cells. *J Virol* 25(1), 331-8.
- Campbell, M.E., Palfreyman, J.W. and Preston, C.M. (1984) Identification of herpes simplex virus DNA sequences which encode a trans-acting polypeptide responsible for stimulation of immediate early transcription. *J Mol Biol* 180(1), 1-19.
- Carter, B.J., Laughlin, C.A., de la Maza, L.M. and Myers, M. (1979) Adeno-associated virus autointerference. *Virology* 92(2), 449-62.

- Casto, B.C., Armstrong, J.A., Atchison, R.W. and Hammon, W.M. (1967a) Studies on the relationship between adeno-associated virus type 1 (AAV-1) and adenoviruses. II. Inhibition of adenovirus plaques by AAV; its nature and specificity. *Virology* 33(3), 452-8.
- Casto, B.C., Atchison, R.W. and Hammon, W.M. (1967b) Studies on the relationship between adeno-associated virus type I (AAV-1) and adenoviruses. I. Replication of AAV-1 in certain cell cultures and its effect on helper adenovirus. *Virology* 32(1), 52-9.
- Chadeuf, G., Favre, D., Tessier, J., Provost, N., Nony, P., Kleinschmidt, J., Moullier, P. and Salvetti, A. (2000) Efficient recombinant adeno-associated virus production by a stable rep-cap HeLa cell line correlates with adenovirus-induced amplification of the integrated rep-cap genome. *J Gene Med* 2(4), 260-8.
- Chang, L.S., Shi, Y. and Shenk, T. (1989) Adeno-associated virus P5 promoter contains an adenovirus E1A-inducible element and a binding site for the major late transcription factor. *J Virol* 63(8), 3479-88.
- Chejanovsky, N. and Carter, B.J. (1989) Mutagenesis of an AUG codon in the adeno-associated virus rep gene: effects on viral DNA replication. *Virology* 173(1), 120-8.
- Chen, C. and Okayama, H. (1987) High-efficiency transformation of mammalian cells by plasmid DNA. *Mol Cell Biol* 7(8), 2745-52.
- Chen, C.A. and Okayama, H. (1988) Calcium phosphate-mediated gene transfer: a highly efficient transfection system for stably transforming cells with plasmid DNA. *Biotechniques* 6(7), 632-8.
- Cherepanov, P.P. and Wackernagel, W. (1995) Gene disruption in Escherichia coli: TcR and KmR cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance determinant. *Gene* 158(1), 9-14.
- Cheung, A.K., Hoggan, M.D., Hauswirth, W.W. and Berns, K.I. (1980) Integration of the adeno-associated virus genome into cellular DNA in latently infected human Detroit 6 cells. *J Virol* 33(2), 739-48.
- Chiaroni, J.A., Kim, F., Yang, L. and Kotin, R.M. (1999) Cloning and characterization of adeno-associated virus type 5. *J Virol* 73(2), 1309-19.
- Chiaroni, J.A., Weitzman, M.D., Owens, R.A., Urcelay, E., Safer, B. and Kotin, R.M. (1994) Biologically active Rep proteins of adeno-associated virus type 2 produced as fusion proteins in Escherichia coli. *J Virol* 68(2), 797-804.
- Chirmule, N., Xiao, W., Truneh, A., Schnell, M.A., Hughes, J.V., Zoltick, P. and Wilson, J.M. (2000) Humoral immunity to adeno-associated virus type 2 vectors following administration to murine and nonhuman primate muscle. *J Virol* 74(5), 2420-5.
- Cigan, A.M., Feng, L. and Donahue, T.F. (1988) tRNA<sub>i</sub>(met) functions in directing the scanning ribosome to the start site of translation. *Science* 242(4875), 93-7.
- Conway, J.E., Rhys, C.M., Zolotukhin, I., Zolotukhin, S., Muzychka, N., Hayward, G.S. and Byrne, B.J. (1999) High-titer recombinant adeno-associated virus production utilizing a recombinant herpes simplex virus type I vector expressing AAV-2 Rep and Cap. *Gene Ther* 6(6), 986-93.
- Crute, J.J. and Lehman, I.R. (1989) Herpes simplex-1 DNA polymerase. Identification of an intrinsic 5'---3' exonuclease with ribonuclease H activity. *J Biol Chem* 264(32), 19266-70.
- Crute, J.J., Tsurumi, T., Zhu, L.A., Weller, S.K., Olivo, P.D., Challberg, M.D., Mocarski, E.S. and Lehman, I.R. (1989) Herpes simplex virus 1 helicase-primase: a complex of three herpes-encoded gene products. *Proc Natl Acad Sci U S A* 86(7), 2186-9.
- Datsenko, K.A. and Wanner, B.L. (2000) One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* 97(12), 6640-5.
- Davis, M.D., Wu, J. and Owens, R.A. (2000) Mutational analysis of adeno-associated virus type 2 Rep68 protein endonuclease activity on partially single-stranded substrates. *J Virol* 74(6), 2936-42.
- De, B., Heguy, A., Leopold, P.L., Wasif, N., Korst, R.J., Hackett, N.R. and Crystal, R.G. (2004) Intrapleural administration of a serotype 5 adeno-associated virus coding for alpha1-antitrypsin mediates persistent, high lung and serum levels of alpha1-antitrypsin. *Mol Ther* 10(6), 1003-10.
- de la Maza, L.M. and Carter, B.J. (1981) Inhibition of adenovirus oncogenicity in hamsters by adeno-associated virus DNA. *J Natl Cancer Inst* 67(6), 1323-6.

- Deiss, L.P., Chou, J. and Frenkel, N. (1986) Functional domains within the a sequence involved in the cleavage-packaging of herpes simplex virus DNA. *J Virol* 59(3), 605-18.
- Delius, H. and Clements, J.B. (1976) A partial denaturation map of herpes simplex virus type 1 DNA: evidence for inversions of the unique DNA regions. *J Gen Virol* 33(1), 125-33.
- DeLuca, N.A., McCarthy, A.M. and Schaffer, P.A. (1985) Isolation and characterization of deletion mutants of herpes simplex virus type 1 in the gene encoding immediate-early regulatory protein ICP4. *J Virol* 56(2), 558-70.
- DeLuca, N.A. and Schaffer, P.A. (1985) Activation of immediate-early, early, and late promoters by temperature-sensitive and wild-type forms of herpes simplex virus type 1 protein ICP4. *Mol Cell Biol* 5(8), 1997-2008.
- Di Pasquale, G. and Chiorini, J.A. (2003) PKA/PrKX activity is a modulator of AAV/adenovirus interaction. *Embo J* 22(7), 1716-24.
- Dolan, A., Jamieson, F.E., Cunningham, C., Barnett, B.C. and McGeoch, D.J. (1998) The genome sequence of herpes simplex virus type 2. *J Virol* 72(3), 2010-21.
- Dubielzig, R., King, J.A., Weger, S., Kern, A. and Kleinschmidt, J.A. (1999) Adeno-associated virus type 2 protein interactions: formation of pre-encapsidation complexes. *J Virol* 73(11), 8989-98.
- Ellison, K.S., Rice, S.A., Verity, R. and Smiley, J.R. (2000) Processing of alpha-globin and ICP0 mRNA in cells infected with herpes simplex virus type 1 ICP27 mutants. *J Virol* 74(16), 7307-19.
- Farson, D., Harding, T.C., Tao, L., Liu, J., Powell, S., Vimal, V., Yendluri, S., Koprivnikar, K., Ho, K., Twitty, C., Husak, P., Lin, A., Snyder, R.O. and Donahue, B.A. (2004) Development and characterization of a cell line for large-scale, serum-free production of recombinant adeno-associated viral vectors. *J Gene Med* 6(12), 1369-81.
- Frenkel, N., Locker, H., Batterson, W., Hayward, G.S. and Roizman, B. (1976) Anatomy of herpes simplex virus DNA. VI. Defective DNA originates from the S component. *J Virol* 20(2), 527-31.
- Geraghty, R.J., Krummenacher, C., Cohen, G.H., Eisenberg, R.J. and Spear, P.G. (1998) Entry of alphaherpesviruses mediated by poliovirus receptor-related protein 1 and poliovirus receptor. *Science* 280(5369), 1618-20.
- Giraud, C., Winocour, E. and Berns, K.I. (1994) Site-specific integration by adeno-associated virus is directed by a cellular DNA sequence. *Proc Natl Acad Sci U S A* 91(21), 10039-43.
- Girod, A., Wobus, C.E., Zadori, Z., Ried, M., Leike, K., Tijssen, P., Kleinschmidt, J.A. and Hallek, M. (2002) The VP1 capsid protein of adeno-associated virus type 2 is carrying a phospholipase A2 domain required for virus infectivity. *J Gen Virol* 83(Pt 5), 973-8.
- Godowski, P.J. and Knipe, D.M. (1986) Transcriptional control of herpesvirus gene expression: gene functions required for positive and negative regulation. *Proc Natl Acad Sci U S A* 83(2), 256-60.
- Gottlieb, J., Marcy, A.I., Coen, D.M. and Challberg, M.D. (1990) The herpes simplex virus type 1 UL42 gene product: a subunit of DNA polymerase that functions to increase processivity. *J Virol* 64(12), 5976-87.
- Green, M.R. and Roeder, R.G. (1980) Transcripts of the adeno-associated virus genome: mapping of the major RNAs. *J Virol* 36(1), 79-92.
- Grimm, D. and Kay, M.A. (2003) From virus evolution to vector revolution: use of naturally occurring serotypes of adeno-associated virus (AAV) as novel vectors for human gene therapy. *Curr Gene Ther* 3(4), 281-304.
- Grimm, D., Kern, A., Rittner, K. and Kleinschmidt, J.A. (1998) Novel tools for production and purification of recombinant adenoassociated virus vectors. *Hum Gene Ther* 9(18), 2745-60.
- Hann, L.E., Cook, W.J., Uprichard, S.L., Knipe, D.M. and Coen, D.M. (1998) The role of herpes simplex virus ICP27 in the regulation of UL24 gene expression by differential polyadenylation. *J Virol* 72(10), 7709-14.
- Hardwicke, M.A. and Sandri-Goldin, R.M. (1994) The herpes simplex virus regulatory protein ICP27 contributes to the decrease in cellular mRNA levels during infection. *J Virol* 68(8), 4797-810.
- Hardy, W.R. and Sandri-Goldin, R.M. (1994) Herpes simplex virus inhibits host cell splicing, and regulatory protein ICP27 is required for this effect. *J Virol* 68(12), 7790-9.

- Hauswirth, W.W. and Berns, K.I. (1977) Origin and termination of adeno-associated virus DNA replication. *Virology* 78(2), 488-99.
- Hauswirth, W.W. and Berns, K.I. (1979) Adeno-associated virus DNA replication: nonunit-length molecules. *Virology* 93(1), 57-68.
- Hayward, G.S., Jacob, R.J., Wadsworth, S.C. and Roizman, B. (1975) Anatomy of herpes simplex virus DNA: evidence for four populations of molecules that differ in the relative orientations of their long and short components. *Proc Natl Acad Sci U S A* 72(11), 4243-7.
- Heilbronn, R., Burkle, A., Stephan, S. and zur Hausen, H. (1990) The adeno-associated virus rep gene suppresses herpes simplex virus-induced DNA amplification. *J Virol* 64(6), 3012-8.
- Heilbronn, R., Engstler, M., Weger, S., Krahn, A., Schetter, C. and Boshart, M. (2003) ssDNA-dependent colocalization of adeno-associated virus Rep and herpes simplex virus ICP8 in nuclear replication domains. *Nucleic Acids Res* 31(21), 6206-13.
- Heine, J.W., Honess, R.W., Cassai, E. and Roizman, B. (1974) Proteins specified by herpes simplex virus. XII. The virion polypeptides of type 1 strains. *J Virol* 14(3), 640-51.
- Hermonat, P.L. (1989) The adeno-associated virus Rep78 gene inhibits cellular transformation induced by bovine papillomavirus. *Virology* 172(1), 253-61.
- Hermonat, P.L. (1992) Inhibition of bovine papillomavirus plasmid DNA replication by adeno-associated virus. *Virology* 189(1), 329-33.
- Hermonat, P.L. (1994) Adeno-associated virus inhibits human papillomavirus type 16: a viral interaction implicated in cervical cancer. *Cancer Res* 54(8), 2278-81.
- Hernandez, Y.J., Wang, J., Kearns, W.G., Loiler, S., Poirier, A. and Flotte, T.R. (1999) Latent adeno-associated virus infection elicits humoral but not cell-mediated immune responses in a nonhuman primate model. *J Virol* 73(10), 8549-58.
- Hirt, B. (1967) Selective extraction of polyoma DNA from infected mouse cell cultures. *J Mol Biol* 26(2), 365-9.
- Honess, R.W. and Roizman, B. (1974) Regulation of herpesvirus macromolecular synthesis. I. Cascade regulation of the synthesis of three groups of viral proteins. *J Virol* 14(1), 8-19.
- Hoque, M., Ishizu, K., Matsumoto, A., Han, S.I., Arisaka, F., Takayama, M., Suzuki, K., Kato, K., Kanda, T., Watanabe, H. and Handa, H. (1999) Nuclear transport of the major capsid protein is essential for adeno-associated virus capsid formation. *J Virol* 73(9), 7912-5.
- Hörer, M. (1992) Untersuchung der AAV-2 Rep-Genprodukte und deren Einfluß auf die HPV18 URR getriebene Genexpression, Universität Stuttgart.
- Im, D.S. and Muzyczka, N. (1990) The AAV origin binding protein Rep68 is an ATP-dependent site-specific endonuclease with DNA helicase activity. *Cell* 61(3), 447-57.
- Im, D.S. and Muzyczka, N. (1992) Partial purification of adeno-associated virus Rep78, Rep52, and Rep40 and their biochemical characterization. *J Virol* 66(2), 1119-28.
- Janik, J.E., Huston, M.M. and Rose, J.A. (1981) Locations of adenovirus genes required for the replication of adenovirus-associated virus. *Proc Natl Acad Sci U S A* 78(3), 1925-9.
- Jean, S., LeVan, K.M., Song, B., Levine, M. and Knipe, D.M. (2001) Herpes simplex virus 1 ICP27 is required for transcription of two viral late (gamma 2) genes in infected cells. *Virology* 283(2), 273-84.
- Johnson, D.C. and Spear, P.G. (1982) Monensin inhibits the processing of herpes simplex virus glycoproteins, their transport to the cell surface, and the egress of virions from infected cells. *J Virol* 43(3), 1102-12.
- Jooss, K., Yang, Y., Fisher, K.J. and Wilson, J.M. (1998) Transduction of dendritic cells by DNA viral vectors directs the immune response to transgene products in muscle fibers. *J Virol* 72(5), 4212-23.
- Kashiwakura, Y., Tamayose, K., Iwabuchi, K., Hirai, Y., Shimada, T., Matsumoto, K., Nakamura, T., Watanabe, M., Oshimi, K. and Daida, H. (2005) Hepatocyte growth factor receptor is a coreceptor for adeno-associated virus type 2 infection. *J Virol* 79(1), 609-14.
- Kay, M.A., Manno, C.S., Ragni, M.V., Larson, P.J., Couto, L.B., McClelland, A., Glader, B., Chew, A.J., Tai, S.J., Herzog, R.W., Arruda, V., Johnson, F., Scallan, C., Skarsgard, E., Flake, A.W. and High, K.A. (2000) Evidence for gene transfer and expression of factor IX in haemophilia B patients treated with an AAV vector. *Nat Genet* 24(3), 257-61.

- Kearns, W.G., Afione, S.A., Fulmer, S.B., Pang, M.C., Erikson, D., Egan, M., Landrum, M.J., Flotte, T.R. and Cutting, G.R. (1996) Recombinant adeno-associated virus (AAV-CFTR) vectors do not integrate in a site-specific fashion in an immortalized epithelial cell line. *Gene Ther* 3(9), 748-55.
- Khleif, S.N., Myers, T., Carter, B.J. and Trempe, J.P. (1991) Inhibition of cellular transformation by the adeno-associated virus rep gene. *Virology* 181(2), 738-41.
- King, J.A., Dubielzig, R., Grimm, D. and Kleinschmidt, J.A. (2001) DNA helicase-mediated packaging of adeno-associated virus type 2 genomes into preformed capsids. *Embo J* 20(12), 3282-91.
- Kirschstein, R.L., Smith, K.O. and Peters, E.A. (1968) Inhibition of adenovirus 12 oncogenicity by adeno-associated virus. *Proc Soc Exp Biol Med* 128(3), 670-3.
- Klein, R.L., Meyer, E.M., Peel, A.L., Zolotukhin, S., Meyers, C., Muzyczka, N. and King, M.A. (1998) Neuron-specific transduction in the rat septohippocampal or nigrostriatal pathway by recombinant adeno-associated virus vectors. *Exp Neurol* 150(2), 183-94.
- Kleinschmidt, J.A., Mohler, M., Weindler, F.W. and Heilbronn, R. (1995) Sequence elements of the adeno-associated virus rep gene required for suppression of herpes-simplex-virus-induced DNA amplification. *Virology* 206(1), 254-62.
- Kotin, R.M., Linden, R.M. and Berns, K.I. (1992) Characterization of a preferred site on human chromosome 19q for integration of adeno-associated virus DNA by non-homologous recombination. *Embo J* 11(13), 5071-8.
- Kotin, R.M., Siniscalco, M., Samulski, R.J., Zhu, X.D., Hunter, L., Laughlin, C.A., McLaughlin, S., Muzyczka, N., Rocchi, M. and Berns, K.I. (1990) Site-specific integration by adeno-associated virus. *Proc Natl Acad Sci U S A* 87(6), 2211-5.
- Kozak, M. (1983) Translation of insulin-related polypeptides from messenger RNAs with tandemly reiterated copies of the ribosome binding site. *Cell* 34(3), 971-8.
- Kozak, M. (1986) Point mutations define a sequence flanking the AUG initiator codon that modulates translation by eukaryotic ribosomes. *Cell* 44(2), 283-92.
- Kozak, M. (1987) An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs. *Nucleic Acids Res* 15(20), 8125-48.
- Kozak, M. (1991a) An analysis of vertebrate mRNA sequences: intimations of translational control. *J Cell Biol* 115(4), 887-903.
- Kozak, M. (1991b) Structural features in eukaryotic mRNAs that modulate the initiation of translation. *J Biol Chem* 266(30), 19867-70.
- Kozak, M. (1995) Adherence to the first-AUG rule when a second AUG codon follows closely upon the first. *Proc Natl Acad Sci U S A* 92(15), 7134.
- Kozak, M. (1997) Recognition of AUG and alternative initiator codons is augmented by G in position +4 but is not generally affected by the nucleotides in positions +5 and +6. *Embo J* 16(9), 2482-92.
- Kozak, M. (1999) Initiation of translation in prokaryotes and eukaryotes. *Gene* 234(2), 187-208.
- Kozak, M. (2001) Constraints on reinitiation of translation in mammals. *Nucleic Acids Res* 29(24), 5226-32.
- Kuddus, R., Gu, B. and DeLuca, N.A. (1995) Relationship between TATA-binding protein and herpes simplex virus type 1 ICP4 DNA-binding sites in complex formation and repression of transcription. *J Virol* 69(9), 5568-75.
- Kyostio, S.R., Wonderling, R.S. and Owens, R.A. (1995) Negative regulation of the adeno-associated virus (AAV) P5 promoter involves both the P5 rep binding site and the consensus ATP-binding motif of the AAV Rep68 protein. *J Virol* 69(11), 6787-96.
- Labow, M.A. and Berns, K.I. (1988) The adeno-associated virus rep gene inhibits replication of an adeno-associated virus/simian virus 40 hybrid genome in cos-7 cells. *J Virol* 62(5), 1705-12.
- Labow, M.A., Graf, L.H., Jr. and Berns, K.I. (1987) Adeno-associated virus gene expression inhibits cellular transformation by heterologous genes. *Mol Cell Biol* 7(4), 1320-5.
- Labow, M.A., Hermonat, P.L. and Berns, K.I. (1986) Positive and negative autoregulation of the adeno-associated virus type 2 genome. *J Virol* 60(1), 251-8.
- Lamartina, S., Roscilli, G., Rinaudo, D., Delmastro, P. and Toniatti, C. (1998) Lipofection of purified adeno-associated virus Rep68 protein: toward a chromosome-targeting nonviral particle. *J Virol* 72(9), 7653-8.

- Laughlin, C.A., Cardellicchio, C.B. and Coon, H.C. (1986) Latent infection of KB cells with adeno-associated virus type 2. *J Virol* 60(2), 515-24.
- Laughlin, C.A., Myers, M.W., Risin, D.L. and Carter, B.J. (1979) Defective-interfering particles of the human parvovirus adeno-associated virus. *Virology* 94(1), 162-74.
- Li, J., Samulski, R.J. and Xiao, X. (1997) Role for highly regulated rep gene expression in adeno-associated virus vector production. *J Virol* 71(7), 5236-43.
- Lindberg, A. and Kreivi, J.P. (2002) Splicing inhibition at the level of spliceosome assembly in the presence of herpes simplex virus protein ICP27. *Virology* 294(1), 189-98.
- Linden, R.M., Winocour, E. and Berns, K.I. (1996) The recombination signals for adeno-associated virus site-specific integration. *Proc Natl Acad Sci U S A* 93(15), 7966-72.
- Liu, X., Vougaropoulou, F., Chen, R., Johnson, P.R. and Clark, K.R. (2000) Selective Rep-Cap gene amplification as a mechanism for high-titer recombinant AAV production from stable cell lines. *Mol Ther* 2(4), 394-403.
- Locker, H., Frenkel, N. and Halliburton, I. (1982) Structure and expression of class II defective herpes simplex virus genomes encoding infected cell polypeptide number 8. *J Virol* 43(2), 574-93.
- Lusby, E., Fife, K.H. and Berns, K.I. (1980) Nucleotide sequence of the inverted terminal repetition in adeno-associated virus DNA. *J Virol* 34(2), 402-9.
- Lusby, E.W. and Berns, K.I. (1982) Mapping of the 5' termini of two adeno-associated virus 2 RNAs in the left half of the genome. *J Virol* 41(2), 518-26.
- Malik, A.K. and Weller, S.K. (1996) Use of transdominant mutants of the origin-binding protein (UL9) of herpes simplex virus type 1 to define functional domains. *J Virol* 70(11), 7859-66.
- Manno, C.S., Chew, A.J., Hutchison, S., Larson, P.J., Herzog, R.W., Arruda, V.R., Tai, S.J., Ragni, M.V., Thompson, A., Ozelo, M., Couto, L.B., Leonard, D.G., Johnson, F.A., McClelland, A., Scallan, C., Skarsgard, E., Flake, A.W., Kay, M.A., High, K.A. and Glader, B. (2003) AAV-mediated factor IX gene transfer to skeletal muscle in patients with severe hemophilia B. *Blood* 101(8), 2963-72.
- Mayor, H.D., Houlditch, G.S. and Mumford, D.M. (1973) Influence of adeno-associated satellite virus on adenovirus-induced tumours in hamsters. *Nat New Biol* 241(106), 44-6.
- McCarthy, A.M., McMahan, L. and Schaffer, P.A. (1989) Herpes simplex virus type 1 ICP27 deletion mutants exhibit altered patterns of transcription and are DNA deficient. *J Virol* 63(1), 18-27.
- McCarty, D.M., Christensen, M. and Muzyczka, N. (1991) Sequences required for coordinate induction of adeno-associated virus p19 and p40 promoters by Rep protein. *J Virol* 65(6), 2936-45.
- McCarty, D.M., Monahan, P.E. and Samulski, R.J. (2001) Self-complementary recombinant adeno-associated virus (scAAV) vectors promote efficient transduction independently of DNA synthesis. *Gene Ther* 8(16), 1248-54.
- McCarty, D.M., Pereira, D.J., Zolotukhin, I., Zhou, X., Ryan, J.H. and Muzyczka, N. (1994a) Identification of linear DNA sequences that specifically bind the adeno-associated virus Rep protein. *J Virol* 68(8), 4988-97.
- McCarty, D.M., Ryan, J.H., Zolotukhin, S., Zhou, X. and Muzyczka, N. (1994b) Interaction of the adeno-associated virus Rep protein with a sequence within the A palindrome of the viral terminal repeat. *J Virol* 68(8), 4998-5006.
- McGregor, F., Phelan, A., Dunlop, J. and Clements, J.B. (1996) Regulation of herpes simplex virus poly (A) site usage and the action of immediate-early protein IE63 in the early-late switch. *J Virol* 70(3), 1931-40.
- McLauchlan, J., Phelan, A., Loney, C., Sandri-Goldin, R.M. and Clements, J.B. (1992) Herpes simplex virus IE63 acts at the posttranscriptional level to stimulate viral mRNA 3' processing. *J Virol* 66(12), 6939-45.
- McLaughlin, S.K., Collis, P., Hermonat, P.L. and Muzyczka, N. (1988) Adeno-associated virus general transduction vectors: analysis of proviral structures. *J Virol* 62(6), 1963-73.
- Mendelson, E., Trempe, J.P. and Carter, B.J. (1986) Identification of the trans-acting Rep proteins of adeno-associated virus by antibodies to a synthetic oligopeptide. *J Virol* 60(3), 823-32.

- Mizukami, H., Okada, T., Ogasawara, Y., Matsushita, T., Urabe, M., Kume, A. and Ozawa, K. (2004) Separate control of Rep and Cap expression using mutant and wild-type loxP sequences and improved packaging system for adeno-associated virus vector production. *Mol Biotechnol* 27(1), 7-14.
- Modrow, S. and Falke, D. (1997) *Molekulare Virologie*. Spektrum Akademischer Verlag.
- Monahan, P.E. and Samulski, R.J. (2000a) AAV vectors: is clinical success on the horizon? *Gene Ther* 7(1), 24-30.
- Monahan, P.E. and Samulski, R.J. (2000b) Adeno-associated virus vectors for gene therapy: more pros than cons? *Mol Med Today* 6(11), 433-40.
- Montgomery, R.I., Warner, M.S., Lum, B.J. and Spear, P.G. (1996) Herpes simplex virus-1 entry into cells mediated by a novel member of the TNF/NGF receptor family. *Cell* 87(3), 427-36.
- Morgan, C., Rose, H.M. and Mednis, B. (1968) Electron microscopy of herpes simplex virus. I. Entry. *J Virol* 2(5), 507-16.
- Muralidhar, S., Becerra, S.P. and Rose, J.A. (1994) Site-directed mutagenesis of adeno-associated virus type 2 structural protein initiation codons: effects on regulation of synthesis and biological activity. *J Virol* 68(1), 170-6.
- Muyrers, J.P., Zhang, Y. and Stewart, A.F. (2001) Techniques: Recombinogenic engineering--new options for cloning and manipulating DNA. *Trends Biochem Sci* 26(5), 325-31.
- Muzyczka, N. (1992) Use of adeno-associated virus as a general transduction vector for mammalian cells. *Curr Top Microbiol Immunol* 158, 97-129.
- Muzyczka, N. and Berns, K.I. (2001) Parvoviridae: The Viruses and Their Replication. In: D.M. Knipe and P.M. Howley (Eds), *Fields Virology*, pp. 2327-2359.
- Nakai, H., Storm, T.A. and Kay, M.A. (2000) Increasing the size of rAAV-mediated expression cassettes in vivo by intermolecular joining of two complementary vectors. *Nat Biotechnol* 18(5), 527-32.
- Nakamura, S., Nakamura, R., Shibata, K., Kobayashi, M., Sahara, N., Shigeno, K., Shinjo, K., Naito, K., Ohnishi, K., Kasahara, N. and Iwaki, Y. (2004) Development of packaging cell lines for generation of adeno-associated virus vectors by lentiviral gene transfer of trans-complementary components. *Eur J Haematol* 73(4), 285-94.
- Newcomb, W.W. and Brown, J.C. (1994) Induced extrusion of DNA from the capsid of herpes simplex virus type 1. *J Virol* 68(1), 433-40.
- O'Hare, P. and Hayward, G.S. (1985) Three trans-acting regulatory proteins of herpes simplex virus modulate immediate-early gene expression in a pathway involving positive and negative feedback regulation. *J Virol* 56(3), 723-33.
- Okada, T., Mizukami, H., Urabe, M., Nomoto, T., Matsushita, T., Hanazono, Y., Kume, A., Tobita, K. and Ozawa, K. (2001) Development and characterization of an antisense-mediated prepackaging cell line for adeno-associated virus vector production. *Biochem Biophys Res Commun* 288(1), 62-8.
- Olivo, P.D., Nelson, N.J. and Challberg, M.D. (1988) Herpes simplex virus DNA replication: the UL9 gene encodes an origin-binding protein. *Proc Natl Acad Sci U S A* 85(15), 5414-8.
- Ostrove, J.M., Duckworth, D.H. and Berns, K.I. (1981) Inhibition of adenovirus-transformed cell oncogenicity by adeno-associated virus. *Virology* 113(2), 521-33.
- Owens, R.A., Weitzman, M.D., Kyostio, S.R. and Carter, B.J. (1993) Identification of a DNA-binding domain in the amino terminus of adeno-associated virus Rep proteins. *J Virol* 67(2), 997-1005.
- Pereira, D.J. and Muzyczka, N. (1997a) The adeno-associated virus type 2 p40 promoter requires a proximal Sp1 interaction and a p19 CArG-like element to facilitate Rep transactivation. *J Virol* 71(6), 4300-9.
- Pereira, D.J. and Muzyczka, N. (1997b) The cellular transcription factor SP1 and an unknown cellular protein are required to mediate Rep protein activation of the adeno-associated virus p19 promoter. *J Virol* 71(3), 1747-56.
- Phelan, A., Dunlop, J. and Clements, J.B. (1996) Herpes simplex virus type 1 protein IE63 affects the nuclear export of virus intron-containing transcripts. *J Virol* 70(8), 5255-65.
- Portis, J.L., Spangrude, G.J. and McAtee, F.J. (1994) Identification of a sequence in the unique 5' open reading frame of the gene encoding glycosylated Gag which influences the incubation period of neurodegenerative disease induced by a murine retrovirus. *J Virol* 68(6), 3879-87.

- Post, L.E., Mackem, S. and Roizman, B. (1981) Regulation of alpha genes of herpes simplex virus: expression of chimeric genes produced by fusion of thymidine kinase with alpha gene promoters. *Cell* 24(2), 555-65.
- Preston, C.M. (1979) Control of herpes simplex virus type 1 mRNA synthesis in cells infected with wild-type virus or the temperature-sensitive mutant tsK. *J Virol* 29(1), 275-84.
- Qiao, C., Li, J., Skold, A., Zhang, X. and Xiao, X. (2002a) Feasibility of generating adeno-associated virus packaging cell lines containing inducible adenovirus helper genes. *J Virol* 76(4), 1904-13.
- Qiao, C., Wang, B., Zhu, X., Li, J. and Xiao, X. (2002b) A novel gene expression control system and its use in stable, high-titer 293 cell-based adeno-associated virus packaging cell lines. *J Virol* 76(24), 13015-27.
- Qing, K., Mah, C., Hansen, J., Zhou, S., Dwarki, V. and Srivastava, A. (1999) Human fibroblast growth factor receptor 1 is a co-receptor for infection by adeno-associated virus 2. *Nat Med* 5(1), 71-7.
- Rabinowitz, J.E., Xiao, W. and Samulski, R.J. (1999) Insertional mutagenesis of AAV2 capsid and the production of recombinant virus. *Virology* 265(2), 274-85.
- Rice, S.A. and Knipe, D.M. (1988) Gene-specific transactivation by herpes simplex virus type 1 alpha protein ICP27. *J Virol* 62(10), 3814-23.
- Rice, S.A. and Knipe, D.M. (1990) Genetic evidence for two distinct transactivation functions of the herpes simplex virus alpha protein ICP27. *J Virol* 64(4), 1704-15.
- Richardson, W.D. and Westphal, H. (1981) A cascade of adenovirus early functions is required for expression of adeno-associated virus. *Cell* 27(1 Pt 2), 133-41.
- Rittner, K., Heilbronn, R., Kleinschmidt, J.A., Oelze, I. and Sczakiel, G. (1991) Replication of the human immunodeficiency virus type 1 is inhibited by the adeno-associated virus rep gene. *Biochem Soc Trans* 19(4), 438S.
- Rittner, K., Heilbronn, R., Kleinschmidt, J.A. and Sczakiel, G. (1992) Adeno-associated virus type 2-mediated inhibition of human immunodeficiency virus type 1 (HIV-1) replication: involvement of p78rep/p68rep and the HIV-1 long terminal repeat. *J Gen Virol* 73 (Pt 11), 2977-81.
- Rixon, F.J. and McLauchlan, J. (1990) Insertion of DNA sequences at a unique restriction enzyme site engineered for vector purposes into the genome of herpes simplex virus type 1. *J Gen Virol* 71 (Pt 12), 2931-9.
- Rixon, F.J. and McLauchlan, J. (1993) Herpes simplex virus vectors. In: A.J. Davidson and R.M. Elliot (Eds), *Molecular Virology: A Practical Approach*, pp. 285-307. Oxford University Press.
- Roizman, B. and Knipe, D.M. (2001) Herpes Simplex Viruses and Their Replication. In: D.M. Knipe and P.M. Howley (Eds), *Fields Virology*, pp. 2399-2459.
- Rose, J.A., Berns, K.I., Hoggan, M.D. and Koczot, F.J. (1969) Evidence for a single-stranded adenovirus-associated virus genome: formation of a DNA density hybrid on release of viral DNA. *Proc Natl Acad Sci U S A* 64(3), 863-9.
- Rose, J.A., Maizel, J.V., Jr., Inman, J.K. and Shatkin, A.J. (1971) Structural proteins of adenovirus-associated viruses. *J Virol* 8(5), 766-70.
- Ruffing, M., Heid, H. and Kleinschmidt, J.A. (1994) Mutations in the carboxy terminus of adeno-associated virus 2 capsid proteins affect viral infectivity: lack of an RGD integrin-binding motif. *J Gen Virol* 75 (Pt 12), 3385-92.
- Ruffing, M., Zentgraf, H. and Kleinschmidt, J.A. (1992) Assembly of viruslike particles by recombinant structural proteins of adeno-associated virus type 2 in insect cells. *J Virol* 66(12), 6922-30.
- Ryan, J.H., Zolotukhin, S. and Muzyczka, N. (1996) Sequence requirements for binding of Rep68 to the adeno-associated virus terminal repeats. *J Virol* 70(3), 1542-53.
- Sacks, W.R., Greene, C.C., Aschman, D.P. and Schaffer, P.A. (1985) Herpes simplex virus type 1 ICP27 is an essential regulatory protein. *J Virol* 55(3), 796-805.
- Saiki, R.K., Gelfand, D.H., Stoffel, S., Scharf, S.J., Higuchi, R., Horn, G.T., Mullis, K.B. and Erlich, H.A. (1988) Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239(4839), 487-91.
- Saiki, R.K., Scharf, S., Falloona, F., Mullis, K.B., Horn, G.T., Erlich, H.A. and Arnheim, N. (1985) Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* 230(4732), 1350-4.

- Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989) Molecular Cloning, A Laboratory Manual. Cold Spring Harbor Laboratory Press.
- Samulski, R.J., Chang, L.S. and Shenk, T. (1989) Helper-free stocks of recombinant adeno-associated viruses: normal integration does not require viral gene expression. *J Virol* 63(9), 3822-8.
- Samulski, R.J., Srivastava, A., Berns, K.I. and Muzyczka, N. (1983) Rescue of adeno-associated virus from recombinant plasmids: gene correction within the terminal repeats of AAV. *Cell* 33(1), 135-43.
- Samulski, R.J., Zhu, X., Xiao, X., Brook, J.D., Housman, D.E., Epstein, N. and Hunter, L.A. (1991) Targeted integration of adeno-associated virus (AAV) into human chromosome 19. *Embo J* 10(12), 3941-50.
- Sandri-Goldin, R.M. (2004) Viral regulation of mRNA export. *J Virol* 78(9), 4389-96.
- Sandri-Goldin, R.M. and Mendoza, G.E. (1992) A herpesvirus regulatory protein appears to act post-transcriptionally by affecting mRNA processing. *Genes Dev* 6(5), 848-63.
- Schlehofer, J.R., Ehrbar, M. and zur Hausen, H. (1986) Vaccinia virus, herpes simplex virus, and carcinogens induce DNA amplification in a human cell line and support replication of a helpervirus dependent parvovirus. *Virology* 152(1), 110-7.
- Schlehofer, J.R., Heilbronn, R., Georg-Fries, B. and zur Hausen, H. (1983) Inhibition of initiator-induced SV40 gene amplification in SV40-transformed Chinese hamster cells by infection with a defective parvovirus. *Int J Cancer* 32(5), 591-5.
- Senapathy, P., Tratschin, J.D. and Carter, B.J. (1984) Replication of adeno-associated virus DNA. Complementation of naturally occurring rep- mutants by a wild-type genome or an ori- mutant and correction of terminal palindrome deletions. *J Mol Biol* 179(1), 1-20.
- Shelling, A.N. and Smith, M.G. (1994) Targeted integration of transfected and infected adeno-associated virus vectors containing the neomycin resistance gene. *Gene Ther* 1(3), 165-9.
- Shiau, A.L., Liu, P.S. and Wu, C.L. (2005) Novel strategy for generation and titration of recombinant adeno-associated virus vectors. *J Virol* 79(1), 193-201.
- Shieh, M.T., WuDunn, D., Montgomery, R.I., Esko, J.D. and Spear, P.G. (1992) Cell surface receptors for herpes simplex virus are heparan sulfate proteoglycans. *J Cell Biol* 116(5), 1273-81.
- Siminoff, P. and Menefee, M.G. (1966) Normal and 5-bromodeoxyuridine-inhibited development of herpes simplex virus. An electron microscope study. *Exp Cell Res* 44(2), 241-55.
- Smith, I.L., Hardwicke, M.A. and Sandri-Goldin, R.M. (1992) Evidence that the herpes simplex virus immediate early protein ICP27 acts post-transcriptionally during infection to regulate gene expression. *Virology* 186(1), 74-86.
- Smith, R.H. and Kotin, R.M. (1998) The Rep52 gene product of adeno-associated virus is a DNA helicase with 3'-to-5' polarity. *J Virol* 72(6), 4874-81.
- Smith, R.H., Spano, A.J. and Kotin, R.M. (1997) The Rep78 gene product of adeno-associated virus (AAV) self-associates to form a hexameric complex in the presence of AAV ori sequences. *J Virol* 71(6), 4461-71.
- Snyder, R.O., Im, D.S. and Muzyczka, N. (1990a) Evidence for covalent attachment of the adeno-associated virus (AAV) rep protein to the ends of the AAV genome. *J Virol* 64(12), 6204-13.
- Snyder, R.O., Samulski, R.J. and Muzyczka, N. (1990b) In vitro resolution of covalently joined AAV chromosome ends. *Cell* 60(1), 105-13.
- Sodeik, B., Ebersold, M.W. and Helenius, A. (1997) Microtubule-mediated transport of incoming herpes simplex virus 1 capsids to the nucleus. *J Cell Biol* 136(5), 1007-21.
- Southern, E.M. (1975) Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* 98(3), 503-17.
- Spear, P.G. and Roizman, B. (1972) Proteins specified by herpes simplex virus. V. Purification and structural proteins of the herpesvirion. *J Virol* 9(1), 143-59.
- Srivastava, A., Lusby, E.W. and Berns, K.I. (1983) Nucleotide sequence and organization of the adeno-associated virus 2 genome. *J Virol* 45(2), 555-64.
- Stackpole, C.W. (1969) Herpes-type virus of the frog renal adenocarcinoma. I. Virus development in tumor transplants maintained at low temperature. *J Virol* 4(1), 75-93.
- Stow, N.D., Hammarsten, O., Arbuckle, M.I. and Elias, P. (1993) Inhibition of herpes simplex virus type 1 DNA replication by mutant forms of the origin-binding protein. *Virology* 196(2), 413-8.

- Straus, S.E., Sebring, E.D. and Rose, J.A. (1976) Concatemers of alternating plus and minus strands are intermediates in adenovirus-associated virus DNA synthesis. *Proc Natl Acad Sci U S A* 73(3), 742-6.
- Summerford, C., Bartlett, J.S. and Samulski, R.J. (1999) AlphaVbeta5 integrin: a co-receptor for adeno-associated virus type 2 infection. *Nat Med* 5(1), 78-82.
- Summerford, C. and Samulski, R.J. (1998) Membrane-associated heparan sulfate proteoglycan is a receptor for adeno-associated virus type 2 virions. *J Virol* 72(2), 1438-45.
- Sun, L., Li, J. and Xiao, X. (2000) Overcoming adeno-associated virus vector size limitation through viral DNA heterodimerization. *Nat Med* 6(5), 599-602.
- Surosky, R.T., Urabe, M., Godwin, S.G., McQuiston, S.A., Kurtzman, G.J., Ozawa, K. and Natsoulis, G. (1997) Adeno-associated virus Rep proteins target DNA sequences to a unique locus in the human genome. *J Virol* 71(10), 7951-9.
- Tattersall, P. and Ward, D.C. (1976) Rolling hairpin model for replication of parvovirus and linear chromosomal DNA. *Nature* 263(5573), 106-9.
- Tenenbaum, L., Lehtonen, E. and Monahan, P.E. (2003) Evaluation of risks related to the use of adeno-associated virus-based vectors. *Curr Gene Ther* 3(6), 545-65.
- Towbin, H., Staehelin, T. and Gordon, J. (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci U S A* 76(9), 4350-4.
- Trempe, J.P. and Carter, B.J. (1988) Regulation of adeno-associated virus gene expression in 293 cells: control of mRNA abundance and translation. *J Virol* 62(1), 68-74.
- Uprichard, S.L. and Knipe, D.M. (1996) Herpes simplex ICP27 mutant viruses exhibit reduced expression of specific DNA replication genes. *J Virol* 70(3), 1969-80.
- Urabe, M., Ding, C. and Kotin, R.M. (2002) Insect cells as a factory to produce adeno-associated virus type 2 vectors. *Hum Gene Ther* 13(16), 1935-43.
- van Genderen, I.L., Brandimarti, R., Torrisi, M.R., Campadelli, G. and van Meer, G. (1994) The phospholipid composition of extracellular herpes simplex virions differs from that of host cell nuclei. *Virology* 200(2), 831-6.
- Varmuza, S.L. and Smiley, J.R. (1985) Signals for site-specific cleavage of HSV DNA: maturation involves two separate cleavage events at sites distal to the recognition sequences. *Cell* 41(3), 793-802.
- Vincent, K.A., Piraino, S.T. and Wadsworth, S.C. (1997) Analysis of recombinant adeno-associated virus packaging and requirements for rep and cap gene products. *J Virol* 71(3), 1897-905.
- Vincent-Lacaze, N., Snyder, R.O., Gluzman, R., Bohl, D., Lagarde, C. and Danos, O. (1999) Structure of adeno-associated virus vector DNA following transduction of the skeletal muscle. *J Virol* 73(3), 1949-55.
- Vlazny, D.A., Kwong, A. and Frenkel, N. (1982) Site-specific cleavage/packaging of herpes simplex virus DNA and the selective maturation of nucleocapsids containing full-length viral DNA. *Proc Natl Acad Sci U S A* 79(5), 1423-7.
- Wadsworth, S., Jacob, R.J. and Roizman, B. (1975) Anatomy of herpes simplex virus DNA. II. Size, composition, and arrangement of inverted terminal repetitions. *J Virol* 15(6), 1487-97.
- Wang, L., Muramatsu, S., Lu, Y., Ikeguchi, K., Fujimoto, K., Okada, T., Mizukami, H., Hanazono, Y., Kume, A., Urano, F., Ichinose, H., Nagatsu, T., Nakano, I. and Ozawa, K. (2002) Delayed delivery of AAV-GDNF prevents nigral neurodegeneration and promotes functional recovery in a rat model of Parkinson's disease. *Gene Ther* 9(6), 381-9.
- Warrington, K.H., Jr., Gorbatyuk, O.S., Harrison, J.K., Opie, S.R., Zolotukhin, S. and Muzyczka, N. (2004) Adeno-associated virus type 2 VP2 capsid protein is nonessential and can tolerate large peptide insertions at its N terminus. *J Virol* 78(12), 6595-609.
- Weger, S., Wistuba, A., Grimm, D. and Kleinschmidt, J.A. (1997) Control of adeno-associated virus type 2 cap gene expression: relative influence of helper virus, terminal repeats, and Rep proteins. *J Virol* 71(11), 8437-47.
- Weindler, F.W. and Heilbronn, R. (1991) A subset of herpes simplex virus replication genes provides helper functions for productive adeno-associated virus replication. *J Virol* 65(5), 2476-83.
- Weitzman, M.D., Kyostio, S.R., Kotin, R.M. and Owens, R.A. (1994) Adeno-associated virus (AAV) Rep proteins mediate complex formation between AAV DNA and its integration site in human DNA. *Proc Natl Acad Sci U S A* 91(13), 5808-12.

- Weller, S.K., Lee, K.J., Sabourin, D.J. and Schaffer, P.A. (1983) Genetic analysis of temperature-sensitive mutants which define the gene for the major herpes simplex virus type 1 DNA-binding protein. *J Virol* 45(1), 354-66.
- Wonderling, R.S., Kyostio, S.R. and Owens, R.A. (1995) A maltose-binding protein/adeno-associated virus Rep68 fusion protein has DNA-RNA helicase and ATPase activities. *J Virol* 69(6), 3542-8.
- Wu, P., Xiao, W., Conlon, T., Hughes, J., Agbandje-McKenna, M., Ferkol, T., Flotte, T. and Muzyczka, N. (2000) Mutational analysis of the adeno-associated virus type 2 (AAV2) capsid gene and construction of AAV2 vectors with altered tropism. *J Virol* 74(18), 8635-47.
- WuDunn, D. and Spear, P.G. (1989) Initial interaction of herpes simplex virus with cells is binding to heparan sulfate. *J Virol* 63(1), 52-8.
- Wustner, J.T., Arnold, S., Lock, M., Richardson, J.C., Himes, V.B., Kurtzman, G. and Peluso, R.W. (2002) Production of recombinant adeno-associated type 5 (rAAV5) vectors using recombinant herpes simplex viruses containing rep and cap. *Mol Ther* 6(4), 510-8.
- Xiao, X., Li, J. and Samulski, R.J. (1998) Production of high-titer recombinant adeno-associated virus vectors in the absence of helper adenovirus. *J Virol* 72(3), 2224-32.
- Xie, Q., Bu, W., Bhatia, S., Hare, J., Somasundaram, T., Azzi, A. and Chapman, M.S. (2002) The atomic structure of adeno-associated virus (AAV-2), a vector for human gene therapy. *Proc Natl Acad Sci U S A* 99(16), 10405-10.
- Yakobson, B., Hryntko, T.A., Peak, M.J. and Winocour, E. (1989) Replication of adeno-associated virus in cells irradiated with UV light at 254 nm. *J Virol* 63(3), 1023-30.
- Yakobson, B., Koch, T. and Winocour, E. (1987) Replication of adeno-associated virus in synchronized cells without the addition of a helper virus. *J Virol* 61(4), 972-81.
- Yalkinoglu, A.O., Heilbronn, R., Burkle, A., Schlehofer, J.R. and zur Hausen, H. (1988) DNA amplification of adeno-associated virus as a response to cellular genotoxic stress. *Cancer Res* 48(11), 3123-9.
- Yan, Z., Zhang, Y., Duan, D. and Engelhardt, J.F. (2000) Trans-splicing vectors expand the utility of adeno-associated virus for gene therapy. *Proc Natl Acad Sci U S A* 97(12), 6716-21.
- Yang, C.C., Xiao, X., Zhu, X., Ansardi, D.C., Epstein, N.D., Frey, M.R., Matera, A.G. and Samulski, R.J. (1997) Cellular recombination pathways and viral terminal repeat hairpin structures are sufficient for adeno-associated virus integration in vivo and in vitro. *J Virol* 71(12), 9231-47.
- Yang, Q., Chen, F., Ross, J. and Trempe, J.P. (1995) Inhibition of cellular and SV40 DNA replication by the adeno-associated virus Rep proteins. *Virology* 207(1), 246-50.
- Yang, Q., Chen, F. and Trempe, J.P. (1994) Characterization of cell lines that inducibly express the adeno-associated virus Rep proteins. *J Virol* 68(8), 4847-56.
- Yang, Q. and Trempe, J.P. (1993) Analysis of the terminal repeat binding abilities of mutant adeno-associated virus replication proteins. *J Virol* 67(7), 4442-7.
- Zadori, Z., Szelei, J., Lacoste, M.C., Li, Y., Gariepy, S., Raymond, P., Allaire, M., Nabi, I.R. and Tijssen, P. (2001) A viral phospholipase A2 is required for parvovirus infectivity. *Dev Cell* 1(2), 291-302.
- Zhang, H.G., Wang, Y.M., Xie, J.F., Liang, X., Hsu, H.C., Zhang, X., Douglas, J., Curiel, D.T. and Mountz, J.D. (2001) Recombinant adenovirus expressing adeno-associated virus cap and rep proteins supports production of high-titer recombinant adeno-associated virus. *Gene Ther* 8(9), 704-12.