

## 7 REFERENCES

- Aebi U., Cohn J., Buhle L. and Gerace, L. (1986)  
The nuclear lamina is a meshwork of intermediate-type filaments  
*Nature* **323**:560-564
- Albagli O., Dhordain P., Deweindt C., Lecocq G. and Leprince D. (1995)  
The BTB/POZ domain : a new protein-protein interaction motif common to DNA- and actin-binding proteins.  
*Cell Growth Differ* **6**:1193-1198
- Altschul S.F., Madden T.L., Schäffer A.A., Zhang J., Zhang Z., Miller W. and Lipman D.J. (1997)  
Gapped BLAST and PSI-BLAST: a new generation of protein database search programs  
*Nucleic Acid Res* **25**:3389-3402
- Apel E.D., Lewis R.M., Grady R.M. and Sanes J.R. (2000)  
Syne-1, a dystrophin- and Klarsicht-related protein associated with synaptic nuclei at the neuromuscular junction.  
*J Biol Chem* **275**:31986-31995
- Ashery-Padan R., Ulitzur N., Arbel A., Goldberg M., Weiss A.M., Maus N., Fisher P.A. and Gruenbaum Y. (1997)  
Localization and posttranslational modifications of otefin, a protein required for vesicle attachment to chromatin, during *Drosophila melanogaster* development.  
*Mol Cell Biol* **17**:4114-4123
- Beaudouin J., Gerlich D., Daigle N., Eils R. and Ellenberg J. (2002)  
Nuclear envelope breakdown proceeds by microtubule-induced tearing of the lamina.  
*Cell* **108**:83-96
- Bell A.C., West A.G. and Felsenfeld G. (2001)  
Insulators and boundaries: versatile regulatory elements in the eukaryotic genome.  
*Science* **291**:447-450
- Belmont A.S., Zhai Y. and Thelenius A. (1993)  
Lamin B distribution and association with peripheral chromatin revealed by optical sectioning and electron microscopy tomography.  
*J Cell Biol* **123**:1671-1685
- Benavente R. and Krohne G. (1986)  
Involvement of nuclear lamins in postmitotic reorganization of chromatin as demonstrated by microinjection of lamin antibodies.  
*J Cell Biol* **103**:184718-184754

Bione, S., Maestrini, E., Rivella, S., Mancini, M., Regis, S., Romeo, G. and Toniolo, D. (1994)  
Identification of a novel X-linked gene responsible for Emery-Dreifuss muscular dystrophy.  
*Nature Genet* **8**:323-327

Bodoor K., Shaikh S., Salina D., Raharjo W.H., Bastos R., Lohka M. and Burke B. (1999)  
Sequential recruitment of NPC proteins to the nuclear periphery at the end of mitosis.  
*J Cell Sci* **112**:2253-2256

Bonne G., Di Barletta M.R., Varnous S., Becane H.M., Hammouda E.H., Merlini L., Muntoni F., Greenberg C.R., Gary F., Urtizberea J.A., Duboc D., Fardeau M., Toniolo D. and Schwartz K. (1999)  
Mutations in the gene encoding lamin A/C cause autosomal dominant Emery-Dreifuss muscular dystrophy.  
*Nat Genet* **21**:285-288

Borden K.L. (2000)  
RING domains: master builders of molecular scaffolds?  
*J Mol Biol* **295**:1103-12

Borden K.L. and Freemont P.S. (1996)  
The RING finger domain: a recent example of a sequence-structure family.  
*Curr Opin Struct Biol* **6**:395-401

Bradford, M.M. (1976)  
A rapid and sensitive method for the quantitation of microgramm quantities of protein utilizing the principle of protein-dye-binding.  
*Anal Biochem* **72**: 248-254

Broers J.L., Machiels B.M., van Eys G.J., Kuijpers H.J., Manders E.M., van Driel R. and Ramaekers F.C. (1999)  
Dynamics of the nuclear lamina as monitored by GFP-tagged A-type lamins.  
*J Cell Sci* **112**:3463-3475

Buendia B., Courvalin J.C. and Collas P. (2001)  
Dynamics of the nuclear envelope at mitosis and during apoptosis.  
*Cell Mol Life Sci* **58**:1781-1789

Busson S., Dujardin D., Moreau A., Dompierre J. and De Mey J.R. (1998)  
Dynein and dynactin are localized to astral microtubules and at cortical sites in mitotic epithelial cells.  
*Curr Biol* **8**:541-544

Carmo-Fonseca M. (2002 )  
The contribution of nuclear compartmentalization to gene regulation.  
*Cell* **108**:513-521

Carmo-Fonseca M., Mendes-Soares L. and Campos I. (2000)  
To be or not to be in the nucleolus.  
*Nat Cell Biol* **2**:E107-112

Chaudhary N. and Courvalin J.C. (1993)  
Stepwise reassembly of the nuclear envelope at the end of mitosis.  
*J Cell Biol* **122**:295-306

Chu A., Rassadi R. und Stochaj U. (1998)  
Velcro in the nuclear envelope: LBR and LAPs.  
*FEBS Lett* **441**:165-169

Clements L., Manilal S., Love D.R. and Morris G.E. (2000)  
Direct interaction between emerin and lamin A.  
*Biochem Biophys Res Commun* **267**:709-714

Collas I. and Courvalin J.C. (2000)  
Sorting nuclear membrane proteins at mitosis.  
*Trends Cell Biol* **10**:5-8

Collas P. (1999)  
Sequential PKC - and cdc2 - mediated phosphorylation events elicit Zebrafish nuclear envelope disassembly.  
*J. Cell Sci* **112**:977-987

Courvalin J.C., Segil N., Blobel G., and Worman H.J. (1992)  
The lamin B receptor of the nuclear membrane undergoes mitosis-specific phosphorylation and is a substrate for p34cdc2-type protein kinase.  
*J Biol Chem* **26**:19035-19038

Cremer T. and Cremer C. (2001)  
Chromosome territories, nuclear architecture and gene regulation in mammalian cells.  
*Nat Rev Genet* **2**:292-301

Croft J-A., Bridger J.M., Boyle S., Perry P. and Bickmore W.A. (1999)  
Differences in the localization and morphology of chromosomes in the human nucleus.  
*J Cell Biol* **145**:1119-1131

Davis L.I. (1995)  
The nuclear pore complex.  
*Ann Rev Biochem* **64**:865-896

de Noronha C.M.C., Sherman M.P., Lin H.W., Cavrois M.V., Moir R.D., Goldman R.D. and Greene W.C. (2001)  
Dynamic disruptions in nuclear envelope architecture and integrity induced by HIV-1 Vpr.  
*Science* **294**:1105-1108

Dechat T., Gotzmann J., Stockinger A., Harris C.A., Talle M.A., Siekierka J.J. and Foisner R. (1998)

Detergent-salt resistance of LAP 2 $\alpha$  in interphase nuclei and phosphorylation dependent association with chromosomes early in nuclear assembly implies functions in nuclear structure dynamics.

*EMBO J* **17**:4887-4902

Dreger M., Bengtsson L., Schoneberg T., Otto H. and Hucho F. (2001)

Nuclear envelope proteomics: novel integral membrane proteins of the inner nuclear membrane.

*Proc Natl Acad Sci USA* **98**:11943-11948

Dreger M., Otto H., Neubauer G., Mann M. and Hucho F. (1999)

Identification of phosphorylation sites in native lamina-associated polypeptide 2 $\beta$ .

*Biochemistry* **38**:9426-9434

Duband-Goulet I. and Courvalin J.C. (2000)

Inner nuclear membrane protein LBR preferentially interacts with DNA secondary structures and nucleosomal linker.

*Biochemistry* **39**:6483-6488

Eide T., Coghlann V., Örstavik S., Holsve C., Solberg R., Skalhegg B.S., Lamb N.J., Langeberg L., Fernandez A., Scott J.D., Jahnson T. and Tasken K. (1998)

Molecular cloning, chromosomal localization and cell cycle dependent subcellular distribution of the A-kinase anchoring protein AKAP95.

*Exp Cell Res* **238**:305-316

Ellenberg J., Siggia E.D., Moreira J.E., Smith C.L., Presley J.F., Worman H.J. and Lippincott-Schwartz J. (1997)

Nuclear membrane dynamics and reassembly in living cells: targeting of an inner nuclear membrane protein in interphase and mitosis.

*J Cell Biol* **138**:1193-1206

Ellis J.A., Craxton M., Yates J.R.W. and Kendrick-Jones J (1998)

Aberrant intracellular targeting and cell cycle-dependent phosphorylation of emerin contribute to the Emery-Dreifuss muscular dystrophy phenotype.

*J Cell Sci* **111**:781-792

Enoch,T., Peter M., Nurse P., and Ngg E.A. (1991)

p34cdc2 acts as a lamin kinase in fission yeast.

*J Cell Biol* **112**:797-807

Fackelmeyer F.O. (2000)

Die Architektur des Zellkerns.

*Biospektrum* **6**:441-444

Fasshauer D, Otto H, Eliason WK, Jahn R, Brunger AT. (1997)  
Structural changes are associated with soluble N-ethylmaleimide-sensitive fusion protein attachment protein receptor complex formation.  
*J Biol Chem* **272**:28036-28041

Favreau C., Worman H.J., Wozniak R.W., Frappier T., and Courvalin J.C. (1996)  
Cell cycle-dependent phosphorylation of nucleoporins and nuclear pore membrane protein gp210  
*Biochemistry* **35**:8035-8044

Fischer N., Boudreau E., Hippel M., Drepper F., Haehnel W. and Rochaix J.D. (1999)  
A large fraction of PsaF is nonfunctional in photosystem I complexes lacking the PsaJ subunit.  
*Biochemistry* **38**:5546-5552

Foisner R. and Gerace L. (1993)  
Integral membrane proteins of the nuclear envelope interact with lamins and chromosomes, and binding is modulated by mitotic phosphorylation.  
*Cell* **73**:1267-1279

Fricke M., Hollinshead M., White N. and Vaux D. (1997)  
Interphase nuclei of many mammalian cell types contain deep, dynamic, tubular membrane – bound invaginations of the nuclear envelope  
*J Cell Biol* **136**:531-544

Furukawa K. (1999)  
LAP 2 binding protein 1 (L2BP1/BAF) is a candidate mediator of LAP 2-chromatin interaction.  
*J Cell Sci* **112**:2485-2492

Furukawa K. and Kondo T. (1998)  
Identification of lamina-associated-polypeptide-2 binding domain of B-type lamin.  
*Eur J Biochem* **251**:729-733

Furukawa K., Glass C. and Kondo T. (1997)  
Characterization of the chromatin binding activity of lamina-associated polypeptide (LAP) 2.  
*Biochem Biophys Res Commun* **238**:240-246

Gall J.G. (2000)  
Cajal bodies: the first 100 years.  
*Annu Rev Cell Dev Biol* **16**:273-300

Gant T.M. and Wilson K.L. (1997)  
Nuclear assembly.  
*Ann Rev Cell Dev Biol* **13**:669-695

Gant T.M., Harris C.A. and Wilson K.L. (1999)  
Roles of LAP 2 proteins in nuclear assembly and DNA replication: truncated LAP 2 $\beta$  proteins alter lamina assembly, envelope formation, nuclear size, and DNA replication efficiency in *Xenopus laevis* extracts.  
*J Cell Biol* **144**:1083-1096

Georgatos S.D., Meier J. and Simos G. (1994)  
Lamins and lamin-associated protein.  
*Curr Opinion Cell Biol* **6**:347-353

Gerace L. and Blobel G. (1980)  
The nuclear envelope lamina is reversibly depolymerized during mitosis.  
*Cell* **19**:277-287

Glass J.R. und Gerace L. (1990)  
Lamins A and C bind and assemble at the surface of mitotic chromosomes.  
*J Cell Biol* **111**:1047-1057

Goldberg M., Lu H., Stuurman N., Ashery-Padan R., Weiss A.M., Yu J., Bhattacharyya D., Fisher P.A., Gruenbaum Y. and Wolfner M.F. (1998)  
Interactions among *Drosophila* nuclear envelope proteins lamin, otefin, and YA.  
*Mol Cell Biol* **18**:4315-4323

Goldberg M.W. and Allen T.D. (1996)  
The nuclear pore complex and lamina: three dimensional structures and interactions determined by field emission in-lens scanning electron microscopy.  
*J Mol Biol* **257**:848-865

Hagan L. and Yanmagida M. (1995)  
The product of spindle formation gene Sad1+ associates with the fission yeast spindle pole body and is essential for viability.  
*J Cell Biol* **129**:1033-1047

Haraguchi T., Koujin T., Hayakawa T., Kaneda T., Tsutsumi C., Imamoto N., Akazawa C., Sukegawa J., Yoneda Y. and Hiraoka Y. (2000)  
Live fluorescence imaging reveals early recruitment of emerin, LBR, RanBP2, and Nup153 to reforming functional nuclear envelopes.  
*J Cell Sci* **113**:779-794

Haraguchi T., Koujin T., Segura-Totten M., Lee K.K., Matsuoka Y., Yoneda Y., Wilson K.L. and Hiraoka Y. (2001)  
BAF is required for emerin assembly into the reforming nuclear envelope.  
*J Cell Sci* **114**:4575-4585

- Harel A., Zlotkin E., Nainudel-Epszteyn S., Feinstein N., Fisher P.A. and Gruenbaum Y. (1989) Persistence of major nuclear envelope antigens in an envelope-like structure during mitosis in *Drosophila melanogaster* embryos. *J Cell Sci* **94**:463-470
- Harris C.A., Andryuk P.J., Cline S.W., Mathew S., Siekierka J.J. and Goldstein G. (1995) Structure and mapping of the human thymopoietin (TMPO) gene and relationship of human TMPO  $\beta$  to rat Lamin-Associated Polypeptide 2. *Genomics* **28**:198-205
- Hartinger J., Stenius K., Hogemann D. and Jahn R. (1996) 16-BAC/SDS-PAGE: a two-dimensional gel electrophoresis system for the separation of integral membrane proteins. *Anal Biochem* **240**:126-133
- Hayward-Lester A., Hewetson A., Beale E.G., Oefner P.J., Doris P.A. and Chilton B.S. (1996) Cloning, characterization, and steroid-dependent posttranscriptional processing of RUSH-1 alpha and beta, two uteroglobin promoter-binding proteins. *Mol Endocrinol* **10**:1335-49
- Hennig W. (1999) Heterochromatin. *Chromosoma* **108**:1-9
- Holmer L., Pezhman A. and Worman H.J. (1998) The human Lamin B Receptor/Sterol Reductase multigene family. *Genomics* **54**:469-476
- Inoue H., Nojima H. and Okayama H. (1990) High efficiency transformation of *Escherichia coli* with plasmids. *Gene* **96**:23-28
- Jenkins H., Holman T., Lyon C., Lane B., Stick R. and Hutchison C. (1993) Nuclei that lack a lamina accumulate karyophilic proteins and assemble a nuclear matrix. *J Cell Sci* **106**:275-285
- Karas M. and Hillenkamp F. (1988) Laser desorption ionization of proteins with molecular masses exceeding 10000 daltons. *Anal Chem* **60**:2299-2301
- Karcher R.L., Deacon S.W. and Gelfand V.I (2002) Motor-cargo interactions: the key to transport specificity. *Trends Cell Biol* **12**: 21-27

La Luna S., Allen K.E., Mason S.L. and La Thangue N.B. (1999)  
Integration of growth-suppressing BTB/POZ domain protein with the DP component of the E2F transcription factor.  
*EMBO J* **18**:212-228

Lee K.K., Haraguchi T., Lee R.S., Koujin T., Hiraoka Y. and Wilson K.L. (2001)  
Distinct functional domains in emerin bind lamin A and DNA-bridging protein BAF.  
*J Cell Sci* **114**:4567-4573

Lichter P., Cremer T., Borden J., Manuelidis L. and Ward D.C. (1988)  
Delineation of individual human chromosomes in metaphase and interphase cells by in situ suppression hybridization using recombinant DNA libraries.  
*Hum Genet* **80**:224-234

Lin F., Blake D.L., Callebaut I., Skerjanc I.S., Holmer L., McBurney M.W., Paulin-Levasseur M. and Worman H.J. (2000)  
MAN1, an inner nuclear membrane protein that shares the LEM domain with lamina-associated polypeptide 2 and emerin.  
*J Biol Chem* **275**:4840-4847

Lin F., Noyer C.M., Ye Q., Courvalin J.C. and Worman H.J. (1996)  
Autoantibodies from patients with primary biliary cirrhosis recognize a region within the nucleoplasmic domain of inner nuclear membrane protein LBR.  
*Hepatology* **23**:57-61

Mackay J.P. and Crossley M. (1998)  
Zinc fingers are sticking together.  
*Trends Biochem Sci* **23**:1-4

Maison C., Pyrpasopoulou A., Theodoropoulos P.A. and Georgatos S.D. (1997)  
The inner nuclear membrane protein LAP 1 forms a native complex with B-type lamins and partitions with spindle-associated mitotic vesicles.  
*EMBO J* **16**:4839-4850

Malone C.J., Fixsen W.D., Horvitz H.R. and Han M. (1999)  
UNC-84 localizes to the nuclear envelope and is required for nuclear migration and anchoring during *C. elegans* development.  
*Development* **126**:3171-3181

Manilal S., Nguyen T.M., Sewry C.A. and Morris G.E. (1996)  
The Emery-Dreifuss muscular dystrophy protein, emerin, is a nuclear membrane protein.  
*Hum Mol Genet* **5**:801-808

Mansharamani M., Hewetson A. and Chilton B.S. (2001)  
Cloning and characterization of an atypical type IV P-type ATPase that binds to the RING motif of RUSH transcription factors.  
*J Biol Chem* **276**:3641-3649

Martin L., Crimando C. and Gerace L. (1995)  
cDNA cloning and characterization of lamina associated polypeptide 1C (LAP 1C), an integral protein of the inner nuclear membrane.  
*J Biol Chem* **270**:8822-8828

Martins S.B, Eide T., Steen R.L., Jahnsen T., Skalhegg B.S. and Collas P. (2000)  
HA95 is a protein of the chromatin and nuclear matrix regulating nuclear envelope dynamics.  
*J Cell Sci* **113**:3703-3713

Matsuoka Y., Takagi M., Ban T., Miyazaki M., Yamamoto T., Kondo Y., and Yoneda Y. (1999)  
Identification and characterization of nuclear pore subcomplexes in mitotic extract of human somatic cells.  
*Biochem Biophys Res Comm* **254**:417-423

Matynia A., Salus S.S. and Sazer S. (2002)  
Three proteins required for early steps in the protein secretory pathway also affect nuclear envelope structure and cell cycle progression in fission yeast.  
*J Cell Sci* **115**:421-431

Mical T.I. and Monteiro M.J. (1998)  
The role of sequences unique to nuclear intermediate filaments in the targeting and assembly of human lamin B: evidence for lack of interaction of lamin B with its putative receptor.  
*J Cell Sci* **111**:3471-3485

Mislow J.M., Kim M.S., Davis D.B. and McNally E.M. (2002)  
Myne-1, a spectrin repeat transmembrane protein of the myocyte inner nuclear membrane, interacts with lamin A/C.  
*J Cell Sci* **115**:61-70.

Misteli T. (2001)  
Protein dynamics: implications for nuclear architecture and gene expression.  
*Science* **291**:843-847

Misteli T. and Spector D.L. (1997)  
Protein phosphorylation and the nuclear organization of pre-mRNA splicing.  
*Trends Cell Biol* **7**:135-138

Misteli T. and Spector D.L. (1999)  
RNA polymerase II targets pre-mRNA splicing factors to transcription sites in vivo.  
*Mol Cell* **3**:697-705

Misteli T., Cáceres J.F. and Spector D.L. (1997)  
The dynamics of a pre-mRNA splicing factor in living cells.  
*Nature* **387**:523-527

Moir R.D. and Spann T.P. (2001)  
The structure and function of nuclear lamins: implication for disease.  
*Cell Mol Life Sci* **58**:1748-1757

Morris N.R., Ziang X. and Beckwith S.M. (1995)  
Nuclear migration advances in fungi.  
*Trends Cell Biol* **5**:278-282

Mosley-Bishop K.L., Li Q., Patterson L. and Fischer J.A. (1999)  
Molecular analysis of the klarsicht gene and its role in nuclear migration within  
differentiating cells of the *Drosophila* eye.  
*Cur. Biol.* **9**:1211-1220

Nagano A., Koga R., Ogawa M., Kurano Y., Kawada J., Okada R., Hayashi Y.K., Tsukahara  
T. and Arahata K. (1996)  
Emerin deficiency at the nuclear membrane in patients with Emery-Dreifuss muscular  
dystrophy  
*Nature Genet* **12**:254-259

Nigg, E.A. (1992)  
Assembly and cell cycle dynamics of the nuclear lamina  
*Sem Cell Biol* **3**:245-253

Nigro V., Bruni P., Ciccodicola A., Politano L., Nigro G., Piluso G., Cappa V., Covone A.E.,  
Romeo G. and D'Urso M. (1995)  
SSCP detection of novel mutations in patients with Emery-Dreifuss muscular dystrophy:  
definition of a small C-terminal region required for emerin function.  
*Hum Mol Genet* **4**:2003-2004

Nikolakaki E., Meier J., Simos G., Georgatos S.D. and Giannakouros T. (1997)  
Mitotic phosphorylation of the lamin B receptor by a serine/arginine kinase and p34(cdc2).  
*J Biol Chem* **272**:6208-6213

Nikolakaki E., Simos G., Georgatos S.D. and Giannakouros T. (1996)  
A nuclear envelope-associated kinase phosphorylates arginine-serine motifs and modulates  
interactions between the lamin B receptor and other nuclear proteins.  
*J Biol Chem* **271**:8365-8272

Nili E., Cojocaru G.S., Kalma Y., Ginsberg D., Copeland N.G., Gilbert D.J., Jenkins N.A.,  
Berger R., Shaklai S., Amariglio N., Brok-Simoni F., Simon A.J. and Rechavi G. (2001)  
Nuclear membrane protein LAP 2 $\beta$  mediates transcriptional repression alone and with its  
binding partner GCL (germ-cell-less).  
*J Cell Sci* **114**:3297-3307

Orphanides G. and Reinberg D. (2002)  
A unified theory of gene expression.  
*Cell* **108**:439-451

Örstavik S., Eide T., Collas P., Han I.O., Tasken K., Kieff E., Jahnsen T. and Skalhegg B.S. (2000)  
Identification, cloning and characterization of a novel nuclear protein, HA95, homologous to the A-kinase anchoring protein 95.  
*Biol Cell* **92**: 27-37

Padan R., Nainudel-Epszteyn S., Goitein R., Fainsod A. and Gruenbaum Y. (1990)  
Isolation and characterization of the Drosophila nuclear envelope otefin cDNA.  
*J Biol Chem* **265**:7808-7813

Paddy M.R., Belmeont A.S., Saumweber H., Agard D.A. and Sedat J.W. (1990)  
Interphase nuclear envelope lamins form a discontinuous network that interacts with only a fraction of the chromatin in the nuclear periphery  
*Cell* **62**:89-106

Papoutsopoulou S., Nikolakaki E. and Giannakouros T. (1999)  
SRPK1 and LBR protein kinases show identical substrate specificities.  
*Biochem Biophys Res Commun* **255**:602-607

Paulin-Levasseur M., Blake D.L., Julien M. and Rouleau L. (1996)  
The MAN antigens are non-lamin constituents of the nuclear lamina in vertebrate cells.  
*Chromosoma* **104**:367-379

Peter M. Nakagawa J. Doree M. Labbe J.C. und Nigg E.A. (1990)  
In vitro disassembly of the nuclear lamina and M-phase specific phosphorylation of lamins by cdc2 kinase.  
*Cell* **61**:591-602

Polioudaki H., Kourmouli N., Drosou V., Bakou A., Theodoropoulos P.A., Singh P.B., Giannakouros T. and Georgatos S.D. (2001)  
Histones H3/H4 form a tight complex with the inner nuclear membrane protein LBR and heterochromatin protein 1.  
*EMBO Rep* **2**:920-925

Qumsiyeh M.B. (1999)  
Structure and function of the nucleus: anatomy and physiology of chromatin.  
*Cell Mol Life Sci* **55**:1129-1140

Raff J.W. and Glover D.M. (1989)  
Centrosomes and not nuclei, initiate pole cell formation in Drosophila embryo.  
*Cell* **57**:611-619

Rolls M.M., Stein P.A., Taylor S.S., Ha E., McKeon F. and Rapoport T.A. (1999)  
A visual screen of a GFP-fusion library identifies a new type of nuclear envelope membrane protein.  
*J Cell Biol.* **146**:29-44

Salina D., Bodoor K., Eckley D.M., Schroer T.A., Rattner J.B. and Burke B. (2002)  
Cytoplasmic dynein as a fasciliator of nuclear envelope breakdown.  
*Cell* **108**:97-107

Sambrook J., Fritsch F.R. and Maniatis T. (1989)  
Molecular Cloning: A Laboratory Manual  
New York: Cold Spring Harbor Laboratory Press

Saurin A.J, Borden K.L., Boddy M.N. and Freemont P.S. (1996)  
Does this have a familiar RING?  
*Trends Biochem Sci* **21**:208-14

Schägger H. and Jagow G. (1991)  
Blue native electrophoresis for isolation of membrane protein complexes in enzymatically active form.  
*Anal Biochem* **199**:223-231

Schuler E., Lin F. and Worman H.J. (1994)  
Characterization of the human gene encoding LBR, an integral protein of the nuclear envelope inner membrane.  
*J Biol Chem* **269**:11312-11317.

Schultz J., Copley R.R., Doerks T., Ponting C.P. and Bork P. (2000)  
SMART: a web-based tool for the study of genetically mobile domains.  
*Nucleic Acids Res* **28**:231-234

Schultz J., Milpetz F., Bork P. and Ponting C.P. (1998)  
SMART, a simple modular architecture research tool: identification of signaling domains.  
*Proc Natl Acad Sci USA* **95**:5857-5864

Senior A. and Gerace L. (1988)  
Integral membrane proteins specific to the inner nuclear membrane and associated with the nuclear lamina.  
*J Cell Biol* **107**:2029-2036

Shevchenko A., Wilm M., Vorm O. and Mann M. (1996)  
Mass spectrometric sequencing of proteins from silver-stained polyacrylamide gels.  
*Anal Chem* **68**:850-858

Shumaker D.K., Lee K.K., Tanhehco Y.C., Craigie R. and Wilson K.L. (2001)  
LAP 2 binds to BAF DNA complexes: requirement for the LEM domain and modulation by variable regions.  
*EMBO J* **20**:1754-64

Silve S., Dupuy P.H, Ferrara P. and Loison G. (1998)  
Human lamin B receptor exhibits sterol C14-reductase activity in *Saccharomyces cerevisiae*.  
*Biochim Biophys Acta* **1392**:233-244

Spann T.P., Moir R.D., Goldman A.E., Stick R. and Goldman R.D. (1997)  
Disruption of nuclear lamin organization alters the distribution of replication factors and inhibits DNA synthesis.  
*J Cell Bio.* **136**:1201-1212

Starr D.A., Hermann G.J., Malone C.J., Fixsen W., Priess J.R., Horvitz H.R. and Han M. (2001)  
unc-83 encodes a novel component of the nuclear envelope and is essential for proper nuclear migration.  
*Development* **128**:5039-5050

Stewart M. and Clarkson W.D. (1996)  
Nuclear pores and macromolecular assemblies involved in nucleocytoplasmic transport.  
*Curr Opin Struct Biol* **6**:162-165

Sullivan T., Escalante-Alcalde D., Bhatt H., Anver M., Bhat N., Nagashima K., Stewart C.L. and Burke B. (1999)  
Loss of A-type lamin expression compromises nuclear envelope integrity leading to muscular dystrophy.  
*J Cell Biol* **147**:913-920

Valenzuela d.m., Stitt T.N., DiStefano P.S., Rojas E., Mattsson K., Compton D.L., Nunez L., Park J.S., Stark J.L., Gies D.R., Thomas S., LeBeau M.M., Fernald A.A., Copeland N.G., Jenkins N.A., Burden S.J., Glass D.J. and Yancoupulos G.D. (1995)  
Receptor tyrosine kinase specific for the skeletal muscle lineage: expression in embryonic muscle, at the neuromuscular junction, and after injury.  
*Neuron* **15**:573-584

Vlcek S., Just H., Dechat T. and Foisner R. (1999)  
Functional diversity of LAP 2alpha and LAP 2beta in postmitotic chromosome association is caused by an alpha-specific nuclear targeting domain.  
*EMBO J* **18**:6370-6384

Vuillard L., Marret N. and Rabilloud T. (1995)  
Enhancing protein solubilization with nondetergent sulfobetaines.  
*Electrophoresis* **16**:295-7

Welte M.A., Gross S.P., Postner M., Block S.M., and Wieschaus E.F. (1998)  
Developmental regulation of vesicle transport in Drosophila embryos: forces and kinetics  
*Cell* **92**:547-557

Worman H.J., Evans C.D. and Blobel G. (1990)  
A lamin B receptor of the nuclear envelope inner membrane: a polytopic protein with eight potential transmembrane domains.  
*J Cell Biol* **111**:1535-1542

Worman H.J., Yuan J., Blobel G. and Georgatos S.D. (1988)  
A lamin B receptor in the nuclear envelope.  
*Proc Natl Acad Sci USA* **85**:8531-8534

Yang L., Guan T. and Gerace L. (1997)  
Lamin-binding fragment of LAP 2 inhibits increase in nuclear volume during the cell cycle and progression into S phase.  
*J Cell Biol* **139**:1077-1087

Ye Q. and Worman H.J. (1996)  
Interaction between an integral protein of the nuclear envelope inner membrane and human chromodomain proteins homologous to Drosophila HP1.  
*J Biol Chem* **271**:14653-14656

Ye Q. and Worman H.J. (1994)  
Primary structure analysis and lamin B and DNA binding of human LBR, an integral protein of the nuclear envelope inner membrane.  
*J Biol Chem* **269**:11306-11311

Ye Q., Callebaut I., Pezhman A., Courvalin J.C. and Worman H.J. (1997)  
Domain-specific interactions of human HP1-type chromodomain proteins and inner nuclear membrane protein LBR.  
*J Biol Chem* **272**:14983-14989

Zhang C., Jenkins H., Goldberg M.W., Allen T.D. and Hutchison C.J. (1996)  
Nuclear lamina and nuclear matrix organization in sperm pronuclei assembled in Xenopus egg extract.  
*J Cell Sci* **109**:2275-2286

Zhang Q., Skepper J.N., Yang F., Davies J.D., Hegyi L., Roberts R.G., Weissberg P.L. ,Ellis J.A., Shanahan C.M. (2001)  
Nesprins: a novel family of spectrin-repeat-containing proteins that localize to the nuclear membrane in multiple tissues.  
*J Cell Sci* **114**:4485-4498

Zheng R., Ghirlando R., Soo Lee M., Mizuuchi K., Krause M. and Craigie R. (2000)  
Barrier-to-autointegration factor (BAF) bridges DNA in a discrete, higher-order nucleoprotein complex.  
*Proc Natl Acad Sci USA* **97**:8997-9002

Zhong S., Salomoni P. and Pandolfi P.P. (2000)  
The transcriptional role of PML and the nuclear body.  
*Nat Cell Biol* **2**:E85-90

Zoghbi H.Y. and Orr H.T. (2000)  
Glutamine repeats and neurodegeneration.  
*Annu Rev Neurosci* **23**:217-247