

9 LITERATURVERZEICHNIS

- Abrams, C.S., Zhang, J., Downes, C.P., Tang, X.W., Zhao, W., Rittenhouse, S.E. (1996) Phosphopleckstrin inhibits G $\beta\gamma$ -activable platelet phosphatidylinositol-4,5-bisphosphate 3-kinase. *J. Biol. Chem.*, **271**, 25192-25197.
- Ahnert-Hilger, G., Nürnberg, B., Exner, T., Schäfer, T., Jahn, R. (1998) The heterotrimeric G protein G α_{o2} regulates catecholamine uptake by secretory vesicles. *EMBO J.*, **17**, 406-413.
- Asano, T., Morishita, R., Matsuda, T., Fukada, Y., Yoshizawa, T., Kato, K. (1993) Purification of four forms of the $\beta\gamma$ subunit complex of G proteins containing different γ subunits. *J. Biol. Chem.*, **268**, 20512-20519.
- Auger, K.R., Serunian, L.A., Soltoff, S.P., Libby, P., Cantley, L.C. (1989) PDGF-dependent tyrosine phosphorylation stimulates production of novel polyphosphoinositides in intact cells. *Cell*, **57**, 167-175.
- Bae, Y.S., Cantley, L.G., Chen, C.S., Kim, S.R., Kwon, K.S., Rhee, S.G. (1998) Activation of phospholipase C- γ by phosphatidylinositol 3,4,5-trisphosphate. *J. Biol. Chem.*, **273**, 4465-4469.
- Baier, R., Bondeva, T., Klinger, R., Bondev, A., Wetzker, R. (1999) Retinoic acid induces selective expression of phosphoinositide 3-kinase γ in myelomonocytic U937 cells. *Cell Growth Differ.*, **10**, 447-456.
- Barr, A.J., Ali, H., Haribabu, B., Snyderman, R., Smrcka, A.V. (2000) Identification of a region at the N-terminus of phospholipase C- β_3 that interacts with G protein $\beta\gamma$ subunits. *Biochemistry*, **39**, 1800-1806.
- Bayewitch, M.L., Avidor-Reiss, T., Levy, R., Pfeuffer, T., Nevo, I., Simonds, W.F., Vogel, Z. (1998a) Differential modulation of adenylyl cyclases I and II by various G β subunits. *J. Biol. Chem.*, **273**, 2273-2276.
- Bayewitch, M.L., Avidor-Reiss, T., Levy, R., Pfeuffer, T., Nevo, I., Simonds, W.F., Vogel, Z. (1998b) Inhibition of adenylyl cyclase isoforms V and VI by various G $\beta\gamma$ subunits. *FASEB J.*, **12**, 1019-1025.
- Belluscio, L., Gold, G.H., Nemes, A., Axel, R. (1998) Mice deficient in G $_{olf}$ are anosmic. *Neuron*, **20**, 69-81.
- Bernstein, H.-G., Keilhoff, G., Reiser, M., Freese, S., Wetzker, R. (1998) Tissue distribution and subcellular localization of a G protein activated phosphoinositide 3-kinase. An immunohistochemical study. *Cell. Mol. Biol.*, **44**, 973-983.
- Bi, L., Okabe, I., Bernard, D.J., Wynshaw-Boris, A., Nussbaum, R.L. (1999) Proliferative defect and embryonic lethality in mice homozygous for a deletion in the p110 α subunit of phosphoinositide 3-kinase. *J. Biol. Chem.*, **274**, 10963-10968.
- Blair, L.A., Marshall, J. (1997) IGF-1 modulates N and L calcium channels in a PI 3-kinase-dependent manner. *Neuron*, **19**, 421-429.

- Blank, J.L., Ross, A.H., Exton, J.H. (1991) Purification and characterization of two G-proteins that activate the β_1 isozyme of phosphoinositide-specific phospholipase C. *J. Biol. Chem.*, **266**, 18206-18216.
- Bondeva, T., Pirola, L., Bulgarelli-Leva, G., Rubio, I., Wetzker, R., Wymann, M.P. (1998) Bifurcation of lipid and protein kinase signals of PI3K γ to the protein kinases PKB and MAPK. *Science*, **282**, 293-296.
- Bootman, M.D., Berridge, M.J. (1995) The elemental principles of calcium signaling. *Cell*, **83**, 675-678.
- Breeze, A.L., Kara, B.V., Barrat, D.G., Anderson, M., Smith, J.C., Luke, R.W., Best, J.R., Cartledge, S.A. (1996) Structure of a specific peptide complex of the carboxy-terminal SH2 domain from the p85 α subunit of phosphatidylinositol 3-kinase. *EMBO J.*, **15**, 3579-3589.
- Breitwieser, G.E., Szabo, G. (1985) Uncoupling of cardiac muscarinic and β -adrenergic receptors from ion channels by a guanine nucleotide analogue. *Nature*, **317**, 538-540.
- Brown, R.A., Domin, J., Arcaro, A., Waterfield, M.D. (1999) Insulin activates the α isoform of class II phosphoinositide 3-kinase. *J. Biol. Chem.*, **274**, 14529-14532.
- Brunk, I., Pahner, I., Maier, U., Jenner, B., Veh, R.W., Nürnberg, B., Ahnert-Hilger, G. (1999) Differential distribution of G-protein β -subunits in brain: an immunocytochemical analysis. *Eur. J. Cell. Biol.*, **78**, 311-322.
- Buck, E., Li, J., Chen, Y., Weng, G., Scarlata, S., Iyengar, R. (1999) Resolution of a signal transfer region from a general binding domain in G β for stimulation of phospholipase C- β_2 . *Science*, **283**, 1332-1335.
- Cabrera, J.L., de Freitas, F., Satpaev, D.K., Slepak, V.Z. (1998) Identification of the G β_5 -RGS7 protein complex in the retina. *Biochim. Biophys. Res. Commun.*, **249**, 898-902.
- Camps, M., Carozzi, A., Schnabel, P., Scheer, A., Parker, P.J., Gierschik, P. (1992) Isozyme-selective stimulation of phospholipase C- β_2 by G protein $\beta\gamma$ subunits. *Nature*, **360**, 684-686.
- Carozzi, A.J., Kriz, R.W., Webster, C., Parker, P.J. (1992) Identification, purification and characterization of a novel phosphatidylinositol-specific phospholipase C, a third member of the β subfamily. *Eur. J. Biochem.*, **210**, 521-529.
- Carozzi, A., Camps, M., Gierschik, P., Parker, P.J. (1993) Activation of phosphatidylinositol lipid-specific phospholipase C- β_3 by G protein $\beta\gamma$ subunits. *FEBS Lett.*, **315**, 340-342.
- Carpenter, C.L., Auger, K.R., Duckworth, B.C., Hou, W.M., Schaffhausen, B.S., Cantley, L.C. (1993) A tightly associated serine/threonine protein kinase regulates phosphoinositide 3-kinase activity. *Mol. Cell Biol.*, **13**, 1657-1665.
- Casey, P., Seabra, M.C. (1996) Protein prenyltransferases. *J. Biol. Chem.*, **271**, 5289-5892.
- Chen, W.-J., Andres, D.A., Goldstein, J.L., Russell, D.W., Brown, M.S. (1991) cDNA cloning and expression of the peptide-binding β subunit of rat p21ras farnesyltransferase, the counterpart of yeast DPR1/RAM1. *Cell*, **66**, 327-334.

- Chen, C.-K., Burns, M.E., He, W., Wensel, T.G., Baylor, D.A., Simon, M.I. (2000) Slowed recovery of rod photoresponse in mice lacking the GTPase accelerating protein RGS9-1. *Nature*, **403**, 557-559.
- Chung, J., Grammer, T.C., Lemon, K.P., Kazlauskas, A., Blenis, J. (1994) PDGF- and insulin-dependent pp70S6k activation mediated by phosphatidylinositol-3-OH kinase. *Nature*, **370**, 71-75.
- Clapham, D.E., Neer, E.J. (1997) G protein $\beta\gamma$ subunits. *Annu. Rev. Pharmacol. Toxicol.*, **37**, 167-203.
- Codina, J., Carty, D.J., Birnbaumer, L., Iyengar, R. (1990) Purification of G proteins. *Meth. Enzymol.*, **195**, 177-188.
- Cook, L.A., Schey, K.L., Wilcox, M.D., Dingus, J., Hildebrandt, J.D. (1998) Heterogenous processing of a G protein γ subunit at a site critical for protein and membrane interactions. *Biochemistry*, **37**, 12280-12286.
- Cooper, D.M., Brooker, G. (1993) Ca^{2+} -inhibited adenylyl cyclase in cardiac tissue. *Trends Pharmacol. Sci.*, **14**, 34-36.
- Coso, O.A., Chiariello, M., Yu, J.C., Teramoto, H., Crespo, P., Xu, N., Miki, T., Gutkind, J.S. (1995) The small GTP-binding proteins Rac1 and Cdc42 regulate the activity of the JNK/SAPK signaling pathway. *Cell*, **81**, 1137-1146.
- De Camilli, P., Emr, S.D., McPherson, P.S., Novick, P. (1996) Phosphoinositides as regulators in membrane traffic. *Science*, **271**, 1533-1539.
- Dekker, L.V., Segal, A.W. (2000) Signals to move cells. *Science*, **287**, 982-984.
- Dhanasekaran, N., Vara Prasad, M.V.V.S., Wadsworth, S.J., Dermott, J.M., van Rossum, G. (1994) Protein kinase C-dependent and -independent activation of Na^+/H^+ exchanger by $\text{G}\alpha_{12}$ class of G proteins. *J. Biol. Chem.*, **269**, 11802-11806.
- Dhanasekaran, N., Heasley, L.E., Johnson, G.L. (1995) G protein-coupled receptor systems involved in cell growth and oncogenesis. *Endocrine Rev.*, **16**, 259-270.
- Dhand, R., Hara, K., Hiles, I., Bax, B., Gout, I., Panayotou, G., Fry, M.J., Yonezawa, K., Kasuga, M., Waterfield, M.D. (1994a) PI 3-kinase: structural and functional analysis of intersubunit interactions. *EMBO J.*, **13**, 511-521.
- Dhand, R., Hiles, I., Panayotou, G., Roche, S., Fry, M.J., Gout, I., Totty, N.F., Truong, O., Vicendo, P., Yonezawa, K., Waterfield, M.D. (1994b) PI 3-kinase is a dual specificity enzyme: autoregulation by an intrinsic protein-serine kinase activity. *EMBO J.*, **13**, 522-533.
- Dietrich, A., Meister, M., Spicher, K., Schultz, G., Camps, M., Gierschik, P. (1992) Expression, characterization and purification of soluble G-protein $\beta\gamma$ dimers composed of defined subunits in baculovirus-infected insect cells. *FEBS Lett.*, **313**, 220-224.
- Dobos, G.J., Wu, X.R., Traynor-Kaplan, A. (1993) A product of phosphatidylinositol-3 kinase is elevated in dividing HT29 colonic epithelial cells. *FEBS Lett.*, **324**, 143-146.
- Downward, J. (1998) Mechanisms and consequences of activation of protein kinase B/Akt, *Curr. Opin. Cell Biol.*, **10**, 262-267.

- Evans, R. (1988) The steroid and thyroid hormone receptor subfamily. *Science*, **240**, 889-895.
- Exner, T., Jensen, O.N., Mann, M., Kleuss, C., Nürnberg, B. (1999) Posttranslational modification of G α_{o1} generates G α_{o3} , an abundant G protein in brain. *Proc. Natl. Acad. Sci. U S A*, **96**, 1327-32.
- Exner, T., Nürnberg, B. (1999) Immuno- and gold staining of a single Western blot. *Anal. Biochem.*, **260**, 108-110.
- Exton, J.H. (1997) Cell signaling through guanine-nucleotide-binding regulatory proteins (G proteins) and phospholipases. *Eur. J. Biochem.*, **243**, 10-20.
- Fanger, G.R., Gerwins, P., Widmann, C., Jarpe, M.B., Johnson, G.L. (1997) MEKKs, GCKs, MLKs, PAKs, TAKs, and tpls: upstream regulators of the c-Jun amino-terminal kinases? *Curr. Opin. Genet. Dev.*, **7**, 67-74.
- Fawzi, A.B., Fay, D.S., Murphy, E.A., Tamir, H., Erdos, J.J., Northup, J.K. (1991) Rhodopsin and the retinal G-protein distinguish among G-protein $\beta\gamma$ subunit forms. *J. Biol. Chem.*, **266**, 12194-12200.
- Flanagan, C.A., Schnieders, E.A., Emerick, A.W., Kunisawa, R., Thorner, J.W., Admon, A. (1993) Phosphatidylinositol 4-kinase: gene structure and requirement for yeast cell viability. *Science*, **262**, 1444-1448.
- Fletcher, M.A., Myung, C.-S., Savino, Y., Yasuda, H., Kazhan, R., Garrison, J.C. (1998) Differential activity of the G protein $\beta_5\gamma_2$ subunit at receptors and effectors. *J. Biol. Chem.*, **273**, 34429-34436.
- Ford, C.E., Skiba, N.P., Bae, H., Daaka, Y., Reuveny, E., Shekter, L.R., Rosal, R., Weng, G., Yang, C.S., Iyengar, R., Miller, R.J., Jan, L.Y., Lefkowitz, R.J., Hamm, H.E. (1998) Molecular basis for interactions of G protein $\beta\gamma$ subunits with effectors. *Science*, **280**, 1271-1274.
- Franke, T.F., Cantley, L.C. (1997a) Apoptosis. A Bad kinase makes good. *Nature*, **390**, 116-117.
- Franke, T.F., Kaplan, D.R., Cantley, L.C. (1997b) PI3K: downstream AKTion blocks apoptosis. *Science*, **275**, 665-668.
- Frech, M., Andjelkovich, M., Ingley, E., Reddy, K.K., Falck, J.R., Hemmings, B.A. (1997) High affinity binding of inositol phosphates and phosphoinositides to the pleckstrin homology domain of RAC/protein kinase B and their influence on kinase activity. *J. Biol. Chem.*, **272**, 8474-8481.
- Fruman, D.A., Meyers, R.E., Cantley, L.C. (1998) Phosphoinositide kinases. *Annu. Rev. Biochem.*, **67**, 481-507.
- Fukada, Y., Takao, T., Ohguro, H., Yoshizawa, T., Akino, T., Shimonishi, Y. (1990) Farnesylated γ -subunit of photoreceptor G protein indispensable for GTP-binding. *Nature*, **346**, 658-660.
- Galetic, I., Andjelkovic, M., Meier, R., Brodbeck, D., Park, J., Hemmings, B.A. (1999) Mechanism of protein kinase B activation by insulin/insulin-like growth factor-1 revealed by specific inhibitors of phosphoinositide 3-kinase. Significance for diabetes and cancer. *Pharmacol. Ther.*, **82**, 409-425.

- Garbers, D.L., Koesling, D., Schultz, G. (1994) Guanylyl cyclase receptors. *Mol. Biol. Cell*, **5**, 1-5.
- Garcia, D.E., Li, B., Garcia-Ferreiro, E., Hernandez-Ochoa, E.O., Yan, K., Gautam, N., Catterall, W.A., Mackie, K., Hille, B. (1998) G-protein $\beta\gamma$ specificity in the fast membrane-delimited inhibition of Ca^{2+} channels. *J. Neurosci.*, **18**, 9163-9170.
- Gautam, N., Downes, G.B., Yan, K., Kisselev, O. (1998) The G-protein $\beta\gamma$ -complex. *Cell. Signal.*, **10**, 447-455.
- Gierschik, P., Camps, M. (1994) Stimulation of phospholipase C by G-protein $\beta\gamma$ subunits. *Methods Enzymol.*, **238**, 181-195.
- Gohla, A., Harhammer, R., Schultz, G. (1997) The G-protein G_{13} but not G_{12} mediates signaling from lysophosphatidic acid receptor via epidermal growth factor receptor to Rho. *J. Biol. Chem.*, **273**, 4653-4659.
- Gohla, A., Offermanns, S., Wilkie, T.M., Schultz, G. (1999) Differential involvement of $G\alpha_{12}$ and $G\alpha_{13}$ in receptor-mediated stress fiber formation. *J. Biol. Chem.*, **274**, 17901-17907.
- Graf, R., Mattera, R., Codina, J., Evans, T., Ho, Y.K., Estes, M.K., Birnbaumer, L. (1992) Studies on the interaction of α subunits of GTP-binding proteins with $\beta\gamma$ dimers. *Eur. J. Biochem.*, **210**, 609-619.
- Graness, A., Adomeit, A., Heinze, R., Wetzker, R., Liebmann, C. (1998) A novel mitogenic signaling pathway of bradykinin in the human colon carcinoma cell line SW-480 involves sequential activation of a $G_{q/11}$ protein, phosphatidylinositol 3-kinase β , and protein kinase C ϵ . *J. Biol. Chem.*, **273**, 32016-32022.
- Gudermann, T., Nürnberg, B., Schultz, G. (1995) Receptors and G proteins as primary components of transmembrane signal transduction. Part 1. G-protein-coupled receptors: structure and function. *J. Mol. Med.*, **73**, 51-63.
- Gutkind, J.S. (1998) The pathways connecting G protein coupled receptors to the nucleus through divergent mitogen-activated protein kinase cascades. *J. Biol. Chem.*, **273**, 1839-1842.
- Hamm, H.E. (1998) The many faces of G protein Signaling. *J. Biol. Chem.*, **273**, 669-672.
- Hartley, K.O., Gell, D., Smith, G.C., Zhang, H., Divecha, N., Connelly, M.A., Admon, A., Lees-Miller, S.P., Anderson, C.W., Jackson, S.P. (1995) DNA-dependent protein kinase catalytic subunit: a relative of phosphatidylinositol 3-kinase and the ataxia telangiectasia gene product. *Cell*, **82**, 849-856.
- Hawes, B.E., Luttrell, L.M., van Biesen, T., Lefkowitz, R.J. (1996) Phosphatidylinositol 3-kinase is an early intermediate in the $G\beta\gamma$ -mediated mitogen-activated protein kinase signaling pathway. *J. Biol. Chem.*, **271**, 12133-12136.
- Hawkins, P.T., Jackson, T.R., Stephens, L.R. (1992) Platelet-derived growth factor stimulates synthesis of $\text{PtdIns}(3,4,5)\text{P}_3$ by activating a $\text{PtdIns}(4,5)\text{P}_2$ 3-OH kinase. *Nature*, **358**, 157-159.
- Hazeki, O., Hazeki, K., Katada, T., Ui, M. (1996) Inhibitory effect of wortmannin on phosphatidylinositol 3-kinase-mediated cellular events. *J. Lipid Mediat. Cell. Signal.*, **14**, 259-261.

- He, Z., He, Y.S., Kim, Y., Chu, L., Ohmstede, C., Biron, K.K., Coen, D.M. (1997) The human cytomegalovirus UL97 protein is a protein kinase that autophosphorylates on serines and threonines. *J. Virol.*, **71**, 405–411.
- Hepler, J.R., Kozasa, T., Smrcka, A.V., Simon, M.I., Rhee, S.G., Sternweis, P.C., Gilman, A.G. (1993) Purification from Sf9 cells and characterization of recombinant G α_q and G α_{11} . Activation of purified phospholipase C isozymes by G α subunits. *J. Biol. Chem.*, **268**, 14367-14375.
- Hirsch, E., Katanaev, V.L., Garlanda, C., Azzolino, O., Pirola, L., Silengo, L., Sozzani, S., Mantovani, A., Atruda, F., Wymann, M.P. (2000) Central role for G protein-coupled phosphoinositide 3-kinase γ in inflammation. *Science*, **287**, 1049-1053.
- Hofmann, F., Lacinová, L., Klugbauer, N. (1997) Voltage-dependent calcium channels: from structure to function. *Rev. Physiol. Biochem. Pharmacol.*, **139**, 33-87.
- Hofmann, T., Obukhov, A.G., Schaefer, M., Harteneck, C., Gudermann, T., Schultz, G. (1999) Direct activation of human TRPC6 and TRPC3 channels by diacylglycerol. *Nature*, **297**, 259-263.
- Hu, P., Mondino, A., Skolnik, E.Y., Schlessinger, J. (1993) Cloning of a novel, ubiquitously expressed human phosphatidylinositol 3-kinase and identification of its binding site on p85. *Mol. Cell. Biol.*, **13**, 7677-7688.
- Hu, Z.W., Shi, X.Y., Lin, R.Z., Hoffman, B.B. (1999) Contrasting signaling pathways of α_{1A} - α_{1B} -adrenergic receptor subtype activation of phosphatidylinositol 3-kinase and ras in transfected NIH3T3 cells. *Mol. Endocrinol.*, **13**, 3-14.
- Huang, L., Shanker, Y.G., Dubauskaite, J., Zheng, J.Z., Yan, W., Rosenzweig, S., Spielman, A.I., Max, M., Margolskee, R.F. (1999) G γ_{13} colocalizes with gustducin in taste receptor cells and mediates IP $_3$ responses to bitter denatonium. *Nat. Neurosci.*, **12**, 1055-1062.
- Hunter, T. (1995) When is a lipid kinase not a lipid kinase? When it is a protein kinase. *Cell*, **83**, 1-4.
- Illenberger, D., Schwald, F., Gierschik, P. (1997) Characterization and purification from bovine neutrophils of a soluble guanine-nucleotide-binding protein that mediates isozyme-specific stimulation of phospholipase C β_2 . *Eur. J. Biochem.*, **246**, 71-77.
- Iñiguez-Lluhi, J.A., Simon, M.I., Robishaw, J.D., Gilman, A.G. (1992) G protein $\beta\gamma$ subunits synthesized in Sf9 cells: Functional characterization and the significance of prenylation of γ . *J. Biol. Chem.*, **267**, 23409-23417.
- Jan, L., Jan, Y. (1997) Receptor-regulated ion channels. *Curr. Opin. Cell Biol.*, **9**, 155-160.
- Janmey, P.A., Lamb, J., Allen, P.G., Matsudaira, P.T. (1992) Phosphoinositide-binding peptides derived from the sequences of gelsolin and villin. *J. Biol. Chem.*, **267**, 11818-11823.
- Jelsema, C.L., Axelrod, J. (1987) Stimulation of phospholipase A $_2$ activity in bovine rod outer segments by the $\beta\gamma$ subunits of transducin and its inhibition by the α subunit. *Proc. Natl. Acad. Sci. U S A*, **84**, 3623-3627.

- Jensen, O.N., Shevchenko, A., Mann, M. (1997) Protein analysis by mass spectrometry. In: Creighton, T.E. (Hrsg.): Protein structure – A practical approach. IRL Press, 2.Auflage, 1997.
- Jeong, S.W., Ikeda, S.R. (2000) Effect of G protein heterotrimer composition on coupling of neurotransmitter receptors to N-type Ca²⁺ channel modulation in sympathetic neurons. *Proc. Natl. Acad. Sci. U S A*, **97**, 907-912.
- Ji, T.H., Grossmann, M., Ji, I. (1998) G protein-coupled receptors. I. Diversity of receptor-ligand interactions. *J. Biol. Chem.*, **273**, 17299-17302.
- Jiang, H., Wu, D., Simon, M.I. (1994) Activation of phospholipase C- β_4 by heterotrimeric GTP-binding proteins. *J. Biol. Chem.*, **269**, 7593-7596.
- Jiang, H., Kuang, Y., Wu, Y., Smrcka, A., Simon, M.I., Wu, D. (1996) Pertussis toxin-sensitive activation of phospholipase C by the C5a and fMet-Leu-Phe receptors. *J. Biol. Chem.*, **271**, 13430-13434.
- Johnson, G.L., Gardner, A.M., Lange-Carter, C., Qian, N.X., Russell, M., Winitz, S. (1994) How does the G protein, G_{i2}, transduce mitogenic signals? *J. Cell. Biochem.*, **54**, 415-422.
- Jones, D.T., Reed, R.R. (1989) G_{olf}: An olfactory neuron specific-G protein involved in odorant signal transduction. *Science*, **244**, 790-795.
- Jones, P.G., Lombardi, S.J., Cockett, M.I. (1998) Cloning and tissue distribution of the human G protein β_5 cDNA. *Biochim. Biophys. Acta*, **1402**, 288-291.
- Jones, M.B., Garrison, J.C. (1999) Instability of the G protein β_5 subunit in detergent. *Anal. Biochem.* **268**, 126-133.
- Kalkbrenner, F., Dippel, E., Wittig, B., Schultz, G. (1996) Specificity of interaction between receptor and G protein: Use of antisense techniques to relate G protein subunits to function. *Biochim. Biophys. Acta*, **1314**, 125-139.
- Katada, T. und Ui, M. (1982) ADP ribosylation of the specific membrane protein of C6 cells by islet-activating protein associated with modification of adenylate cyclase activity. *J. Biol. Chem.*, **257**, 7210-7216.
- Katz, A., Wu, D.Q., Simon, M.I. (1993) Subunits $\beta\gamma$ of heterotrimeric G protein activate β_2 isoform of phospholipase C, *Nature*, **360**, 686-689.
- Kaupp, U.B., Koch, K.-W. (1992) Role of cGMP and Ca²⁺ in vertebrate photoreceptor excitation and adaptation. *Annu. Rev. Physiol.*, **54**, 153-175.
- Kaziro, Y., Itoh, H., Kozasa, T., Nakafuku, M., Satoh, T. (1991) Structure and function of signal-transducing GTP-binding proteins. *Annu. Rev. Biochem.*, **60**, 349-400.
- Keely, P.J., Westwick, I.P., Whitehead, I.P., Der, C.J., Parise, L.V. (1997) Cdc42 and Rac1 induce integrin-mediated cell motility and invasiveness through PI(3)K. *Nature*, **390**, 632-636.
- Kisselev, O.G., Ermolaeva, M.V., Gautam, N. (1994) A farnesylated domain in the G protein γ subunit is a specific determinant of receptor coupling. *J. Biol. Chem.*, **269**, 21399-21402.

- Kisselev, O.G., Pronin, A., Ermolaeva, M.V., Gautam, N. (1995) Receptor-G protein coupling is established by a potential conformational switch in the $\beta\gamma$ complex. *Proc. Natl. Acad. Sci. U S A*, **90**, 9102-9106.
- Klippel, A., Reinhard, C., Kavanaugh, W.M., Apell, G., Escobedo, M.A., Williams, L.T. (1996) Membrane localization of phosphatidylinositol 3-kinase is sufficient to activate multiple signal-transducing kinase pathways. *Mol. Cell. Biol.*, **16**, 4117-4127.
- Kotzyba-Hilbert, F., Grutter, T., Goeldner, M. (1999) Molecular investigations on the nicotinic acetylcholine receptor: conformational mapping and dynamic exploration using photoaffinity labeling. *Mol. Neurobiol.*, **20**, 45-59.
- Kovoor, A., Chen, C.K., He, W., Wensel, T.G., Simon, M.I., Lester, H.A. (2000) Co-expression of $G\beta_5$ enhances the function of two $G\gamma$ subunit-like domain-containing regulators of G protein signaling proteins. *J. Biol. Chem.*, **275**, 3397-3402.
- Kozasa, T., Hepler, J.R., Smrcka, A.V., Simon, M.I., Rhee, S.G., Sternweis, P.C., Gilman, A.G. (1993) Purification from Sf9 cells and characterization of recombinant $G\alpha_{16}$. Activation of purified phospholipase C isozymes by $G\alpha$ subunits. *Proc. Natl. Acad. Sci. U S A*, **90**, 9176-9180.
- Kozasa, T., Gilman, A.G. (1995) Purification of recombinant G proteins from Sf9 cells by hexahistidine tagging of associated subunits. Characterization of α_{12} and inhibition of adenylyl cyclase by α_z . *J. Biol. Chem.*, **270**, 1734-1741.
- Krugmann, S., Hawkins, P.T., Pryer, N., Braselmann, S. (1999) Characterizing the interactions between the two subunits of the p101/p110 γ phosphoinositide 3-kinase and their role in the activation of this enzyme by $G\beta\gamma$ subunits. *J. Biol. Chem.*, **274**, 17152-17158.
- Kuang, Y., Wu, Y., Smrcka, A., Jiang, H., Wu, D. (1996) Identification of a phospholipase C β_2 region that interacts with $G\beta\gamma$. *Proc. Natl. Acad. Sci. U S A*, **93**, 2964-2968.
- Kurachi, Y., Nakajima, T., Sugimoto, T. (1986) Acetylcholine activation of potassium channels in cell-free membrane patches of atrial cells. *Am. J. Physiol.*, **251**, H681-H684.
- Kurosu, H., Maehama, T., Okada, T., Yamamoto, T., Hoshino, S., Fukui, Y., Ui, M., Hazeki, O., Katada, T. (1997) Heterodimeric phosphoinositide 3-kinase consisting of p85 and p110 β is synergistically activated by the $\beta\gamma$ subunits of G proteins and phosphotyrosyl peptide. *J. Biol. Chem.*, **272**, 24252-24262.
- Lam, K., Carpenter, C.L., Ruderman, N.B., Friel, J.C., Kelly, K.L. (1994) The phosphatidylinositol 3-kinase serine kinase phosphorylates IRS-1. Stimulation by insulin and inhibition by Wortmannin. *J. Biol. Chem.*, **269**, 20648-20652.
- Lämmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, **227**, 680-685.
- Langhans-Rajasekaran, S.A., Wan, Y., Huang, X.Y. (1995) Activation of Tsk and Btk tyrosin kinases by G protein $\beta\gamma$ subunits. *Proc. Natl. Acad. Sci. U S A*, **92**, 8601-8605.

- Lee, C., Murakami, T., Simonds, W.F. (1995a) Identification of a discrete region of the G protein γ subunit conferring selectivity in $\beta\gamma$ complex formation. *J. Biol. Chem.*, **270**, 8779-8784.
- Lee, S.B., Rhee, S.G. (1995b) Significance of PIP₂ hydrolysis and regulation of phospholipase C isoenzymes. *Curr. Opin. Cell Biol.*, **7**, 183-189.
- Leevers, S.J., Vanhaesebroeck, B., Waterfield, M.D. (1999) Signalling through phosphoinositide 3-kinases: the lipids take centre stage. *Curr. Opin. Cell Biol.*, **2**, 219-225.
- Leopoldt, D., Harteneck, C., Nürnberg, B. (1997) G proteins endogenously expressed in Sf9 cells: interactions with mammalian histamine receptors. *Naunyn-Schmiedeberg's Arch. Pharmacol.*, **350**, 329-338.
- Leopoldt, D., Hanck, T., Exner, T., Maier, U., Wetzker, R., Nürnberg, B. (1998) G $\beta\gamma$ stimulates phosphoinositide 3-kinase γ by direct interaction with two domains of the catalytic p110 subunit. *J. Biol. Chem.*, **273**, 7024-7029.
- Levay, K., Cabrera, J.L., Satpaev, D.K., Slepak, V.Z. (1999) G β_5 prevents the RGS7-G α_0 interaction through binding to a distinct G γ -like domain found in RGS7 and other RGS proteins. *Proc. Natl. Acad. Sci. U S A*, **96**, 2503-2507.
- Li, Y., Sternweis, P.M., Charnecki, S., Smith, T.F., Gilman, A.G., Neer, E.J., Kozasa, T. (1998) Sites for G α binding on the G protein β subunit overlap with sites for regulation of phospholipase C β and adenylyl cyclase. *J. Biol. Chem.*, **273**, 16265-16272.
- Li, Z., Jiang, H., Xie, W., Zhang, Z., Smrcka, A.V., Wu, D. (2000) Roles of PLC- β_2 and - β_3 and PI3K γ in chemoattractant-mediated signal transduction. *Science*, **287**, 1046-1049.
- Lindorfer, M.A., Myung, C.S., Savino, Y., Yasuda, H., Khazan, R., Garrison, J.C. (1998) Differential activity of the G protein $\beta_5\gamma_2$ subunit at receptors and effectors. *J. Biol. Chem.*, **273**, 34429-34436.
- Loew, A., Ho, Y.K., Blundell, T., Bax, B. (1998) Phosducin induces a structural change in transducin $\beta\gamma$. *Structure*, **6**, 1007-1019.
- Logothetis, D.E., Kurachi, Y., Galper, J., Neer, E.J., Clapham, D.E. (1987) The $\beta\gamma$ subunits of GTP-binding proteins activate the muscarinic K⁺ channel in heart. *Nature*, **325**, 321-326.
- Lopez-Illasaca, M., Crespo, P., Pellici, P.G., Gutkind, J.S., Wetzker, R. (1997) Linkage of G-Protein coupled receptors to the MAPK signaling pathway through PI 3-Kinase γ . *Science*, **275**, 394-397.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., Randall, R.J. (1951) Protein measurement with the folin phenol reagent. *J. Biol. Chem.*, **193**, 265-275.
- Lu, P.-J., Chen, C.-S. (1997) Selective recognition of phosphatidylinositol 3,4,5-trisphosphate by a synthetic peptide. *J. Biol. Chem.*, **272**, 466-472.

- Luttrell, L.M., Hawes, B.E., van Biesen, T., Luttrell, D.K., Lansing, T.J., Lefkowitz, R.J. (1996) Role of c-Src tyrosin kinase in G protein coupled receptor- and G $\beta\gamma$ subunit-mediated activation of mitogen-activated protein kinases. *J. Biol. Chem.*, **271**, 19443-19450.
- Luttrell, L.M., Daaka, Y., Della, Rocca. G.J., Lefkowitz, R.J. (1997) G protein-coupled receptors mediate two functionally distinct pathways of tyrosine phosphorylation in rat 1a fibroblasts. Shc phosphorylation and receptor endocytosis correlate with activation of Erk kinases. *J. Biol. Chem.*, **272**, 31648-31656.
- Luttrell, L.M., Daaka, Y., Lefkowitz, R. (1999) Regulation of tyrosine kinase cascades by G-protein-coupled receptors. *Curr. Opin. Cell. Biol.*, **11**, 177-183.
- Ma, A.D., Metjian, A., Bagrodia, S., Taylor, S., Abrams, C.S. (1998) Cytoskeletal reorganization by G protein-coupled receptors is dependent on phosphoinositide 3-kinase γ , a Rac guanine exchange factor, and Rac. *Mol. Cell. Biol.*, **18**, 4744-4751.
- MacDermott, A.B., Role, L.W., Siegelbaum, S.A. (1999) Presynaptic ionotropic receptors and the control of transmitter release. *Annu. Rev. Neurosci.*, **22**, 443-485.
- MacDougall, L.K., Domin, M.D., Waterfield, M.D. (1995) A family of phosphoinositide 3-kinases in Drosophila identifies a new mediator of signal transduction. *Curr. Biol.*, **5**, 1404-1415.
- Macrez, N., Morel, J.L., Kalkbrenner, F., Viard, P., Schultz, G., Mironneau, J (1997) A $\beta\gamma$ dimer derived from G₁₃ transduces the angiotensin AT₁ receptor signal to stimulation of Ca²⁺ channels in rat portal vein myocytes. *J. Biol. Chem.*, **272**, 23180-23185.
- Macrez-Lepretre, N., Kalkbrenner, F., Morel, J.L., Schultz, G., Mironneau, J. (1997) G protein heterotrimer G $\alpha_{13}\beta_1\gamma_3$ couples the angiotensin AT_{1A} receptor to increases in cytoplasmic Ca²⁺ in rat portal vein myocytes. *J. Biol. Chem.*, **272**, 10095-10102.
- Maier, U., Babich, A., Nürnberg, B. (1999) Roles of non-catalytic subunits in G $\beta\gamma$ induced activation of class I phosphoinositide 3-kinase isoforms β and γ . *J. Biol. Chem.*, **274**, 29311-29317.
- Maier, U., Babich, A., Macrez, N., Leopoldt, D., Gierschik, P., Illenberger, D. und Nürnberg, B. (2000) G $\beta_5\gamma_2$ is a highly selective activator of phospholipid-dependent enzymes. *J. Biol. Chem.*, **275**, 13746-13754.
- Matsuda, T., Hashimoto, Y., Ueda, H., Asano, T., Matsuura, Y., Doi, T., Takao, T., Shimonishi, Y., Fukada, Y. (1998) Specific isoprenyl group linked to transducin γ -subunit is a determinant of its unique signaling properties among G-proteins. *Biochemistry*, **37**, 9843-9850.
- Mattingly, R.R., Macara, I.G. (1996) Phosphorylation-dependent activation of the Ras-GRF/CDC25Mm exchange factor by muscarinic receptors and G-protein $\beta\gamma$ subunits. *Nature*, **382**, 268-272.
- Meister, M., Dietrich, A., Gierschik, P. (1995) Identification of a three-amino-acid region in G protein γ_1 as a determinant of selective $\beta\gamma$ heterodimerization. *Eur. J. Biochem.*, **234**, 171-177.

- Metjian, A., Roll, R.L., Ma, A.D., Abrams, C.S. (1999) Agonists cause nuclear translocation of phosphatidylinositol 3-kinase γ . *J. Biol. Chem.*, **274**, 27943-27947.
- Misawa, H., Ohtsubo, M., Copeland, N.G., Gilbert, D.J., Jenkins, N.A., Yoshimura, A. (1998) Cloning and characterization of a novel class II phosphoinositide 3-kinase containing C2 domain. *Biochem. Biophys. Res. Commun.*, **244**, 531-539.
- Molendijk, A.J., Irvine, R.F. (1998) Inositide signalling in Chlamydomonas: characterization of a phosphatidylinositol 3-kinase gene. *Plant Mol. Biol.*, **37**, 53-66.
- Murga, C., Laguinge, L., Wetzker, R., Cuadrado, A., Gutkind, J.S. (1998) Activation of Akt/protein kinase B by G protein-coupled receptors. A role for α and $\beta\gamma$ subunits of heterotrimeric G proteins acting through phosphatidylinositol-3-OH kinase γ . *J. Biol. Chem.*, **273**, 19080-19085.
- Myung, C.S., Yasuda, H., Liu, W.W., Harden, T.K., Garrison, J.C. (1999) Role of isoprenoid lipids on the heterotrimeric G protein γ subunit in determining effector activation. *J. Biol. Chem.*, **274**, 16595-16603.
- Nakanishi, S., Yano, H., Matsuda, Y. (1995) Novel functions of phosphatidylinositol-3-kinase in terminally differentiated cells. *Cell. Signal.*, **7**, 545-557.
- Nolte, R.T., Eck, M.J., Schlessinger, J., Shoelson, S.E., Harrison, S.C. (1996) Crystal structure of the PI 3-kinase p85 amino-terminal SH2 domain and its phosphopeptide complexes. *Nat. Struct. Biol.*, **3**, 364-374.
- Nürnberg, B., Spicher, K., Harhammer, R., Bosserhoff, A., Frank, R., Hilz, H., Schultz, G. (1994) Purification of a novel G protein α_o -subtype from mammalian brain. *Biochem. J.*, **300**, 387-394.
- Nürnberg, B., Gudermann, T., Schultz, G. (1995) Receptors and G proteins as primary components of transmembrane signal transduction. Part 2. G proteins: structure and function. *J. Mol. Med.*, **73**, 123-132. *Corr.* **73**, 379.
- Nürnberg, B. (1997) Pertussis toxin as a cell biology tool. In: Aktories, K. (Hrsg.): Bacterial toxins, tools in cell biology. Chapman & Hall, Weinheim, 47-60.
- Nürnberg, B. (2000) Pertussis toxin as a pharmacological tool. In: Aktories K., Just I., (Hrsg.): Handbook of Experimental Pharmacology, **145**, Bacterial Protein Toxins. Springer Verlag, Berlin, Heidelberg, S. 187-206.
- O'Reilly, D.R., Miller, L.K., Luckow, V.A. (1992) Baculovirus expression vectors. WH Freeman, New York.
- Okada, T., Hazeki, O., Ui, M., Katada, T. (1996) Synergistic activation of PtdIns 3-kinase by tyrosine-phosphorylated peptide and $\beta\gamma$ -subunits of GTP-binding proteins. *Biochem. J.*, **317**, 475-480.
- Panchenko, M.P., Saxena, K., Li, Y., Charnecki, S., Sternweis, P.M., Smith, T.F., Gilman, A.G., Kozasa, T., Neer, E.J. (1998) Sites important for PLC β_2 activation by the G protein $\beta\gamma$ subunit map to the sides of the β propeller structure. *J. Biol. Chem.*, **273**, 28298-28304.

- Parish, C.A., Smrcka, A.V., Rando, R.R. (1995) Functional significance of $\beta\gamma$ -subunit carboxymethylation for the activation of phospholipase C and phosphoinositide 3-kinase. *Biochemistry*, **34**, 7722-7727.
- Park, D., Jhon, D.Y., Lee, C.W., Lee, K.H., Rhee, S.G. (1993) Activation of phospholipase C isozymes by G protein $\beta\gamma$ subunits. *J. Biol. Chem.*, **268**, 4573-4576.
- Parker, M.G. (1993) Steroid and related receptors. *Curr. Opin. Cell. Biol.*, **3**, 499-504.
- Paulssen, R.H., Woodson, J., Liu, Z., Ros, E.M. (1996) Carboxyl-terminal fragments of phospholipase C- β_1 with intrinsic G_q GTPase-activating protein (GAP) activity. *J. Biol. Chem.*, **271**, 26622-26629.
- Peterson, G.L. (1983) Determination of total protein. *Meth. Enzymol.*, **91**, 95-119.
- Pitcher, J.A., Touhara, K., Payne, E.S., Lefkowitz, R.J. (1995) Pleckstrin homology domain-mediated membrane association and activation of the β -adrenergic receptor kinase requires coordinate interaction with $G\beta\gamma$ subunits and lipid. *J. Biol. Chem.*, **270**, 11707-11710.
- Pitcher, J.A., Freedman, N.J., Lefkowitz, R.J. (1998) G protein-coupled receptor kinases. *Annu. Rev. Biochem.*, **67**, 653-692.
- Posner, B.A., Gilman, A.G., Harris B.A. (1999) Regulators of G protein signaling 6 and 7. Purification of complexes with $G\beta_5$ and assessment of their effects on G protein-mediated signaling pathways. *J. Biol. Chem.*, **274**, 31087-31093.
- Possee, L.A., Possee, R.D. (1992) The baculovirus expression system: A laboratory guide. Chapman & Hall.
- Prasad, S.V.N., Esposito, G., Mao, L., Koch, W.J., Rockman, H.A. (2000) $G\beta\gamma$ -dependent phosphoinositide 3-kinase activation in hearts with in vivo pressure overload hypertrophy. *J. Biol. Chem.*, **275**, 4693-4698.
- Pumiglia, K.M., LeVine, H., Haske, T., Habib, T., Jove, R., Decker, S.J. (1995) A direct interaction between G protein $\beta\gamma$ subunits and the Raf-1 protein kinase. *J. Biol. Chem.*, **270**, 14251-14254.
- Rameh, L.E., Cantley, L.C. (1999) The role of phosphoinositide 3-kinase lipid products in cell function. *J. Biol. Chem.*, **274**, 8347-8350.
- Ray, K., Ganguly, R. (1992) The Drosophila G protein γ subunit gene (D- $G\gamma_1$) produces three developmentally regulated transcripts and is predominantly expressed in the central nervous system. *J. Biol. Chem.*, **267**, 6086-6092.
- Ray, K., Kunsch, C., Bonner, L.M., Robishaw, J.D. (1995) Isolation of cDNA clones encoding eight different human G protein γ subunits, including three novel forms designated the γ_4 , γ_{10} , and γ_{11} subunits. *J. Biol. Chem.*, **270**, 21765-21771.
- Rebecchi, M.J., Scarlata, S. (1998) Pleckstrin homology domains: a common fold with diverse functions. *Annu. Rev. Biophys Biomol. Struct.*, **27**, 503-528.
- Roche, S., Downward, J., Raynal, P., Courtneidge, S.A. (1998) A function for phosphatidylinositol 3-kinase β (p85 α -p110 β) in fibroblasts during mitogenesis: requirement for insulin- and lysophosphatidic acid-mediated signal transduction. *Mol. Cell. Biol.*, **18**, 7119-7129.

- Rodriguez-Viciano, P., Warne, P.H., Dhand, R., Vanhaesebroeck, B., Gout., I., Fry, M.J., Waterfield, M.D., Downward, J. (1994) Phosphatidylinositol-3-OH kinase as a direct target of Ras. *Nature*, **370**, 527-532.
- Rosenberg, S.J., Rane, M.J., Dean, W.L., Corpier, C.L., Hoffman, J.L., McLeish, K.R. (1998) Effect of γ subunit carboxyl methylation on the interaction of G protein α subunits with $\beta\gamma$ subunits of defined composition. *Cell. Signal.*, **10**, 131-136.
- Rubio, I., Wittig, U., Meyer, C., Heinze, R., Kadereit, D., Waldmann, H., Downward, J., Wetzker, R. (1999) Farnesylation of Ras is important for the interaction with phosphoinositide 3-kinase γ . *Eur. J. Biochem.*, **266**, 70-82.
- Ruiz-Velasco, V., Ikeda, S.R. (2000) Multiple G-protein $\beta\gamma$ combinations produce voltage-dependent inhibition of N-type calcium channels in rat superior cervical ganglion. *J. Neurosci.*, **20**, 2183-2191.
- Sankaran, B., Osterhout, J., Wu, D., Smrcka, A.V. (1998) Identification of a structural element in phospholipase C β_2 that interacts with G protein $\beta\gamma$ subunits. *J. Biol. Chem.*, **273**, 7148-7154.
- Sasaki, T., Irie-Sasaki, J., Jones, R.G., Oliveira-dos-Santos, A.J., Stanford, W.L., Bolon, B., Wakeham, A., Itie, A., Bouchard, D., Kozieradzki, I., Joza, N., Mak, T.W., Ohashi, P.S., Suzuki, A., Penninger, J.M. (2000) Function of PI3K γ in thymocyte development, T cell activation and neutrophil migration. *Science*, **287**, 1040-1046.
- Scharenberg, A.M., El-Hillal, O., Fruman, D.A., Beitz, L.O., Li, Z., Lin, S., Gout, I., Cantley, L.C., Rawlings, D.J., Kinet, J.P. (1998) Phosphatidylinositol-3,4,5-trisphosphate (PtdIns-3,4,5-P₃)/Tec kinase-dependent calcium signaling pathway: a target for SHIP-mediated inhibitory signals. *EMBO J.*, **17**, 1961-1972.
- Schinkmann, K., Blenis, J. (1997) Cloning and characterization of a human STE20-like protein kinase with unusual cofactor requirements. *J. Biol. Chem.*, **272**, 28695-28703.
- Schlicher, U., Linden, L., Calenberg, M., Kreimer, G. (1996) G proteins and Ca²⁺-modulated protein kinases of a plasma membrane-enriched fraction and isolated eyespot apparatuses of *Spermatozopsis similis* (Chlorophyceae). *Eur. J. Phycol.*, **30**, 319-330.
- Schu, P.V., Takegawa, K., Fry, M.J., Stack, J.H., Waterfield, M.D., Emr, S.D. (1993) Phosphatidylinositol 3-kinase encoded by yeast VPS34 gene essential for protein sorting. *Science*, **260**, 88-91.
- Siffert, W., Roskopf, D., Siffert, G., Busch, S., Moritz, A., Erbel, R., Sharma, A.M., Ritz, E., Wichmann, H.-E., Jakobs, K.H., Horsthemke, B. (1998) Association of a human G-protein β_3 subunit variant with hypertension. *Nature Genet.*, **18**, 45-48.
- Simões, A.P., Reed, J., Schnabel, P., Camps, M, Gierschik, P. (1995) Characterization of putative polyphosphoinositide binding motifs from phospholipase C- β_2 . *Biochemistry*, **34**, 5113-5119.
- Simon, M.I., Strathmann, M.P., Gautam, N. (1991) Diversity of G proteins in signal transduction. *Science*, **252**, 802-808.

- Smrcka, A.V., Hepler, J.R., Brown, K.O., Sternweis, P.C. (1991) Regulation of polyphosphoinositide-specific phospholipase C activity by purified G_q. *Science*, **251**, 804-807.
- Smrcka, A.V., Sternweis, P.C. (1993) Regulation of purified subtypes of phosphatidylinositol-specific phospholipase C β by G protein α and $\beta\gamma$ -subunits. *J. Biol. Chem.*, **268**, 9667-9674.
- Snow, B.E., Krumins, A.M., Brothers, G.M., Lee, S.F., Wall, M.A., Chung, S., Mangion, J., Arya, S., Gilman, A.G., Siderovski, D.P. (1998) A G protein γ subunit-like domain shared between RGS11 and other RGS proteins specifies binding to G β_5 subunits. *Proc. Natl. Acad. Sci. U S A*, **95**, 13307-13312.
- Sondek, J., Bohm, A., Lambright, D.G., Hamm, H.E., Sigler, P.B. (1996) Crystal structure of a G-protein $\beta\gamma$ dimer at 2.1 Å resolution. *Nature*, **379**, 369-374. *Korr.* **379**, 847.
- Songyang, Z., Cantley, L.C. (1995) Recognition and specificity in protein tyrosine kinase-mediated signalling. *Trends Biochem. Sci.*, **20**, 470-475.
- Spicher, K., Kalkbrenner, F., Zobel, A., Harhammer, R., Nürnberg, B., Söling, A., Schultz, G. (1994) G₁₂ and G₁₃ α -subunits are immunochemically detectable in membranes of most tissues of various mammalian species. *Biochem. Biophys. Res. Comm.*, **198**, 906-914.
- Spring, D.J., Neer, E.J. (1994) A 14-amino acid region of the G protein γ subunit is sufficient to confer selectivity of γ binding to the β subunit. *J. Biol. Chem.*, **269**, 22882-22886.
- Srinivasan, N., Waterfield, M.D., Blundell, T.L. (1996) Comparative analysis of the regions binding $\beta\gamma$ -subunits in G α and PH domains. *Biochem. Biophys. Res. Comm.*, **220**, 697-702.
- Stack, J.H., De Wald, D.B., Takegawa, K., Emr, S.D. (1995) Vesicle-mediated protein transport: regulatory interactions between the Vps15 protein kinase and the Vps34 PtdIns 3-kinase essential for protein sorting to the vacuole in yeast. *J. Cell Biol.*, **129**, 321-334.
- Stephens, L.R., Hughes, K.T., Irvine, R.F. (1991) Pathway of phosphatidylinositol(3,4,5)-trisphosphate synthesis in activated neutrophils. *Nature*, **351**, 33-39
- Stephens, L., Smrcka, A., Cooke, F.T., Jackson, T.R., Sternweis, P.C., Hawkins, P.T. (1994) A novel phosphoinositide 3 kinase activity in myeloid-derived cells is activated by G protein $\beta\gamma$ subunits. *Cell*, **77**, 83-89.
- Stephens, L.R., Eguinoa, A., Erdjument-Bromage, H., Lui, M., Cooke, F., Coadwell, J., Smrcka, A.S., Thelen, M., Cadwallader, K., Tempst, P., Hawkins, P.T. (1997) The G $\beta\gamma$ sensitivity of a PI3K is dependent upon a tightly associated adaptor, p101. *Cell*, **89**, 105-114.
- Stephens, L.R., Anderson, K., Stokoe, D., Erdjument-Bormage, H., Painter, G.F., Holmes, A.B., Gaffney, P.R., Reese, C.B., McCormick, F., Tempst, P., Coadwell, J., Hawkins, P.T. (1998) Protein kinase B kinases that mediate phosphatidylinositol 3,4,5-trisphosphate-dependent activation of protein kinase B. *Science*, **279**, 710-714.

- Sternweis, P.C. (1986) The purified α subunits of G_o and G_i from bovine brain require $\beta\gamma$ for association with phospholipid vesicles. *J. Biol. Chem.*, **261**, 631-637.
- Sternweis, P.C., Pang, I.-H. (1990). Preparation of G-proteins and their subunits. In: Receptor-effector coupling. A Practical Approach. Chapter I. Ed. E.J. Hulme, S.1-30, Oxford University Press.
- Stokoe, D., Stephens, L.R., Copeland, T., Gaffney, P.R., Reese, C.B., Painter, G.F., Holmes, A.B., McCormick, F., Hawkins, P.T. (1997) Dual role of phosphatidylinositol-3,4,5-trisphosphate in the activation of protein kinase B. *Science*, **277**, 567-570.
- Stoyanov, V., Volinia, S., Hanck, T., Rubio, I., Loubtchenkov, M., Malek, D., Stoyanova, S., Vanhaesebroeck, B., Dhand, R., Nürnberg, B., Gierschik, P., Seedorf, K., Hsuan, J.J., Waterfield, M.D., Wetzker, R. (1995) Cloning and characterization of a G protein-activated human phosphatidylinositol-3 kinase. *Science*, **269**, 690-693.
- Stoyanova, S., Bulgarelli-Leva, G., Kirsch, C., Hanck, T., Klinger, R., Wetzker, R., Wymann, M.P. (1997) Lipid and protein kinase activities of G-protein-coupled phosphoinositide 3-kinase γ : structure-activity analysis and interactions with wortmannin. *Biochem. J.*, **324**, 489-495.
- Strathmann, M.P., Simon, M.I. (1991) $G\alpha_{12}$ and $G\alpha_{13}$ subunits define a fourth class of G protein α subunits. *Proc. Natl. Acad. Sci. U S A*, **88**, 5582-5586.
- Sunahara, R.K., Dessauer, C.W., Gilman, A.G. (1996) Complexity and diversity of mammalian adenylyl cyclases. *Annu. Rev. Pharmacol. Toxicol.*, **36**, 461-480.
- Takahashi, T., Taniguchi, T., Konishi, H., Kikkawa, U., Ishikawa, Y., Yokoyama, M. (1999) Activation of Akt/protein kinase B after stimulation with angiotensin II in vascular smooth muscle cells. *Am. J. Physiol.*, **276**, H1927-H1934.
- Tang, W., Gilman, A.G. (1991) Type-specific regulation of adenylyl cyclase by G protein $\beta\gamma$ subunits. *Science*, **254**, 1500-1503.
- Tang, X., Downes, C.P. (1997) Purification and characterization of $G\beta\gamma$ -responsive phosphoinositide 3-kinases from pig platelet cytosol. *J. Biol. Chem.*, **272**, 14193-14199.
- Taniguchi, T. (1995) Cytokine signalling through nonreceptor protein tyrosin kinases. *Science*, **268**, 251-255.
- Tanti, J.F., Gremeaux, T., Van Obberghen, E., Le Marchand-Brustel, Y. (1994) Insulin receptor substrate 1 is phosphorylated by the serine kinase activity of phosphatidylinositol 3-kinase. *Biochem. J.*, **304**, 17-21.
- Taussig, R., Gilman, A.G. (1995) Mammalian membrane-bound adenylyl cyclases. *J. Biol. Chem.*, **270**, 1-4.
- Taylor, S.S., Knighton, D.R., Zheng, J., Sowadski, J.M., Gibbs, C.S., Zoller, M.J. (1993) A template for the protein kinase family. *Trends Biochem. Sci.*, **18**, 84-89.
- Thomason, P.A., James, S.R., Casey, P.J., Downes, C.P. (1994) A G-protein $\beta\gamma$ -subunit-responsive phosphoinositide 3-kinase activity in human platelet cytosol. *J. Biol. Chem.*, **269**, 16525-16528.

- Tolias, K.F., Cantley, L.C., Carpenter, C.L. (1995) Rho family GTPases bind to phosphoinositide kinases. *J. Biol. Chem.*, **270**, 17656-17659.
- Towbin, H., Staehelin, T., Gordon, J. (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets. *Proc. Natl. Acad. Sci. U S A*, **76**, 4350-4354.
- Traynor-Kaplan, A.E., Harris, A.L., Thompson, B.L., Taylor, P., Sklar, L.A. (1988) An inositol tetrakisphosphate-containing phospholipid in activated neutrophils. *Nature*, **334**, 353-356.
- Traynor-Kaplan, A.E., Thompson, B.L., Harris, A.L., Taylor, P., Sklar, L.A. (1989) Transient increase in phosphatidylinositol 3,4-bisphosphate and phosphatidylinositol trisphosphate during activation of human neutrophils. *J. Biol. Chem.*, **264**, 15668-15673.
- Tsukada, S., Simon, M.I., Witte, O.N., Katz, A. (1994) Binding of $\beta\gamma$ subunits of heterotrimeric G proteins to the PH domain of Bruton tyrosine kinase. *Proc. Natl. Acad. Sci. U S A*, **91**, 11256-11260.
- Turner, S.J., Domin, J., Waterfield, M.D., Ward, S.G., Westwick, J. (1998) The CC chemokine monocyte chemoattractant peptide-1 activates both the class I p85/p110 phosphatidylinositol 3-kinase and the class II PI3K-C2 α . *J. Biol. Chem.*, **273**, 25987-25995.
- Uddin, S., Fish, E.N., Sher, D.A., Gardziola, C., White, M.F., Platanius, L.C. (1997) Activation of the phosphatidylinositol 3-kinase serine kinase by IFN- α . *J. Immunol.*, **158**, 2390-2397.
- Ui, M., Okada, T., Hazeki, K., Hazeki, O. (1995) Wortmannin as a unique probe for an intracellular signalling protein, phosphoinositide 3-kinase. *Trends Biochem. Sci.*, **20**, 303-307.
- Unger, V.M., Hargrave, P.A., Baldwin, J.M., Schertler, G.F.X. (1997) Arrangement of rhodopsin transmembrane α -helices. *Nature*, **389**, 203-206.
- Vanhaesebroeck, B., Higashi, K., Raven, C., Welham, M., Anderson, S., Brennan, P., Ward, S.G., Waterfield, M.D. (1999) Autophosphorylation of p110 δ phosphoinositide 3-kinase: a new paradigm for the regulation of lipid kinases in vitro and in vivo. *EMBO J.*, **18**, 1292-1302.
- Vanhaesebroeck, B., Waterfield, M.D. (1999) Signaling by distinct classes of phosphoinositide 3-kinases. *Experim. Cell Res.*, **253**, 239-254.
- Viard, P., Exner, T., Maier, U., Mironneau, J., Nürnberg, B., Macrez, N. (1999) G $\beta\gamma$ dimers stimulate vascular L-type Ca²⁺ channels via phosphoinositide 3-kinase. *FASEB J.*, **13**, 685-694.
- Virchow, S., Ansorge, N., Roskopf, D., Rübber, H., Siffert, W. (1999) The G protein β_3 subunit splice variant G β_{3-s} causes enhanced chemotaxis of human neutrophils in response to interleukin-8. *Naunyn-Schmiedeberg's Arch. Pharmacol.*, **360**, 27-32.
- Walker, E.H., Perisic, O., Ried, C., Stephens, L., Williams, R.L. (1999) Structural insights into phosphoinositide 3-kinase catalysis and signalling. *Nature*, **402**, 313-320.

- Wang, T., Pentylala, S., Rebecchi, M.J., Scarlata, S. (1999) Differential association of the pleckstrin homology domains of phospholipases-C β_1 , C β_2 , and C δ_1 with lipid bilayers and the $\beta\gamma$ subunits of heterotrimeric G proteins. *Biochemistry*, **38**, 1517-1524.
- Watson, A.J., Katz, A., Simon, M.I. (1994) A fifth member of the mammalian G protein β -subunit family. *J. Biol. Chem.*, **269**, 22150-22156.
- Watson, A.J., Aragay, A.M., Slepak, V.Z., Simon, M.I. (1996) A novel form of the G protein β subunit G β_5 is specifically expressed in the vertebrate retina. *J. Biol. Chem.*, **271**, 28154-28160.
- Whiting, P.J., Bonnert, T.P., McKernan, R.M., Farrar, S., Le Bourdelles, B., Heavens, R.P., Smith, D.W., Hewson, L., Rigby, M.R., Sirinathsinghji, D.J., Thompson, S.A., Wafford, K.A. (1999) Molecular and functional diversity of the expanding GABA_A receptor gene family. *Ann. N Y Acad. Sci.*, **868**, 645-653.
- Wilkie, T.M., Gilbert, D.J., Olsen, A.S., Chen, X.N., Amatruda, T.T., Korenberg, J.R., Trask, B.J., de Jong, P., Reed, R.R., Simon, M.I., Jenkins, N.A., Copeland, N.G. (1992) Evolution of the mammalian G protein α subunit multigene family. *Nature Genet.*, **1**, 85-91.
- Wilson, M., Burt, A.R., Milligan, G., Anderson, N.G. (1996) Wortmannin-sensitive activation of p70^{s6k} by endogenous and heterologously expressed G_i-coupled receptors. *J. Biol. Chem.*, **271**, 8537-8540.
- Wong, G.T., Gannon, K.S., Margolskee, R.F. (1996) Transduction of bitter and sweet taste by gustducin. *Nature*, **381**, 796-800.
- Woscholski, R., Parker, P.J. (1997) Inositol lipid 5-phosphatases – traffic signals and signal traffic. *Trends Biochem. Sci.*, **22**, 427-431.
- Wurmser, A.E., Gary, J.D., Emr, S.D. (1999) Phosphoinositide 3-kinases and their FYVE domain-containing effectors as regulators of vacuolar/lysosomal membrane trafficking pathways. *J. Biol. Chem.*, **274**, 9129-9132.
- Wymann, M.P., Bulgarelli-Leva, G., Zvelebil, M.J., Pirola, L., Vanhaesebroeck, B., Waterfield, M.D., Panayotou, G. (1996) Wortmannin inactivates phosphoinositide 3-kinase by covalent modification of Lys-802, a residue involved in the phosphate transfer reaction. *Mol. Cell. Biol.* **16**, 1722-1733.
- Wymann, M.P., Pirola, L. (1998) Structure and function of phosphoinositide 3-kinases. *Biochim. Biophys. Acta*, **1436**, 127-150.
- Yan, K., Kalyanaraman, V., Gautam, N. (1996) Differential ability to form the G protein $\beta\gamma$ complex among members of the β and γ subunit families. *J. Biol. Chem.*, **271**, 7141-7146.
- Yan, K., Gautam, N. (1997) Structural determinants for interaction with three different effectors on the G protein β subunit. *J. Biol. Chem.*, **272**, 2056-2059.
- Yarfitz, S., Provost, N.M., Hurley, J.B. (1988) Cloning of a drosophila melanogaster guanine nucleotide regulatory protein β -subunit gene and characterization of its expression during development. *Proc. Natl. Acad. Sci. U S A*, **85**, 7134-7138.

- Yasuda, H., Lindorfer, M.A., Woodfork, K.A., Fletcher, J.E., Garrison, J.C. (1996) Role of the prenyl group on the G protein γ subunit in coupling trimeric G proteins to A1 adenosine receptors. *J. Biol. Chem.*, **271**, 18588-18595.
- Yu, F.-X., Sun, H.-Q., Janmey, P.A., Yin, H.L. (1992) Identification of a polyphosphoinositide-binding sequence in an actin monomer-binding domain of gelsolin. *J. Biol. Chem.*, **267**, 14616-14621.
- Yu, J., Wjasow, C., Backer, J.M. (1998a) Regulation of the p85/p110 α phosphatidylinositol 3'-kinase. Distinct roles for the N-terminal and C-terminal SH2 domains. *J. Biol. Chem.*, **273**, 30199-30203.
- Yu, J., Zhang, Y., McIlroy, J., Rordorf-Nikolic, T., Orr, G.A., Backer, J.M. (1998b) Regulation of the p85/p110 phosphatidylinositol 3'-kinase: stabilization and inhibition of the p110 α catalytic subunit by the p85 regulatory subunit. *Mol. Cell. Biol.*, **18**, 1379-1387.
- Zakian, V.A. (1995) ATM-related genes: What do they tell us about functions of the human gene? *Cell*, **82**, 685-687.
- Zhang, J., Zhang, J., Shattil, S.J., Cunningham, M.C., Rittenhouse, S.E. (1996) Phosphoinositide 3-kinase γ and p85/phosphoinositide 3-kinase in platelets. Relative activation by thrombin receptor or β -phorbol myristate acetate and roles in promoting the ligand-binding function of $\alpha_{IIb}\beta_3$ integrin. *J. Biol. Chem.*, **271**, 6265-6572.
- Zhang, S., Coso, O.A., Lee, C., Gutkind, J.S., Simonds, W.F. (1996) Selective activation of effector pathways by brain-specific G protein β_5 . *J. Biol. Chem.*, **271**, 33575-33579.
- Zheng, Y., Bagrodia, S., Cerione, R.A. (1994) Activation of phosphoinositide 3-kinase activity by Cdc42Hs binding to p85. *J. Biol. Chem.*, **269**, 18727-18730.
- Zvelebil, M.J., MacDougall, L., Leever, S., Volinia, S., Vanhaesebroeck, B., Gout, I., Panayotou, G., Domin, J., Stein, R., Pages, F., Koga, H., Salim, K., Linacre, J., Das, P., Panaretou, C., Wetzker, R., Waterfield, M. (1996) Structural and functional diversity of phosphoinositide 3-kinases. *Phil. Trans. R. Soc. Lond. B*, **351**, 217-223.