

11. Literaturverzeichnis

- 1) Effects of enalapril on mortality in severe congestive heart failure. Results of the Cooperative North Scandinavian Enalapril Survival Study (CONSENSUS). The CONSENSUS Trial Study Group. *N Engl J Med* 1987; 316: 1429-35
- 2) Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. The SOLVD Investigators. *N Engl J Med* 1991; 325: 293-302
- 3) Effectiveness of spironolactone added to an angiotensin-converting enzyme inhibitor and a loop diuretic for severe chronic congestive heart failure (the Randomized Aldactone Evaluation Study [RALES]). *Am J Cardiol* 1996; 78: 902-7
- 4) The Cardiac Insufficiency Bisoprolol Study II (CIBIS-II): a randomised trial. *Lancet* 1999; 353: 9-13
- 5) Effect of metoprolol CR/XL in chronic heart failure: Metoprolol CR/XL Randomised Intervention Trial in Congestive Heart Failure (MERIT-HF). *Lancet* 1999; 353: 2001-7
- 6) Anand IS, Fisher LD, Chiang YT, et al. Changes in brain natriuretic peptide and norepinephrine over time and mortality and morbidity in the Valsartan Heart Failure Trial (Val-HeFT). *Circulation* 2003; 107: 1278-83
- 7) Antoni DH, Beuckelmann D, Hambrecht R, et al. Treatment of chronic heart failure with beta-adrenoceptor antagonists. Guideline-based optimization with ERICA, a quality register. *Dtsch Med Wochenschr* 2002; 127: 2459-63
- 8) Artinian NT, Harden JK, Kronenberg MW, et al. Pilot study of a Web-based compliance monitoring device for patients with congestive heart failure. *Heart Lung* 2003; 32: 226-33
- 9) Baer CA, Di Salvo TG, Cail MI, et al. Electronic home monitoring of congestive heart failure patients: design and feasibility. *Congest Heart Fail* 1999; 5: 105-113
- 10) Bauriedel G, Skowasch D, Lüderitz B Die chronische Herzinsuffizienz. *Dtsch Arztebl* 2005; 102: A 592-601
- 11) Benatar D, Bondmass M, Ghitelman J, et al. Outcomes of chronic heart failure. *Arch Intern Med* 2003; 163: 347-52
- 12) Benedict CR, Johnstone DE, Weiner DH, et al. Relation of neurohumoral activation to clinical variables and degree of ventricular dysfunction: a report from the Registry of Studies of Left Ventricular Dysfunction. SOLVD Investigators. *J Am Coll Cardiol* 1994; 23: 1410-20

- 13) Berger R Telematik im Gesundheitswesen - Perspektiven der Telemedizin in Deutschland. Im Auftrag für das Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie und Gesundheitsministerium für Gesundheit. München 1997: <http://hcp.katalog-procurement.de>, Abruf 10.01.2006
- 14) Blue L, Lang E, McMurray JJ, et al. Randomised controlled trial of specialist nurse intervention in heart failure. *Bmj* 2001; 323: 715-8
- 15) Bondmass MD, Benatar JD, Castro G, et al. A Prospective Randomized Study Comparing Outcomes and Outpatient Care Delivery Methods for Chronic Heart Failure. *J Am Coll Cardiol* 2001; 37 Suppl 2: 196A, Abstract 1234-57
- 16) Bondmass MD, Malhotra V, Castro G, et al. The Long Term Effect of a Telemedicine Intervention on Heart Failure Admission and Length of Stay. *J Card Fail* 1999; 5 Suppl 1: 78, Abstract 293
- 17) Brater DC Benefits and risks of torasemide in congestive heart failure and essential hypertension. *Drug Saf* 1996; 14: 104-20
- 18) Capomolla S, Pinna GD, La Rovere MT, et al. Heart failure case disease management program: a pilot study of home telemonitoring versus usual care. *Eur Heart J* 2004; 6 Suppl F: F91-F98
- 19) Cheng V, Kazanagra R, Garcia A, et al. A rapid bedside test for B-type peptide predicts treatment outcomes in patients admitted for decompensated heart failure: a pilot study. *J Am Coll Cardiol* 2001; 37: 386-91
- 20) Chryseogolos ET, Gemme D, Coleman K, et al. Telemonitoring Devices Further Improve Outcomes of a Multidisciplinary Heart Failure Outpatient Program. *J Card Fail* 1999; 5 Suppl 1: 73, Abstract 274
- 21) Cleland JG Health economic consequences of the pharmacological treatment of heart failure. *Eur Heart J* 1998; 19 Suppl P: P32-9
- 22) Cleland JG, Cohen-Solal A, Aguilar JC, et al. Management of heart failure in primary care (the IMPROVEMENT of Heart Failure Programme): an international survey. *Lancet* 2002; 360: 1631-9
- 23) Cleland JG, Louis AA, Rigby AS, et al. Noninvasive home telemonitoring for patients with heart failure at high risk of recurrent admission and death: the Trans-European Network-Home-Care Management System (TEN-HMS) study. *J Am Coll Cardiol* 2005; 45: 1654-64
- 24) Cleland JG, Swedberg K, Follath F, et al. The EuroHeart Failure survey programme - a survey on the quality of care among patients with heart failure in Europe. Part 1: patient characteristics and diagnosis. *Eur Heart J* 2003; 24: 442-63
- 25) Cohn JN, Johnson G, Ziesche S, et al. A comparison of enalapril with hydralazine-isosorbide dinitrate in the treatment of chronic congestive heart failure. *N Engl J Med*

1991; 325: 303-10

- 26) Cohn JN, Tognoni G A randomized trial of the angiotensin-receptor blocker valsartan in chronic heart failure. *N Engl J Med* 2001; 345: 1667-75
- 27) Cordisco ME, Benjaminovitz A, Hammond K, et al. Use of telemonitoring to decrease the rate of hospitalization in patients with severe congestive heart failure. *Am J Cardiol* 1999; 84: 860-2, A8
- 28) Cosin J, Diez J Torasemide in chronic heart failure: results of the TORIC study. *Eur J Heart Fail* 2002; 4: 507-13
- 29) Cowie MR, Mosterd A, Wood DA, et al. The epidemiology of heart failure. *Eur Heart J* 1997; 18: 208-25
- 30) Cowie MR, Wood DA, Coats AJ, et al. Incidence and aetiology of heart failure; a population-based study. *Eur Heart J* 1999; 20: 421-8
- 31) Dao Q, Krishnaswamy P, Kazanegra R, et al. Utility of B-type natriuretic peptide in the diagnosis of congestive heart failure in an urgent-care setting. *J Am Coll Cardiol* 2001; 37: 379-85
- 32) de Lusignan S, Meredith K, Wells S, et al. A controlled pilot study in the use of telemedicine in the community on the management of heart failure - a report of the first three months. *Stud Health Technol Inform* 1999; 64: 126-37
- 33) de Lusignan S, Wells S, Johnson P, et al. Compliance and effectiveness of 1 year's home telemonitoring. The report of a pilot study of patients with chronic heart failure. *Eur J Heart Fail* 2001; 3: 723-30
- 34) Deering M, Baines B, Christianson C, et al. Patients and providers evaluate daily home weight and symptom monitoring for CHF management. *J Card Fail* 2002; 8 Suppl 1: 97, Abstract 361
- 35) Dzau VJ Renal and circulatory mechanisms in congestive heart failure. *Kidney Int* 1987; 31: 1402-15
- 36) Ertle DA, Litmann GI Hospital outpatient disease management initiative achieves marked reduction inpatient admission and costs for congestive heart failure. *J Card Fail* 2002; 9 Suppl 1: 90, Abstract 334
- 37) Ertle DA, Litmann GI Hospital Outpatient Heart Failure Disease Management Initiative Achieves 54 % Reduction in HF Inpatient Admissions and 24 % Reduction in All Cause (Non-HF) Admissions. *J Card Fail* 2003; 9: 111, Abstract 419
- 38) Finkelstein SM, Speedie SM, Demiris G, et al. Telehomecare: quality, perception, satisfaction. *Telemed J E Health* 2004; 10: 122-8

- 39) Flather MD, Shibata MC, Coats AJ, et al. Randomized trial to determine the effect of nebivolol on mortality and cardiovascular hospital admission in elderly patients with heart failure (SENIORS). *Eur Heart J* 2005; 26: 215-25
- 40) Fonarow GC, Stevenson LW, Walden JA, et al. Impact of a comprehensive heart failure management program on hospital readmission and functional status of patients with advanced heart failure. *J Am Coll Cardiol* 1997; 30: 725-32
- 41) Fox KF, Cowie MR, Wood DA, et al. Coronary artery disease as the cause of incident heart failure in the population. *Eur Heart J* 2001; 22: 228-36
- 42) Francis GS, Goldsmith SR, Levine TB, et al. The neurohumoral axis in congestive heart failure. *Ann Intern Med* 1984; 101: 370-7
- 43) Galbreath AD, Krasuski RA, Smith B, et al. Long-term healthcare and cost outcomes of disease management in a large, randomized, community-based population with heart failure. *Circulation* 2004; 110: 3518-26
- 44) Goldberg LR, Piette JD, Walsh MN, et al. Randomized trial of a daily electronic home monitoring system in patients with advanced heart failure: the Weight Monitoring in Heart Failure (WHARF) trial. *Am Heart J* 2003; 146: 705-12
- 45) Gonseth J, Guallar-Castillon P, Banegas JR, et al. The effectiveness of disease management programmes in reducing hospital re-admission in older patients with heart failure: a systematic review and meta-analysis of published reports. *Eur Heart J* 2004; 25: 1570-95
- 46) Grancelli H, Varini S, Ferrante D, et al. Randomized Trial of Telephone Intervention in Chronic Heart Failure (DIAL): study design and preliminary observations. *J Card Fail* 2003; 9: 172-9
- 47) Granger CB, McMurray JJ, Yusuf S, et al. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function intolerant to angiotensin-converting-enzyme inhibitors: the CHARM-Alternative trial. *Lancet* 2003; 362: 772-6
- 48) Hanumanthu S, Butler J, Chomsky D, et al. Effect of a heart failure program on hospitalization frequency and exercise tolerance. *Circulation* 1997; 96: 2842-8
- 49) Heidenreich PA, Ruggerio CM, Massie BM. Effect of a home monitoring system on hospitalization and resource use for patients with heart failure. *Am Heart J* 1999; 138: 633-40
- 50) Hense HW. Epidemiologie der Herzinsuffizienz. *Med Welt* 2004; 55: 374-379
- 51) Hoppe UC, Böhm M, Dietz R, et al. Leitlinien zur Therapie der chronischen Herzinsuffizienz - Herausgegeben vom Vorstand der Deutschen Gesellschaft für Kardiologie-Herz-und Kreislaufforschung e.V. - Bearbeitet im Auftrag der Kommission für klinische Kardiologie - in Zusammenarbeit mit der Arzneimittelkommission der

Deutschen Ärzteschaft. 2005:

- 52) Hunt SA, Baker DW, Chin MH, et al. ACC/AHA guidelines for the evaluation and management of chronic heart failure in the adult: executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to revise the 1995 Guidelines for the Evaluation and Management of Heart Failure). *J Am Coll Cardiol* 2001; 38: 2101-13
- 53) Jäckel A, Schollmayer A, Dudeck J (Hrsg.) Einführung in die Chancen und Voraussetzungen von Telematikanwendungen im Gesundheitswesen. *Telemedizin-führer Deutschland 2000: Ausgabe 2000*: Bad Nauheim:10-13
- 54) Jerant AF, Azari R, Nesbitt TS Reducing the cost of frequent hospital admissions for congestive heart failure: a randomized trial of a home telecare intervention. *Med Care* 2001; 39: 1234-45
- 55) Johnston B, Wheeler L, Deuser J, et al. Outcomes of the Kaiser Permanente Tele-Home Health Research Project. *Arch Fam Med* 2000; 9: 40-5
- 56) Kesinger T, Gilani S, S.H. J Electronic Home Monitoring Reduces Hospiatl Admission, Length of Stay and Readmission Frequency in a Selected Heart Failure Population. *Journal of Cardiac Failure* 2002; 8 Suppl: 94, Abstract 352
- 57) Knox DA, Mueller TM, Vuckovic KM, et al. Remote titration of a beta blocker therapy for heart failure of advanced practice nurse, titration protocols an daily patient telemanagement. *J Card Fail* 2002; 8 Suppl: 83, Abstract 305
- 58) Komajda M, Follath F, Swedberg K, et al. The EuroHeart Failure Survey programme - a survey on the quality of care among patients with heart failure in Europe. Part 2: treatment. *Eur Heart J* 2003; 24: 464-74
- 59) Lapworth DJ, DiBiase A Decreased Hopitalizations Using a Home Based Electronic Weight Monitoring System as a Part of a Comprehensive Heart Failure Program. *J Card Fail* 2000; 6 Suppl: 69, Abstract 258
- 60) Levy D, Kenchaiah S, Larson MG, et al. Long-term trends in the incidence of and survival with heart failure. *N Engl J Med* 2002; 347: 1397-402
- 61) Lloyd-Jones DM, Larson MG, Leip EP, et al. Lifetime risk for developing congestive heart failure: the Framingham Heart Study. *Circulation* 2002; 106: 3068-72
- 62) Louis AA, Turner T, Gretton M, et al. A systematic review of telemonitoring for the management of heart failure. *Eur J Heart Fail* 2003; 5: 583-90
- 63) Macropoulos LR, Selna MJ CHF Hospital Admissions Reduced by 48 % (to 0.272 PPPY at 360 days) in Large Medicare Population Using Advanced Home Monitoring Program. *J Card Fail* 2002; 8 Suppl: 97, Abstract 362
- 64) Maggioni AP, Anand I, Gottlieb SO, et al. Effects of valsartan on morbidity and

mortality in patients with heart failure not receiving angiotensin-converting enzyme inhibitors. J Am Coll Cardiol 2002; 40: 1414-21

- 65) Malek M Health economics of heart failure. Heart 1999; 82 Suppl 4: IV11-3
- 66) Massie BM, West J, Van Ostaeyen D, et al. A controlled trial of heart failure management programs. J Am Coll Cardiol 2001; 37 Suppl 2: A196, Abstract 1234-57
- 67) McMurray JJ, Ostergren J, Swedberg K, et al. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function taking angiotensin-converting-enzyme inhibitors: the CHARM-Added trial. Lancet 2003; 362: 767-71
- 68) McMurray JJ, Petrie MC, Murdoch DR, et al. Clinical epidemiology of heart failure: public and private health burden. Eur Heart J 1998; 19 Suppl P: P9-16
- 69) McMurray JJ, Stewart S Epidemiology, aetiology, and prognosis of heart failure. Heart 2000; 83: 596-602
- 70) Meystre S The current state of telemonitoring: a comment on the literature. Telemed J E Health 2005; 11: 63-9
- 71) Ministerium für Arbeit, Soziales, Gesundheit und Frauen Brandenburg Statistik der stationär behandelten Patienten im Land Brandenburg. Potsdam 2003: http://www.mbj.s.brandenburg.de/media/1336/sgb_2_gender.pdf, Abruf 25.11.2005
- 72) Mortara A, Pinna GD, Capomolla S, et al. A multi-country randomised trial of the role of a new telemonitoring system in CHF: the HHH (Home or Hospital in Heart Failure). Preliminary data. ESC Kongress 2005: Poster 358: <http://www.escardio.org/knowledge/congresses/abol/presentation?id=22806>, Abruf 25.02.2006
- 73) Mortara A, Pinna GD, Johnson P, et al. A multi-country randomised trial of the role of a new telemonitoring system in CHF: the HHH study (Home or Hospoital in Heart Failure). Rational, study design and protocol. Eur Heart J 2004; 6 Suppl F: F99-F102
- 74) Narang R, Cleland JG, Erhardt L, et al. Mode of death in chronic heart failure. A request and proposition for more accurate classification. Eur Heart J 1996; 17: 1390-403
- 75) Osterziel KJ Medikamentöse Therapie der Herzinsuffizienz. Med Welt 2004; 55: 367-373
- 76) Packer M, Coats AJ, Fowler MB, et al. Effect of carvedilol on survival in severe chronic heart failure. N Engl J Med 2001; 344: 1651-8
- 77) Packer M, Fowler MB, Roecker EB, et al. Effect of carvedilol on the morbidity of patients with severe chronic heart failure: results of the carvedilol prospective randomized cumulative survival (COPERNICUS) study. Circulation 2002; 106: 2194-9

- 78) Patterson JH, Adams KF, Jr., Applefeld MM, et al. Oral torsemide in patients with chronic congestive heart failure: effects on body weight, edema, and electrolyte excretion. Torsemide Investigators Group. *Pharmacotherapy* 1994; 14: 514-21
- 79) Pfister R, Scholz M, Wielckens K, et al. The value of natriuretic peptides NT-pro-BNP and BNP for the assessment of left-ventricular volume and function. A prospective study of 150 patients]. *Dtsch Med Wochenschr* 2002; 127: 2605-9
- 80) Pitt B, Remme W, Zannad F, et al. Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 2003; 348: 1309-21
- 81) Pitt B, Zannad F, Remme WJ, et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldactone Evaluation Study Investigators. *N Engl J Med* 1999; 341: 709-17
- 82) Rauchhaus M Pathophysiologie der Herzinsuffizienz. *Med Welt* 2004; 55: 380-385
- 83) Rich MW, Beckham V, Wittenberg C, et al. A multidisciplinary intervention to prevent the readmission of elderly patients with congestive heart failure. *N Engl J Med* 1995; 333: 1190-5
- 84) Richards AM, Troughton R, Lainchbury J, et al. Guiding and monitoring of heart failure therapy with NT-ProBNP: concepts and clinical studies. *J Card Fail* 2005; 11: S34-7
- 85) Rickenbauer P Herzinsuffizienz: Epidemiologie, Pathophysiologie. *Schweiz Med Forum* 2001; 1: 4-9
- 86) Rihal CS, Davis KB, Kennedy JW, et al. The utility of clinical, electrocardiographic, and roentgenographic variables in the prediction of left ventricular function. *Am J Cardiol* 1995; 75: 220-3
- 87) Roger VL, Weston SA, Redfield MM, et al. Trends in heart failure incidence and survival in a community-based population. *Jama* 2004; 292: 344-50
- 88) Roth A, Kajiloti I, Elkayam I, et al. Telecardiology for patients with chronic heart failure: the 'SHL' experience in Israel. *Int J Cardiol* 2004; 97: 49-55
- 89) Ryan KN Results of a clinical-based, automated weight and symptom monitoring pilot program for CHF disease management. *J Card Fail* 2001; 7 Suppl 2: Abstract 370
- 90) Scalfini S, Martinelli G, Baratti D, et al. Telecardiology: one-lead electrocardiogram monitoring and nurse triage in chronic heart failure. *J Telemed Telecare* 2005; 11 Suppl 1: 18-20
- 91) Scalfini S, Zanelli E, Volterrani M, et al. A pilot study of nurse-led, home-based

- telecardiology for patients with chronic heart failure. *J Telemed Telecare* 2004; 10: 113-7
- 92) Shah NB, Der E, Ruggerio C, et al. Prevention of hospitalizations for heart failure with an interactive home monitoring program. *Am Heart J* 1998; 135: 373-8
- 93) Stewart S, Blue L, Walker A, et al. An economic analysis of specialist heart failure nurse management in the UK; can we afford not to implement it? *Eur Heart J* 2002; 23: 1369-78
- 94) Swedberg K, Cleland J, Dargie H, et al. Guidelines for the diagnosis and treatment of chronic heart failure: executive summary (update 2005): The Task Force for the Diagnosis and Treatment of Chronic Heart Failure of the European Society of Cardiology. *Eur Heart J* 2005; 26: 1115-40
- 95) Thompson DR, Roebuck A, Stewart S Effects of a nurse-led, clinic and home-based intervention on recurrent hospital use in chronic heart failure. *Eur J Heart Fail* 2005; 7: 377-84
- 96) Valli N, Georges A, Corcuff JB, et al. Assessment of brain natriuretic peptide in patients with suspected heart failure: comparison with radionuclide ventriculography data. *Clin Chim Acta* 2001; 306: 19-26
- 97) Wang L, Yu CM, Chau E, et al. Feasibility of predicting CHF hospitalization using pacemaker - based impedance sensor in CHF patients. *J Card Fail* 2002; 8 Suppl: 81, Abstract 297
- 98) Werdan K, J. TH, Zerkowski HR (Hrsg.) *Das Herz Buch: Praktische Herz-Kreislauf-Medizin.* 2003: München, Jena: 605-633
- 99) WHO A Health Telematics Policy: Report of the WHO Group Consultation on Health Telematics 11-16 December, Geneva 1997. 1998:
- 100) Willenheimer R, van Veldhuisen DJ, Silke B, et al. Effect on survival and hospitalization of initiating treatment for chronic heart failure with bisoprolol followed by enalapril, as compared with the opposite sequence: results of the randomized Cardiac Insufficiency Bisoprolol Study (CIBIS) III. *Circulation* 2005; 112: 2426-35
- 101) Williams RE, Acker K, Cashy J Prospective, Randomized, Controlled Trial (RCT) of an Automated Daily Heart Failure Telemonitoring Program. *J Card Fail* 2003; 9 Suppl 1: 63, Abstract 227
- 102) Woodend AK, Fraser M, H. S, et al. Readmission and quality of life: the impact of telehome care in heart failure. *J Card Fail* 2002; 8 Suppl 1: 98, Abstract 367
- 103) Zugck C, Nelles M, Frankenstein L, et al. Telemonitoring in chronic heart failure patients - which diagnostic finding prevents hospital readmission? *Herzschr Elektrophys* 2005; 16: 176-82