

5 Literatur

- Adams, M.D., Celniker, S.E., Holt, R.A., et al. (2000) The genome sequence of *Drosophila melanogaster*. *Science*, **287**, 2185-95.
- Afanassiev, V., Hanemann, V. und Wolfl, S. (2000) Preparation of DNA and protein micro arrays on glass slides coated with an agarose film. *Nucleic Acids Res*, **28**, E66.
- Altschul, S.F., Gish, W., Miller, W., et al. (1990) Basic local alignment search tool. *J Mol Biol*, **215**, 403-10.
- Altschul, S.F., Madden, T.L., Schaffer, A.A., et al. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*, **25**, 3389-402.
- Andersen, J.S., Svensson, B. und Roepstorff, P. (1996) Electrospray ionization and matrix assisted laser desorption/ionization mass spectrometry: powerful analytical tools in recombinant protein chemistry. *Nat Biotechnol*, **14**, 449-57.
- Andrade, M.A., Gonzalez-Guzman, M., Serrano, R., et al. (2001) A combination of the F-box motif and kelch repeats defines a large Arabidopsis family of F-box proteins. *Plant Mol Biol*, **46**, 603-14.
- Angenendt, P., Glokler, J., Murphy, D., et al. (2002) Toward optimized antibody microarrays: a comparison of current microarray support materials. *Analytical Biochemistry*, **309**, 253-60.
- Angenendt, P., Glokler, J., Sobek, J., et al. (2003b) Next generation of protein microarray support materials: evaluation for protein and antibody microarray applications. *J Chromatogr A*, **1009**, 97-104.
- Arabidopsis-Genome-Initiative. (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature*, **408**, 796-815.
- Arenkov, P., Kukhtin, A., Gemmell, A., et al. (2000) Protein microchips: use for immunoassay and enzymatic reactions. *Analytical Biochemistry*, **278**, 123-31.
- Asai, T., Tena, G., Plotnikova, J., et al. (2002) MAP kinase signalling cascade in *Arabidopsis* innate immunity. *Nature*, **415**, 977-83.
- Bernard, P. und Couturier, M. (1992) Cell killing by the F plasmid CcdB protein involves poisoning of DNA-topoisomerase II complexes. *J Mol Biol*, **226**, 735-45.
- Bogre, L., Ligterink, W., Heberle-Bors, E., et al. (1996) Mechanosensors in plants. *Nature*, **383**, 489-90.

- Bonfield, J.K. und Staden, R. (1996) Experiment files and their application during large-scale sequencing projects. *DNA Seq*, **6**, 109-17.
- Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*, **72**, 248-54.
- Braun, P., Hu, Y., Shen, B., et al. (2002) Proteome-scale purification of human proteins from bacteria. *Proceedings of the National Academy of Sciences of the United States of America*, **99**, 2654-9.
- Burke, J., Davison, D. und Hide, W. (1999) d2_cluster: a validated method for clustering EST and full-length cDNA sequences. *Genome Res*, **9**, 1135-42.
- Burton, G.R. und McGehee, R.E., Jr. (2004) Identification of candidate genes involved in the regulation of adipocyte differentiation using microarray-based gene expression profiling. *Nutrition*, **20**, 109-14.
- Büssow, K., Cahill, D., Nietfeld, W., et al. (1998) A method for global protein expression and antibody screening on high - density filters of an arrayed cDNA library. *Nucleic Acids Research*, **26**, 5007-5008.
- Büssow, K., Nordhoff, E., Lübbert, C., et al. (2000) A human cDNA library for high-throughput protein expression screening. *Genomics*, **65**, 1-8.
- Byzova, M.V., Franken, J., Aarts, M.G., et al. (1999) Arabidopsis STERILE APETALA, a multifunctional gene regulating inflorescence, flower, and ovule development. *Genes Dev*, **13**, 1002-14.
- C.elegans-Sequencing-Consortium. (1998) Genome sequence of the nematode *C. elegans*: a platform for investigating biology. The *C. elegans* Sequencing Consortium. *Science*, **282**, 2012-8.
- Cahill, D.J. (2001) Protein and antibody arrays and their medical applications. *Journal of Immunological Methods*, **250**, 81-91.
- Cardinale, F., Jonak, C., Ligterink, W., et al. (2000) Differential activation of four specific MAPK pathways by distinct elicitors. *J Biol Chem*, **275**, 36734-40.
- Champion, A., Kreis, M., Mockaitis, K., et al. (2004) Arabidopsis kinome: after the casting. *Funct Integr Genomics*, **4**, 163-87. Epub 2004 Jan 22.
- Chien, C.T., Bartel, P.L., Sternglanz, R., et al. (1991) The two-hybrid system: a method to identify and clone genes for proteins that interact with a protein of interest. *Proc Natl Acad Sci U S A*, **88**, 9578-82.

- Clark, M.D., Panopoulou, G.D., Cahill, D.J., et al. (1999) Construction and analysis of arrayed cDNA libraries [Review]. *Methods in Enzymology*, **303**, 205-233.
- Cordonnier, M.M., Greppin, H. und Pratt, L.H. (1986) Identification of a highly conserved domain on phytochrome from angiosperms to algae. *Plant Physiology*, **80**, 982-987.
- DeRisi, J.L., Iyer, V.R. und Brown, P.O. (1997) Exploring the metabolic and genetic control of gene expression on a genomic scale. *Science*, **278**, 680-686.
- Diezel, W., Kopperschlager, G. und Hofmann, E. (1972) An improved procedure for protein staining in polyacrylamide gels with a new type of Coomassie Brilliant Blue. *Anal Biochem*, **48**, 617-20.
- Dove, S.L. und Hochschild, A. (2004) A bacterial two-hybrid system based on transcription activation. *Methods Mol Biol*, **261**, 231-46.
- Droillard, M.J., Thibivilliers, S., Cazale, A.C., et al. (2000) Protein kinases induced by osmotic stresses and elicitor molecules in tobacco cell suspensions: two crossroad MAP kinases and one osmoregulation-specific protein kinase. *FEBS Lett*, **474**, 217-22.
- Ewing, B. und Green, P. (1998) Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res*, **8**, 186-94.
- Ewing, B., Hillier, L., Wendl, M.C., et al. (1998) Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res*, **8**, 175-85.
- Falsey, J.R., Renil, M., Park, S., et al. (2001) Peptide and small molecule microarray for high throughput cell adhesion and functional assays. *Bioconjug Chem*, **12**, 346-53.
- Fields, S., Kohara, Y. und Lockhart, D.J. (1999) Functional genomics. *Proc Natl Acad Sci U S A*, **96**, 8825-6.
- Fields, S. und Sternglanz, R. (1994) The two-hybrid system: an assay for protein-protein interactions. *Trends Genet*, **10**, 286-92.
- Ge, H. (2000) UPA, a universal protein array system for quantitative detection of protein-protein, protein-DNA, protein-RNA and protein-ligand interactions. *Nucleic Acids Research*, **28**, e3.
- Glinski, M., Romeis, T., Witte, C.P., et al. (2003) Stable isotope labeling of phosphopeptides for multiparallel kinase target analysis and identification of phosphorylation sites. *Rapid Commun Mass Spectrom*, **17**, 1579-84.
- Glöckler, J. und Angenendt, P. (2003) Protein and antibody microarray technology. *J Chromatogr B Analyt Technol Biomed Life Sci*, **797**, 229-40.

- Goff, S.A., Ricke, D., Lan, T.H., et al. (2002) A draft sequence of the rice genome (*Oryza sativa* L. ssp. japonica). *Science*, **296**, 92-100.
- Gómez-Gómez, L. und Boller, T. (2000) FLS2: an LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in Arabidopsis. *Mol Cell*, **5**, 1003-11.
- Guschin, D., Yershov, G., Zaslavsky, A., et al. (1997) Manual manufacturing of oligonucleotide, DNA, and protein microchips. *Anal Biochem*, **250**, 203-11.
- Gutjahr, C. (2003) Generierung und Applikation von Hochdichte-Proteinarrays für die Identifizierung von Autoantigenen bei systemischen Erkrankungen. *Dissertation, Freien Universität Berlin*.
- Haab, B.B., Dunham, M.J. und Brown, P.O. (2001) Protein microarrays for highly parallel detection and quantitation of specific proteins and antibodies in complex solutions. *Genome Biology*, **2**, RESEARCH0004.
- Harashima, S. (1994) Heterologous protein production by yeast host-vector systems. *Bioprocess Technol*, **19**, 137-58.
- Hardie, D.G. (1999) PLANT PROTEIN SERINE/THREONINE KINASES: Classification and Functions. *Annu Rev Plant Physiol Plant Mol Biol*, **50**, 97-131.
- Heß, D. (1991) Pflanzenphysiologie. *UTB für Wissenschaft*, **9. Auflage**.
- Heukeshoven, J. und Dernick, R. (1985) Simplified method for silverstaining of proteins in polyacrylamide gels and the mechanism of silver staining. *Electrophoresis*, **6**, 103-12.
- Heyman, J.A., Cornthwaite, J., Foncerrada, L., et al. (1999) Genome-scale cloning and expression of individual open reading frames using topoisomerase I-mediated ligation. *Genome Research*, **9**, 383-392.
- Heyman, J.A., Cornthwaite, J., Foncerrada, L., et al. (1999) Genome-scale cloning and expression of individual open reading frames using topoisomerase I-mediated ligation. *Genome Research*, **9**, 383-392.
- Holz, C., Lueking, A., Bovekamp, L., Gutjahr, C., Bolotina, N., Lehrach, H., Cahill, D. J., et al. (2001) A human cDNA expression library in yeast enriched for open reading frames. *Genome Research*, **11**; 1730-5.
- Hillenkamp, F. und Karas, M. (1990) Mass spectrometry of peptides and proteins by matrix-assisted ultraviolet laser desorption/ionization. *Methods Enzymol*, **193**, 280-95.
- Hink, M.A., Bisselin, T. und Visser, A.J. (2002) Imaging protein-protein interactions in living cells. *Plant Mol Biol*, **50**, 871-83.

- Houseman, B.T., Huh, J.H., Kron, S.J., et al. (2002) Peptide chips for the quantitative evaluation of protein kinase activity. *Nat Biotechnol*, **20**, 270-4.
- Ichimura, K., Mizoguchi, T., Yoshida, R., et al. (2000) Various abiotic stresses rapidly activate Arabidopsis MAP kinases ATMPK4 and ATMPK6. *Plant J*, **24**, 655-65.
- Jonak, C., Kiegerl, S., Ligterink, W., et al. (1996) Stress signaling in plants: a mitogen-activated protein kinase pathway is activated by cold and drought. *Proc Natl Acad Sci U S A*, **93**, 11274-9.
- Jonak, C., Okresz, L., Bogre, L., et al. (2002) Complexity, cross talk and integration of plant MAP kinase signalling. *Curr Opin Plant Biol*, **5**, 415-24.
- Joos, T.O., Schrenk, M., Hopfl, P., et al. (2000) A microarray enzyme-linked immunosorbent assay for autoimmune diagnostics. *Electrophoresis*, **21**, 2641-50.
- Joung, J.K., Ramm, E.I. und Pabo, C.O. (2000) A bacterial two-hybrid selection system for studying protein-DNA and protein-protein interactions. *Proc Natl Acad Sci U S A*, **97**, 7382-7.
- Kallus, B. (2001) cAMP-abhängige Phosphorylierung der Cytochrom c Oxidase in Rattenhepatozyten. *Dissertation, Heinrich-Heine-Universität Düsseldorf*.
- Kersten, B., Burkle, L., Kuhn, E.J., et al. (2002) Large-scale plant proteomics. *Plant Mol Biol*, **48**, 133-41.
- Kersten, B., Feilner, T., Angenendt, P., et al. (2004a) Proteomic Approaches in Plant Biology. *Current Proteomics*, **1**, 131-144.
- Kersten, B., Feilner, T., Kramer, A., et al. (2003) Generation of Arabidopsis protein chips for antibody and serum screening. *Plant Molecular Biology*, **52**, 999-1010.
- Kersten, B., Possling, A., Blaesing, F., et al. (2004b) Protein microarray technology and ultraviolet crosslinking combined with mass spectrometry for the analysis of protein-DNA interactions. *Anal Biochem*, **331**, 303-13.
- Keyse, S.M. (2000) Protein phosphatases and the regulation of mitogen-activated protein kinase signalling. *Curr Opin Cell Biol*, **12**, 186-92.
- Kimura, M., Yamamoto, Y.Y., Seki, M., et al. (2003) Identification of Arabidopsis genes regulated by high light-stress using cDNA microarray. *Photochem Photobiol*, **77**, 226-33.
- Kingsmore, S.F. und Patel, D.D. (2003) Multiplexed protein profiling on antibody-based microarrays by rolling circle amplification. *Curr Opin Biotechnol*, **14**, 74-81.

- Klose, J. (1975) Protein mapping by combined isoelectric focusing and electrophoresis of mouse tissues. A novel approach to testing for induced point mutations in mammals. *Humangenetik*, **26**, 231-43.
- Klose, J., Nock, C., Herrmann, M., et al. (2002) Genetic analysis of the mouse brain proteome. *Nat Genet*, **30**, 385-93.
- Kramer, A. (2003) Herstellung von Gerste-Protein-Chips mit Proteinen aus cDNA-Expressions-Bibliotheken und Testung ihrer Anwendbarkeit für Phosphorylierungsstudien *in vitro*. *Diplomarbeit, Freie Universität Berlin*.
- Kramer, A., Feilner, T., Possling, A., et al. (2004) Identification of barley CK2alpha targets by using the protein microarray technology. *Phytochemistry*, **65**, 1777-84.
- Kreps, J.A., Wu, Y., Chang, H.S., et al. (2002) Transcriptome changes for Arabidopsis in response to salt, osmotic, and cold stress. *Plant Physiol*, **130**, 2129-41.
- Kroj, T., Rudd, J.J., Nurnberger, T., et al. (2003) Mitogen-activated protein kinases play an essential role in oxidative burst-independent expression of pathogenesis-related genes in parsley. *J Biol Chem*, **278**, 2256-64. Epub 2002 Nov 7.
- Lamb, C. und Dixon, R.A. (1997) The Oxidative Burst in Plant Disease Resistance. *Annu Rev Plant Physiol Plant Mol Biol*, **48**, 251-275.
- Lander, E.S., Linton, L.M., Birren, B., et al. (2001) Initial sequencing and analysis of the human genome. *Nature*, **409**, 860-921.
- Lee, J., Klessig, D.F. und Nurnberger, T. (2001) A harpin binding site in tobacco plasma membranes mediates activation of the pathogenesis-related gene HIN1 independent of extracellular calcium but dependent on mitogen-activated protein kinase activity. *Plant Cell*, **13**, 1079-93.
- Lehrach, H., Bancroft, D. und Maier, E. (1997) Robotics, computing, and biology. *Interdisciplinary Science Review*, **22**, 37-44.
- Lesaicherre, M.L., Uttamchandani, M., Chen, G.Y., et al. (2002) Antibody-based fluorescence detection of kinase activity on a peptide array. *Bioorg Med Chem Lett*, **12**, 2085-8.
- Lizcano, J.M., Deak, M., Morrice, N., et al. (2002) Molecular basis for the substrate specificity of NIMA-related kinase-6 (NEK6). Evidence that NEK6 does not phosphorylate the hydrophobic motif of ribosomal S6 protein kinase and serum- and glucocorticoid-induced protein kinase *in vivo*. *J Biol Chem*, **277**, 27839-49.

- Lu, C., Han, M.H., Guevara-Garcia, A., et al. (2002) Mitogen-activated protein kinase signaling in postgermination arrest of development by abscisic acid. *Proc Natl Acad Sci U S A*, **99**, 15812-7. Epub 2002 Nov 14.
- Lüking, A., Horn, M., Eickhoff, H., et al. (1999) Protein microarrays for gene expression and antibody screening. *Analytical Biochemistry*, **270**, 103-11.
- Lüking, A., Horn, S., Lehrach, H., et al. (2003a) A dual-expression vector allowing expression in *E. coli* and *P. pastoris*, including new modifications. *Methods Mol Biol*, **205**, 31-42.
- Lüking, A., Possling, A., Huber, O., et al. (2003b) A Nonredundant Human Protein Chip for Antibody Screening and Serum Profiling. *Mol Cell Proteomics*, **2**, 1342-1349.
- MacBeath, G. und Schreiber, S.L. (2000) Printing proteins as microarrays for high-throughput function determination. *Science*, **289**, 1760-1763.
- MAPKgroup. (2002) Mitogen-activated protein kinase cascades in plants: a new nomenclature. *Trends Plant Sci*, **7**, 301-8.
- Martinsky, T. (2004) Protein Microarray Manufacturing. *PharmaGenomics*, **3**, 42-48.
- Mayer, K. (2001) DAS ARABIDOPSIS GENOM— Eine Blaupause für die Genome höherer Pflanzen. *GenomXPress*, **1**, 3-5.
- Meri, S. und Baumann, M. (2001) Proteomics: posttranslational modifications, immune responses and current analytical tools. *Biomol Eng*, **18**, 213-20.
- Michaud, G.A., Salcius, M., Zhou, F., et al. (2003) Analyzing antibody specificity with whole proteome microarrays. *Nat Biotechnol*, **21**, 1509-12. Epub 2003 Nov 9.
- Mizoguchi, T., Irie, K., Hirayama, T., et al. (1996) A gene encoding a mitogen-activated protein kinase kinase kinase is induced simultaneously with genes for a mitogen-activated protein kinase and an S6 ribosomal protein kinase by touch, cold, and water stress in *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A*, **93**, 765-9.
- Mundy, J. und Schneitz, K. (2002) Protein phosphorylation in and around signal transduction. *Trends Plant Sci*, **7**, 54-5.
- Neill, S.J., Desikan, R., Clarke, A., et al. (2002) Nitric oxide is a novel component of abscisic acid signaling in stomatal guard cells. *Plant Physiol*, **128**, 13-6.
- Nilsson, J., Stahl, S., Lundeberg, J., et al. (1997) Affinity fusion strategies for detection, purification, and immobilization of recombinant proteins. *Protein Expr Purif*, **11**, 1-16.
- Nühse, T.S., Peck, S.C., Hirt, H., et al. (2000) Microbial elicitors induce activation and dual phosphorylation of the *Arabidopsis thaliana* MAPK 6. *J Biol Chem*, **275**, 7521-6.

- Nürnbergger, T. und Scheel, D. (2001) Signal transmission in the plant immune response. *Trends Plant Sci*, **6**, 372-9.
- O'Farrell, P.H. (1975) High resolution two-dimensional electrophoresis of proteins. *J Biol Chem*, **250**, 4007-21.
- Ouaked, F., Rozhon, W., Lecourieux, D., et al. (2003) A MAPK pathway mediates ethylene signaling in plants. *Embo J*, **22**, 1282-8.
- Pandey, A. und Mann, M. (2000) Proteomics to study genes and genomes. *Nature*, **405**, 837-46.
- Parekh, R.B. und Rohlff, C. (1997) Post-translational modification of proteins and the discovery of new medicine. *Curr Opin Biotechnol*, **8**, 718-23.
- Patton, W.F. (2000) A thousand points of light: the application of fluorescence detection technologies to two-dimensional gel electrophoresis and proteomics. *Electrophoresis*, **21**, 1123-44.
- Perruc, E., Charpentreau, M., Ramirez, B.C., et al. (2004) A novel calmodulin-binding protein functions as a negative regulator of osmotic stress tolerance in *Arabidopsis thaliana* seedlings. *Plant J*, **38**, 410-20.
- Petricoin, E.F., 3rd, Hackett, J.L., Lesko, L.J., et al. (2002a) Medical applications of microarray technologies: a regulatory science perspective. *Nature Genetics*, **32 Suppl**, 474-9.
- Petricoin, E.F., Zoon, K.C., Kohn, E.C., et al. (2002b) Clinical proteomics: translating benchside promise into bedside reality. *Nature Reviews. Drug Discovery*, **1**, 683-95.
- Phizicky, E., Bastiaens, P.I., Zhu, H., et al. (2003) Protein analysis on a proteomic scale. *Nature*, **422**, 208-15.
- Rice, P., Longden, I. und Bleasby, A. (2000) EMBOSS: the European Molecular Biology Open Software Suite. *Trends Genet*, **16**, 276-7.
- Romeis, T., Piedras, P., Zhang, S., et al. (1999) Rapid Avr9- and Cf-9 -dependent activation of MAP kinases in tobacco cell cultures and leaves: convergence of resistance gene, elicitor, wound, and salicylate responses. *Plant Cell*, **11**, 273-87.
- Rubin, G.M., Yandell, M.D., Wortman, J.R., et al. (2000) Comparative genomics of the eukaryotes. *Science*, **287**, 2204-15.
- Sayed, M., Kim, S.O., Salh, B.S., et al. (2000) Stress-induced activation of protein kinase CK2 by direct interaction with p38 mitogen-activated protein kinase. *J Biol Chem*, **275**, 16569-73.

- Schatz, P.J. (1993) Use of peptide libraries to map the substrate specificity of a peptide-modifying enzyme: a 13 residue consensus peptide specifies biotinylation in *Escherichia coli*. *Biotechnology (N Y)*, **11**, 1138-43.
- Scheich, C., Sievert, V. und Bussow, K. (2003) An automated method for high-throughput protein purification applied to a comparison of His-tag and GST-tag affinity chromatography. *BMC Biotechnol*, **3**, 12.
- Schena, M., Shalon, D., Davis, R.W., et al. (1995) Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science*, **270**, 467-70.
- Schweitzer, B., Roberts, S., Grimwade, B., et al. (2002) Multiplexed protein profiling on microarrays by rolling-circle amplification. *Nature Biotechnology*, **20**, 359-65.
- Seki, M., Ishida, J., Narusaka, M., et al. (2002) Monitoring the expression pattern of around 7,000 Arabidopsis genes under ABA treatments using a full-length cDNA microarray. *Funct Integr Genomics*, **2**, 282-91. Epub 2002 Aug 21.
- Snappyan, M., Lecocq, M., Guevel, L., et al. (2003) Dissecting DNA-protein and protein-protein interactions involved in bacterial transcriptional regulation by a sensitive protein array method combining a near-infrared fluorescence detection. *Proteomics*, **3**, 647-57.
- Staden, R., Beal, K.F. und Bonfield, J.K. (2000) The Staden package, 1998. *Methods Mol Biol*, **132**, 115-30.
- Teige, M., Scheickl, E., Eulgem, T., et al. (2004) The MKK2 pathway mediates cold and salt stress signaling in Arabidopsis. *Mol Cell*, **15**, 141-52.
- Templin, M.F., Stoll, D., Schrenk, M., et al. (2002) Protein microarray technology. *Drug Discov Today*, **7**, 815-22.
- Templin, M.F., Stoll, D., Schwenk, J.M., et al. (2003) Protein microarrays: promising tools for proteomic research. *Proteomics*, **3**, 2155-66.
- Tena, G., Asai, T., Chiu, W.L., et al. (2001) Plant mitogen-activated protein kinase signaling cascades. *Curr Opin Plant Biol*, **4**, 392-400.
- Tena, G. und Renaudin, J.P. (1998) Cytosolic acidification but not auxin at physiological concentration is an activator of MAP kinases in tobacco cells. *Plant J*, **16**, 173-82.
- Ulm, R., Ichimura, K., Mizoguchi, T., et al. (2002) Distinct regulation of salinity and genotoxic stress responses by Arabidopsis MAP kinase phosphatase 1. *Embo J*, **21**, 6483-93.
- Verwoerd, T.C., Dekker, B.M. und Hoekema, A. (1989) A small-scale procedure for the rapid isolation of plant RNAs. *Nucleic Acids Res*, **17**, 2362.

- Walhout, A.J., Temple, G.F., Brasch, M.A., et al. (2000) GATEWAY recombinational cloning: application to the cloning of large numbers of open reading frames or ORFeomes. *Methods in Enzymology*, **328**, 575-92.
- Walter, G., Bussow, K., Lueking, A., et al. (2002) High-throughput protein arrays: prospects for molecular diagnostics. *Trends in Molecular Medicine*, **8**, 250-3.
- Wan, J., Zhang, S. und Stacey, G. (2004) Activation of a mitogen-activated protein kinase pathway in Arabidopsis by chitin. *Molecular Plant Pathology*, **5**, 125.
- Willems, A.R., Goh, T., Taylor, L., et al. (1999) SCF ubiquitin protein ligases and phosphorylation-dependent proteolysis. *Philos Trans R Soc Lond B Biol Sci*, **354**, 1533-50.
- Yamagata, H., Saka, K., Tanaka, T., et al. (2001) Light activates a 46-kDa MAP kinase-like protein kinase in soybean cell culture. *FEBS Lett*, **494**, 24-9.
- Yu, J., Hu, S., Wang, J., et al. (2002) A draft sequence of the rice genome (*Oryza sativa* L. ssp. indica). *Science*, **296**, 79-92.
- Zhang, S., Du, H. und Klessig, D.F. (1998) Activation of the tobacco SIP kinase by both a cell wall-derived carbohydrate elicitor and purified proteinaceous elicitors from *Phytophthora* spp. *Plant Cell*, **10**, 435-50.
- Zhang, S. und Klessig, D.F. (2001) MAPK cascades in plant defense signaling. *Trends Plant Sci*, **6**, 520-7.
- Zhu, H., Bilgin, M., Bangham, R., et al. (2001) Global analysis of protein activities using proteome chips. *Science*, **293**, 2101-5.
- Zhu, H., Bilgin, M. und Snyder, M. (2003b) Proteomics. *Annu Rev Biochem*, **72**, 783-812.
- Zhu, H., Klemic, J.F., Chang, S., et al. (2000) Analysis of yeast protein kinases using protein chips. *Nature Genetics*, **26**, 283-289.
- Zhu, H. und Snyder, M. (2003a) Protein chip technology. *Current Opinion in Chemical Biology*, **7**, 55-63.
- Zwerger, K. und Hirt, H. (2001) Recent advances in plant MAP kinase signalling. *Biol Chem*, **382**, 1123-31.