

## 8. Literaturverzeichnis

1. Alward WLM: Color atlas of gonioscopy. Mosby-Verlag, Iowa, 1994, 52-57
2. Anders N, Wollensak J: Biometrische Daten von Augen nach Glaukomanfall. Klin Monatsbl Augenheilk 1992; 201:155-158
3. Bailey IL, Bullimore MA, Raasch TW, Taylor HR: Clinical grading and the effects of scaling. Invest Ophthalmol Vis Sci 1991; 32:422-432
4. Baumann M, Gentile RC, Liebmann JM, Ritch R: Reproducibility of retinal thickness measurements in normal eyes using optical coherence tomography. Ophthalmic Surg Lasers 1998; 29:280-285
5. Bengtsson B: The prevalence of glaucoma. Br J Ophthalmol 1981;65:46-9.
6. Bland JM, Altman DG: Statistical methods for assessing agreement between two methods of clinical measurement. Lancet 1986;I:307-310
7. Böcker T, Sheqem J, Rauwolf M, Wegener A: Anterior chamber angle biometry: A comparison of Scheimpflug photography and ultrasound biomicroscopy. Ophthalmic Res 1995; 27:104-109
8. Broman AT: Biometric gonioscopy and the effects of age, race, and sex on the anterior chamber angle. Br J Ophthalmol 2002; 86:18-22
9. Brosius F: SPSS 11 mitpt Verlag, Bonn 2002
10. Chan R, Smith JA, Richardson KT: Anterior segment configuration correlated with Shaffer's grading of anterior chamber angle. Arch Ophthalmol 1981;99:104-107
11. Chandler PA: Narrow-angle glaucoma. Arch Ophthalmol 1952; 47:695-716
12. Coleman AL, Brigatti L: The glaucomas. Minerva Med 2001; 92:365-379
13. Coleman AL: Glaucoma. Lancet 1999; 354:1803-10
14. Collins JF, Augustin AJ: Augenheilkunde, Springer Verlag 1997, S.251-279
15. Congdon NG, Wang F, Tielsch JM: Issues in the epidemiology and population-based screening of primary angle-closure glaucoma. Surv Ophthalmol 1992;36:411-23
16. Congdon NG, Quigley HA, Huang PT, Wang TH: The impact of age, cataract and visual acuity on wholefield scotopic sensitivity screening for glaucoma in rural Taiwan. Arch Ophthalmol 1995;113:1138-43

17. Congdon NG, Quigley HA, Huang PT, Wang TH: Screening technique for angle closure glaucoma in rural Taiwan. *Acta Ophthalmol Scand* 1996; 74:113-119
18. Congdon NG, Youlin Q, Quigley H, Hung PT, Wang TH, Ho TC, Tielsch JM: Biometry and primary angle-closure glaucoma among chinese, white and black populations. *Ophthalmology* 1997;104:1489-1495
19. Congdon NG, Spaeth GL, Augsburger J, Klancnik J, Ketan Patel, David G: A proposed simple method for measurement in the anterior chamber angle. Biometric gonioscopy. *Ophthalmology* 1999; 106:2161-2167
20. Douthwaite WA, Spence D: Slit-lamp measurement of the anterior chamber depth. *Br J Ophthalmol* 1986;70:205-208
21. Drexler W, Baumgartner A, Findl O, Hitzemberger CK, Sattmann H, Fercher AF: Submicrometer precision biometry of the anterior segment of the human eye. *Invest Ophthalmol Vis Sci* 1997; 38:1304-1313
22. Forbes M: Gonioscopy with corneal indentation. *Arch Ophthalmol* 1966; 76:488-492
23. Foster PJ, Johnson GJ: Glaucoma in China: how big is the problem. *Br J Ophthalmol* 2001; 85:1277-1282
24. Fujimoto JG, Brezinski ME, Tearney GJ, Boppart SA, Bouma B, Hee MR, Southern JF, Swanson EA: Optical biopsy and imaging using optical coherence tomography. *Nature Med* 1995; 1:970-972
25. Grehn F: Diagnostik und Therapie des primären Winkelblockglaukoms. *Fortschr Ophthalmol* 1988;85:437-444
26. Grehn F, Mackensen G: Die Glaukome. Verlag W. Kohlhammer, Stuttgart Berlin Köln 1993 S.36-46
27. Guggenmoos-Holzmann I, Wernecke KD: Medizinische Statistik 1996 Backwell Wissenschafts-Verlag Berlin-Wien: S.160-4
28. Hee MR, Izatt JA, Swanson EA, Huang D, Schumann JS, Fujimoto JG: Optical coherence tomography of the human retina. *Arch Ophthalmol* 1995;113:325-332

29. Hirano K, Ito Y, Suzuki T, Kojima T, Kaji S, Miyake Y: Optical coherence tomography for the noninvasive evaluation of the cornea. *Cornea* 2001; 20:281-289
30. Hoerauf H, Gordes RS, Scholz C, Wirbelauer C, Koch P, Engelhardt R, Winkler J, Laqua H, Birngruber R: First experimental and clinical results with transscleral optical coherence tomography. *Ophthalmic Surg Lasers*. 2000; 31:218-222
31. Hoerauf H, Wirbelauer C, Scholz C, Engelhardt R, Koch P, Laqua H, Birngruber R: Slit-lamp-adapted optical coherence tomography of the anterior segment. *Graefe's Arch Clin Exp Ophthalmol* 2000;238:8-18
32. Hoerauf H, Scholz C, Koch P, Engelhart R, Laqua H, Birngruber R: Transscleral optical coherence tomography. *Arch Ophthalmol* 2002; 120:816-819
33. Hoerauf H, Winkler J, Scholz C, Wirbelauer C, Gordes RS, Koch P, Engelhardt R, Laqua H, Birngruber R: Transscleral optical coherence tomography- An experimental study in ex-vivo human eyes. *Lasers Surg Med* 2002; 30:209-215
34. Huang D, Swanson EA, Lin CP, Schuman JS, Stinson WG, Chang W, Hee MR, Flotte T, Gregory K, Puliafito CA, Fujimoto JG: Optical coherence tomography. *Science* 1991; 254:1178-1181
35. Ishikawa H, Esaki K, Liebermann JM, Uji Y, Ritch R: Ultrasound biomicroscopy dark room provocative testing: A quantitative method for estimating anterior chamber angle width. *Jpn J Ophthalmol* 1999; 43:526-534
36. Izatt JA, Hee MR, Swanson EA, Lin CP, Huang D, Schumann JS, Puliafito CA, Fujimoto JG: Micrometer-scale resolution imaging of the anterior eye in vivo with optical coherence tomography. *Arch Ophthalmol*. 1994;112:1584-1589
37. Kim YY, Jung HR: Clarifying the nomenclature for primary angle-closure glaucoma. *Surv Ophthalmol* 1997; 42:125-136
38. Klemm M, Rumberger E, Walter A, Richard G: Reproduzierbarkeit von Messungen der retinalen Nervenfaserschichtdicke. *Ophthalmologe* 2002; 99:345-351

39. Koop N, Brinkmann R, Lankeau E, Flache S, Engelhardt R, Birngruber R: Optische Kohärenztomographie der Kornea und des vorderen Augenabschnitts. *Ophthalmologie* 1997; 94:481-486
40. Krumpaszky HG, Ludtke R, Mickler A, Klauss V, Selbmann HK: Blindness incidence in Germany. A population-based study from Württemberg-Hohenzollern. *Ophthalmologica* 1999; 213:176-82
41. Liebmann JM, Ritch R: Ultrasound biomicroscopy of the anterior segment. *J Am Optom Assoc* 1996;67:469-479
42. Lowe RF: A history of primary angle-closure glaucoma. *Surv Ophthalmol* 1995;40:163-170
43. Lütjen E, Rohen JW: Histometrische Untersuchungen über die Kammerwinkelregion des menschlichen Auges bei verschiedenen Altersstufen und Glaukomformen. *Graefes Arch Clin Exp Ophthalmol* 1968; 176:1-12
44. Makabe R: Vergleichende Untersuchung der Kammerwinkelweite mit Echographie und Gonioskopie. *Klin Monatsbl Augenheilkd* 1989;194:6-9
45. Marchini G, Pagliaruso A, Toscano A, Tosi R, Brunelli C, Bonomi L: Ultrasound biomicroscopic and conventional ultrasonographic study of ocular dimensions in primary angle-closure glaucoma. *Ophthalmology* 1998; 105:2091-2098
46. Markowitz SN, Morin D: The ratio of lens thickness to axial length for biometric standardization in angle-closure glaucoma. *Am J Ophthalmol* 1985;99:400-402
47. Müller R, Büttner P: A critical discussion of intraclass correlation coefficients. *Stat Med* 1994; 13:1-12
48. Oh YG, Minelli S, Spaeth GL, Steinman WC: The anterior chamber angle is different in different racial groups: a gonioscopic study. *Eye* 1994;8:104-108
49. Pavlin CJ, Sherar MJ, Foster FS: Subsurface ultrasound microscopic imaging of the intact eye. *Ophthalmology* 1990; 97:244-250
50. Pavlin CJ, Harasiewicz K, Sherar MD, Foster FS: Clinical use of ultrasound biomicroscopy. *Ophthalmology* 1991; 98:287-295

51. Pavlin CJ, Harasiewicz K, Foster FS: Ultrasound biomicroscopy of anterior segment structures in normal and glaucomatous eyes. *Am J Ophthalmol* 1992;113:381-389
52. Pavlin CJ, Ritch R, Foster FS: Ultrasound biomicroscopy in plateau iris syndrome. *Am J Ophthalmol* 1992; 113:390-395
53. Pham DT, Wollensak J. Ciliary Block (malignant glaucoma) following posterior chamber lens implantation. *Ophthalmic Surg.* 1987;18:741-744
54. Potash SD, Tello C, Liebmann J, Ritch R: Ultrasound biomicroscopy in pigment dispersion syndrome. *Ophthalmology* 1994; 101:332-339
55. Puliafito CA, Hee MR, Lin CR, Reichel E, Schuman JS, Ducker JS, Izatt JA, Swanson EA, Fujimoto JG: Imaging of macular diseases with optical coherence tomography. *Ophthalmology* 1995; 102:217-219
56. Quigley HA, Nickels RW, Kerrigan LA: Retinal ganglion cell apoptosis. *Invest Ophthal Vis Sci* 1995; 36:774-86
57. Quigley HA: Number of people with glaucoma worldwide. *Br J Ophthalmol* 1996; 80:389-393
58. Radhakrishnan S, Rollins AM, Roth JE, Yazdanfar S, Westphal V, Bardenstein DS, Izatt JA: Real-time optical coherence tomography of the anterior segment at 1310 nm. *Arch Ophthalmol.* 2001; 119:1179-1185
59. Rait J: Seven million too many. *Br J Ophthalmol* 1996; 80:385-386
60. Riley SF, Nairn JP, Maestre FA, Smith TJ: Analysis of the anterior chamber angle by gonioscopy and by ultrasound biomicroscopy. *Int Ophthalmol Clin* 1994; 34:271-282
61. Rohen JW, Lutjen-Drecoll E, Flugel C, Meyer M, Grierson: Ultrastructure of the trabecular meshwork in untreated cases of primary open-angle glaucoma. *Exp Eye Res* 1993; 56:683-92
62. Scheie HG: Width and pigmentation of the angle of the anterior chamber a system of grading by gonioscopy. *Arch Ophthalmol* 1957;4:509-513
63. Schuman JS, Pedut-Kloizman T, Hertzmark E, Hee MR, Wilkins JR, Coker JG, Puliafito CA, Fujimoto JG, Swanson EA: Reproducibility of nerve fiber layer thickness measurements using optical coherence tomography. *Ophthalmology* 1996; 103:1889-1898

64. Schirmer KE: Gonioscopy and artefacts. *Br J Ophthalmol* 1967;51:50-53
65. Shaffer R, Schwartz A: Gonioscopy. *Surv Ophthalmol* 1957;2:389-409
66. Shaffer RN: A suggested anatomic classification to define pupillary block glaucomas. *Invest Ophthalmol Vis Sci* 1973; 12:540-5
67. Shaffer RN: The centennial history of glaucoma (1896-1996) American Academy of Ophthalmology. *Ophthalmology* 1996; 103(suppl8):40-50
68. Shaffer RN, Hoskins HD Jr: Symposium, glaucoma, ciliary block (malignant) glaucoma. *Ophthalmology* 1978; 85:215-220
69. Shields MB, Krieglstein GK: *Glaukom*, Springer Verlag, 1993, 191-210
70. Smith RJH: A new method of estimating the depth of the anterior chamber. *Br J Ophthalmol* 1979; 63:215
71. Sommer A, Tielsch JM, Quigley HA, Gottsch JD, Javitt J, Singh K: Relationship between intraocular pressure and primary open angle glaucoma among white and black Americans. *Arch Ophthalmol* 1991; 109:1090-5
72. Spaeth GL: The normal development of the human anterior chamber angle: a new system of descriptive grading. *Trans Ophthalmol Soc U K* 1971;91:709-739
73. Spaeth GL, Aruajo S, Azuara A: Comparison of the configuration of the human anterior chamber angle, as determined by the Spaeth gonioscopic grading system and ultrasound biomicroscopy. *Trans Am Ophthalmol Soc* 1995;93:337-347
74. Spaeth GL, Azuara-Blanco A, Araujo SV, Augsburger JJ: Intraobserver and interobserver agreement in evaluating the anterior chamber angle configuration by ultrasound biomicroscopy. *J Glaucoma* 1997; 6:13-7
75. Tello C , Liebmann J, Potash SD, Cohen H, Ritch R: Measurement of ultrasound biomicroscopy images: Intraobserver and interobserver reliability. *Invest Ophthalmol Vis Sci* 1994; 35:3549-3552
76. Thylefore B, Negrel AD: The global impact of glaucoma. *Bulletin of World Health Organization* 1994; 72:323-326
77. Unger HH: Zur Kammerwinkelanatomie. *Klin Monatsbl Augenheilkd* 1959; 135:161-183

78. Van Beuningen E: Atlas der Spaltlampen gonioskopie, 1955, Thieme Verlag, Seite 2-19
79. Van Herick W, Shaffer RN, Schwarz A: Estimation of width of angle of anterior chamber, incidence and significance of the narrow angle. *Am J Ophthalmol* 1969; 68:626-629
80. Wiederholt M, Thieme H, Stumpff F: The regulation of trabecular meshwork and ciliary muscle contractility. *Prog Retin Eye Res.* 2000;19:271-95
81. Wirbelauer C, Scholz C, Hoerauf H, Engelhardt R, Birngruber R, Laqua H: Corneal optical coherence tomography before and immediately after excimer laser photorefractive keratectomy. *Am J Ophthalmol* 2000; 130:693-699
82. Wirbelauer C, Scholz C, Hoerauf H, Bastian GO, Engelhardt R, Birngruber R, Laqua H: Untersuchungen der Hornhaut mittels optischer Kohärenztomographie. *Ophthalmologe* 2001; 98:151-156
83. Wirbelauer C, Scholz C, Engelhardt R, Laqua H, Pham DT: Biomorphometrie des Hornhautepithels mittels spaltlampenadaptierter optischer Kohärenztomographie. *Ophthalmologe* 2001; 98:848-852
84. Wirbelauer C, Scholz C, Hoerauf H, Pham DT, Laqua H, Birngruber R: Noncontact corneal pachymetry with slitlamp-adapted optical coherence tomography. *Am J Ophthalmol* 2002; 133:444-450
85. Wirbelauer C, Scholz C, Häberle H, Laqua H, Pham DT: Corneal optical coherence tomography before and after excimer laser phototherapeutic keratectomy for recurrent epithelial erosions. *J Cataract Refract Surg* 2002; 28:1631-1637
86. Wirbelauer C, Winkler J, Bastian GO, Häberle H, Pham DT: Histopathological correlation of corneal diseases with optical coherence tomography. *Graefe's Arch Clin Exp Ophthalmol* 2002; 240:727-734
87. Wirbelauer C, Karandish A, Häberle H, Pham DT: Optical coherence tomography in malignant glaucoma following filtration surgery. *Br J Ophthalmol* 2003; 87:952-955
88. Wirbelauer C, Karandish A, Häberle H, Pham DT: Imaging of scleral expansion bands for presbyopia with optical coherence tomography. *J Cataract Refract Surg* 2003;29:2435-2438

89. Woo EK, Pavlin CJ, Slomovic A, Taback N, Buys YM: Ultrasound biomicroscopic quantitative analysis of light-dark changes associated with pupillary block. *Am J Ophthalmol* 1999; 127:43-47
90. Zadnik K, Mutti DO, Adams AJ: The repeatability of measurement of the ocular components. *Invest Ophthalmol Vis Sci* 1992; 33:2325-2333