

7 Literaturverzeichnis

- Adams JC. Thrombospondins: multifunctional regulators of cell interactions. *Annu Rev Cell Dev Biol*, 2001;17:25-51.
- Asch AS, Silbiger S, Heimer E, Nachman RL. Thrombospondin sequence motif (CSVTCG) is responsible for CD36 binding. *Biochem Biophys Res Commun*, 1992;82(3):1208-17.
- Baenziger NL, Brodie GN, Majerus PW. A thrombin-sensitive protein of human platelet membranes. *Proc Natl Acad Sci USA*, 1971;68:240-9.
- Barakat AI, Davies PF. Mechanisms of shear stress transmission and transduction in endothelial cells. *Chest*, 1998;114(1 Suppl):58S-63S.
- Bartling B, Tostlebe H, Darmer D, Holtz J, Silber RE, Morawietz H. Shear stress-dependent expression of apoptosis-regulating genes in endothelial cells. *Biochem Biophys Res Comm*, 2000;278 (3):740-6
- Bein K, Simons M. Thrombospondin type 1 repeats interact with matrix metalloproteinase 2. Regulation of metalloproteinase activity. *J Biol Chem*, 2000;275:32167-73
- Berk BC, Abe JI, Min W, Surapisitchat J, Yan C. Endothelial atheroprotective and anti-inflammatory mechanisms. *Ann N Y Acad Sci*, 2001;947:93-109.
- Berthiaume F, Frangos JA. Flow-induced prostacyclin production is mediated by a pertussis toxin-sensitive G protein. *FEBS Lett*, 1992;308:272-9
- Bhakdi S. Complement and atherogenesis: the unknown connection. *Ann Med*, 1998;30:503-7.
- Bjarnason JB, Fox JW. Snake venom metalloendopeptidases: reprotolysins. *Methods Enzymol*, 1995;248:345-68.
- Bongrazio M, Baumann C, Zakrzewicz A, Pries AR, Gaehtgens P. Evidence for modulation of genes involved in vascular adaptation by prolonged exposure of endothelial cells to shear stress. *Cardiovasc Res*, 2000;47:384-93.

- Bornstein P, O'Rourke K, Wikstrom K, Wolf FW, Katz R, Li P, Dixit VM. A second, expressed thrombospondin gene (Thbs2) exists in the mouse genome. *J Biol Chem*, 1991;266(20):12821-4.
- Bornstein P, Armstrong LC, Hankenson KD, Kyriakides TR, Yang Z. Thrombospondin 2, a matricellular protein with diverse functions. *Matrix Biol*, 2000;19(7):557-68.
- Braddock M, Schwachtgen JL, Houston P, Dickson M. C, Lee MJ, Campbell CJ. Fluid shear stress modulation of gene expression in endothelial cells. *News Physiol Sci*, 1998;13:241-6
- Brooimans, RA, Hiemstra, PS, van der Ark, AA, Sim, RB, van Es, LA, Daha MR. Biosynthesis of complement factor H by human umbilical vein endothelial cells. Regulation by T cell growth factor and IFN-gamma. *J Immunol*, 1989;142:2024-30.
- Brooks PC, Clark R, Cheresh DA. Requirement of vascular integrin $\alpha_v\beta_3$ for angiogenesis. *Science*, 1994;264:569-71
- Burri PH, Djonov V. Intussusceptive angiogenesis – the alternative to capillary sprouting. *Molecular Aspects of Medicine*, 2002;23:S2-S27.
- Buschmann I, Schaper W. Arteriogenesis versus Angiogenesis: Two mechanisms of vessel growth. *News Physiol Sci*, 1999;14:121-5
- Busse R, Edwards G, Feletou M, Fleming I, Vanhoutte PM, Weston AH. EDHF: bringing the concepts together. *Trends Pharmacol Sci*, 2002;23(8):374-80.
- Bussolari SR, Dewey CF Jr, Gimbrone MA Jr. Apparatus for subjecting living cells to fluid shear stress. *Rev Sci Instrum*, 1982;53(12):1851-4.
- Carmeliet P, Jain RK. Angiogenesis in cancer and other diseases. *Nature*, 2000;407:249-57
- Castle VP, Dixit VM, Polverini PJ. Thrombospondin-1 suppresses tumorigenesis and angiogenesis in serum- and anchorage-independent NIH 3T3 cells. *Lab Invest*, 1997;77(1):51-61.

- Chen H, Sottile J, O'Rourke KM, Dixit VM, Mosher DF. Properties of recombinant mouse thrombospondin 2 expressed in *Spodoptera* cells. *J Biol Chem*, 1994;269(51):32226-32.
- Chen H, Strickland DK, Mosher DF. Metabolism of thrombospondin 2. Binding and degradation by 3t3 cells and glycosaminoglycan-variant Chinese hamster ovary cells. *J Biol Chem*, 1996;271(27):15993-9.
- Chun TH, Itoh H, Ogawa Y, Tamura N, Takaya K, Igaki T, Yamashita J, Doi K, Inoue M, Masatsugu K, Korenaga R, Ando J, Nakao K. Shear stress augments expression of C-type natriuretic peptide and adrenomedullin. *Hypertension*, 1997;29:1296-302
- Colige A, Li SW, Sieron AL, Nusgens BV, Prockop DJ, Lapiere CM. cDNA cloning and expression of bovine procollagen I N-proteinase: a new member of the superfamily of zinc-metalloproteinases with binding sites for cells and other matrix components. *Proc Natl Acad Sci U S A*, 1997;94(6):2374-9.
- Colige A, Sieron AL, Li SW, Schwarze U, Petty E, Wertelecki W, Wilcox W, Krakow D, Cohn DH, Reardon W, Byers PH, Lapiere CM, Prockop DJ, Nusgens BV. Human Ehlers-Danlos syndrome type VII C and bovine dermatosparaxis are caused by mutations in the procollagen I N-proteinase gene. *Am J Hum Genet*, 1999;65(2):308-17.
- Conway EM, Collen D, Carmeliet P. Molecular mechanisms of blood vessel growth. *Cardiovasc Res*, 2001;49:507-21.
- Cowan DB, Langille BL. Cellular and molecular biology of vascular remodeling. *Curr Opin Lipidol*, 1996;7(2):94-100.
- Dameron KM, Volpert OV, Tainsky MA, Bouck N. Control of angiogenesis in fibroblasts by p53 regulation of thrombospondin-1. *Science*, 1994;265:1582-4
- Davies PF, Tripathi SC. Mechanical stress mechanisms and the cell. An endothelial paradigm. *Circ Res*, 1993;72(2):239-45.

- Davies PF Flow-mediated endothelial mechanotransduction. *Physiol Rev*, 1995;75(3):519-60.
- Dawson DW, Pearce SFA, Zhong R, Silverstein RL, Frazier WA, Bouck N. CD36 mediates the in vitro inhibitory effects of thrombospondin-1 on endothelial cells. *J Cell Biol*, 1997;183:707-17
- Dawson DW, Bouck NP. Thrombospondin as an inhibitor of angiogenesis. *Antiangiogenic Agents in Cancer Therapy*, edited by Teicher BA, Totowa NJ. Human Press Inc, 1999;185-203
- Dawson JM, Hudlicka O. The effects of long term administration of prazosin on the microcirculation in skeletal muscles. *Cardiovasc Res*, 1989;23(11):913-20.
- Diamond SL, Sharefkin JB, Dieffenbach C, Frasier-Scott K, McIntire LV, Eskin SG. Tissue plasminogen activator messenger RNA levels increase in cultured human endothelial cells exposed to laminar shear stress. *J Cell Physiol*, 1990;143(2):364-71.
- Dobrin PB, Littooy FN, Endean ED Mechanical factors predisposing to intimal hyperplasia and medial thickening in autogenous vein grafts. *Surgery*, 1989;105(3):393-400.
- Duda DG, Sunamura M, Lozonschi L, Yokoyama T, Yatsuoka T, Motoi F, Horii A, Tani K, Asano S, Nakamura Y, Matsuno S. Overexpression of the p53-inducible brain-specific angiogenesis inhibitor 1 suppresses efficiently tumour angiogenesis. *Br J Cancer*, 2002;86(3):490-6.
- Nerem RM, Alexander RW, Chappell DC, Medford RM, Varner SE, Taylor WR. The study of the influence of flow on vascular endothelial biology. *Am J Med Sci*, 1998;316(3):169-75.
- Egginton S, Zhou A-L, Brown MD, Hudlicka O. Unorthodox angiogenesis in skeletal muscle. *Cardiovasc Res*, 2001;49:634-46

- Febbraio M, Hajjar DP, Silverstein RL. CD36: A class B scavenger receptor involved in angiogenesis, atherosclerosis, inflammation, and lipid metabolism. *J Clin Invest*, 2001;108:785-91
- Ferrara N. Molecular and biological properties of vascular endothelial growth factor. *J Mol Med*. 1999;77(7):527-43.
- Ferrara N. Role of vascular endothelial growth factor in regulation of physiological angiogenesis. *Am J Physiol Cell Physiol*, 2001;280(6):C1358-66.
- Fijen CA, van den Bogaard R, Schipper M, Mannens M, Schlesinger M, Nordin FG, Dankert J, Daha MR, Sjöholm AG, Truedsson L, Kuijper EJ. Properdin deficiency: molecular basis and disease association. *Mol Immunol*, 1999;36, 863-7.
- Forsythe JA, Jiang BH, Iyer NV, Agani F, Leung SW, Koos RD, Semenza GL. Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1. *Mol Cell Biol*, 1996;16(9):4604-13.
- Freyberg MA, Kaiser D, Graf R, Vischer P, Friedl P. Integrin-associated protein and thrombospondin-1 as endothelial mechanosensitive death mediators. *Biochem Biophys Res Commun*, 2000;271(3):584-8.
- Freyberg MA, Kaiser D, Graf R, Buttenbender J, Friedl P. Proatherogenic flow conditions initiate endothelial apoptosis via thrombospondin-1 and the integrin-associated protein. *Biochem Biophys Res Commun*, 2001;286(1):141-9.
- Gimbrone MA Jr, Topper JN, Nagel T, Anderson KR, Garcia-Cardena G. Endothelial dysfunction, hemodynamic forces, and atherogenesis. *Ann N Y Acad Sci*, 2000;902,230-9.
- Good DJ, Polverini PJ, Rastinejad F, LeBeau MM, Lemons RS, Frazier WA, Bouck NP. A tumor suppressor-dependent inhibitor of angiogenesis is immunologically and functionally indistinguishable from a fragment of thrombospondin. *Proc Natl Acad Sci*, 1990;87:6624-8.

- Greenwalt DE, Lipsky RH, Ockenhouse CF, Ikeda H, Tandon HH, Jamieson GA. Membrane glycoprotein CD36: A review of its roles in adherence, signal transduction, and transfusion medicine. *Blood*, 1992;80:1005-15.
- Grossfeld GD, Ginsberg DA, Stein JP, Bochner BH, Esrig D, Groshen S, Dunn M, Nichols PW, Taylor CR, Skinner DG, Cote RJ. Thrombospondin-1 expression in bladder cancer: association with p53 alterations, tumor angiogenesis, and tumor progression. *J Natl Cancer Inst*, 1997;89(3):219-27.
- Guc D, Gulati P, Lemercier C, Lappin D, Birnie GD, Whaley K. Expression of the components and regulatory proteins of the alternative complement pathway and the membrane attack complex in normal and diseased synovium. *Rheumatol Int*, 1993;13, 39-46.
- Guo N, Krutzsch HC, Inman JK, Roberts DD. Thrombospondin 1 and type I repeat peptides of thrombospondin 1 specifically induce apoptosis of endothelial cells. *Cancer Res*, 1997;57(9):1735-42
- Gupta K, Gupta P, Wild R, Ramakrishnan S, Hebbel RP. Binding and displacement of vascular endothelial growth factor (VEGF) by thrombospondin: effect on human microvascular endothelial cell proliferation and angiogenesis. *Angiogenesis*, 1999;3(2):147-58.
- Hanahan D. Signaling vascular morphogenesis and maintenance. *Science*, 1997;277:48-50.
- Heissig B, Hattori K, Friedrich M, Rafii S, Werb Z. Angiogenesis: vascular remodeling of the extracellular matrix involves metalloproteinases. *Curr Opin Hematol*, 2003;10(2):136-41.
- Hochleitner BW, Hochleitner EO, Obrist P, Eberl T, Amberger A, Xu Q, Margreiter R, Wick G. Fluid shear stress induces heat shock protein 60 expression in endothelial cells in vitro and in vivo. *Arterioscler Thromb Vasc Biol*, 2000;20(3):617-23.

- Hsieh H-J, Li N-Q, Frangos JA. Shear stress increases endothelial platelet-derived growth factor mRNA levels. *Am J Phys*, 1991;260:642-4.
- Hsieh H-J, Li NQ, Frangos JA. Pulsatile and steady flow induces c-fos expression in human endothelial cells. *J Cell Physiol*, 1993;154:143-51.
- Hudlicka O, Brown M, Egginton S. Angiogenesis in skeletal and cardiac muscle. *Physiol Rev*, 1992;72:369-417.
- Ingber DE, Dike L, Hansen L, Karp S, Liley H, Maniotis A, McNamee H, Mooney D, Plopper G, Sims J. Cellular tensegrity: exploring how mechanical changes in the cytoskeleton regulate cell growth, migration, and tissue pattern during morphogenesis. *Int Rev Cytol*, 1994;150:173-224.
- Iruela-Arispe ML, Porter P, Bornstein P, Sage EH. Thrombospondin-1, an Inhibitor of Angiogenesis, is regulated by Progesterone in the Human Endometrium. *J Clin Invest*, 1996;97:403-12.
- Iruela-Arispe ML, Vazquez F, Ortega MA. Antiangiogenic domains shared by thrombospondins and metallopondins, a new family of angiogenic inhibitors. *Ann N Y Acad Sci*, 1999;886:58-66.
- Iruela-Arispe ML, Carpizo D, Luque A. ADAMTS1: a matrix metalloprotease with angioinhibitory properties. *Ann N Y Acad Sci*, 2003;995:183-90.
- Ishida T, Takahashi M, Corson MA, Berk BC. Fluid shear stress-mediated signal transduction: how do endothelial cells transduce mechanical force into biological responses ? *Ann N Y Acad Sci*, 1997;811:12-3.
- Jaffe EA, Nachman RL, Becker CG, Minick RC. Culture of human endothelial cells derived from umbilical veins. Identification by morphologic and immunologic criteria. *J Clin Invest*, 1973;52:2745-56.
- Jimenez B, Volpert OV, Crawford SE, Febbraio M, Silverstein RL, Bouck N. Signals leading to apoptosis-dependent inhibition of neovascularization by thrombospondin-1. *Nat Med*, 2000;6:41-8.

- Julen N, Dauchel H, Lemerrier C, Sim RB, Fontaine M, Ripoche J. In vitro biosynthesis of complement factor I by human endothelial cells. *Eur J Immunol*, 1992;22, 213-7.
- Kaiser D, Freyberg MA, Schrimpf G, Friedl P. Apoptosis induced by lack of hemodynamic forces is a general endothelial feature even occurring in immortalized cell lines. *Endothelium*, 1999;6(4):325-34.
- Kawakami Y, Watanabe Y, Yamaguchi M, Sakaguchi H, Kono I, Ueki A. TNF- α stimulates the biosynthesis of complement C3 and factor B by human umbilical cord vein endothelial cells. *Cancer Lett*, 1997;116:21-6.
- Khachigian LM, Anderson KR, Halnon NJ, Gimbrone MA Jr, Resnick N, Collins T. Egr-1 is activated in endothelial cells exposed to fluid shear stress and interacts with a novel shear-stress-response element in the PDGF-A chain promoter. *Arterioscler Thromb Vasc Biol*, 1997;17:2280-6.
- Klegeris A, Bissonnette CJ, Dorovini-Zis K, McGeer PL. Expression of complement messenger RNAs by human endothelial cells. *Brain Res*, 2000;871:1-6.
- Kuchan MJ, Jo H, Frangos JA. Role of G proteins in shear stress-mediated nitric oxide production by endothelial cells. *Am J Physiol*, 1994;267(3):C753-8.
- Kuno K, Kanada N, Nakashima E, Fujiki F, Ichimura F, Matsushima K. Molecular cloning of a gene encoding a new type of metalloproteinase-disintegrin family protein with thrombospondin motifs as an inflammation associated gene. *J Biol Chem*, 1997;272(1):556-62.
- Kyriakides TR, Zhu YH, Smith LT, Bain SD, Yang Z, Lin MT, Danielson KG, Iozzo RV, LaMarca M, McKinney CE, Ginns EI, Bornstein P. Mice that lack thrombospondin 2 display connective tissue abnormalities that are associated with disordered collagen fibrillogenesis, an increased vascular density, and a bleeding diathesis, *J Cell Biol*, 1998;140(2):419-30.

- Lan Q, Mercurius KO, Davies PF. Stimulation of transcription factors NF kappa B and AP 1 in endothelial cells subjected to shear stress. *Biochem Biophys Res Commun*, 1994;201:950-6.
- Langeggen H, Pausa M, Johnson E, Casarsa C, Tedesco F. The endothelium is an extrahepatic site of synthesis of the seventh component of the complement system. *Clin Exp Immunol*, 2000;121:69-76.
- Langille BL. Remodeling of developing and mature arteries: endothelium, smooth muscle, and matrix. *J Cardiovasc Pharmacol*, 1993;21(suppl 1):S11-S17.
- Lapiere CM, Lenaers A, Kohn LD. Procollagen peptidase: an enzyme excising the coordination peptides of procollagen. *Proc Natl Acad Sci USA*, 1971;68(12):3054-8.
- Lawler J. The functions of thrombospondin-1 and-2. *Curr Opin Cell Biol*, 2000;12(5):634-40.
- Lawler J. Thrombospondin-1 as an endogenous inhibitor of angiogenesis and tumor growth. *J Cell Mol Med*, 2002;6(1):1-12
- Lelièvre E, Lionneton F, Soncin F, Vandebunder B. The Ets family contains transcriptional activators and repressors involved in angiogenesis. *Int J Biochem Cell Biol*, 2001;33:391-407.
- Li SW, Arita M, Fertala A, Bao Y, Kopen GC, Langsjö TK, Hyttinen MM, Helminen HJ, Prockop DJ. Transgenic mice with inactive alleles for procollagen N-proteinase (ADAMTS-2) develop fragile skin and male sterility. *Biochem J*, 2001;355(Pt 2):271-8.
- Lin MC, Almus-Jacobs F, Chen HH, Parry GCN, Mackman N, Shyy JYJ, Chien S. Shear stress induction of the tissue factor gene. *J Clin Invest*, 1997;99:737-44.
- Lindahl P, Bostrom H, Karlsson L, Hellstrom M, Kalen M, Betsholtz C. Role of platelet-derived growth factors in angiogenesis and alveogenesis. *Curr Top Pathol*, 1999;93:27-33.

- Luque A, Carpizo DR, Iruela-Arispe ML. ADAMTS1/METH1 inhibits endothelial cell proliferation by direct binding and sequestration of VEGF165. *J Biol Chem*, 2003;278(26):23656-65.
- Lymn JS, Rao SJ, Clunn GF, Gallagher KL, O'Neil CO, Thompson NT, Hughes AD. Phosphatidylinositol 3-kinase and focal adhesion kinase are early signals in the growth factor-like responses to thrombospondin-1 seen in human vascular smooth muscle. *Arterioscler Thromb Vasc Biol*, 1999;19:2133-40.
- Maeda K, Nishiguchi Y, Yashiro M, Yamada S, Onoda N, Sawada T, Kang SM, Hirakawa K. Expression of vascular endothelial growth factor and thrombospondin-1 in colorectal carcinoma. *Int J Mol Med*, 2000;5(4):373-8.
- Maves K.K, Weiler J.M. Properdin: approaching four decades of research. *Immunol Res*, 1993;12, 233-43.
- Malek A, Izumo S. Physiological fluid shear stress causes downregulation of endothelin-1 mRNA in bovine aortic endothelium. *Am J Phys*, 1992;263:C389-C396.
- Malek AM, Jackman R, Rosenberg RD, Izumo S. Endothelial expression of thrombomodulin is reversibly regulated by fluid shear stress. *Circ Res*, 1994;74:852-60.
- Metivier F, Marchais SJ, Guerin AP, Pannier B, London GM. Pathophysiology of anaemia: focus on the heart and blood vessels. *Nephrol Dial Transplant*, 2000;15(Suppl 3):14-8.
- Mettouchi A, Cabon F, Montreau N, Vernier P, Mercier G, Bangy D, Tricoire H, Vigier P, Binetruy B. SPARC and thrombospondin genes are repressed by the c-jun oncogene in rat embryo fibroblasts. *EMBO J*, 1994;13:5668-78.
- Mo M, Eskin SG, Schilling WP. Flow-induced changes in Ca²⁺ signaling of vascular endothelial cells: effect of shear stress and ATP. *Am J Physiol*, 1991;260(5 Pt 2):H1698-707.

- Morandi V, Fauvel-Lafeve F, Legrand C, Legrand YJ. Role of thrombospondin in the adhesion of human endothelial cells in primary culture. *In Vitro Cell Dev Biol*, 1993;29 A:585-91.
- Nagel T, Atkinson WJ, Dewey CF, Gimbrone MA. Shear stress selectively upregulates intercellular adhesion molecule-1 expression in cultured human vascular endothelial cells. *J Clin Invest*, 1994;94:885-91.
- Nesselroth SM, Willis AI, Fuse S, Olson ET, Lawler J, Sumpio BE, Gahtan V. The C-terminal domain of thrombospondin-1 induces vascular smooth muscle cell chemotaxis. *J Vasc Surg*, 2001;33(3):595-600.
- Nicosia RF, Tuszynski GP. Matrix-bound thrombospondin promotes angiogenesis in vitro. *J Cell Biol*, 1994;124:183-93.
- Niculescu F, Rus H. Complement activation and atherosclerosis. *Mol Immunol*, 1999;36, 949-55.
- Niewiarowski S, McLane MA, Kloczewiak M, Stewart GJ. Disintegrins and other naturally occurring antagonists of platelet fibrinogen receptors. *Semin Hematol*, 1994;31(4):289-300.
- Nishida K, Harrison DG, Navas JP, Fisher AA, Dockery SP, Uematsu M, Nerem RM, Alexander RW, Murphy TJ. Molecular cloning and characterization of the constitutive bovine aortic endothelial cell nitric oxide synthase. *J Clin Invest*, 1992;90/5:2092-6.
- Nishimori H, Shiratsuchi T, Urano T, Kimura Y, Kiyono K, Tatsumi K, Yoshida S, Ono M, Kuwano M, Nakamura Y, Tokino T. A novel brain-specific p53-target gene, BAI1, containing thrombospondin type 1 repeats inhibits experimental angiogenesis. *Oncogene*, 1997;15(18):2145-50.
- Nolan KF, Reid KB. Properdin. *Methods Enzymol*, 1993;223, 35-46.
- Nor JE, Mitra RS, Sutorik MM, Mooney DJ, Castle VP, Polverini PJ. Thrombospondin-1 induces endothelial cell apoptosis and inhibits

- angiogenesis by activating the caspase death way. *J Vasc Res*, 2000;37(3):209-18.
- Ohno M, Cooke JP, Dzau VL, Gibbons GH. Fluid shear stress induces endothelial transforming growth factor beta-1 transcription and production. Modulation by potassium channel blockade. *J Clin Invest*, 1995;95:1363-9.
- Ohtsuka A, Ando J, Korenaga R, Kamiya A, Toyama-Sorimachi N, Miyasaka M. The effect of flow on the expression of vascular adhesion molecule-1 by cultured mouse endothelial cells. *Biochem Biophys Res Commun*, 1993;193:303-10.
- Okahara K, Sun B, Kawasaki T, Monden M, Kambayashi J. Expression of platelet-activating factor receptor transcript-2 is induced by shear stress in HUVEC. *Prostaglandins Other Lipid Mediat*, 1998;55:323-9.
- Olesen SP, Clapham DE, Davies PF. Haemodynamic shear stress activates a K⁺ current in vascular endothelial cells. *Nature*, 1988;331:168-70.
- Papadaki M, Eskin SG. Effects of fluid shear stress on gene regulation of vascular cells. *Biotechnol Prog*, 1997;13(3):209-21.
- Pangburn MK. Analysis of the natural polymeric forms of human properdin and their functions in complement activation. *J Immunol*, 1989;142:202-7.
- Panetti TS, Chen H, Misenheimer TM, Getzler SB, Mosher DF. Endothelial cell mitogenesis induced by LPA: inhibition by thrombospondin-1 and thrombospondin-2. *Lab Clin Med*, 1997;129(2):208-16.
- Phelan MW, Forman LW, Perrine SP, Faller DV. Hypoxia increases thrombospondin-1 transcript and protein in cultured endothelial cells. *J Lab Clin Med*, 1998;132(6):519-29.
- Poiseuille JLM. *Recherches sur la force du cœur aortique [Dissertation]*". Paris: 1828, Didot le Jeune
- Pries AR, Neuhaus D, Gaehtgens P. Blood viscosity in tube flow: dependence on diameter and hematocrit. *Am J Physiol*, 1992;263(6 Pt 2):H1770-8.

- Pries AR, Secomb TW, Gaetgens P. Design principles of vascular beds. *Circ Res*, 1995;77:1017-1023.
- Pries AR, Reglin B, Secomb TW. Structural adaptation of microvascular networks: functional roles of adaptive responses. *Am J Physiol Heart Circ Physiol*, 2000;281:H1015-25.
- Pries AR, Reglin B, Secomb TW. Structural response of microcirculatory networks to changes in demand: information transfer by shear stress. *Am J Physiol Heart Circ Physiol*, 2003;284(6):H2204-12.
- Qian X, Wang TN, Rothman VL, Nicosia RF, Tuszynski GP. Thrombospondin-1 modulates angiogenesis in vitro by up-regulation of matrixmetalloproteinase-9 in endothelial cells. *Exp Cell Res*, 1997;235:403-12.
- Resnick N, Collins T, Atkinson W, Bonthron DT, Dewey CF Jr, Gimbron MA Jr. Platelet-derived growth factor B chain promoter contains a cis-acting fluid shear-stress-responsive element. *Proc Natl Acad Sci USA*, 1993; 90:4591-5.
- Resnick N, Yahav H, Shay-Salit A, Shushy M, Schubert S, Zilberman LC, Wofovitz E. Fluid shear stress and the vascular endothelium: for better and for worse. *Prog Biophys Mol Biol*, 2003;81(3):177-99.
- Rieder MJ, Carmona R, Krieger JE, Pritchard KA, Greene AS. Suppression of angiotensin-converting enzyme expression and activity by shear stress. *Circ Res*, 1997;80:312-9.
- Rieger H, Gaetgens P. Pathophysiologie der Gewebsdurchblutung und ihre Regulation: Durchblutungsstörungen in der Pathophysiologie des Menschen, Chapman & Hall, 1994;17.3.
- Risau W. Mechanisms of angiogenesis. *Nature*, 1997;386(6626):671-4.

- Rodriguez-Manzaneque JC, Milchanowski AB, Dufour EK, Leduc R, Iruela-Arispe ML. Characterization of METH-1/ADAMTS1 processing reveals two distinct active forms. *J Biol Chem*, 2000;275(43):33471-9.
- Rundhaug JE. Matrix metalloproteinases, angiogenesis, and cancer: commentary re: AC. Lockhart et al, Reduction of wound angiogenesis in patients treated with BMS-275291, a broad spectrum matrix metalloproteinase inhibitor. *Clin Cancer Res*, 2003;9(2):551-4.
- Sargiannidou I, Zhou J, Tuszynski GP. The role of thrombospondin-1 in tumor progression. *Exp Biol Med*, 2001;226(8):726-33.
- Schultz-Cherry S, Chen H, Mosher DF, Misenheimer TM, Kruttsch HC, Roberts DD, Murphy-Ullrich JE. Regulation of transforming growth factor-beta activation by discrete sequences of thrombospondin 1. *Biol Chem*, 1995;270(13):7304-10.
- Schwachtgen J.-L, Houston P, Campbell C, Sukhatme V, Braddock M. Fluid shear stress activation of egr-1 transcription in cultured human endothelial and epithelial cells is mediated via the extracellular signal-related kinase 1/2 mitogen-activated protein kinase pathway. *J Clin Invest*, 1998;101:2540-9.
- Schwaeble W, Huemer HP, Most J, Dierich MP, Strobel M, Claus C, Reid KB, Ziegler-Heitbrock HW. Expression of properdin in human monocytes. *Eur J Biochem*, 1994;219:759-64.
- Sdougos HP, Bussolari SR, Dewey CF. Secondary flow and turbulence in a cone-and-plate device. *J Fluid Mech*, 1984;138:397-404.
- Shyy YJ, Hsieh J-H, Usami S, Chien S. Fluid shear stress induces a biphasic response of human monocyte chemotactic protein 1 gene expression in vascular endothelium. *Proc Natl Acad Sci USA*, 1994;91:4678-82.
- Simon MI, Strathmann MP, Gautam N. Diversity of G proteins in signal transduction. *Science*, 1991;252:802-8.
- Skalak TC, Price RJ. The role of mechanical stresses in microvascular remodeling. *Microcirculation*, 1996;3:143-65.

- Streit M, Riccardi L, Velasco P, Brown LF, Hawighorst T, Bornstein P, Detmar M. Thrombospondin-2: a potent endogenous inhibitor of tumor growth and angiogenesis. *Proc Natl Acad Sci USA*, 1999;96:14888-93.
- Suzuma K, Takagi H, Otani A, Oh A, Honda Y. Expression of thrombospondin-1 in ischemia-induced retinal neovascularization. *Am J Pathol*, 1999;154(2):343-54.
- Swerlick RA, Lee KH, Wick TM, Lawley TJ. Human dermal microvascular endothelial but not human umbilical vein endothelial cells express CD 36 in vivo and in vitro. *J Immunol*, 1992;148:78-83.
- Takahashi M, Berk BC. Mitogen-activated protein kinase (ERK 1/2) activation by shear stress and adhesion in endothelial cells – essential role for a herbimycin-sensitive kinase. *J Clin Invest*, 1996;98:2623-31.
- Tang BL. ADAMTS: a novel family of extracellular matrix proteases. *Int J Biochem Cell Biol* 2001;33(1):33-44.
- Taraboletti G, Roberts D, Liotta LA, Giavazzi R. Platelet thrombospondin modulates endothelial cell adhesion, motility, and growth: a potential angiogenesis regulatory factor. *J Cell Biol*, 1990;111:765-72.
- Tolsma SS, Volpert OV, Good DJ, Frazier DA, Poverini PJ, Bouck N. Peptides derived from two separate domains of the matrix protein thrombospondin-1 have anti-angiogenic activity. *J Cell Biol*, 1993;122:497-511.
- Topper JN, Cai J, Falb D, Gimbrone MA. Identification of vascular endothelial genes differentially responsive to fluid mechanical stimuli: cyclooxygenase-2, manganese superoxide dismutase, and endothelial cell nitric oxide synthase are selectively up-regulated by steady laminar shear stress. *Proc Natl Acad Sci USA*, 1996;93(19):10417-22.
- Topper JN, Cai J, Qiu Y, Anderson KR, Xu YY, Deeds JD, Feeley R, Gimeno CJ, Woolf EA, Tayber O, Mays GG, Sampson BA, Schoen FJ, Gimbrone MA Jr, Falb D. Vascular MADs: two novel MAD-related genes selectively inducible

- by flow in human vascular endothelium. *Proc Natl Acad Sci USA*, 1997;94 (17):9314-9.
- Topper JN, Gimbrone MA Jr. Blood flow and vascular gene expression: fluid shear stress as a modulator of endothelial phenotype. *Mol Med Today*, 1999;5:40-6.
- Torzewski J, Bowyer DE, Waltenberger J, Fitzsimmons C. Processes in atherogenesis: complement activation. *Atherosclerosis*, 1997;132:131-8.
- Tsuboi H, Ando J, Korenaga R, Takada Y, Kamiya A. Flow stimulates ICAM-1 expression time and shear stress dependently in cultured human endothelial cells. *Biochem Biophys Res Commun*, 1995;206(3):988-96.
- Tuszynski GP, Rothman V, Murphy A, Siegler K, Smith L, Smith S, Karczewski J, Knudsen KA. Thrombospondin promotes cell-substratum adhesion. *Science*, 1987;236:1570-3.
- Tuszynski GP, Nicosia RF. Localization of thrombospondin and its cysteine-serine-valine-threonine-cysteine-glycine-specific receptor in human breast carcinoma. *Lab Invest*, 1994;70(2):228-33.
- Tuszynski GP, Nicosia RF. The role of thrombospondin-1 in tumor progression and angiogenesis. *Bioessays*, 1996;18(1):71-6.
- Urbich C, Fritzenwanger M, Zeiher AM, Dimmeler S. Laminar shear stress upregulates the complement-inhibitory protein clusterin: a novel potent defense mechanism against complement-induced endothelial cell activation. *Circulation*, 2000;101:352-5.
- Vazquez F, Hastings G, Ortega MA, Lane T.F, Oikemus S, Lombardo M, Iruela-Arispe ML. METH-1, a human ortholog of ADAMTS-1, and METH-2 are members of a new family of proteins with angio-inhibitory activity. *J Biol.Chem*, 1999;274 (33):23349-57.

- Volpert OV, Stellmach V.;Bouck N. The modulation of thrombospondin and other naturally occurring inhibitors of angiogenesis during tumor progression. *Breast Cancer Res Treat*, 1995;36(2):119-26.
- Wang XQ, Frazier WA. The thrombospondin receptor CD47(IAP) modulates and associates with alpha 2 beta 1 integrin in vascular smooth muscle cells. *Mol Biol Cell*, 1998;9:865-74.
- Wang X-Q, Lindberg F, Frazier W. Integrin-associated protein stimulates $\alpha_2\beta_1$ -dependent chemotaxis via Gi-mediated inhibition of adenylate-cyclase and extracellular-regulated kinases. *J Cell Biol*, 1999;147:389-99.
- Wirthmueller U, Dewald B, Thelen M, Schafer MK, Stover C, Whaley K, North J, Eggleton P, Reid KB, Schwaeble WJ. Properdin, a positive regulator of complement activation, is released from secondary granules of stimulated peripheral blood neutrophils. *J Immunol*, 1997;158:4444-51.
- Witzenbichler B, Maisonpierre PC, Jones P, Yancopoulos GD, Isner JM. Chemotactic properties of Angiopoietin-1 and-2, ligands for the endothelial-specific receptor tyrosine kinase Tie2. *J Biol Chem*, 1998;273:18514-24.
- Wolfsberg TG, Primakoff P, Myles DG, White JM. ADAM, a novel family of membrane proteins containing A Disintegrin And Metalloprotease domain: multipotential functions in cell-cell and cell-matrix interactions. *J Cell Biol*, 1995;131(2):275-8.
- Yang Z, Kyriakides TR, Bornstein P. Matricellular proteins as modulators of cell-matrix interactions: adhesive defect in thrombospondin 2-null fibroblasts is a consequence of increased levels of matrix metalloproteinase-2. *Mol Biol Cell*, 2000;11(10):3353-64.
- Yao L, Zhao YL, Itoh S, Wada S, Yue L, Furuta I. Thrombospondin-1 expression in oral squamous cell carcinomas: correlations with tumor vascularity, clinicopathological features and survival. *Oral Oncol*, 2000;36(6):539-44.
- Zarins CK, Giddens DP, Bharadvaj BK, Sottiurai VS, Mabon RF, Glagov S. Carotid bifurcation atherosclerosis. Quantitative correlation of plaque localization

with flow velocity profiles and wall shear stress. *Circ Res*, 1983;53(4):502-14.

Zakrzewicz A, Grafe M, Terbeek D, Bongrazio M, Auch-Schwelk W, Walzog B, Graf K, Fleck E, Ley K, Gaehtgens P. L-selectin-dependent leukocyte adhesion to microvascular but not to macrovascular endothelial cells of the human coronary system. *Blood*, 1997;89:3228-35.

Zakrzewicz A, Secomb TW, Pries AR. Angioadaptation: Keeping the vascular system in shape. *News Physiol Sci*, 2002;17:197-201.

Zhou A, Egginton S, Hudlicka O, Brown MD. Internal division of capillaries in rat skeletal muscle in response to chronic vasodilator treatment with alpha1-antagonist prazosin. *Cell Tissue Res*, 1998;293(2):293-303.