

VI. References

- Andersen, O.M., Christensen, L.L., Christensen, P.A., Sorensen, E.S., Jacobsen, C., Moestrup, S.K., Etzerodt, M. and Thogersen, H.C. (2000a) Identification of the minimal functional unit in the low density lipoprotein receptor-related protein for binding the receptor-associated protein (RAP). A conserved acidic residue in the complement-type repeats is important for recognition of RAP. *J Biol Chem*, **275**, 21017-21024.
- Andersen, O.M., Christensen, P.A., Christensen, L.L., Jacobsen, C., Moestrup, S.K., Etzerodt, M. and Thogersen, H.C. (2000b) Specific binding of alpha-macroglobulin to complement-type repeat CR4 of the low-density lipoprotein receptor-related protein. *Biochemistry*, **39**, 10627-10633.
- Anderson, L.E., Walsh, K.A. and Neurath, H. (1977) Bovine enterokinase. Purification, specificity, and some molecular properties. *Biochemistry*, **16**, 3354-3360.
- Arden, K.C., Viars, C.S., Weiss, S., Argentin, S. and Nemer, M. (1995) Localization of the human B-type natriuretic peptide precursor (NPPB) gene to chromosome 1p36. *Genomics*, **26**, 385-389.
- Bhanot, P., Brink, M., Samos, C.H., Hsieh, J.C., Wang, Y., Macke, J.P., Andrew, D., Nathans, J. and Nusse, R. (1996) A new member of the frizzled family from Drosophila functions as a Wingless receptor. *Nature*, **382**, 225-230.
- Bieker, J.J. (2001) Kruppel-like factors: three fingers in many pies. *J Biol Chem*, **276**, 34355-34358.
- Bleyl, S., Nelson, L., Odelberg, S.J., Ruttenberg, H.D., Otterud, B., Leppert, M. and Ward, K. (1995) A gene for familial total anomalous pulmonary venous return maps to chromosome 4p13-q12. *Am J Hum Genet*, **56**, 408-415.
- Bliss, D.P., Jr., Battey, J.F., Linnoila, R.I., Birrer, M.J., Gazdar, A.F. and Johnson, B.E. (1990) Expression of the atrial natriuretic factor gene in small cell lung cancer tumors and tumor cell lines. *J Natl Cancer Inst*, **82**, 305-310.

- Bravo, D.A., Gleason, J.B., Sanchez, R.I., Roth, R.A. and Fuller, R.S. (1994) Accurate and efficient cleavage of the human insulin proreceptor by the human proprotein-processing protease furin. Characterization and kinetic parameters using the purified, secreted soluble protease expressed by a recombinant baculovirus. *J Biol Chem*, **269**, 25830-25837.
- Brenner, B.M., Ballermann, B.J., Gunning, M.E. and Zeidel, M.L. (1990) Diverse biological actions of atrial natriuretic peptide. *Physiol Rev*, **70**, 665-699.
- Brown, J., Chen, Q. and Hong, G. (1997) An autocrine system for C-type natriuretic peptide within rat carotid neointima during arterial repair. *Am J Physiol*, **272**, H2919-2931.
- Brown, M.S. and Goldstein, J.L. (1986) A receptor-mediated pathway for cholesterol homeostasis. *Science*, **232**, 34-47.
- Brown, M.S., Herz, J., and Goldstein, J.L. (1997) LDL-receptor structure. Calcium cages, acid baths and recycling receptors. *Nature*, **388**, 629-30.
- Brown, S.D., Twells, R.C., Hey, P.J., Cox, R.D., Levy, E.R., Soderman, A.R., Metzker, M.L., Caskey, C.T., Todd, J.A. and Hess, J.F. (1998) Isolation and characterization of LRP6, a novel member of the low density lipoprotein receptor gene family. *Biochem Biophys Res Commun*, **248**, 879-888.
- Bruneau, B.G., Nemer, G., Schmitt, J.P., Charron, F., Robitaille, L., Caron, S., Conner, D.A., Gessler, M., Nemer, M., Seidman, C.E. and Seidman, J.G. (2001) A murine model of Holt-Oram syndrome defines roles of the T-box transcription factor Tbx5 in cardiogenesis and disease. *Cell*, **106**, 709-721.
- Bucher, P. (1990) Weight matrix descriptions of four eukaryotic RNA polymerase II promoter elements derived from 502 unrelated promoter sequences. *J Mol Biol*, **212**, 563-578.
- Burnett, J.C., Jr., Kao, P.C., Hu, D.C., Heser, D.W., Heublein, D., Granger, J.P., Opgenorth, T.J. and Reeder, G.S. (1986) Atrial natriuretic peptide elevation in congestive heart failure in the human. *Science*, **231**, 1145-1147.
- Cal, S., Quesada, V., Garabaya, C. and Lopez-Otin, C. (2003) Polyserase-I, a human polyprotease with the ability to generate independent serine protease domains from a single translation product. *Proc Natl Acad Sci U S A*, **100**, 9185-9190.
- Campbell, I.D. (2003) Modular proteins at the cell surface. *Biochem Soc Trans*, **31**, 1107-1114.
- Campling, B.G., Sarda, I.R., Baer, K.A., Pang, S.C., Baker, H.M., Lofters, W.S. and Flynn, T.G. (1995) Secretion of atrial natriuretic peptide and vasopressin by small cell lung cancer. *Cancer*, **75**, 2442-2451.

- Chan, J., Knudson, O., Morser, J., Dole, W., and Wu, Q. (2004) Salt-sensitive and pregnancy-induced hypertension in mice deficient in corin, the pro-atrial natriuretic peptide convertase. *Cardiovascular Pathology 13 Meeting*, Oral presentation.
- Charron, F. and Nemer, M. (1999) GATA transcription factors and cardiac development. *Semin Cell Dev Biol*, **10**, 85-91.
- Chen, C.Y. and Schwartz, R.J. (1995) Identification of novel DNA binding targets and regulatory domains of a murine tinman homeodomain factor, nkx-2.5. *J Biol Chem*, **270**, 15628-15633.
- Chen, H.H. and Burnett, J.C., Jr. (1998) C-type natriuretic peptide: the endothelial component of the natriuretic peptide system. *J Cardiovasc Pharmacol*, **32 Suppl 3**, S22-28.
- Chinkers, M., Garbers, D.L., Chang, M.S., Lowe, D.G., Chin, H.M., Goeddel, D.V. and Schulz, S. (1989) A membrane form of guanylate cyclase is an atrial natriuretic peptide receptor. *Nature*, **338**, 78-83.
- Cho, E.G., Kim, M.G., Kim, C., Kim, S.R., Seong, I.S., Chung, C., Schwartz, R.H. and Park, D. (2001) N-terminal processing is essential for release of epithin, a mouse type II membrane serine protease. *J Biol Chem*, **276**, 44581-44589.
- Chusho, H., Tamura, N., Ogawa, Y., Yasoda, A., Suda, M., Miyazawa, T., Nakamura, K., Nakao, K., Kurihara, T., Komatsu, Y., Itoh, H., Tanaka, K., Saito, Y. and Katsuki, M. (2001) Dwarfism and early death in mice lacking C-type natriuretic peptide. *Proc Natl Acad Sci U S A*, **98**, 4016-4021.
- Claycomb, W.C., Lanson, N.A., Jr., Stallworth, B.S., Egeland, D.B., Delcarpio, J.B., Bahinski, A. and Izzo, N.J., Jr. (1998) HL-1 cells: a cardiac muscle cell line that contracts and retains phenotypic characteristics of the adult cardiomyocyte. *Proc Natl Acad Sci U S A*, **95**, 2979-2984.
- Cogan, E., Debieve, M.F., Philipart, I., Pepersack, T. and Abramow, M. (1986) High plasma levels of atrial natriuretic factor in SIADH. *N Engl J Med*, **314**, 1258-1259.
- Colucci, W.S., Elkayam, U., Horton, D.P., Abraham, W.T., Bourge, R.C., Johnson, A.D., Wagoner, L.E., Givertz, M.M., Liang, C.S., Neibaur, M., Haught, W.H. and LeJemtel, T.H. (2000) Intravenous nesiritide, a natriuretic peptide, in the treatment of decompensated congestive heart failure. Nesiritide Study Group. *N Engl J Med*, **343**, 246-253.
- Corthorn, J., Cantin, M. and Thibault, G. (1991) Rat atrial secretory granules and pro-ANF processing enzyme. *Mol Cell Biochem*, **103**, 31-39.

- Cromlish, J.A., Seidah, N.G. and Chretien, M. (1986a) A novel serine protease (IRCM-serine protease 1) from porcine neurointermediate and anterior pituitary lobes. Isolation, polypeptide chain structure, inhibitor sensitivity, and substrate specificity with fluorogenic peptide substrates. *J Biol Chem*, **261**, 10850-10858.
- Cromlish, J.A., Seidah, N.G. and Chretien, M. (1986b) Selective cleavage of human ACTH, beta-lipotropin, and the N-terminal glycopeptide at pairs of basic residues by IRCM-serine protease 1. Subcellular localization in small and large vesicles. *J Biol Chem*, **261**, 10859-10870.
- Croy, J.E., Shin, W.D., Knauer, M.F., Knauer, D.J. and Komives, E.A. (2003) All three LDL receptor homology regions of the LDL receptor-related protein bind multiple ligands. *Biochemistry*, **42**, 13049-13057.
- Currie, M.G., Geller, D.M., Chao, J., Margolius, H.S. and Needleman, P. (1984a) Kallikrein activation of a high molecular weight atrial peptide. *Biochem Biophys Res Commun*, **120**, 461-466.
- Currie, M.G., Geller, D.M., Cole, B.R. and Needleman, P. (1984b) Proteolytic activation of a bioactive cardiac peptide by *in vitro* trypsin cleavage. *Proc Natl Acad Sci U S A*, **81**, 1230-1233.
- Currie, M.G. and Newman, W.H. (1986) Evidence for alpha-1 adrenergic receptor regulation of atriopeptin release from the isolated rat heart. *Biochem Biophys Res Commun*, **137**, 94-100.
- Dann, C.E., Hsieh, J.C., Rattner, A., Sharma, D., Nathans, J. and Leahy, D.J. (2001) Insights into Wnt binding and signalling from the structures of two Frizzled cysteine-rich domains. *Nature*, **412**, 86-90.
- Davie, E.W., Fujikawa, K. and Kisiel, W. (1991) The coagulation cascade: initiation, maintenance, and regulation. *Biochemistry*, **30**, 10363-10370.
- de Bold, A.J. (1979) Heart atria granularity effects of changes in water-electrolyte balance. *Proc Soc Exp Biol Med*, **161**, 508-511.
- de Bold, A.J. (1985) Atrial natriuretic factor: a hormone produced by the heart. *Science*, **230**, 767-770.
- de Bold, A.J. and Bencosme, S.A. (1973) Studies on the relationship between the catecholamine distribution in the atrium and the specific granules present in atrial muscle cells. 1. Isolation of a purified specific granule subfraction. *Cardiovasc Res*, **7**, 351-363.
- de Bold, A.J., Borenstein, H.B., Veress, A.T. and Sonnenberg, H. (1981) A rapid and potent natriuretic response to intravenous injection of atrial myocardial extract in rats. *Life Sci*, **28**, 89-94.

- de Bold, A.J., Raymond, J.J. and Bencosme, S.A. (1978) Atrial specific granules of the rat heart: light microscopic staining and histochemical reactions. *J Histochem Cytochem*, **26**, 1094-1102.
- de Winther, M.P., van Dijk, K.W., Havekes, L.M. and Hofker, M.H. (2000) Macrophage scavenger receptor class A: A multifunctional receptor in atherosclerosis. *Arterioscler Thromb Vasc Biol*, **20**, 290-297.
- Denault, J.B., Claing, A., D'Orleans-Juste, P., Sawamura, T., Kido, T., Masaki, T. and Leduc, R. (1995) Processing of proendothelin-1 by human furin convertase. *FEBS Lett*, **362**, 276-280.
- Donckier, J.E., Meunier, H.T., Ketelslegers, J.M. and Lambert, A.E. (1986) Atrial natriuretic peptide and the syndrome of inappropriate secretion of antidiuretic hormone. *Ann Intern Med*, **105**, 795-796.
- Dong, Y., Lathrop, W., Weaver, D., Qiu, Q., Cini, J., Bertolini, D. and Chen, D. (1998) Molecular cloning and characterization of LR3, a novel LDL receptor family protein with mitogenic activity. *Biochem Biophys Res Commun*, **251**, 784-790.
- Doolittle, R.F. (1995) The multiplicity of domains in proteins. *Annu Rev Biochem*, **64**, 287-314.
- Durocher, D. and Nemer, M. (1998) Combinatorial interactions regulating cardiac transcription. *Dev Genet*, **22**, 250-262.
- Edman, P. (1950) Method for determination of the amino acid sequence in peptides. *Acta Chem Scand*, **4**, 283-293
- Esser, V., Limbird, L.E., Brown, M.S., Goldstein, J.L. and Russell, D.W. (1988) Mutational analysis of the ligand binding domain of the low density lipoprotein receptor. *J Biol Chem*, **263**, 13282-13290.
- Fass, D., Blacklow, S., Kim, P.S. and Berger, J.M. (1997) Molecular basis of familial hypercholesterolaemia from structure of LDL receptor module. *Nature*, **388**, 691-693.
- Field, L.J. (1988) Atrial natriuretic factor-SV40 T antigen transgenes produce tumors and cardiac arrhythmias in mice. *Science*, **239**, 1029-1033.
- Finch, P.W., He, X., Kelley, M.J., Uren, A., Schaudies, R.P., Popescu, N.C., Rudikoff, S., Aaronson, S.A., Varmus, H.E. and Rubin, J.S. (1997) Purification and molecular cloning of a secreted, Frizzled-related antagonist of Wnt action. *Proc Natl Acad Sci U S A*, **94**, 6770-6775.
- Flynn, T.G., de Bold, M.L. and de Bold, A.J. (1983) The amino acid sequence of an atrial peptide with potent diuretic and natriuretic properties. *Biochem Biophys Res Commun*, **117**, 859-865.

- Ford, C.F., Suominen, I. and Glatz, C.E. (1991) Fusion tails for the recovery and purification of recombinant proteins. *Protein Expr Purif*, **2**, 95-107.
- Fox, K.F., Cowie, M.R., Wood, D.A., Coats, A.J., Gibbs, J.S., Underwood, S.R., Turner, R.M., Poole-Wilson, P.A., Davies, S.W. and Sutton, G.C. (2001) Coronary artery disease as the cause of incident heart failure in the population. *Eur Heart J*, **22**, 228-236.
- Francis, G.S., McDonald, K., Chu, C. and Cohn, J.N. (1995) Pathophysiologic aspects of end-stage heart failure. *Am J Cardiol*, **75**, 11A-16A.
- Friedrich, R., Fuentes-Prior, P., Ong, E., Coombs, G., Hunter, M., Oehler, R., Pierson, D., Gonzalez, R., Huber, R., Bode, W. and Madison, E.L. (2002) Catalytic domain structures of MT-SP1/matriptase, a matrix-degrading transmembrane serine proteinase. *J Biol Chem*, **277**, 2160-2168.
- Fuller, F., Porter, J.G., Arfsten, A.E., Miller, J., Schilling, J.W., Scarborough, R.M., Lewicki, J.A. and Schenk, D.B. (1988) Atrial natriuretic peptide clearance receptor. Complete sequence and functional expression of cDNA clones. *J Biol Chem*, **263**, 9395-9401.
- Furie, B. and Furie, B.C. (1988) The molecular basis of blood coagulation. *Cell*, **53**, 505-518.
- Furuya, M., Yoshida, M., Hayashi, Y., Ohnuma, N., Minamino, N., Kangawa, K. and Matsuo, H. (1991) C-type natriuretic peptide is a growth inhibitor of rat vascular smooth muscle cells. *Biochem Biophys Res Commun*, **177**, 927-931.
- Gliemann, J. (1998) Receptors of the low density lipoprotein (LDL) receptor family in man. Multiple functions of the large family members via interaction with complex ligands. *Biol Chem*, **379**, 951-964.
- Goldstein, J.L., Brown, M.S., Anderson, R.G., Russell, D.W. and Schneider, W.J. (1985) Receptor-mediated endocytosis: concepts emerging from the LDL receptor system. *Annu Rev Cell Biol*, **1**, 1-39.
- Grant, S.G., Jesse, J., Bloom, F.R. and Hanahan, D. (1990) Differential plasmid rescue from transgenic mouse DNAs into Escherichia coli methylation-restriction mutants. *Proc Natl Acad Sci U S A*, **87**, 4645-4649.
- Grepin, C., Dagnino, L., Robitaille, L., Haberstroh, L., Antakly, T. and Nemer, M. (1994) A hormone-encoding gene identifies a pathway for cardiac but not skeletal muscle gene transcription. *Mol Cell Biol*, **14**, 3115-3129.
- Gross, A.J., Steinberg, S.M., Reilly, J.G., Bliss, D.P., Jr., Brennan, J., Le, P.T., Simmons, A., Phelps, R., Mulshine, J.L., Ihde, D.C. and et al. (1993) Atrial natriuretic factor and arginine vasopressin production in tumor cell lines from patients with lung cancer and their relationship to serum sodium. *Cancer Res*, **53**, 67-74.

- Hartmann, E., Rapoport, T.A. and Lodish, H.F. (1989) Predicting the orientation of eukaryotic membrane-spanning proteins. *Proc Natl Acad Sci U S A*, **86**, 5786-5790.
- Hemker, H.C. (1983) Handbook of synthetic substrates for the coagulation and fibrinolytic system. *Martinus Nijhoff Publ*, The Hague
- Henry, J.P., Gauer, O.H. and Reeves, J.L. (1956) Evidence of the atrial location of receptors influencing urine flow. *Circ Res*, **4**, 85-90.
- Herz, J. and Strickland, D.K. (2001) LRP: a multifunctional scavenger and signaling receptor. *J Clin Invest*, **108**, 779-784.
- Hey, P.J., Twells, R.C., Phillips, M.S., Yusuke, N., Brown, S.D., Kawaguchi, Y., Cox, R., Guochun, X., Dugan, V., Hammond, H., Metzker, M.L., Todd, J.A. and Hess, J.F. (1998) Cloning of a novel member of the low-density lipoprotein receptor family. *Gene*, **216**, 103-111.
- Hobbs, H.H., Brown, M.S. and Goldstein, J.L. (1992) Molecular genetics of the LDL receptor gene in familial hypercholesterolemia. *Hum Mutat*, **1**, 445-466.
- Hobson, J.P., Netzel-Arnett, S., Szabo, R., Rehault, S.M., Church, F.C., Strickland, D.K., Lawrence, D.A., Antalis, T.M. and Bugge, T.H. (2004) Mouse DESC1 is located within a cluster of seven DESC1-like genes and encodes a type II transmembrane serine protease that forms serpin inhibitory complexes. *J Biol Chem*.
- Hohenester, E., Sasaki, T. and Timpl, R. (1999a) Crystal structure of a scavenger receptor cysteine-rich domain sheds light on an ancient superfamily. *Nat Struct Biol*, **6**, 228-232.
- Hooper, J.D., Clements, J.A., Quigley, J.P. and Antalis, T.M. (2001) Type II transmembrane serine proteases. Insights into an emerging class of cell surface proteolytic enzymes. *J Biol Chem*, **276**, 857-860.
- Hooper, J.D., Scarman, A.L., Clarke, B.E., Normyle, J.F. and Antalis, T.M. (2000) Localization of the mosaic transmembrane serine protease corin to heart myocytes. *Eur J Biochem*, **267**, 6931-6937.
- Horn, I.R., van den Berg, B.M., van der Meijden, P.Z., Pannekoek, H. and van Zonneveld, A.J. (1997) Molecular analysis of ligand binding to the second cluster of complement-type repeats of the low density lipoprotein receptor-related protein. Evidence for an allosteric component in receptor-associated protein-mediated inhibition of ligand binding. *J Biol Chem*, **272**, 13608-13613.
- Hosoda, K., Nakao, K., Mukoyama, M., Saito, Y., Jougasaki, M., Shirakami, G., Suga, S., Ogawa, Y., Yasue, H. and Imura, H. (1991) Expression of brain natriuretic peptide gene in human heart. Production in the ventricle. *Hypertension*, **17**, 1152-1155.

- Hsieh, J.C., Kodjabachian, L., Rebbert, M.L., Rattner, A., Smallwood, P.M., Samos, C.H., Nusse, R., Dawid, I.B. and Nathans, J. (1999a) A new secreted protein that binds to Wnt proteins and inhibits their activities. *Nature*, **398**, 431-436.
- Hsieh, J.C., Rattner, A., Smallwood, P.M. and Nathans, J. (1999b) Biochemical characterization of Wnt-frizzled interactions using a soluble, biologically active vertebrate Wnt protein. *Proc Natl Acad Sci U S A*, **96**, 3546-3551.
- Huang, Y., Dai, J., Tang, R., Zhao, W., Zhou, Z., Wang, W., Ying, K., Xie, Y. and Mao, Y. (2001) Cloning and characterization of a human lysyl oxidase-like 3 gene (hLOXL3). *Matrix Biol*, **20**, 153-157.
- Huntington, J.A., Olson, S.T., Fan, B. and Gettins, P.G. (1996) Mechanism of heparin activation of antithrombin. Evidence for reactive center loop preinsertion with expulsion upon heparin binding. *Biochemistry*, **35**, 8495-8503.
- Hussain, M.M., Strickland, D.K. and Bakillah, A. (1999) The mammalian low-density lipoprotein receptor family. *Annu Rev Nutr*, **19**, 141-172.
- Hutchinson, H.G., Trindade, P.T., Cunanan, D.B., Wu, C.F. and Pratt, R.E. (1997) Mechanisms of natriuretic-peptide-induced growth inhibition of vascular smooth muscle cells. *Cardiovasc Res*, **35**, 158-167.
- Igaki, T., Itoh, H., Suga, S., Hama, N., Ogawa, Y., Komatsu, Y., Mukoyama, M., Sugawara, A., Yoshimasa, T., Tanaka, I. and Nakao, K. (1996) C-type natriuretic peptide in chronic renal failure and its action in humans. *Kidney Int Suppl*, **55**, S144-147.
- Imada, T., Takayanagi, R. and Inagami, T. (1987) Identification of a peptidase which processes atrial natriuretic factor precursor to its active form with 28 amino acid residues in particulate fractions of rat atrial homogenate. *Biochem Biophys Res Commun*, **143**, 587-592.
- Imada, T., Takayanagi, R. and Inagami, T. (1988) Atrioactivase, a specific peptidase in bovine atria for the processing of pro-atrial natriuretic factor. Purification and characterization. *J Biol Chem*, **263**, 9515-9519.
- John, S.W., Krege, J.H., Oliver, P.M., Hagaman, J.R., Hodgin, J.B., Pang, S.C., Flynn, T.G. and Smithies, O. (1995) Genetic decreases in atrial natriuretic peptide and salt-sensitive hypertension. *Science*, **267**, 679-681.
- Kambayashi, Y., Nakao, K., Mukoyama, M., Saito, Y., Ogawa, Y., Shiono, S., Inouye, K., Yoshida, N. and Imura, H. (1990) Isolation and sequence determination of human brain natriuretic peptide in human atrium. *FEBS Lett*, **259**, 341-345.
- Kangawa, K. and Matsuo, H. (1984) Purification and complete amino acid sequence of alpha-human atrial natriuretic polypeptide (alpha-hANP). *Biochem Biophys Res Commun*, **118**, 131-139.

- Kangawa, K., Tawaragi, Y., Oikawa, S., Mizuno, A., Sakuragawa, Y., Nakazato, H., Fukuda, A., Minamino, N. and Matsuo, H. (1984) Identification of rat gamma atrial natriuretic polypeptide and characterization of the cDNA encoding its precursor. *Nature*, **312**, 152-155.
- Katsube, N., Schwartz, D. and Needleman, P. (1985) Release of atriopeptin in the rat by vasoconstrictors or water immersion correlates with changes in right atrial pressure. *Biochem Biophys Res Commun*, **133**, 937-944.
- Kazama, Y., Hamamoto, T., Foster, D.C. and Kisiel, W. (1995) Hepsin, a putative membrane-associated serine protease, activates human factor VII and initiates a pathway of blood coagulation on the cell surface leading to thrombin formation. *J Biol Chem*, **270**, 66-72.
- Kim, D.R., Sharmin, S., Inoue, M. and Kido, H. (2001) Cloning and expression of novel mosaic serine proteases with and without a transmembrane domain from human lung. *Biochim Biophys Acta*, **1518**, 204-209.
- Kisch, B. (1956) Electron microscopy of the atrium of the heart. I. Guinea pig. *Exp Med Surg*, **14**, 99-112.
- Kishimoto, I., Dubois, S.K. and Garbers, D.L. (1996) The heart communicates with the kidney exclusively through the guanylyl cyclase-A receptor: acute handling of sodium and water in response to volume expansion. *Proc Natl Acad Sci U S A*, **93**, 6215-6219.
- Kitamoto, Y., Veile, R.A., Donis-Keller, H. and Sadler, J.E. (1995) cDNA sequence and chromosomal localization of human enterokinase, the proteolytic activator of trypsinogen. *Biochemistry*, **34**, 4562-4568.
- Knappe, S. (1999) Biochemische Charakterisierung einer neu entdeckten im Herz exprimierten Serinprotease. *Diplomarbeit*. Technische Fachhochschule Berlin
- Kodama, T., Freeman, M., Rohrer, L., Zabrecky, J., Matsudaira, P. and Krieger, M. (1990) Type I macrophage scavenger receptor contains alpha-helical and collagen-like coiled coils. *Nature*, **343**, 531-535.
- Koller, K.J. and Goeddel, D.V. (1992) Molecular biology of the natriuretic peptides and their receptors. *Circulation*, **86**, 1081-1088.
- Komatsu, Y., Itoh, H., Suga, S., Ogawa, Y., Hama, N., Kishimoto, I., Nakagawa, O., Igaki, T., Doi, K., Yoshimasa, T. and Nakao, K. (1996) Regulation of endothelial production of C-type natriuretic peptide in coculture with vascular smooth muscle cells. Role of the vascular natriuretic peptide system in vascular growth inhibition. *Circ Res*, **78**, 606-614.

- Kone, B.C. (2001) Molecular biology of natriuretic peptides and nitric oxide synthases. *Cardiovasc Res*, **51**, 429-441.
- Kraut, J. (1977) Serine proteases: structure and mechanism of catalysis. *Annu Rev Biochem*, **46**, 331-358.
- Kurachi, K., Torres-Rosado, A. and Tsuji, A. (1994) Hepsin. *Methods Enzymol*, **244**, 100-114.
- Kyte, J. and Doolittle, R.F. (1982) A simple method for displaying the hydropathic character of a protein. *J Mol Biol*, **157**, 105-132.
- Ladenson, P.W., Sherman, S.I., Baughman, K.L., Ray, P.E. and Feldman, A.M. (1992) Reversible alterations in myocardial gene expression in a young man with dilated cardiomyopathy and hypothyroidism. *Proc Natl Acad Sci U S A*, **89**, 5251-5255.
- Lämmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, **227**, 680-685.
- Lang, J.C. and Schuller, D.E. (2001) Differential expression of a novel serine protease homologue in squamous cell carcinoma of the head and neck. *Br J Cancer*, **84**, 237-243.
- Lang, R.E., Tholken, H., Ganten, D., Luft, F.C., Ruskoaho, H. and Unger, T. (1985) Atrial natriuretic factor--a circulating hormone stimulated by volume loading. *Nature*, **314**, 264-266.
- Langenickel, T., Pagel, I., Hohnel, K., Dietz, R. and Willenbrock, R. (2000) Differential regulation of cardiac ANP and BNP mRNA in different stages of experimental heart failure. *Am J Physiol Heart Circ Physiol*, **278**, H1500-1506.
- Langenickel, T.H., Pagel, I., Buttgereit, J., Tenner, K., Lindner, M., Dietz, R., Willenbrock, R. and Bader, M. (2004) Rat corin gene: molecular cloning and reduced expression in experimental heart failure. *Am J Physiol Heart Circ Physiol*, **287**, H1516-H1521.
- Ledsome, J.R., Wilson, N., Rankin, A.J. and Courneya, C.A. (1986) Time course of release of atrial natriuretic peptide in the anaesthetized dog. *Can J Physiol Pharmacol*, **64**, 1017-1022.
- Lee, J.S., Ishimoto, A. and Yanagawa, S. (1999) Characterization of mouse dishevelled (Dvl) proteins in Wnt/Wingless signaling pathway. *J Biol Chem*, **274**, 21464-21470.
- Lee, M.E., Miller, W.L., Edwards, B.S. and Burnett, J.C., Jr. (1989) Role of endogenous atrial natriuretic factor in acute congestive heart failure. *J Clin Invest*, **84**, 1962-1966.
- Lee, S.L., Dickson, R.B. and Lin, C.Y. (2000) Activation of hepatocyte growth factor and urokinase/plasminogen activator by matriptase, an epithelial membrane serine protease. *J Biol Chem*, **275**, 36720-36725.

- Levin, E.R., Gardner, D.G. and Samson, W.K. (1998) Natriuretic peptides. *N Engl J Med*, **339**, 321-328.
- Levy, D., Larson, M.G., Vasan, R.S., Kannel, W.B. and Ho, K.K. (1996) The progression from hypertension to congestive heart failure. *Jama*, **275**, 1557-1562.
- Leyns, L., Bouwmeester, T., Kim, S.H., Piccolo, S. and De Robertis, E.M. (1997) Frzb-1 is a secreted antagonist of Wnt signaling expressed in the Spemann organizer. *Cell*, **88**, 747-756.
- Leytus, S.P., Loeb, K.R., Hagen, F.S., Kurachi, K. and Davie, E.W. (1988) A novel trypsin-like serine protease (hepsin) with a putative transmembrane domain expressed by human liver and hepatoma cells. *Biochemistry*, **27**, 1067-1074.
- Light, A. and Janska, H. (1989) Enterokinase (enteropeptidase): comparative aspects. *Trends Biochem Sci*, **14**, 110-112.
- Lin, C.Y., Anders, J., Johnson, M., Sang, Q.A. and Dickson, R.B. (1999) Molecular cloning of cDNA for matriptase, a matrix-degrading serine protease with trypsin-like activity. *J Biol Chem*, **274**, 18231-18236.
- Lohse, P., Mann, W.A., Stein, E.A. and Brewer, H.B., Jr. (1991) Apolipoprotein E-4Philadelphia (Glu13----Lys,Arg145----Cys). Homozygosity for two rare point mutations in the apolipoprotein E gene combined with severe type III hyperlipoproteinemia. *J Biol Chem*, **266**, 10479-10484.
- Lopez, M.J., Garbers, D.L. and Kuhn, M. (1997) The guanylyl cyclase-deficient mouse defines differential pathways of natriuretic peptide signaling. *J Biol Chem*, **272**, 23064-23068.
- Lopez, M.J., Wong, S.K., Kishimoto, I., Dubois, S., Mach, V., Friesen, J., Garbers, D.L. and Beuve, A. (1995) Salt-resistant hypertension in mice lacking the guanylyl cyclase-A receptor for atrial natriuretic peptide. *Nature*, **378**, 65-68.
- Lu, D., Futterer, K., Korolev, S., Zheng, X., Tan, K., Waksman, G. and Sadler, J.E. (1999) Crystal structure of enteropeptidase light chain complexed with an analog of the trypsinogen activation peptide. *J Mol Biol*, **292**, 361-373.
- Lu, D., Yuan, X., Zheng, X. and Sadler, J.E. (1997) Bovine proenteropeptidase is activated by trypsin, and the specificity of enteropeptidase depends on the heavy chain. *J Biol Chem*, **272**, 31293-31300.
- Lucas, K.A., Pitari, G.M., Kazerounian, S., Ruiz-Stewart, I., Park, J., Schulz, S., Chepenik, K.P. and Waldman, S.A. (2000) Guanylyl cyclases and signaling by cyclic GMP. *Pharmacol Rev*, **52**, 375-414.

- Madison, E.L., Kobe, A., Gething, M.J., Sambrook, J.F. and Goldsmith, E.J. (1993) Converting tissue plasminogen activator to a zymogen: a regulatory triad of Asp-His-Ser. *Science*, **262**, 419-421.
- Maki, M., Parmentier, M. and Inagami, T. (1984) Cloning of genomic DNA for human atrial natriuretic factor. *Biochem Biophys Res Commun*, **125**, 797-802.
- Mann, K.G., Nesheim, M.E., Church, W.R., Haley, P. and Krishnaswamy, S. (1990) Surface-dependent reactions of the vitamin K-dependent enzyme complexes. *Blood*, **76**, 1-16.
- Marchioli, C.C. and Graziano, S.L. (1997) Paraneoplastic syndromes associated with small cell lung cancer. *Chest Surg Clin N Am*, **7**, 65-80.
- Maroux, S., Baratti, J. and Desnuelle, P. (1971) Purification and specificity of porcine enterokinase. *J Biol Chem*, **246**, 5031-5039.
- Masiakowski, P. and Yancopoulos, G.D. (1998) The Wnt receptor CRD domain is also found in MuSK and related orphan receptor tyrosine kinases. *Curr Biol*, **8**, R407.
- Matsumoto, A., Naito, M., Itakura, H., Ikemoto, S., Asaoka, H., Hayakawa, I., Kanamori, H., Aburatani, H., Takaku, F., Suzuki, H. and et al. (1990) Human macrophage scavenger receptors: primary structure, expression, and localization in atherosclerotic lesions. *Proc Natl Acad Sci U S A*, **87**, 9133-9137.
- Mattingly, M.T., Brandt, R.R., Heublein, D.M., Wei, C.M., Nir, A. and Burnett, J.C., Jr. (1994) Presence of C-type natriuretic peptide in human kidney and urine. *Kidney Int*, **46**, 744-747.
- Michener, M.L., Gierse, J.K., Seetharam, R., Fok, K.F., Olins, P.O., Mai, M.S. and Needleman, P. (1986) Proteolytic processing of atriopeptin prohormone. *Mol Pharmacol*, **30**, 552-557.
- Mikhailenko, I., Considine, W., Argraves, K.M., Loukinov, D., Hyman, B.T. and Strickland, D.K. (1999) Functional domains of the very low density lipoprotein receptor: molecular analysis of ligand binding and acid-dependent ligand dissociation mechanisms. *J Cell Sci*, **112 (Pt 19)**, 3269-3281.
- Minamino, N., Aburaya, M., Ueda, S., Kangawa, K. and Matsuo, H. (1988) The presence of brain natriuretic peptide of 12,000 daltons in porcine heart. *Biochem Biophys Res Commun*, **155**, 740-746.
- Minamino, N., Kangawa, K. and Matsuo, H. (1990) N-terminally extended form of C-type natriuretic peptide (CNP-53) identified in porcine brain. *Biochem Biophys Res Commun*, **170**, 973-979.
- Moeller, C., Swindell, E.C., Kispert, A. and Eichele, G. (2003) Carboxypeptidase Z (CPZ) modulates Wnt signaling and regulates the development of skeletal elements in the chicken. *Development*, **130**, 5103-5111.

- Mukoyama, M., Nakao, K., Hosoda, K., Suga, S., Saito, Y., Ogawa, Y., Shirakami, G., Jougasaki, M., Obata, K., Yasue, H. and et al. (1991) Brain natriuretic peptide as a novel cardiac hormone in humans. Evidence for an exquisite dual natriuretic peptide system, atrial natriuretic peptide and brain natriuretic peptide. *J Clin Invest*, **87**, 1402-1412.
- Nakayama, K., Ohkubo, H., Hirose, T., Inayama, S. and Nakanishi, S. (1984) mRNA sequence for human cardiodilatin-atrial natriuretic factor precursor and regulation of precursor mRNA in rat atria. *Nature*, **310**, 699-701.
- Neels, J.G., van Den Berg, B.M., Lookene, A., Olivecrona, G., Pannekoek, H. and van Zonneveld, A.J. (1999) The second and fourth cluster of class A cysteine-rich repeats of the low density lipoprotein receptor-related protein share ligand-binding properties. *J Biol Chem*, **274**, 31305-31311.
- Neurath, H. (1984) Evolution of proteolytic enzymes. *Science*, **224**, 350-357.
- North, C.L. and Blacklow, S.C. (1999) Structural independence of ligand-binding modules five and six of the LDL receptor. *Biochemistry*, **38**, 3926-3935.
- North, C.L. and Blacklow, S.C. (2000) Evidence that familial hypercholesterolemia mutations of the LDL receptor cause limited local misfolding in an LDL-A module pair. *Biochemistry*, **39**, 13127-13135.
- Novikova, E.G., Reznik, S.E., Varlamov, O. and Fricker, L.D. (2000) Carboxypeptidase Z is present in the regulated secretory pathway and extracellular matrix in cultured cells and in human tissues. *J Biol Chem*, **275**, 4865-4870.
- Nusse, R. and Varmus, H.E. (1992) Wnt genes. *Cell*, **69**, 1073-1087.
- Nykjaer, A. and Willnow, T.E. (2002) The low-density lipoprotein receptor gene family: a cellular Swiss army knife? *Trends Cell Biol*, **12**, 273-280.
- Oberst, M.D., Williams, C.A., Dickson, R.B., Johnson, M.D. and Lin, C.Y. (2003) The activation of matriptase requires its noncatalytic domains, serine protease domain, and its cognate inhibitor. *J Biol Chem*, **278**, 26773-26779. Epub 22003 May 26778.
- Ogawa, Y., Nakao, K., Nakagawa, O., Komatsu, Y., Hosoda, K., Suga, S., Arai, H., Nagata, K., Yoshida, N. and Imura, H. (1992) Human C-type natriuretic peptide. Characterization of the gene and peptide. *Hypertension*, **19**, 809-813.
- Oikawa, S., Imai, M., Ueno, A., Tanaka, S., Noguchi, T., Nakazato, H., Kangawa, K., Fukuda, A. and Matsuo, H. (1984) Cloning and sequence analysis of cDNA encoding a precursor for human atrial natriuretic polypeptide. *Nature*, **309**, 724-726.

- Oliver, P.M., Fox, J.E., Kim, R., Rockman, H.A., Kim, H.S., Reddick, R.L., Pandey, K.N., Milgram, S.L., Smithies, O. and Maeda, N. (1997) Hypertension, cardiac hypertrophy, and sudden death in mice lacking natriuretic peptide receptor A. *Proc Natl Acad Sci U S A*, **94**, 14730-14735.
- Oliver, P.M., John, S.W., Purdy, K.E., Kim, R., Maeda, N., Goy, M.F. and Smithies, O. (1998) Natriuretic peptide receptor 1 expression influences blood pressures of mice in a dose-dependent manner. *Proc Natl Acad Sci U S A*, **95**, 2547-2551.
- Pan, J., Hinzmann, B., Yan, W., Wu, F., Morser, J. and Wu, Q. (2002) Genomic structures of the human and murine corin genes and functional GATA elements in their promoters. *J Biol Chem*, **277**, 38390-38398.
- Pandey, K.N., Inagami, T. and Misono, K.S. (1987) Three distinct forms of atrial natriuretic factor receptors: kidney tubular epithelium cells and vascular smooth muscle cells contain different types of receptors. *Biochem Biophys Res Commun*, **147**, 1146-1152.
- Pandey, K.N. and Orgebin-Crist, M.C. (1991) Atrial natriuretic factor in mammalian testis: immunological detection in spermatozoa. *Biochem Biophys Res Commun*, **180**, 437-444.
- Paoloni-Giacobino, A., Chen, H., Peitsch, M.C., Rossier, C. and Antonarakis, S.E. (1997) Cloning of the TMPRSS2 gene, which encodes a novel serine protease with transmembrane, LDLRA, and SRCR domains and maps to 21q22.3. *Genomics*, **44**, 309-320.
- Patel, A.M., Davila, D.G. and Peters, S.G. (1993) Paraneoplastic syndromes associated with lung cancer. *Mayo Clin Proc*, **68**, 278-287.
- Potter, L.R. and Hunter, T. (2001) Guanylyl cyclase-linked natriuretic peptide receptors: structure and regulation. *J Biol Chem*, **276**, 6057-6060.
- Proba, K., Gschwend, T.P. and Sonderegger, P. (1998) Cloning and sequencing of the cDNA encoding human neurotrypsin. *Biochim Biophys Acta*, **1396**, 143-147.
- Rattner, A., Hsieh, J.C., Smallwood, P.M., Gilbert, D.J., Copeland, N.G., Jenkins, N.A. and Nathans, J. (1997) A family of secreted proteins contains homology to the cysteine-rich ligand-binding domain of frizzled receptors. *Proc Natl Acad Sci U S A*, **94**, 2859-2863.
- Rehn, M. and Pihlajaniemi, T. (1995) Identification of three N-terminal ends of type XVIII collagen chains and tissue-specific differences in the expression of the corresponding transcripts. The longest form contains a novel motif homologous to rat and *Drosophila* frizzled proteins. *J Biol Chem*, **270**, 4705-4711.

- Resnick, D., Chatterton, J.E., Schwartz, K., Slayter, H. and Krieger, M. (1996) Structures of class A macrophage scavenger receptors. Electron microscopic study of flexible, multidomain, fibrous proteins and determination of the disulfide bond pattern of the scavenger receptor cysteine-rich domain. *J Biol Chem*, **271**, 26924-26930.
- Resnick, D., Pearson, A. and Krieger, M. (1994) The SRCR superfamily: a family reminiscent of the Ig superfamily. *Trends Biochem Sci*, **19**, 5-8.
- Rettenberger, P.M., Oka, K., Ellgaard, L., Petersen, H.H., Christensen, A., Martensen, P.M., Monard, D., Etzerodt, M., Chan, L. and Andreasen, P.A. (1999) Ligand binding properties of the very low density lipoprotein receptor. Absence of the third complement-type repeat encoded by exon 4 is associated with reduced binding of Mr 40,000 receptor-associated protein. *J Biol Chem*, **274**, 8973-8980.
- Riddell, D.R., Vinogradov, D.V., Stannard, A.K., Chadwick, N. and Owen, J.S. (1999) Identification and characterization of LRP8 (apoER2) in human blood platelets. *J Lipid Res*, **40**, 1925-1930.
- Rosenzweig, A. and Seidman, C.E. (1991) Atrial natriuretic factor and related peptide hormones. *Annu Rev Biochem*, **60**, 229-255.
- Ruskoaho, H., Tholken, H. and Lang, R.E. (1986) Increase in atrial pressure releases atrial natriuretic peptide from isolated perfused rat hearts. *Pflugers Arch*, **407**, 170-174.
- Ruskoaho, H. (2003) Cardiac hormones as diagnostic tools in heart failure. *Endocr Rev*, **24**, 341-356
- Sato, H., Kinoshita, T., Takino, T., Nakayama, K. and Seiki, M. (1996) Activation of a recombinant membrane type 1-matrix metalloproteinase (MT1-MMP) by furin and its interaction with tissue inhibitor of metalloproteinases (TIMP)-2. *FEBS Lett*, **393**, 101-104.
- Sawada, Y., Suda, M., Yokoyama, H., Kanda, T., Sakamaki, T., Tanaka, S., Nagai, R., Abe, S. and Takeuchi, T. (1997) Stretch-induced hypertrophic growth of cardiocytes and processing of brain-type natriuretic peptide are controlled by proprotein-processing endopeptidase furin. *J Biol Chem*, **272**, 20545-20554.
- Schechter, I. and Berger, A. (1967) On the size of the active site in proteases. I. Papain. *Biochem Biophys Res Commun*, **27**, 157-162.
- Schiebinger, R.J. and Gomez-Sanchez, C.E. (1990) Endothelin: a potent stimulus of atrial natriuretic peptide secretion by superfused rat atria and its dependency on calcium. *Endocrinology*, **127**, 119-125.
- Schiebinger, R.J. and Linden, J. (1986) The influence of resting tension on immunoreactive atrial natriuretic peptide secretion by rat atria superfused *in vitro*. *Circ Res*, **59**, 105-109.

- Schwartz, D., Geller, D.M., Manning, P.T., Siegel, N.R., Fok, K.F., Smith, C.E. and Needleman, P. (1985) Ser-Leu-Arg-Arg-atriopeptin III: the major circulating form of atrial peptide. *Science*, **229**, 397-400.
- Schweitz, H., Vigne, P., Moinier, D., Frelin, C. and Lazdunski, M. (1992) A new member of the natriuretic peptide family is present in the venom of the green mamba (*Dendroaspis angusticeps*). *J Biol Chem*, **267**, 13928-13932.
- Scott, H.S., Kudoh, J., Wattenhofer, M., Shibuya, K., Berry, A., Chrast, R., Guipponi, M., Wang, J., Kawasaki, K., Asakawa, S., Minoshima, S., Younus, F., Mehdi, S.Q., Radhakrishna, U., Papasavvas, M.P., Gehrig, C., Rossier, C., Korostishevsky, M., Gal, A., Shimizu, N., Bonne-Tamir, B. and Antonarakis, S.E. (2001) Insertion of beta-satellite repeats identifies a transmembrane protease causing both congenital and childhood onset autosomal recessive deafness. *Nat Genet*, **27**, 59-63.
- Sei, C.A., Hand, G.L., Murray, S.F. and Glembotski, C.C. (1992) The cosecretional maturation of atrial natriuretic factor by primary atrial myocytes. *Mol Endocrinol*, **6**, 309-319.
- Seidah, N.G. and Chretien, M. (1999) Proprotein and prohormone convertases: a family of subtilases generating diverse bioactive polypeptides. *Brain Res*, **848**, 45-62.
- Seidah, N.G., Cromlish, J.A., Hamelin, J., Thibault, G. and Chretien, M. (1986) Homologous IRCM-serine protease 1 from pituitary, heart atrium and ventricle: a common prohormone maturation enzyme? *Biosci Rep*, **6**, 835-844.
- Seidman, C.E., Bloch, K.D., Klein, K.A., Smith, J.A. and Seidman, J.G. (1984) Nucleotide sequences of the human and mouse atrial natriuretic factor genes. *Science*, **226**, 1206-1209.
- Shields, P.P. and Glembotski, C.C. (1988) The post-translational processing of rat pro-atrial natriuretic factor by primary atrial myocyte cultures. *J Biol Chem*, **263**, 8091-8098.
- Shimomura, T., Denda, K., Kitamura, A., Kawaguchi, T., Kito, M., Kondo, J., Kagaya, S., Qin, L., Takata, H., Miyazawa, K. and Kitamura, N. (1997) Hepatocyte growth factor activator inhibitor, a novel Kunitz-type serine protease inhibitor. *J Biol Chem*, **272**, 6370-6376.
- Shinomiya, M., Tashiro, J., Saito, Y., Yoshida, S., Furuya, M., Oka, N., Tanaka, S., Kangawa, K. and Matsuo, H. (1994) C-type natriuretic peptide inhibits intimal thickening of rabbit carotid artery after balloon catheter injury. *Biochem Biophys Res Commun*, **205**, 1051-1056.
- Shiono, S., Nakao, K., Morii, N., Yamada, T., Itoh, H., Sakamoto, M., Sugawara, A., Saito, Y., Katsuura, G. and Imura, H. (1986) Nature of atrial natriuretic polypeptide in rat brain. *Biochem Biophys Res Commun*, **135**, 728-734.

- Silberbach, M. and Roberts, C.T., Jr. (2001) Natriuretic peptide signalling: molecular and cellular pathways to growth regulation. *Cell Signal*, **13**, 221-231.
- Song, L. and Fricker, L.D. (1997) Cloning and expression of human carboxypeptidase Z, a novel metallocarboxypeptidase. *J Biol Chem*, **272**, 10543-10550.
- Stanley, K.K., Kocher, H.P., Luzio, J.P., Jackson, P. and Tschopp, J. (1985) The sequence and topology of human complement component C9. *Embo J*, **4**, 375-382.
- Steiner, D.F. (1998) The proprotein convertases. *Curr Opin Chem Biol*, **2**, 31-39.
- Steinhelper, M.E. (1993) Structure, expression, and genomic mapping of the mouse natriuretic peptide type-B gene. *Circ Res*, **72**, 984-992.
- Steinhelper, M.E., Cochrane, K.L. and Field, L.J. (1990) Hypotension in transgenic mice expressing atrial natriuretic factor fusion genes. *Hypertension*, **16**, 301-307.
- Stroud, R.M. (1974) A family of protein-cutting proteins. *Sci Am*, **231**, 74-88.
- Stubbs, M.T., Renatus, M. and Bode, W. (1998) An active zymogen: unravelling the mystery of tissue-type plasminogen activator. *Biol Chem*, **379**, 95-103.
- Sudoh, T., Kangawa, K., Minamino, N. and Matsuo, H. (1988) A new natriuretic peptide in porcine brain. *Nature*, **332**, 78-81.
- Sudoh, T., Minamino, N., Kangawa, K. and Matsuo, H. (1990) C-type natriuretic peptide (CNP): a new member of natriuretic peptide family identified in porcine brain. *Biochem Biophys Res Commun*, **168**, 863-870.
- Suga, S., Nakao, K., Hosoda, K., Mukoyama, M., Ogawa, Y., Shirakami, G., Arai, H., Saito, Y., Kambayashi, Y., Inouye, K. and et al. (1992a) Receptor selectivity of natriuretic peptide family, atrial natriuretic peptide, brain natriuretic peptide, and C-type natriuretic peptide. *Endocrinology*, **130**, 229-239.
- Suga, S., Nakao, K., Itoh, H., Komatsu, Y., Ogawa, Y., Hama, N. and Imura, H. (1992b) Endothelial production of C-type natriuretic peptide and its marked augmentation by transforming growth factor-beta. Possible existence of "vascular natriuretic peptide system". *J Clin Invest*, **90**, 1145-1149.
- Sugiyama, T., Kumagai, H., Morikawa, Y., Wada, Y., Sugiyama, A., Yasuda, K., Yokoi, N., Tamura, S., Kojima, T., Nosaka, T., Senba, E., Kimura, S., Kadowaki, T., Kodama, T. and Kitamura, T. (2000) A novel low-density lipoprotein receptor-related protein mediating cellular uptake of apolipoprotein E-enriched beta-VLDL *in vitro*. *Biochemistry*, **39**, 15817-15825.
- Suzuki, E., Hirata, Y., Hayakawa, H., Omata, M., Kojima, M., Kangawa, K., Minamino, N. and Matsuo, H. (1993) Evidence for C-type natriuretic peptide production in the rat kidney. *Biochem Biophys Res Commun*, **192**, 532-538.

- Suzuki, H., Kurihara, Y., Takeya, M., Kamada, N., Kataoka, M., Jishage, K., Ueda, O., Sakaguchi, H., Higashi, T., Suzuki, T., Takashima, Y., Kawabe, Y., Cynshi, O., Wada, Y., Honda, M., Kurihara, H., Aburatani, H., Doi, T., Matsumoto, A., Azuma, S., Noda, T., Toyoda, Y., Itakura, H., Yazaki, Y., Kodama, T. (1997) A role for macrophage scavenger receptors in atherosclerosis and susceptibility to infection. *Nature*, **386**, 292-296.
- Suzuki, T., Yamazaki, T. and Yazaki, Y. (2001) The role of the natriuretic peptides in the cardiovascular system. *Cardiovasc Res*, **51**, 489-494.
- Szabo, R., Wu, Q., Dickson, R.B., Netzel-Arnett, S., Antalis, T.M. and Bugge, T.H. (2003) Type II transmembrane serine proteases. *Thromb Haemost*, **90**, 185-193.
- Takeuchi, T., Harris, J.L., Huang, W., Yan, K.W., Coughlin, S.R. and Craik, C.S. (2000) Cellular localization of membrane-type serine protease 1 and identification of protease-activated receptor-2 and single-chain urokinase-type plasminogen activator as substrates. *J Biol Chem*, **275**, 26333-26342.
- Takeuchi, T., Shuman, M.A. and Craik, C.S. (1999) Reverse biochemistry: use of macromolecular protease inhibitors to dissect complex biological processes and identify a membrane-type serine protease in epithelial cancer and normal tissue. *Proc Natl Acad Sci U S A*, **96**, 11054-11061.
- Tamura, N., Ogawa, Y., Chusho, H., Nakamura, K., Nakao, K., Suda, M., Kasahara, M., Hashimoto, R., Katsuura, G., Mukoyama, M., Itoh, H., Saito, Y., Tanaka, I., Otani, H. and Katsuki, M. (2000) Cardiac fibrosis in mice lacking brain natriuretic peptide. *Proc Natl Acad Sci U S A*, **97**, 4239-4244.
- Tawaragi, Y., Fuchimura, K., Tanaka, S., Minamino, N., Kangawa, K. and Matsuo, H. (1991) Gene and precursor structures of human C-type natriuretic peptide. *Biochem Biophys Res Commun*, **175**, 645-651.
- Thibault, G., Garcia, R., Gutkowska, J., Bilodeau, J., Lazure, C., Seidah, N.G., Chretien, M., Genest, J. and Cantin, M. (1987) The propeptide Asn1-Tyr126 is the storage form of rat atrial natriuretic factor. *Biochem J*, **241**, 265-272.
- Thibault, G., Lazure, C., Schiffrin, E.L., Gutkowska, J., Chartier, L., Garcia, R., Seidah, N.G., Chretien, M., Genest, J. and Cantin, M. (1985) Identification of a biologically active circulating form of rat atrial natriuretic factor. *Biochem Biophys Res Commun*, **130**, 981-986.
- Thibault, G., Nemer, M., Drouin, J., Lavigne, J.P., Ding, J., Charbonneau, C., Garcia, R., Genest, J., Jasmin, G., Sole, M. and et al. (1989) Ventricles as a major site of atrial natriuretic factor synthesis and release in cardiomyopathic hamsters with heart failure. *Circ Res*, **65**, 71-82.

- Thomas, C.A., Li, Y., Kodama, T., Suzuki, H., Silverstein, S.C. and El Khoury, J. (2000) Protection from lethal gram-positive infection by macrophage scavenger receptor-dependent phagocytosis. *J Exp Med*, **191**, 147-156.
- Tomita, Y., Kim, D.H., Magoori, K., Fujino, T. and Yamamoto, T.T. (1998) A novel low-density lipoprotein receptor-related protein with type II membrane protein-like structure is abundant in heart. *J Biochem (Tokyo)*, **124**, 784-789.
- Towbin, H., Staehelin, T. and Gordon, J. (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci U S A*, **76**, 4350-4354.
- Tran, K.L., Lu, X., Lei, M., Feng, Q. and Wu, Q. (2004) Upregulation of corin gene expression in hypertrophic cardiomyocytes and failing myocardium. *Am J Physiol Heart Circ Physiol*, **287**, H1625-H1631.
- Ueno, H., Haruno, A., Morisaki, N., Furuya, M., Kangawa, K., Takeshita, A. and Saito, Y. (1997) Local expression of C-type natriuretic peptide markedly suppresses neointimal formation in rat injured arteries through an autocrine/paracrine loop. *Circulation*, **96**, 2272-2279.
- Velasco, G., Cal, S., Quesada, V., Sanchez, L.M. and Lopez-Otin, C. (2002) Matriptase-2, a membrane-bound mosaic serine proteinase predominantly expressed in human liver and showing degrading activity against extracellular matrix proteins. *J Biol Chem*, **277**, 37637-37646.
- Vinson, C.R., Conover, S. and Adler, P.N. (1989) A Drosophila tissue polarity locus encodes a protein containing seven potential transmembrane domains. *Nature*, **338**, 263-264.
- Vu, T.K., Liu, R.W., Haaksma, C.J., Tomasek, J.J. and Howard, E.W. (1997) Identification and cloning of the membrane-associated serine protease, hepsin, from mouse preimplantation embryos. *J Biol Chem*, **272**, 31315-31320.
- Wada, A., Tsutamoto, T., Matsuda, Y. and Kinoshita, M. (1994) Cardiorenal and neurohumoral effects of endogenous atrial natriuretic peptide in dogs with severe congestive heart failure using a specific antagonist for guanylate cyclase-coupled receptors. *Circulation*, **89**, 2232-2240.
- Wallin, R., Stanton, C. and Ross, R.P. (1994) Intracellular proteolytic processing of the two-chain vitamin K-dependent coagulation factor X. *Thromb Res*, **73**, 395-403.
- Wallrapp, C., Hahnel, S., Muller-Pillasch, F., Burghardt, B., Iwamura, T., Ruthenburger, M., Lerch, M.M., Adler, G. and Gress, T.M. (2000) A novel transmembrane serine protease (TMPRSS3) overexpressed in pancreatic cancer. *Cancer Res*, **60**, 2602-2606.

- Wardell, M.R., Rall, S.C., Jr., Schaefer, E.J., Kane, J.P. and Weisgraber, K.H. (1991) Two apolipoprotein E5 variants illustrate the importance of the position of additional positive charge on receptor-binding activity. *J Lipid Res*, **32**, 521-528.
- Wei, C.M., Heublein, D.M., Perrella, M.A., Lerman, A., Rodeheffer, R.J., McGregor, C.G., Edwards, W.D., Schaff, H.V. and Burnett, J.C., Jr. (1993) Natriuretic peptide system in human heart failure. *Circulation*, **88**, 1004-1009.
- Wei, C.M., Kim, C.H., Khraibi, A.A., Miller, V.M. and Burnett, J.C., Jr. (1994) Atrial natriuretic peptide and C-type natriuretic peptide in spontaneously hypertensive rats and their vasorelaxing actions *in vitro*. *Hypertension*, **23**, 903-907.
- White, S.M., Constantin, P.E. and Claycomb, W.C. (2004) Cardiac physiology at the cellular level: use of cultured HL-1 cardiomyocytes for studies of cardiac muscle cell structure and function. *Am J Physiol Heart Circ Physiol*, **286**, H823-829.
- Whitney, G.S., Starling, G.C., Bowen, M.A., Modrell, B., Siadak, A.W. and Aruffo, A. (1995) The membrane-proximal scavenger receptor cysteine-rich domain of CD6 contains the activated leukocyte cell adhesion molecule binding site. *J Biol Chem*, **270**, 18187-18190.
- Wilson, I.A. and Jolliffe, L.K. (1999) The structure, organization, activation and plasticity of the erythropoietin receptor. *Curr Opin Struct Biol*, **9**, 696-704.
- Wodarz, A. and Nusse, R. (1998) Mechanisms of Wnt signaling in development. *Annu Rev Cell Dev Biol*, **14**, 59-88.
- Wu, C., Wu, F., Pan, J., Morser, J. and Wu, Q. (2003) Furin-mediated processing of Pro-C-type natriuretic peptide. *J Biol Chem*, **278**, 25847-25852.
- Wu, F. and Wu, Q. (2003) Corin-mediated processing of pro-atrial natriuretic peptide in human small cell lung cancer cells. *Cancer Res*, **63**, 8318-8322.
- Wu, F., Yan, W., Pan, J., Morser, J. and Wu, Q. (2002) Processing of pro-atrial natriuretic peptide by corin in cardiac myocytes. *J Biol Chem*, **277**, 16900-16905.
- Wu, Q. (2001) Gene targeting in hemostasis. Hepsin. *Front Biosci*, **6**, D192-200.
- Wu, Q. (2003) Type II transmembrane serine proteases. *Curr Top Dev Biol*, **54**, 167-206
- Wu, Q.Y., Sheehan, J.P., Tsiang, M., Lentz, S.R., Birktoft, J.J. and Sadler, J.E. (1991) Single amino acid substitutions dissociate fibrinogen-clotting and thrombomodulin-binding activities of human thrombin. *Proc Natl Acad Sci U S A*, **88**, 6775-6779.
- Wypij, D.M. and Harris, R.B. (1988) Atrial granules contain an amino-terminal processing enzyme of atrial natriuretic factor. *J Biol Chem*, **263**, 7079-7086.
- Xu, Y.K. and Nusse, R. (1998) The Frizzled CRD domain is conserved in diverse proteins including several receptor tyrosine kinases. *Curr Biol*, **8**, R405-406.

- Yamaguchi, N., Okui, A., Yamada, T., Nakazato, H. and Mitsui, S. (2002) Spinesin/TMPRSS5, a novel transmembrane serine protease, cloned from human spinal cord. *J Biol Chem*, **277**, 6806-6812.
- Yamahara, K., Itoh, H., Chun, T.H., Ogawa, Y., Yamashita, J., Sawada, N., Fukunaga, Y., Sone, M., Yurugi-Kobayashi, T., Miyashita, K., Tsujimoto, H., Kook, H., Feil, R., Garbers, D.L., Hofmann, F. and Nakao, K. (2003) Significance and therapeutic potential of the natriuretic peptides/cGMP/cGMP-dependent protein kinase pathway in vascular regeneration. *Proc Natl Acad Sci U S A*, **100**, 3404-3409.
- Yamamoto, T., Davis, C.G., Brown, M.S., Schneider, W.J., Casey, M.L., Goldstein, J.L. and Russell, D.W. (1984) The human LDL receptor: a cysteine-rich protein with multiple Alu sequences in its mRNA. *Cell*, **39**, 27-38.
- Yamanaka, M., Greenberg, B., Johnson, L., Seilhamer, J., Brewer, M., Friedemann, T., Miller, J., Atlas, S., Laragh, J., Lewicki, J. and et al. (1984) Cloning and sequence analysis of the cDNA for the rat atrial natriuretic factor precursor. *Nature*, **309**, 719-722.
- Yamaoka, K., Masuda, K., Ogawa, H., Takagi, K., Umemoto, N. and Yasuoka, S. (1998) Cloning and characterization of the cDNA for human airway trypsin-like protease. *J Biol Chem*, **273**, 11895-11901.
- Yan, W., Sheng, N., Seto, M., Morser, J. and Wu, Q. (1999) Corin, a mosaic transmembrane serine protease encoded by a novel cDNA from human heart. *J Biol Chem*, **274**, 14926-14935.
- Yan, W., Wu, F., Morser, J. and Wu, Q. (2000) Corin, a transmembrane cardiac serine protease, acts as a pro-atrial natriuretic peptide-converting enzyme. *Proc Natl Acad Sci U S A*, **97**, 8525-8529.
- Yandle, T.G. (1994) Biochemistry of natriuretic peptides. *J Intern Med*, **235**, 561-576.
- Yang-Feng, T.L., Floyd-Smith, G., Nemer, M., Drouin, J. and Francke, U. (1985) The pronatriodilatin gene is located on the distal short arm of human chromosome 1 and on mouse chromosome 4. *Am J Hum Genet*, **37**, 1117-1128.
- Yasue, H., Yoshimura, M., Sumida, H., Kikuta, K., Kugiyama, K., Jougasaki, M., Ogawa, H., Okumura, K., Mukoyama, M. and Nakao, K. (1994) Localization and mechanism of secretion of B-type natriuretic peptide in comparison with those of A-type natriuretic peptide in normal subjects and patients with heart failure. *Circulation*, **90**, 195-203.
- Zheng, X. and Sadler, J.E. (2002) Mucin-like domain of enteropeptidase directs apical targeting in Madin-Darby canine kidney cells. *J Biol Chem*, **277**, 6858-6863.

- Zhou, A., Webb, G., Zhu, X. and Steiner, D.F. (1999) Proteolytic processing in the secretory pathway. *J Biol Chem*, **274**, 20745-20748.
- Zhukov, A., Hellman, U. and Ingelman-Sundberg, M. (1997) Purification and characterization of hepsin from rat liver microsomes. *Biochim Biophys Acta*, **1337**, 85-95.