

6 Literaturverzeichnis

Abd-Alrahman N., Dokmak A.S., Abou-Madawi A. (1994). Anterior cervical discectomy (ACD) versus cervical fusion (ACF): Clinical and radiological outcome study. *Acta Neurochirurgica*; 141:1089-1092.

Alexander J.T, Branch C.L., Subach B.R., Haid R.W. (2002). Application of a resorbable interbody spacer in posterior lumbar interbody fusion. *J Neurosurg; (Spine 4)* 97: 468-472.

An H. S., Simpson J.M., Glover J.M., Stephany J. (1995). Comparison between allograft plus demineralised bone matrix versus autograft in anterior cervical fusion. A prospective multicenter study. *Spine*; 20 (20): 2211-2216.

Arrington E.D, Smith W. J., Chambers H.G., Bucknell A.L., Davino N. (1996). Complications of iliac bone graft harvesting. *Clin Orthop*; 329: 300-309.

Bagby G.W. (1988). Arthrodesis by the distraction-compression method using a stainless steel implant. *Orthopedics* 11: 931-34.

Banwart J.C., Asher M.A., Hassanein R.S. (1995). Iliac crest bone graft harvest donor site morbidity. A statistical evaluation. *Spine*; 20 (9): 1055-1060.

Bergsma E.J., De Bruijn W.C., Rozema F.R. and Boering G. (1995). Late degradation tissue response to poly (L -lactid) bone plates and screws. *Biomaterials* 16:25-31.

Bergsma E.J., Rozema, F.R. Bos R.R.M., and Bruijn W.C. (1993). Foreign body reaction to resorbable poly (L-lactid) bone plates and screws used for the fixation of unstable zygomatic fractures. *J Oral Maxillofac Surg*; 51(6): 666-70.

Biovision GmbH Ilmenau, Gespräch mit Herrn Günther

Bishop R.C., Moore K.A., Hadley M.N. (1996). Anterior cervical interbody fusion using autogeneic and allogeneic bone graft substrate: A prospective comparative analysis. *J Neurosurg*; 85: 206-210.

- Blauth, M., Schmidt, U., Bastin, L., Knop, C.,Tscherne, H.** (1998). Die ventrale interkorporelle Spondylodese bei Verletzungen der Halswirbelsäule. Indikation, Operationstechnik und Ergebnisse. Zentralblatt für Chirurgie; 123: 919-929.
- Boden S.D. and Summer D.R.** (1995). Biological factors affecting spinal fusion and bone regeneration. Spine; 20 (24Suppl): 102-12.
- Borke D.S., Rorke A.W., King A.T., Gullan R.W.** (1997). Preliminary experience of carbon fibre cage prosthesis for treatment of cervical spine disorders. J Neurosurg Br; 11: 221-227
- Böstman O.** (1991 a). Current concepts review. Absorbable implants for the fixation of fractures. J Bone Joint Surg Am.; 73 (1): 148-152.
- Böstman O.** (1991 b). Osteolytic changes accompanying degradation of absorbable fracture fixation implants. J Bone Joint Surg Br.; 73: 679-682.
- Böstman O., Hirvensalo E., Vainionpää S., et al.** (1989). Ankle fractures treated using biodegradable internal fixation. Clin Orthop Rela Res; 238: 195-203.
- Böstman O., Hirvensalo E., Makinen J. and Rokkanen P.** (1990). Foreign-body reaction to fracture fixation implants of biodegradable synthetic polymers. J Bone Joint Surg; 65B (1): 89-96.
- Böstman O., Pihlajamäki H.** (2000). Clinical biocompatibility of biodegradable orthopaedic implants for internal fixation: A review. Biomaterials; 21: 2615-2621.
- Böstman O., Pihlajamäki H., Partio E., Rokkanen P.** (1995). Clinical biocomparbility and degradation of polylevolactides Screws in the ankle. Clin Orthop and Rela Res; 320: 101-109.
- Brady J. M., Cutright D.E., Miller R.A. and Batistone G.C.** (1973). Resorption rate, route of elimination, and ultra structure of the implant site of polylactic acid in the abdominal wall of the rat. J Biomed Mater Res; 7: 155-166.
- Brantigan J.W. and Steffee A.D.** (2000). Lumbal interbody fusion using the brantigan I/F cage for posterior lumbar interbody fusion and the variable pedicle screw placement system: Two-Year results from a food drug administration inverstigational device exemption clinical trail. Spine; 25 (11): 1437-1446.

Brantigan J.W. and Steffee A.D. (1993). A carbon fibre implantat to aid interbody lumbar fusion. Two-Year clinical result in the first 26 patients. *Spine*; 18 (14): 2106-2117.

Brantigan J.W., Mc Affee P.C., Cunningham B.W., Wang H., Orbegoso C.M. (1994). Interbody lumbar fusion using a carbon fibre cage implantat versus allograft bone. *Spine*; 19 (13): 1436-1444.

Bremer A.M. and Nguyen T.Q. (1983). Internal metal plate fixation combined with anterior interbody fusion in cases of cervical spine injury. *Neurosurg*; 12 (6): 649-653.

Brigham C.D. and Tsahakis P.J. (1995). Anterior cervical foraminotomy and fusion. Surgical technique and results. *Spine*; 20 (7): 766-770.

Brodke D.S., Dick J.C., Kurz D.N., McCabe R.B.S., Zdeblick T.A. (1997). Posterior lumbar interbody fusion: A biomechanical comparison, including a new threaded cage. *Spine*; 22 (1): 26-31.

Brown M.D., Malinin T.I., Davis P.B. (1976). A roentgenographic evaluation of frozen allografts versus autograft in anterior cervical spine fusions. *Clin Orthop and Rela Res*; 119: 231-236.

Cahill D.W., Martin G.J., Hajjar M.V. et al (2003). Suitability of bioresorbable cages for anterior cervical fusion. *J Neuosurg (Spine 2)*; 98: 195-201.

Claes L. (1992). Mechanical characterization of biodegradable implants. *Clin Mater* 10 (1-2): 41-46.

Claes L., Schultheiss M., Wolf S., Wilke H-J., Arand M., Kinzl, L. (1999). A new radiolucent system for vertebral body replacement: It stability in comparison to other systems. *J Biomech Mater Res*; 48: 82-89.

Cornwall G.B., Ames C.P., Crawford N.R., Chamberlain R.H. (2004). In vivo evaluation of bioresorbable polylactide implants for cervical graft containment in an ovine spinal fusion model. *Neurosurg Focus*; 16 (3): E5

Crock H. (1993). Surgery of the cervical spine. A short practice of spinal surgery. Wien New York. Springer-Verlag: Kapitel 7.

Cunningham B.W., Kanayama M., Parker L.M. et al. (1999). Osteogenic protein versus autologous interbody arthrodesis in the sheep thoracic spine: A comparative endoscopic study using the Bagby and Kulich interbody fusion device. *Spine*; 24 (6): 509-518.

Cutright D.E., Hunsuck E.E. and Beasley J.D. (1971). Fracture reduction using a biodegradable material, polylactic acid. *J.Oral Surgery*; 129: 393-397.

David A., Müller D., Eitenmüller J. und Muhr G. (1998). Einjahresergebnisse der osteosynthese distaler Radiusfrakturen mit resorbierbaren Stiften. *Heft Unfallchir*; 265: 203-206.

Debrunner A. M. (2002). Orthopädie, orthopädische Chirurgie: Patientenorientierte Diagnostik und Therapie des Bewegungsapparates. 4.,vollst. neu bearb. Aufl. Bern, Göttingen, Toronto, Seattle : Hubertus: 91-96.

Di Angelo D.J., Scifert J.L., Kitchel S., Cornwall G.B., McVay B.J. (2002). Bioabsorbable anterior lumbar plate fixation in conjunction with cage-assisted anterior interbody fusion. *J Neurosurg (Spine 4)*; 97: 447-455.

Duane E. (1971). Fracture reduction using a biodegradable material, polylactic acid. *Oral Surgery*; 29: 393-397.

Ebara S., Harada T., Hosono N. et al. (1992). Intraoperativ measurement of lumbar spinal instability. *Spine*; 17 (3S): 44-50.

Eindorf T. (2005). Histologische Evaluation eines biodegradierbaren Implantats zur Spondylodese der Halswirbelsäule. (Eine tierexperimentelle Studie im Schafmodell). Inaugural-Dissertation zur Erlangung des Grades eines Doktors der Veterinärmedizin an der Freien Universität Berlin.

Eitenmüller J., David A., Pommer A. und Muhr G. (1996). Operative Behandlung von Sprunggelenksfrakturen mit biodegradierbaren Schrauben und Platten aus Poly-L-Lactid. *Chirurg*; 67: 413-418.

Gautier S.E., Oudega M., Frago M. et al. (1998). Poly (a-hydroxyacids) for application in the spinal cord: Resorbility and biocompatibility with adult Schwann cell and spinal cord. *J Biomed Mater Res*; 42: 642-654.

- Gerlach R.L., Eitenmüller J., Schmitz H.** (1987). In-vivo-Untersuchung der Festigkeitseigenschaften biologisch abbaubarer Polymere zur Anwendung als Osteosynthesematerialien. *Dtsch Z Mund Kiefer GesichtsChir*; 11: 211-216.
- Gogolewski S.** (2000). Bioresorbable polymer in trauma and bone surgery. *Injury, Int J Care Injured*; 31: S-D28-S-D32.
- Göpferich A.** (1996). Mechanisms of polymer degradation and erosion. *Biomaterials*; 17: 103-114.
- Gore D.R., Sepic S.B.** (1984). Anterior cervical fusion for degenerated or protruded discs. A review of one hundred forty-six patients. *Spine*; 9 (7): 667-671.
- Goulet J.A., Senunas L.E., DeSilva G.L., Greenfield M.L.** (1997). Autogenous iliac crest bone graft: Complications and functional assessment. *Clin Orth Relat Res*; 339: 76-81.
- Gourlay S.J., Rice R.M., Hegyeli A.F. et al.** (1978). Biocomparability testing of polymers: In vivo implantation studies. *J Biomed Mater Res*; 12: 219-232.
- Green D.L., Crawford N.R., Chamberlain R.H., Park S.C., Cransall D.** (2003). Biomechanical comparison of cervical interbody cages versus structural bone graft. *Spine*; 3 (4): 262-269.
- Hacker R.J.** (2000 b). A randomized prospective study of an anterior cervical interbody fusion device with a minimum of 2 years of follow-up results. *J Neurosurg (Spine 2)*; 93: 222-226.
- Hacker R.J., Cauthen J.C., Thomas G.J. and Griffith S.L.** (2000 a). A prospective randomized multicenter clinical evaluation of an anterior cervical fusion cage. *Spine*; 25 (20): 2646-2655.
- Heidemann W., Jeschkeit-Schubbert S., Ruffieux K. et al.** (2001). Degradation of poly (D, L)-lactide implants with and without addition of calciumphosphat in vivo. *Biomaterials*; 22: 2371-2381.
- Heidemann W., Jeschkeit-Schubbert S., Ruffieux K. et al.** (2002). PH-stabilization of predegraded PDLA by an admixture of water-soluble sodium hydrogenphosphat-results of an in vitro- and in vivo- study. *Biomaterials*; 23: 3567-3574.

- Hirsch C.** (1960). Cervical disc rupture. Diagnosis and therapy. *Acta Orthop Scan*; 30: 172-186.
- Hoffmann R., Weiler A., Helling H-J., Krettek C., Rehm K.E.** (1997). Lokale Fremdkörperreaktion auf biodegradierbare Implantate. Eine Klassifikation. *Unfallchirurg*; 100: 658-666.
- Hofman G.O.** (1992). Biodegradable implants in orthopaedic surgery-A review on the state-of-the-art. *Clin Mater*; 10: 75-80.
- Hollowell J.P., Vollmer D.G., Wilson C.R., Pintar F., Yogananda N.** (1996). Biomechanical analysis of thoracolumbal interbody constructs: How important is the Endplate? *Spine*; 21(9): 1032-36.
- Horowitz S.M, Gautsch T.L., Frondoza C.G, Riley L. jr.** (1991). Macrophage exposure to polymethylmethacrylat leads to mediator release and injury. *J Orthop Res*; 9: 406-13.
- Ignatus A.A. and Cleas L.E.** (1996). In vitro biocompatibility of bioresorbable polymers: Poly (L, DL-lactide) and poly (L-lactide-co-glycolide). *Biomaterials*; 17: 831-839.
- Jost B., Cripton P.A., Lund T. et al.** (1998). Compressive strength of interbody cage in the lumbar spine: The effect of cage shape, posterior instrumentation and bone density. *Euro Spine J*; 7: 132-141.
- Kanayama M., Cunningham B.W., Haggerty C.J, Abumi K., Kaneda K., McAfee P.C.** (2000). In vitro biomechanical investigation of the stability and stress-shielding effect of interbody fusion devices. *J Neurosurg (Spine 2)*; 93: 259-265.
- Kandziora F., Pflugmacher R., Schäfer J., Duda G., Haas N.P., Mittelmeier T.** (2001a). Biomechanical comparison of cervical interbody fusion Cages. *Spine 26 (17)*: 1850-57.
- Kandziora F., Pflugmacher R., Scholz M., et al.** (2001 b). Comparison between sheep and human cervical spine. *Spine 26 (9)*: 1028-1037.
- Kandziora F., Schollmeier G., Scholz M. et al. (2002 a):** Influences of cage design on interbody fusion in a sheep cervical spine model. *J. Neurosurg (Spine 3)*; 96: 321-332.

- Kandziora F., Schmidmaier G., Schollmeier G., et al.** (2002 b). IGF-I and TG- β I application by a Poly - (D, L-lactid)-coated cage promotes intervertebral bone matrix formation in the sheep cervical spine. *Spine*; 27 (16): 1710-1723.
- Kandziora F., Pflugmacher R., Scholz, M. et al.** (2002 c). Experimentelle Spondylodese der Schafswirbelsäule. Teil 1: Der Effekt des Cagedesigns auf die intervertebrale Fusion. *Der Chirurg*, 73: 909-917.
- Khodadadyan-Klostermann C., Kandziora F., Schnaake K.J. et al.** (2001). Mechanischer Vergleich biodegradierbarer intervertebraler lumbaler Cages. *Chirurg*; 72: 1431-1438
- Krall C.** Radiologische Evaluation eines biodegradierbaren Implantats zur Spondylodese der Halswirbelsäule. (Dissertation in Vorbereitung). Berlin Zentrum für Muskuloskeletale Chirurgie, Charite-Universitätsmedizin Berlin
- Kulich S.D., Danielson G., Dowdle J.D. et al.** (2000). Four-year-follow-up results of lumbar spine arthrodesis using the Bagby and Kulich lumbar fusion cage. *Spine*; 225 (20): 2656-2662.
- Kulkarni R.K.** (1966). Polylactid acid for surgical implants. *Arch Surg*; 93: 839.
- Kulkarni R.K., Moore E.G., Hegyeli A.F. and Leonard F.** (1971). Biodegradable Poly (lactic acid) Polymers. *J Biomes. Mater. Res*; 5: 169-181.
- Kummer B.** (1959). Bauprinzipien des Säugerskeletts. Georg Thieme-Verlag Stuttgart: 36-76.
- Kurz L.T., Garfin S.T., Booth R.E.** (1989). Harvesting autogenous iliac bone grafts. A review of complication and techniques. *Spine*; 14 (12): 1324-1331.
- Lee E-J, Hung Y-H, Lee M-Y et al.** (1999). Kinematics of cervical spine discectomy with and without bone grafting: Quantitative evaluation of late fusion in a sheep model. *Neurosurg*; 44 (1): 139-147.
- Li S.** (1998). Hydrolytic degradation characteristics of aliphatic polyester derived from lactic and glycolic acids. *J Biomed Mater Res*; 48 (3): 342-353.
- Li S. and Vert M.** (1994). Morphological changes resulting from the hydrolytic degradation of stereocopolymers derived from L and DL-lactides. *Macromolecules*; 27: 3107-3110.

- Li H., Laursen M., Egmund N., Lind M., Bungler C.** (2002). The influence of intervertebral disc tissue on anterior spinal interbody fusion: an experimental study on pigs. *Euro Spine J*; 11:476-481.
- Lippmann C.R., Hajjar M., Abshire B., Martin G., Engelman R.W., Cahill D.W.** (2004). Cervical spine fusion with biodegradable cages. *Neurosurg Focus*; 16 (3). Article 4: 1-4.
- Lowe T.G. and Coe J.D.** (2002). Resorbable polymer implants in unilateral transforaminal lumbar interbody fusion. *J Neurosurg*; (Spine 4) 97: 464-467.
- Lowery G.L., Swank L. and Mc Donough R.F.** (1995). Surgical revision for failed anterior cervical fusion. *Spine*; 20 (22): 2436-2441.
- Mainil-Varlet P., Curtis R. and Gogolewski S.** (1997 b). Effects of in vitro and in vivo degradation on molecular and mechanical properties of various polylactides. *J Biomed Mater Res*; 36:360-380.
- Mainil-Varlet P., Rahn B., Gogolewski S.** (1997 a). Long-term in vivo degradation and bone reaction to various bone polylactides. One-year results. *Biomaterials*, 18: 257.
- Majd M.E., Vadhva M., Holt R.T.** (1999). Anterior cervical reconstruction using titanium cage with anterior plating. *Spine*; 24 (15): 1604-1610.
- Majola A., Vainionpää S., Vihtonen K. et al.** (1991). Absorption, biocompatibility and fixation properties of polylactic acid in bone tissue: An experimental study in rats. *Clin Orth*; 268: 260-269.
- Majola A., Vainionpää S., Vihtonen K. et al.** (1992). Intramedullar fixation of cortical bone osteotomies with self-reinforced polylactic rods in rabbits. *Int Orthop*, 16: 101-108.
- Marchesi D.G.** (2000). Spinal fusion: Bone and bone substitutes. *Euro Spine J*; 9: 372-378.
- Martins A.N.** (1976). Anterior cervical discectomy with and without interbody bone graft. *J Neurosurg*; 44 : 290-295.
- Marzischewski S., Helling H.J. und Rehm K.E.** (1998). Klinische Langzeitergebnisse der Pilotanwendung neuer Polylactidstifte. Gibt es Biocompatibilitätsprobleme? *Heft Unfallchir.* 265: 252-257.

- Mc Afee PC, Cunnungham B.W., Lee G.A. et al.** (1999). Revision strategies for salvation or improving failed cylindrical Cages. *Spine*; 24 (20): 2147-2153.
- Mc Afee, P.C., Lee, G.A., Fedder, I.L., Cunningham, B.W.** (2002). Anterior BAK instrumentation and fusion: Complete versus incomplete discectomy. *Clin Orth*; 394: 55-63.
- Miller R.A., Brady J.M. and Cutright D.E.** (1977). Degradation rates of oral resorbable Implants (polylactates and polyglycolates): Rate modifications with changes in PLA/PGA copolymer ration. *J. Biomes. Mater. Res*; 11: 711-719.
- Minkin C. and Shapiro I.M.** (1986). Osteoclasts, mononuclear phagocytes, and physiological bone resorption. *Calcif Tissue Int*; 39 (6): 357-59.
- Murphy M. A., Trimble, M.B., Piedmonte, M.R., Kalfas, I.H.** (1994). Changes in the cervical foraminal area after anterior discectomy with and without a graft. *Neurosurg* 34 (1): 93-96.
- Murphy M.G. and Gado M.** (1972). Anterior Cervical discectomy without interbody bone graft. *J Neurosurg*; 37: 71-74.
- Murray D.W. and Rushton N.** (1990). Macrophages stimulate bone resorption when they phagocyte particles. *J Bone Joint Surg Br.*; 72 (6): 988-992.
- Niethard F. U., Pfeil J.** (1992). *MLP Duale Reihe Orthopädie. 2. Auflage.* Hippokrates Verlag GmbH Stuttgart: 328-36.
- O'Laoire S.A. and Thomas D.G.H.** (1983). Spinal cord compression due to prolaps of cervical intervertebral disc (herniation of nucleus pulposus): Treatment of 26 cases by discectomy without interbody graft. *J Neurosurg*; 59: 847-853.
- Panjabi M.M.** (1988 a). Biomechanical evaluation of spinal fixation devices: I. A conceptual framework. *Spine*; 13 (10): 1129-1134.
- Panajabi M.M., Tech D., Abumi K., Durancenau J. and Crisci J.J.** (1988 b). Biomechanical evaluation of spinal fixation devices: II. Stability provided by eight internal fixation devices. *Spine*; 13 (10): 1135-1140.
- Panjabi M.M.** (1991). Dreidimensionale Testung der Stabilität von Wirbelsäulen. *Orthopäde*; 20: 106-111.

- Peltier L. F.** (1999). Anteriore and posteriore cervical spine fusion. *Clin. Orthop*; 359: 4-17.
- Perren S.M.** (2002). Evolution of the internal fixation of long bone fractures. The scientific basis of biological internal fixation: Choosing a new balance between stability and biology. *J Bone Joint Surg Br.*; 84 (8):1093-110.
- Pflugmacher R., Schleicher P., Gumnior S. et al.** (2004). Biomechanical comparison of bioabsorbable cervical Spine Interbody Fusion Cages. *Spine*; 29 (16): 1717-1722.
- Pistner H, Gutwald, R Schwartz, G. Mühling, J. und Reuther J.** (1998). Vollständige Biodegradation von spritzgegossenem Polylactid-Osteosynthesematerial in Hart- und Weichgewebe in vivo im Langzeitversuch bis zu 36 Monaten. *Heft Unfallchir.* 265: 133-145.
- Profeta G., de Falco R., Ianniciello G., Profeta L., Cigliano A., Raja A.I.** (2000). Preliminary experience with anterior cervical microdiscectomy and interbody titanium cage fusion (Novus CT-TI) in patients with cervical disc disease. *Surg Neurol*; 53: 417-26.
- Rapoff A.J., Ghanayem A.J., Zedeblick T.A.** (1997). Biomechanical comparison of posterior lumbar interbody fusion cages. *Spine*; 22 (20): 2375-2379.
- Robertson J.T., Johnson S.D** (1980). Anterior cervical discectomy without fusion: Long-term results. *Clin Neurosurg*; 27: 440-449.
- Robinson R. A., Walker A. E., Ferlinc D. C., Wiecking D. K.** (1962). The result of anterior interbody fusion of the cervical spine. *The Journal of bone and joint surgery*; 44 (8): 1569-1586.
- Rokkanen P.U., Böstman O., Vainionpää S. et.al.** (1996). Absorbable devices in the fixation of fractures. *J Trauma*; 40 (Suppl 3): S123-127.
- Sandhu H. S., Turner S., Kabo M. et al.** (1996). Distractive properties of a threaded interbody fusion device: An in vivo model. *Spine*; 21 (10): 1201-10.
- Sandhu H.S., Kanim L.E.A., Diwan A.D. et al.** (1995). Evaluation of rh BMP 2 with an OPLA carrier in a canine posterolateral (transversal process) spinal fusion model. *Spine*; 20 (24): 2669-2682.

- Schmidmaier G., Wildemann B., Beil H. et al.** (2000). Die lokale Freisetzung von IGF-1 und TGF- β 1 aus einer biodegradierbaren Poly (D, L-Lactid) Beschichtung von Implantaten beschleunigt die Frakturheilung. *Chirurg*; 71: 1016-1022.
- Schnee C.L., Frees A., Weil R.J., Marcotte P.J.** (1997). Analysis of harvest morbidity and radiographic outcome using autograft for anterior cervical fusion. *Spine*; 22 (19): 2222-2227.
- Schulte B.S., Clark C.R., Goel V.K.** (1989). Kinematics of the cervical spine following discectomy and stabilization. *Spine*; 14 (10): 1116-1121.
- Shono Y., McAfee P.C., Cunningham B.S., Brantigan J.W.** (1993). A biomechanical analysis of decompression and reconstruction methods in the cervical spine. *J Bone Joint Surg Am.*; 75 (11): 1674-1684.
- Silber J.S., Anderson D.G., Daffner S.D. et al.** (2002). Donor site morbidity after iliac crest bone harvest for single-level anterior cervical discectomy and fusion. *Spine*, 28 (2): 134-139.
- Smit T.** (2002). The use of quadruped as an in vivo model for the study of the spine-biomechanical considerations. *Eur Spine J*; 11: 137-44.
- Södergard A., Stolt M.** (2002). Properties of lactic acid based polymers and their correlation with composition. *Prog Polym. Sci*; 27: 1123-1163.
- Stählin A.C., Weiler A., Rüfenach H., Hoffmann R., Geissmann A. and Feinstein R.** (1997). Clinical degradation and biocompatibility of different bioabsorbable interference screws: A report of six cases. *J Arthroscopic Relat Surg*; 13 (2): 238-244.
- Statistisches Bundesamt Deutschland** (2005 a).
<http://www.destatis.de/basis/d/bevoe/bevoetab5.php>
- Statistisches Bundesamt Deutschland** (2005 b).
http://www.destatis.de/themen/d/thm_gesundheit.php
- Suuronen R., Laine P., Pohjonen T. and Linqvist C.** (1994). Sagittal Ramus Osteotomie fixed with biodegradable screws: A preliminary report. *J Oral Maxillofac Surg*; 52: 715-20.
- Suuronen R., Pohjonen T., Hietanen J. et a.** (1998). A 5-Year in vitro and in vivo study of the biodegradation of polylactid plate. *J oral Maxillofac Surg*; 56: 604-14.

- Taylor M.S., Daniel U.A., Andriano K.P. and Heller J.** (1994). Six bioabsorbable polymers: In vitro acute toxicity of accumulated degradation products. *J Appl Biomater*, 5: 151-157.
- Togawa D, Bauer T, Liebermann I. and Sakai H.** (2004). Lumbar intervertebral body fusion cage: Histological evaluation of clinically failed cages retrieved from Humans. *J Bone Joint Surg Am.*; 86 (1): 70-79.
- Toth J.M., Bradley T.E., Wang M. et al.** (2002). Evaluation of 70/30 poly (L-lactid-co-D-L-lactide) for use as a resorbable interbody fusion cage. *J Neurosurg; (Spine 4)* 97: 423-432.
- Tsantrizos A., Andreou A., Aebi A., Steffen T.** (2000). Biomechanical stability of five stand-alone anterior lumbar interbody fusions constructs. *Euro Spine J*; 9: 14-22.
- Türeyen K.** (2003). Disc height loss after anterior cervical microdiscectomy with titanium intervertebral cage fusion. *Acta Neurochir*; 145: 565-570.
- Vaccaro A.R., Dingh K., Haid R. et al.** (2003). The use of biodegradable implants in the spine. *Spine J*; 3: 227-237.
- Van Dijk M., Smit TH., Burger E.H., Wuisman P.I.** (2002 a). Bioabsorbable poly-L-lactic acid cages for lumbar interbody fusion: Three-year follow-up radiographic, histologic, and histomorphometric analysis in goats. *Spine*, 27 (23): 2706-2714.
- Van Dijk M., Smit TH., Sugihara S., Burger E.H., Wuisman P.I.** (2002 b). The effect of cage stiffness on the rate of lumbar interbody fusion. *Spine* 27 (7): 682-688.
- Van Dijk M., Tunc D.C., Smit T.H., Higham P., Burger E.H., Wuisman P.I.** (2002 c). In vitro and in vivo Degradation of biodegradable Pdllla spine fusion cages. *J Biomed Mater Res*, 63: 752-759.
- Van Dijk M., Smit T.H., Arnoe M.F., Burger E.H., Wuisman P.I.** (2003). The use of poly-L-lactic acid in lumbar interbody cages: Design and biomechanical evaluation in vitro. *Eur Spine J*; 12: 34-40.
- Weiler A., Helling H.J., Kirch U., Ziberes T.K., Rehm K.E.** (1996). Foreignbody reaction and the course of osteolysis after polyglycolid implants for fracture fixation: Experimental study in sheep. *J Bone Joint Surgery Br.*; 78 (3): 369-76.

- Weiler A., Hoffmann R. F.G., Stähelin A.C., Helling H-J., Südkamp N.P.** (2002). Biodegradable implants in sports medicine: The biomechanical base. *J Arthrosc Rel Surgery*; 16 (3): 305-321.
- White A.A., III, Southwick W.O., De Ponte R.J., Gainor J.W. and Hardy R.** (1973). Relife of pain by anterior cervical-spine fusion for spondylosis. A report of sixty-five patients. *J Bone Joint Surg. Am.*; 55: 425-534.
- Whitecloud T.S., Sherk H.** (1999). Modern alternatives and techniques for one-level discectomy and fusion. *Clin Orthop*; 359: 67-76.
- Wilke H-J., Keller A., Wenger K.H. and Claes L.** (1997). Anatomy of the sheep spines and its comparison to the human spine. *The Anatom Rec*; 247: 342-555.
- Wilke H-J., Keller A., Claes L.** (1997). Are sheep spines a valid biomechanical model for human spine? *Spine* 22 (20): 2365-2374.
- Wilke H-J., Wenger K., Claes L.** (1998). Testing criteria for spinal implants: Recommendations for the standardization of in vitro stability testing of spinal implants. *Eur Spine J*; 7: 148-154.
- Wilson D.H., Campbell D.D.** (1977). Anterior cervical discectomy without bone graft. *J Neurosurg*; 47:551-555.
- Wissing H., Stümer K.M., Breidenstein G.** (1990). Die Wertigkeit verschiedener Versuchstierspezies für experimentelle Untersuchungen am Knochen. *Heft Unfallheilk*; 212: 479-488.
- Wuismann P.I., van Dijk M., Smith T.H.** (2002). Resorbable cages for spinal fusion: An experimental goat model. *J Neurosurg. Spine*; 4 (97): 433-439.
- Yamamoto I., Ikeda A., Shibuya N., Tsugane R., Osamu S.** (1991). Clinical long-term results of anterior discectomy without interbody fusion for cervical disc disease. *Spine*; 16 (3): 272-279.
- Younger E.M, Chapman M.W.**(1989). Morbidity at bone graft donor sites. *J Orthop Trauma*; 3: 192-195.

Zdeblick T., Coole M.E., Wilson D., Kunz D.N. and Mc Cabe R. (1993). Anterior cervical discectomy, fusion and plating: A comparative animal study. *Spine*; 18 (14): 1974-1983.

Zdeblick T.A., Ducker T.B. (1991). The use of freeze-dried allograft bone for anterior cervical fusion. *Spine*; 16 (7): 726-9

Zdeblick T., Hughes S.S., Riew M.D. and Bohlman H. (1997). Failed anterior discectomy and arthrodesis. *J Bone Joint Surge Am.* 79 (4):523-32.