

## Literaturverzeichnis

- 
- <sup>1</sup> Weinstock MA, Horn JW: Mycosis fungoïdes in the USA: Increasing incidence and descriptive epidemiology. JAMA 260:42, 1988
- <sup>2</sup> Willemze R, Kerl H, Sterry W, et al.: EORTC Classification for primary cutaneous lymphomas. Blood, Vol 90, No 1, 1997: pp 354-371
- <sup>3</sup> Willemze R, Meijer, C. J. L. M: WHO-EORTC classification for cutaneous lymphomas. Blood, 15 May 2005, Vol. 105, No. 10, pp. 3768-3785
- <sup>4</sup> J.L.M. Alibert: Descriptions des maladies de peau observées à l'Hôpital St. Louis, et exposition des meilleurs méthodes suivies pour leur traitement, p. 157 Tableau du plan fongoïde. Barrois l'Aine et Fils, Paris 1806
- <sup>5</sup> P.A.E. Bazin: Leçons sur le traitement des maladies chroniques en général, affections de la peau en particulier, p 425, Adrein Delahaye, Paris 1870
- <sup>6</sup> Nickoloff BJ: Light-microscopic assessment of 100 patients with patch/plaque stage mycosis fungoïdes. Am J Dermatopathol. 10:469, 1988
- <sup>7</sup> Gellrich S, Lukowsky A, Schilling T, Rutz S, Muche JM, Jahn S et al.: Microanatomical compartments of clonal and reactive T cells in mycosis fungoïdes: molecular demonstration by single cell polymerase chain reaction of T cell receptor gene rearrangements. Journal Invest Dermatol. Vol. 115, No. 4, October 2000, pp. 620-624
- <sup>8</sup> Muche JM Lukowsky A, Asadullah K, Gellrich S, Sterry W: Demonstration of frequent occurrence of clonal T-cells in the peripheral blood of patients with primary cutaneous T-cell lymphoma. Blood, Vol 90, No 4 (August 15), 1997: pp 1636-1642
- <sup>9</sup> Sylke Gellrich, Anke Wilks et. al.: T-Cell-Receptor- $\gamma$  Gene Analysis of CD30<sup>+</sup> Large Atypical Cells in CD30<sup>+</sup> Large Primary CTCL. Soc. I. Dermatology, Vol. 120, April 2003
- <sup>10</sup> Washington et al.: A stable aberrant immunophenotype characterizes nearly all cases of cutaneous T-cell lymphoma. BMC Clin Pathol 2002 Dec 10;2(1):5
- <sup>11</sup> Ginaldi et al.: Differential expression of CD3 and CD7 in T-cell malignancies: a quantitative study by flow cytometry. Br J Haematol 1996 Jun;93(4):921-7
- <sup>12</sup> Kukel et al.: Progressive increase of CD7- T cells in human blood lymphocytes with ageing. Clin Exp Immunol 1994 Oct;98(1):163-8
- <sup>13</sup> Wojnowska et al.: Expression of selected T-cell antigens in the skin infiltration in primary cutaneous T-cell lymphomas. Ann Univ Mariae Curie Skłodowska [Med] 2001;56:237-42
- <sup>14</sup> T. Kakinuma et al.: Thymus and activation-regulated chemokine (TARC/CCL17) in MF: Serum TARC reflects the disease activity of MF. J Am Acad Dermatol. 2003 Jan;48(1):23-30
- <sup>15</sup> Querfeld C, Rosen ST, Guitart J, Kuzel T: The spectrum of cutaneous T-cell lymphomas: new insights into biology and therapy. Current Opinion in Hematology 2005, 12:273-278
- <sup>16</sup> Poszepczynska-Guigne E, Schiavon V, D'Incan M et al.: CD158k/KIR3DL2 is a new phenotypic marker of Sézary cells: relevance for the diagnosis and follow-up of Sézary syndrome. J Invest Dermatol 2004; 122: 820-823
- <sup>17</sup> Schöpp S: Molekularbiologische Identifizierung und immunphänotypische Charakterisierung der malignen Zellen bei Sézary Syndrom unter Berücksichtigung von Klinik und Prognose. Dissertation, Klinik für Dermatologie der Charité, Berlin 2004
- <sup>18</sup> P. Sugerman, M. Bigby: Preliminary functional analysis of human epidermal T cells. Arch Dermatology Res (2000) 292: 9-15

---

<sup>19</sup> Tu et al.: The Cutaneous Lymphocyte Antigen is an essential Component of the L-selectin Ligand Induced on Human Vascular Endothelial Cells. *J. Exp. Med.*, Volume 189, Number 2, January 18, 1999 241-252

<sup>20</sup> Armerding D, Kupper TS: Functional CLA can be induced in essentially all peripheral blood T lymphocytes. *Int Arch Allergy Immunol* 1999 Jul; 119(3): 212-22

<sup>21</sup> Nakayama F, Teraki Y, Kudo T, Togayachi A, Iwasaki H, Tamatani T et al.: Expression of CLA regulated by a set of Glucosyltransferases in human T-cells: Involvement of  $\alpha$ 1,3-Fucosyltransferase VII and  $\beta$ 1,4-Galactosyltransferase I. *Journal Invest Dermatol*, 115: 299-306, 2000

<sup>22</sup> Borowitz MJ, Weidner A, Olsen EA, Picker LJ: Abnormalities of circulating T-cell subpopulations in patients with CTCL: CLA-expression on T-cells correlates with extent of disease. *Leukemia*, Vol. 7, No. 6 (June), 1993: pp 859-863

<sup>23</sup> A. Sarris et al.: Cytokine loops involving IFN- $\gamma$  and IP-10, a cytokine chemotactic for CD-4 $^{+}$  Lymphocytes: an explanation for the epidermotropism of CTCL? *Blood*, Vol 86, No 2 (July) 1995: pp 651 – 658

<sup>24</sup> A. Janeway, P. Travers: *Lehrbuch der Immunologie*, Spektrum Verlag 1997

<sup>25</sup> J:E: Gootenberg et al.: Human Cutaneous T-cell Lymphoma and Leukemia cell lines produce and respond to T-cell growth factor. *J Exp Med* 154: 1403-1418, 1982

<sup>26</sup> Preesman et al.: Cell-bound IgE and increased expression of Fc epsilon-receptors on dendritic cells in cutaneous infiltrates of mycosis fungoides. *Clin Exp Immunol* 1991 Nov; 86(2): 246-251

<sup>27</sup> Eklund JW, Kuzel TM: Denileukin diftitox: a concise clinical review. *Expert Review Anticancer Therapy*, 5(1), 2005, pp. 33-38

<sup>28</sup> E. Papadavid et al.: The relevance of peripheral blood T-helper 1 and 2 cytokine pattern in the evaluation of patients with MF and SS. *Br J Dermatol* 2003 Apr;148(4):709-18

<sup>29</sup> R. Dummer et al.: Peripheral blood mononuclear cells in patients with nonleukemic cutaneous T-cell lymphoma: reduced proliferation and preferential secretion of a a T helper-2-like cytokine pattern on stimulation. *Arch Dermatology* 129: 433–436, 1992

<sup>30</sup> Van Dongen JJM, Comans-Bitter WM, Wolvers-Tettero ILM, Borst J : Developement of human T lymphocytes and their thymus-dependency. *Thymus* 26: 235-254, 1990

<sup>31</sup> Van Dongen JJM, Wolvers-Tettero ILM, Seidman JG, Ang SL, Van de Griend RJ, De Vries EFR, Borst J: Two types of gamma T cell receptors expressed by T cell acute lymphoblastic leukemias. *Eur J Immunol* 1987; 17: 1719-1728

<sup>32</sup> Arden B, Clark SP et al.: Human T-cell receptor variable gene segment families. *Immunogenetics* 1995; 42: 455-500

<sup>33</sup> Toyonaga B et al.: Organization and sepuences of the diversity, joining and constant region genes of the human T-cell-receptor beta chain. *Proc Natl Acad Sci USA* 1985; 82: 8624-8628

<sup>34</sup> Blom B et al.: TCR gene arrangements and expression of the pre-T-cell receptor complex during human T-cell differentiation. *Blood* 1999; 93: 3033-3043

<sup>35</sup> Alaibac M et al.: Molecular Analysis of the gamma T-cell receptor repertoire in normal human skin and in oriental cutaneous leishmaniasis. *Exp Dermatol* 1993; 2: 106-112

<sup>36</sup> Chen Z et al.: The human t-cell V gamma gene locus: cloning of new segments and study of V gamma rearrangements in neoplastic T and B cells. *Blood* 1988; 72: 776-783

<sup>37</sup> van Dongen et al.: Immunobiology of leukemia. In: Henderson ES Lister TA, Greaves MF (eds) *Leukemia*. Philadelphia: WB Saunders Company, 2002, pp 85-129

<sup>38</sup> Modifiziert nach: T.M. Breit, I.L.M. Wolverst-Tetero, J.J.M. van Dongen: Receptor diversity of human T-cell receptor  $\gamma\delta$  expressing cells. Aus: *Progress in Histo- and Cytochemistry*, Vol. 26, 1992

---

<sup>39</sup> Modifiziert nach: Report of the BIOMED-2 Concerted Action BMH4-CT98-3936 Aus: Leukemia 17 (2003), 2257-2317

<sup>40</sup> Juarez T, Isenhardt SN, Polissar NL, Sabath DE, Wood B et al.: Analysis of T-cell receptor gene rearrangement for predicting clinical outcome in patients with cutaneous T-cell lymphoma. Arch Dermatol (2005) 141, 1107-1113

<sup>41</sup> Thangavelu M, Finn WG, Yelavarthi KK et al.: Recurring structural chromosomal abnormalities in peripheral blood lymphocytes of patients with mycosis fungoides/Sézary syndrome. Blood 89: 3371-3377, 1997

<sup>42</sup> Espinet B, Salido M, Pujol RM et al.: Genetic characterization of Sézary's syndrome by conventional cytogenetics and cross-species color banding fluorescent in situ hybridization. Haematologica 2004;89: 165-173

<sup>43</sup> Wain EM, Mitchell TJ, Russel-Jones R, Whittaker S: Fine Mapping of chromosome 10q deletions in mycosis fungoides and Sézary syndrome: identification of two discrete regions of deletion at 10q23.33-24.1 and 10q24.33-25.1. Genes chromosomes Cancer 2005; 42: 184-192

<sup>44</sup> Scarisbrick JJ, Woolford AJ, Russel-Jones R, Whittaker SJ: Allelotyping in mycosis fungoides and Sézary syndrome: common regions of allelic loss identified on 9p, 10q and 17p. J Invest Dermatol 117: 663-670, 2001

<sup>45</sup> Gallardo F, Esteller M, Pujol RM et al.: Methylation status of the p15, p16 and MGMT promoter genes in primary T-cell lymphomas. Haematologica 2004; 89: 1401-1403

<sup>46</sup> Sommer VH; Clemmensen OJ, Nielsen O et al.: In vivo activation of STAT3 in cutaneous T-cell lymphoma: evidence for an antiapoptotic function of STAT3 Leukemia 2004; 18: 1288-1295

<sup>47</sup> Dummer R, Willers U, Döbbeling U, Burg G: Aktuelle Aspekte zur Pathogenese von Sézary-Syndrom und Mycosis fungoides. Hautarzt 2001, 52:189-192

<sup>48</sup> Samoszuk MK, Sholly S, Gupta S, Lukes RJ: Enzyme immunoassay for detection of monoclonal immunoglobulin in lymph nodes. A comparison with histologic and DNA gene arrangement studies. Cancer, 1987 Dec 1; 60(11): 2721-2726

<sup>49</sup> Posnett DN, Sinha R, Kabak S, Russo C: Clonal populations of T cells in normal elderly humans: the T cell equivalent to „benign clonal gammopathy“. J Exp Med 1994 Mar 1; 179(3): 1077

<sup>50</sup> J.J. Lafaille et al.: Junctional sequences of T-cell receptor  $\gamma$ : $\delta$  genes: Implications for  $\gamma$ : $\delta$  T-cell lineages and for a novel intermediate of V-(D)-J joining. Cell 59; 859-870, 1989

<sup>51</sup> J.M. Weinberg et al.: Molecular Diagnosis of Lymphocytic Infiltrates of the Skin. Archives in Dermatology 129: 1491-1500, 1993

<sup>52</sup> Guitart J, Camisa C, Ehrlich M, Bergfeld WF: Long-term implications of T-cell receptor gene rearrangement analysis by Southern blot in patients with cutaneous T-cell lymphoma. J Am Acad Dermatol 2003; 48:775-779

<sup>53</sup> Volkenandt M, Soyer HP, Cerroni L, Koch OM, Atzpodien J, Kerl H: Molecular detection of clone-specific DNA in hypopigmented lesions of a patient with early evolving mycosis fungoides. British Journal of Dermatology (1993) 128, 423-428

<sup>54</sup> Chan DW, Liang R, Kwong YL, Chan V: Detection of T-cell receptor delta gene rearrangement in T-cell malignancies by clonal specific PCR and its application to detect minimal residual disease. American Journal of Hematology 52: 171-177 (1996)

<sup>55</sup> Breit TM et al.: Limited combinatorial repertoire of  $\gamma$ : $\delta$ -T-cell-receptors expressed by T-cell acute lymphoblastic leukemias. Leukemia 1991; 5: 116-124

<sup>56</sup> Thomas Schild: Die PCR als Grundlage in der molekularen DNA-Analyse. Aus: Automatische genetische Analytik. Wiley-VCH, 1997

<sup>57</sup> Dummer R, Haeffner AC, Burg G: Cutaneous T-cell lymphomas (CTCL): new aspects in PCR-based diagnostic tools, immunology and treatment. Eur J Dermatol 1994; 4: 281-286

---

<sup>58</sup> Meyer JC, Hassam S, Dummer R, Muletta S, Dobbeling U, Dommann SN, Burg G: A realistic approach to the sensitivity of PCR-DGGE and its application as a sensitive tool for the detection of clonality in cutaneous T-cell proliferations. *Exp Dermatol* 1997 Jun; 6(3): 122-127

<sup>59</sup> Lukowsky A, Audring H, Heiduk U, Muche JM, Richter S, Sterry W: Nachweis von monoklonalen T-Zellen mittels TCRγ-PCR bei Mycosis Fungoidea (MF). *Hautarzt* 1998; 49:641-645

<sup>60</sup> Lukowsky A, Richter S, Dijkstal K, Sterry W, Muche M: A- T-cell receptor γ polymerase chain reaction assay using capillary electrophoresis for the diagnosis of cutaneous T-cell lymphomas. *Diagn. Mol. Pathol.* 11, 59-66

<sup>61</sup> Lessin SR, Rook AH, Rovera G: Molecular diagnosis of cutaneous T-cell lymphoma: polymerase chain reaction amplification of T-cell receptor β-chain rearrangements. *J Invest Dermatol* 96: 299-302, 1991

<sup>62</sup> Dagmar Schuster: Die DNA-Präparation für die automatische DNA-Analyse. Aus: Automatische genetische Analytik. Wiley-VCH, 1997

<sup>63</sup> M. Kneba et al.: Characterization of clone-specific Rearranged T-Cell Receptor γ-Chain Genes in Lymphomas and Leukemias by PCR and DNA Sequencing. *Blood*, Vol 84, No 2(July 15), 1994: pp 574-581

<sup>64</sup> Beylot-Berry M, Sibaud V, Thiebaut R et al.: Evidence, that an identical T cell clone in skin and peripheral blood lymphocytes is an independent prognostical marker in primary cutaneous T cell lymphoma. *J Invest Dermatol* 2001; 117: 920-926

<sup>65</sup> Fraser-Andrews EA, Woolford AJ, Russel-Jones R, Seed PT, Whittaker SJ: Detection of a peripheral blood T-cell clone is an independent prognostic marker in mycosis fungoidea. *J Invest Dermatol* 2000; 114:114-117

<sup>66</sup> Ahnhudt C, Muche M, Dijkstal K, Sterry W et al.: An approach to the sensitivity of temperature-gradient gel electrophoresis in the detection of clonally expanded T-cells in CTCL. *Electrophoresis* 2001, 22, 33-38

<sup>67</sup> Schäfer T: DNA-Sequenzanalyse. Automatische genetische Analytik. Wiley-VCH, 1997

<sup>68</sup> Westermeier R: Gelelektrophorese, in Analytiker-Taschenbuch. Band 7, Springer-Verlag Berlin, Heidelberg 295-344, 1988

<sup>69</sup> Orita M, Iwahana H, Kanaqawa H, Hayashi K, Sekiya T: Detection of polymorphisms of human DNA by gel electrophoresis as single-strand conformation polymorphism. *Proc. Natl. Acad. Sci. USA* 86 (1989), 2766-2770

<sup>70</sup> Nagamine CM et al.: A PCR artifact: Generation of heteroduplexes. *Am J hum Genet*, 45 (1989), 337-339

<sup>71</sup> Witzens M, Möhler T, Willhauck M, Scheibenbogen C, Lee KH, Keilholz U: Detection of clonally rearranged T-cell-receptor gamma chain genes from T-cell malignancies and acute inflammatory disease using PCR amplification, PAGE and automated analysis. *Ann Hematol* (1997) 74: 123-130

<sup>72</sup> Wood GS, Tung RM, Haefner AC, Crooks CF, Liao S, Veelken H et al.: Detection of clonal T-cell-receptor gamma gene rearrangements in early mycosis fungoidea, Sézary Syndrome by polymerase chain reaction and denaturing gradient gel electrophoresis. *J Invest Dermatol* 103: 34

<sup>73</sup> Scheller U, Muche JM, Sterry W, Lukowsky A: Detection of clonal cells in cutaneous T cell Lymphoma by PCR: Comparison of mutation detection enhancement-polyacrylamide gel electrophoresis, temperature gradient gel electrophoresis and fragment analysis of sequencing gels. *Electrophoresis* 1998, 19, 653-658

<sup>74</sup> de Carmejane O, Yamaguchi Y, Todorov TI, Morris MD: Three-dimensional observation of electrophoretic migration of dsDNA in semidilute hydroxyethylcellulose solution. *Electrophoresis*. 2001 Aug;22(12):2433-41.

<sup>75</sup> Chan DW, Liang R, Kwong YL, Chan V: Detection of T-cell Receptor Delta Gene Rearrangement in T-cell malignancies by Clonal Specific PCR and Its Application to Detect Minimal Residual Disease. *Am J Hema* 52: 171-177 (1996)

---

<sup>76</sup> Gutzmer R, Mommert S, Kiehl P, Wittmann M, Kapp A, Werfel T: Detection of Clonal T Cell Receptor  $\gamma$  Gene Rearrangements in Cutaneous T Cell Lymphoma by LightCycler-Polymerase Chain Reaction. *J Invest Dermatol* Vol. 116, No. 6 June 2001

<sup>77</sup> Wickham CL, Lynas C, Ellard S: Detection of Clonal T cell populations by high resolution PCR using fluorescently labelled nucleotides; evaluation using conventional LOS-SSCP. *J Clin Pathol; Mol Pathol* 2000; 53: 150-154

<sup>78</sup> Ayling JF, Iland HJ: High-resolution analysis of gene rearrangements in lymphoid malignancies. *Pathology* 1999 31, pp. 252-256

<sup>79</sup> Sprouse JT, Werling R, Hanke D, Lakey C, McDonnel L, Wood BL et al.: T-cell clonality determination using PCR amplification of the T-cell receptor gamma-chain gene and capillary electrophoresis of fluorescent labelled PCR products. *Am J Clin Pathol* 2000; 113:838-850

<sup>80</sup> Lynas C, Howe D: Simple, reliable detection of t cell clones by PCR-LIS-SSCP analysis of TCR $\gamma$  rearrangement. *Mol and Cell Probes* 1998, 12, 41-48

<sup>81</sup> Lukowsky A: Clonality Analysis by T-cell receptor  $\gamma$ -PCR and high-resolution electrophoresis in the diagnosis of cutaneous T-cell lymphoma. Aus: Methods in molecular biology, vol. 218: Cancer cell signaling: Methods and protocols. Edited by: Terrian DM, © Humana Press Inc. Totowa, NJ

<sup>82</sup> Dippel E, Assaf C, Hummel M, Schrag HJ, Stein H, Goerdt S, Orfanos CE: Clonal T-cell receptor gamma-chain gene rearrangement by PCR-based GeneScan analysis in advanced cutaneous T-cell lymphoma: a critical evaluation. *J Pathol.* 1999 Jun;188(2):146-54.

<sup>83</sup> Simon M, Kind P, Kaudewitz P, Krokowski M, Graf A, Prinz J et al.: Automated high resolution polymerase chain reaction fragment analysis. *Am J Pathol* 1998, 152: 29-33

<sup>84</sup> Beaubier NT, Hart AP, Bartolo C, Willmann CL, Viswanatha DS: Comparison of capillary electrophoresis and polyacrylamid gel electrophoresis for the evaluation of T and B cell clonality by polymerase chain reaction. *Diagn Mol Pathol* 9 (3): 121-131, 2000

<sup>85</sup> Mansfield ES, Kronick MN: Alternative labeling techniques for automated fluorescence based analysis of PCR products. *Bio Techniques* 15 (1993), 20-22

<sup>86</sup> McCollum C, Chakerian V, Kaufman J, Wenz M, Andrus A. *Biomed. Peptides Proteins & Nucleic Acids* 1 (1994) 25-30

<sup>87</sup> Schmidt G: Die DNA-Synthese als grundlegendes Werkzeug in der molekularen DNA-Analyse. Aus: Automatische genetische Analytik. Wiley-VCH, 1997

<sup>88</sup> Vogelstein B, Gillespie D. (1979) *Proc. Natl. Acad. Sci. USA* 76, 615-619

<sup>89</sup> Chen, Loiseau P, Bories JC, Bensussan A, Sigaux F : Rearranging sequence located in the intron of the human T-cell-receptor  $\gamma$  chain gene constant region. *European Journal of Immunology* 19: 637-642, 1989

<sup>90</sup> Delfau-Larue MH, Petrella T, Lahet C et al.: Value of clonality studies of cutaneous T lymphocytes in the diagnosis and follow-up of patients with mycosis fungoides. *J Pathol* 1998; 184: 185-190

<sup>91</sup> Volkenandt M, Soyer HP, Kerl H, Bertino JR: Developement of a highly specific and sensitive molecular probe for detection of cutaneous lymphoma. *J Invest Dermatol* 97: 137-140, 1991

<sup>92</sup> Theodorou I, Bigorgne C, Delfau MH, Lahet C, Cochet G, Vidaud M, Raphael M, Gaulard P, Farset JP: VJ rearrangements of the TCR $\gamma$  locus in peripheral T-cell lymphomas: Analysis by polymerase chain reacton and denaturing gradient gel electrophoresis. *J Pathol* 178:303, 1996

<sup>93</sup> Födinger M, Buchmayer H, Schwarzinger I, Simonitsch I, Winkler K: Multiplex PCR for rapid detection of t-cell receptor-gamma chain gene rearrangements in patients with lymphoproliferative disease. *Br J Haem*, 1996, 94, 136-139

<sup>94</sup> Wood GS: The benign and malignant cutaneous lymphoproliferative disorders including mycosis fungoides. Aus: *Neoplastic Hematology*. Baltimore, MD, Williams and Wilkins p 917, 1992

- 
- <sup>95</sup> Weiss LM, Wood GS, Trella M: Clonal T-cell populations in lymphomatoid papulosis: Evidence of a lymphoproliferative origin for a clinically benign disease. *New England Journal of Medicine* 315: 475-479, 1986
- <sup>96</sup> Lee SC, Berg KD, Racke FK, Griffin CA, Eshleman JR: Pseudo-spikes are common in histologically benign lymphoid tissues. *Journal of Molecular Diagnostics*. Vol 2, No. 3, August 2000, pp.145-149
- <sup>97</sup> Shie S, Mikkola DL, Wood GS: Differentiation and clonality of lesional lymphocytes in pityriasis lichenoides chronica. *Arch Dermatol* 2001; 137: 305-308
- <sup>98</sup> Smith P: Histologic criteria for early diagnosis of CTCL. *Dermatologic clinics*, Vol. 12/2: pp. 315-322, 1994
- <sup>99</sup> Shaikh ZI, Rahman SB: Clinicopathological spectrum of mycosis fungoides type cutaneous T-cell lymphoma. *J Coll Physicians Surg Pak*. 2006 Mar;16(3):171-4
- <sup>100</sup> Olerud EO et al.: Cutaneous T-cell lymphoma. Evaluation of pretreatment skin biopsy specimens by panel of pathologists. *Arch Dermatol* 128: 501, 1998
- <sup>101</sup> Marschalko M: Diagnosis of cutaneous lymphomas. *Orv Hetil*. 2001 Dec 30;142(52):2903-6
- <sup>102</sup> Guitart J, Kennedy J, Ronan S, Chmiel JS, Hsiegh YC, Variakojis D: Histologic criteria for the diagnosis of mycosis fungoides: proposal for a grading system to standardize pathology reporting. *J Cutan Pathol*. 2001 Apr;28(4):174-83
- <sup>103</sup> Bushmarina MS, Bogdanova MS, Selivanova GV, Rodionov AN, Raznatovskii IM: Cytophotometric and electron microscopic studies of chronic human skin lymphomas. II. Research on lymph node involvement. *Tsitolgiia*. 1987 Apr;29(4):421-5
- <sup>104</sup> Imayama S: Differential cell surface distribution of adhesion molecules demonstrated by immuno-scanning electron microscopy. *J Dermatol*. 1994 Nov;21(11):855-9
- <sup>105</sup> Murphy M, Fullen D, Carlson JA: Low CD7 expression in benign and malignant cutaneous lymphocytic infiltrates: experience with an antibody reactive with paraffin-embedded tissue. *Am J Dermatopathol*. 2002 Feb;24(1):6-16
- <sup>106</sup> Ortonne N, Buyukbabani N, Delfau-Larue MH, Bagot M, Wechsler J: Value of the CD8-CD3 ratio for the diagnosis of mycosis fungoides. *Mod Pathol*. 2003 Sep;16(9):857-62
- <sup>107</sup> Florell SR, Cessna M, Lundell RB, Boucher KM, Bowen GM, Harris RM, Petersen MJ, Zone JJ, Tripp S, Perkins SL: Usefulness (or lack thereof) of immunophenotyping in atypical cutaneous T-cell infiltrates. *Am J Clin Pathol*. 2006 May;125(5):727-36
- <sup>108</sup> Talpur R, Jones DM, Alencar AJ, Apisarnthanarak N, Herne KL, Yang Y, Duvic M: CD25 expression is correlated with histological grade and response to denileukin diftitox in cutaneous T-cell lymphoma. *J Invest Dermatol*. 2006 Mar;126(3):575-83
- <sup>109</sup> Jones D, Ibrahim S, Patel K, Luthra R, Duvic M, Medeiros LJ: Degree of CD25 expression in T-cell lymphoma is dependent on tissue site: implications for targeted therapy. *Clin Cancer Res*. 2004 Aug 15;10(16):5587-94
- <sup>110</sup> Yao JL, Cangiarella JF, Cohen JM, Chhieng DC: Fine-needle aspiration biopsy of peripheral T-cell lymphomas. A cytologic and immunophenotypic study of 33 cases. *Cancer*. 2001 Apr 25;93(2):151-9
- <sup>111</sup> Sigurdsson V, Toonstra J, Bihari IC, Bruijnzeel-Koomen CA, van Vloten WA, Thepen T: Interleukin 4 and interferon-gamma expression of the dermal infiltrate in patients with erythroderma and mycosis fungoides. An immuno-histochemical study. *J Cutan Pathol*. 2000 Oct;27(9):429-35
- <sup>112</sup> Shohat M, Hodak E, Sredni B, Shohat B, Sredni D, David M: Cytokine profile of patients with mycosis fungoides and the immunomodulatory effect of AS101. *Acta Derm Venereol*. 2001 Aug-Sep;81(4):255-7
- <sup>113</sup> Leroy S, Dubois S, Tenaud I, Chebassier N, Godard A, Jacques Y, Dreno B: Interleukin-15 expression in cutaneous T-cell lymphoma (mycosis fungoides and Sezary syndrome). *Br J Dermatol*. 2001 May;144(5):1016-23

---

<sup>114</sup> Sugaya M, Nakamura K, Tamaki K: Serum interleukin-15 levels are not elevated in patients with stage I and II mycosis fungoides. *Acta Derm Venereol.* 2000 Nov-Dec;80(6):455

<sup>115</sup> Jones D, O'Hara C, Kraus MD, Perez-Atayde AR, Shahsafaei A, Wu L et al.: Expression pattern of T-cell-associated chemokine receptors and their chemokines correlates with specific subtypes of T-cell non-Hodgkin lymphoma. *Blood.* 2000 Jul 15;96(2):685-90

<sup>116</sup> Poenitz N, Simon-Ackermann J, Gratchev A, Qadoumi M, Klemke CD, Stadler R et al.: Overexpression of c-myb in leukaemic and non-leukaemic variants of cutaneous T-cell lymphoma. *Dermatology.* 2005;211(2):84-92

<sup>117</sup> Smoller BR, Santucci M, Wood GS, Whittaker SJ: Histopathology and genetics of cutaneous T-cell lymphoma. *Hematol Oncol Clin North Am.* 2003 Dec;17(6):1277-311

<sup>118</sup> Munson D, Bunn PA, Schechter GP: Fc receptor-defined T-lymphocyte subpopulations in patients with cutaneous T-cell lymphoma. *J Invest Dermatol.* 1982 Nov;79(5):317-21

<sup>119</sup> Eros N, Karolyi Z, Matolcsy A: Complex histologic, immunophenotypic and molecular genetic investigation in cutaneous T-cell lymphoproliferative diseases. *Orv Hetil.* 2004 Jan 11;145(2):75-80

<sup>120</sup> Smith BD, Wilson LD: Management of mycosis fungoides. Part 1. Diagnosis, staging, and prognosis. *Oncology (Williston Park).* 2003 Sep;17(9):1281-8

<sup>121</sup> Tomasini D, Zampatti C, Palmedo G, Bonfacini V, Sangalli G, Kutzner H: Cytotoxic mycosis fungoides evolving from pityriasis lichenoides chronica in a seventeen-year-old girl. Report of a case. *Dermatology.* 2002;205(2):176-9

<sup>122</sup> Shieh S, Mikkola DL, Wood GS: Differentiation and clonality of lesional lymphocytes in pityriasis lichenoides chronica. *Arch Dermatol.* 2001 Mar;137(3):305-8

<sup>123</sup> Nihal M, Mikkola D, Horvath N, Gilliam AC, Stevens SR, Spiro TP: Cutaneous lymphoid hyperplasia: a lymphoproliferative continuum with lymphomatous potential. *Hum Pathol.* 2003 Jun;34(6):617-22

<sup>124</sup> Gilliam AC, Wood GS: Cutaneous lymphoid hyperplasias. *Semin Cutan Med Surg.* 2000 Jun;19(2):133-41

<sup>125</sup> Kempf W, Haeffner AC, Zepter K, Sander CA, Flraig MJ, Mueller B: Angiolymphoid hyperplasia with eosinophilia: evidence for a T-cell lymphoproliferative origin. *Hum Pathol.* 2002 Oct;33(10):1023-9

<sup>126</sup> Klemke CD, Dippel E, Dembinski A, Ponitz N, Assaf C, Hummel M et al.: Clonal T cell receptor gamma-chain gene rearrangement by PCR-based GeneScan analysis in the skin and blood of patients with parapsoriasis and early-stage mycosis fungoides. *J Pathol.* 2002 Jul;197(3):348-54

<sup>127</sup> Willemze R, Meijer CJ: Classification of cutaneous T-cell lymphoma: from Alibert to WHO-EORTC. *J Cutan Pathol.* 2006 Feb;33 Suppl 1:18-26

<sup>128</sup> Dadej K, Gaboury L, Lamarre L, Petorin C, Seguin C, Cadotte M et al: The value of clonality in the diagnosis and follow-up of patients with cutaneous T-cell infiltrates. *Diagn Mol Pathol.* 2001 Jun;10(2):78-88

<sup>129</sup> Khalil SH, Hamadah IR: The applicability of T-cell receptor gamma gene rearrangement as an adjuvant diagnostic tool in skin biopsies for cutaneous T-cell lymphoma. *Saudi Med J.* 2006 Jul;27(7):951-954

<sup>130</sup> Ponti R, Quaglino P, Novelli M, Fierro MT, Comessatti A, Peroni A, et al.: T-cell receptor gamma gene rearrangement by multiplex polymerase chain reaction/heteroduplex analysis in patients with cutaneous T-cell lymphoma (mycosis fungoides/Sezary syndrome) and benign inflammatory disease: correlation with clinical, histological and immunophenotypical findings. *Br J Dermatol.* 2005 Sep;153(3):565-73

<sup>131</sup> Morgan SM, Hodges E, Mitchell TJ, Harris S, Whittaker SJ, Smith JL: Molecular analysis of T-cell receptor Beta genes in cutaneous T-cell lymphoma reveals jbeta1 bias. *J Invest Dermatol.* 2006 Aug;126(8):1893-9. Epub 2006 Jun 1

<sup>132</sup> Kneba M, Bolz I, Linke B, Hidemann W: Analysis of rearranged TCR $\beta$ -chain genes by PCR, DNA sequencing and automated high resolution PCR fragment analysis. *Blood* Vol 86/10: 3930-3937, 1995

- 
- <sup>133</sup> Murphy KM, Berg KD, Geiger T, Hafez M, Flickinger KA, Cooper L et al.: Capillary electrophoresis artifact due to eosin: implications for the interpretation of molecular diagnostic assays. *J Mol Diagn.* 2005 Feb;7(1):143-148
- <sup>134</sup> Johansson G, Isaksson R, Harang V: Migration time and peak area artifacts caused by systemic effects in voltage controlled capillary electrophoresis. *J Chromatogr A.* 2003 Jul 4;1004(1-2):91-8
- <sup>135</sup> Moreira BG, You Y, Behlke MA, Owczarzy R: Effects of fluorescent dyes, quenchers, and dangling ends on DNA duplex stability. *Biochem Biophys Res Commun.* 2005 Feb 11;327(2):473-84
- <sup>136</sup> Hahn M, Wilhelm J, Pingoud A: Influence of fluorophor dye labels on the migration behavior of polymerase chain reaction-amplified short tandem repeats during denaturing capillary electrophoresis. *Electrophoresis.* 2001 Aug;22(13):2691-700
- <sup>137</sup> Delfau MH, Hance AJ, Lecossier D, Vilmer E, Grandchamp B: Restricted diversity of V gamma 9-JP rearrangements in unstimulated human gamma/delta T lymphocytes. *Eur J Immunol.* 1992 Sep;22(9):2437-43
- <sup>138</sup> Lu D, Patel KA, Duvic M, Jones D: Clinical and pathological spectrum of CD8-positive cutaneous T-cell lymphomas. *J Cutan Pathol.* 2003 Mar;30(3):222
- <sup>139</sup> Chen M, Deng A, Crowson AN, Srinivasan M, Yearsley KH, Jewell S et al.: Assessment of T-cell clonality via T-cell receptor-gamma rearrangements in cutaneous T-cell-dominant infiltrates using polymerase chain reaction and single-stranded DNA conformational polymorphism assay. *Appl Immunohistochem Mol Morphol.* 2004 Dec;12(4):373-9
- <sup>140</sup> Langerak AW, Molina TJ, Lavender FL, Pearson D, Flohr T et al.: Polymerase chain reaction-based clonality testing in tissue samples with reactive lymphoproliferations: usefulness and pitfalls A Report of the BIOMED-2 Concerted Action BHM4-CT98-3936. *Leukemia* (2007) 21, 222-229
- <sup>141</sup> Van Krieken JHJM, Langerak AW, Macintyre EA, Kneba M, Hodges E, Garcia Sanz R et al.: Improved reliability of lymphoma diagnostics via PCR-based clonality-testing: Report of the BIOMED-2 Concerted Action BHM4-CT98-3936. *Leukemia* (2007) 21, 201-206
- <sup>142</sup> Brüggemann M, White H, Gaulard P, Garcia-Sanz P, Gameiro P: Powerful strategy for polymerase chain reaction-based clonality assessment in T-cell malignancies Report of the Biomed-2 Concerted Action BHM4 CT 98-3936. *Leukemia* (2007) 21, 215-221
- <sup>143</sup> Mitchell D, Willerslev E, Hansen A: Damage and repair of ancient DNA. *Mutat Res.* 2005 Apr 1;571(1-2):265-76
- <sup>144</sup> Ross KS, Haites NE, Kelly KF: Repeated freezing and thawing of peripheral blood and DNA in suspension: effects on DNA yield and integrity. *J Med Genet.* 1990 Sep;27(9):569-70
- <sup>145</sup> Gustafson S, Proper JA, Bowie EJ, Sommer SS: Parameters affecting the yield of DNA from human blood. *Anal Biochem.* 1987 Sep;165(2):294-9
- <sup>146</sup> Cushwa WT, Medrano JF: Effects of blood storage time and temperature on DNA yield and quality. *Biotechniques.* 1993 Feb;14(2):204-7
- <sup>147</sup> Farkas DH, Drevon AM, Kiechle FL, DiCarlo RG, Heath EM, Crisan D: Specimen stability for DNA-based diagnostic testing. *Diagn Mol Pathol.* 1996 Dec;5(4):227-35
- <sup>148</sup> Kilpatrick CW: Noncryogenic preservation of mammalian tissues for DNA extraction: an assessment of storage methods. *Biochem Genet.* 2002 Feb;40(1-2):53-62
- <sup>149</sup> Stanta G, Mucelli SP, Petrera F, Bonin S, Bussolati G: A novel fixative improves opportunities of nucleic acids and proteomic analysis in human archive's tissues. *Diagn Mol Pathol.* 2006 Jun;15(2):115-23