

8 Literaturverzeichnis

- Acampora D, Merlo GR, Paleari L, Zerega B, Postiglione MP, Mantero S, Bober E, Barbieri O, Simeone A, Levi G (1999).
Craniofacial, vestibular and bone defects in mice lacking the Distal-less-related gene Dlx5.
Development 1999; **126**, 3795-809.
- Adams JC. (1992).
Biotin amplification of biotin and horseradish peroxidase signals in histochemical stains.
J. Histochem. Cytochem. 1992 Oct; **40(10)**:1457-63.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990).
Basic local alignment search tool.
J. Mol. Biol. 1990; **215**:403-410.
- Alvarez J, Horton J, Sohn P, Serra R (2001).
The perichondrium plays an important role in mediating the effects of TGF-beta1 on endochondral bone formation.
Dev. Dyn. 2001 Jul; **221(3)**:311-21.
- Anderson MK, Pant R, Miracle AL, Sun X, Luer CA, Walsh CJ, Telfer JC, Litman GW, Rothenberg EV (2004).
Evolutionary origins of lymphocytes: ensembles of T cell and B cell transcriptional regulators in a cartilaginous fish.
J. Immunol. 2004 May 15; **172(10)**:5851-60.
- Angelastro JM, Töröcsik B, Greene LA (2002).
Nerve growth factor selectively regulates expression of transcripts encoding ribosomal proteins.
BMC Neurosci. 2002; **3**:3.
- Apte SS, Fukai N, Beier DR, Olsen BR (1997).
The matrix metalloproteinase-14 (MMP-14) gene is structurally distinct from other MMP genes and is co-expressed with the TIMP-2 gene during mouse embryogenesis.
J. Biol. Chem. 1997; **272(41)**:25511-25517.
- Armes N, Fried M (1995).
The genomic organization of the region containing the Drosophila melanogaster rpL7a (Surf-3) gene differs from those of the mammalian and avian Surf-3 loci.
Mol. Cell. Biol. 1995; **15(5)**:2367-2373.
- Aubin I, Adams CP, Opsahl S, Septier D, Bishop CE, Auge N, Salvayre R, Negre-Salvayre A, Goldberg M, Guenet JL, Poirier C.
A deletion in the gene encoding sphingomyelin phosphodiesterase 3 (Smpd3) results in osteogenesis and dentinogenesis imperfecta in the mouse.
Nat Genet. 2005 Aug; **37(8)**:803-5.
- Bangsow C, Rubins N, Glusman G, Bernstein Y, Negreanu V, Goldenberg D, Lotem J, Ben-Asher E, Lancet D, Levanon D, Groner Y (2001).
The RUNX3 gene-sequence, structure and regulated expression.
Gene 2001 Nov 28; **279(2)**:221-32.
- Bendall AJ, Hu G, Levi G, Abate-Shen C (2003).
Dlx5 regulates chondrocyte differentiation at multiple stages.
Int. J. Dev. Biol. 2003; **47**, 335-344.
- Benjamini Y, Hochberg Y (1995).
Controlling the false discovery rate – A practical and powerful approach to multiple testing.
J. Roy. Stat. Soc. B 1995; **57**:289–300.
- Bennett JH, Carter DH, Alavi AL, Beresford JN, Walsh S (2001).
Patterns of integrin expression in a human mandibular explant model of osteoblast differentiation.
Arch. Oral. Biol. 2001; **46(3)**:229-238.

- Blair JE, Hedges SB (2005).
Molecular phylogeny and divergence times of deuterostome animals.
Mol. Biol. Evol. 2005; **22**:2275-2284,.
- Bland YS, Critchlow MA, Ashhurst DE (1999).
The expression of the fibrillar collagen genes during fracture healing: heterogeneity of the matrices and differentiation of the osteoprogenitor cells.
Histochem. J. 1999; **31**(12):797-809.
- Brage M, Abrahamson M, Lindström V, Grubb A, Lerner UH (2005).
Different cysteine proteinases involved in bone resorption and osteoclast formation.
Calcif. Tissue Int. 2005; **76**(6):439-447.
- Brandenberger R, Schmidt A, Linton J, Wang D, Backus C, Denda S, Muller U, Reichardt LF (2001).
Identification and characterization of a novel extracellular matrix protein nephronectin that is associated with integrin alpha8beta1 in the embryonic kidney.
J. Cell. Biol. 2001; **154**:447-458.
- Brown TL, Fischer WC, Collins MD, De BK, Scott WJ (1994).
Identification of a 100-kDa phosphoprotein in developing murine embryos as elongation factor 2.
Arch Biochem Biophys 1994; **309**:105-110.
- Brusca GJ, Brusca RC, Burness TJ (2003).
Invertebrates.
Sunderland, MA, USA: Sinauer Associates Inc.; 2003.
- Cameron CB, Garey JR, Swalla BJ (2000).
Evolution of the chordate body plan: new insights from phylogenetic analyses of deuterostome phyla.
Proc. Natl. Acad. Sci. USA 2000; **97**:4469-4474
- Castresana J.
Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis.
Mol Biol Evol. 2000; **17**(4):540-52.
- Chang DF, Belaguli NS, Iyer D, Roberts WB, Wu SP, Dong XR, Marx JG, Moore MS, Beckerle MC, Majesky MW, Schwartz RJ (2003).
Cysteine-rich LIM-only proteins CRP1 and CRP2 are potent smooth muscle differentiation cofactors.
Dev. Cell 2003; **4**:107-118.
- Chang DJ, Ji C, Kim KK, Casinighino S, McCarthy TL, Centrella M (1998).
Reduction in transforming growth factor beta receptor I expression and transcription factor CBFa1 on bone cells by glucocorticoid.
J. Biol. Chem. 1998 Feb 27; **273**(9):4892-6.
- Chellaiah M, Kizer N, Silva M, Alvarez U, Kwiatkowski D, Hruska KA (2000).
Gelsolin deficiency blocks podosome assembly and produces increased bone mass and strength.
J. Cell. Biol. 2000; **148**:665-78.
- Cho EA, Dressler GR (1998).
TCF-4 binds beta-catenin and is expressed in distinct regions of the embryonic brain and limbs.
Mech. Dev. 1998; **77**:9-18.
- Chu K, Miyazaki M, Man WC, Ntambi JM (2006).
Stearoyl-coenzyme A desaturase 1 deficiency protects against hypertriglyceridemia and increases plasma high-density lipoprotein cholesterol induced by liver X receptor activation.
Mol Cell Biol. 2006; **26**(18):6786-98.
- Coggins KG, Latour A, Nguyen MS, Audoly L, Coffman TM, Koller BH (2002).
Metabolism of PGE2 by prostaglandin dehydrogenase is essential for remodeling the ductus arteriosus.
Nat. Med. 2002; **8**:91-2.

- Cohen MM, Jr. (2003).
The hedgehog signaling network.
Am. J. Med. Genet. A 2003; **123**:5-28.
- Constam DB, Robertson EJ (2000).
SPC4/PACE4 regulates a TGFbeta signaling network during axis formation.
Genes Dev. 2000; **14**:1146-1155.
- Corradetti MN, Inoki K, Guan KL (2005).
The stress-induced proteins RTP801 and RTP801L are negative regulators of the mammalian target of rapamycin pathway.
J Biol Chem. 2005; **280**(11):9769-72.
- D'Alonzo RC, Selvamurugan N, Karsenty G, Partridge NC (2002).
Physical interaction of the activator protein-1 factors c-Fos and c-Jun with Cbfa1 for collagenase-3 promoter activation.
J. Biol. Chem. 2002; **277**:816-822.
- Dang CM, Beanes SR, Lee H, Zhang X, Soo C, Ting K (2003).
Scarless fetal wounds are associated with an increased matrix metalloproteinase-to-tissue-derived inhibitor of metalloproteinase ratio.
Plast. Reconstr. Surg. 2003, **111**(7):2273–2285.
- de Crombrughe B, Lefebvre V, Nakashima K (2001).
Regulatory mechanisms in the pathways of cartilage and bone formation.
Curr. Opin. Cell Biol. 2001; **13**:721-727.
- de Jong DS, Vaes BL, Decherig KJ, Feijen A, Hendriks JM, Wehrens R, Mummery CL, van Zoelen EJ, Olijve W, Steegenga WT (2004).
Identification of novel regulators associated with early-phase osteoblast differentiation.
J. Bone Miner. Res. 2004; **19**:947-958.
- Delaisse JM, Andersen TL, Engsig MT, Henriksen K, Troen T, Blavier L (2003).
Matrix metalloproteinases (MMP) and cathepsin K contribute differently to osteoclastic activities.
Microsc. Res. Tech. 2003; **61**:504-513.
- Deleersnijder W, Hong G, Cortvrindt R, Poirier C, Tylzanowski P, Pittois K, Marck EV, Merregaert J (1996).
Isolation of markers for chondro-osteogenic differentiation using cDNA library subtraction. Molecular cloning and characterization of a gene belonging to a novel multigene family of integral membrane proteins.
J. Biol. Chem. 1996, **271**(32):19475-19482.
- Delsuc F, Brinkmann H, Chourrout D, Philippe H (2006).
Tunicates and not cephalochordates are the closest living relatives of vertebrates.
Nature 2006; **439**:965-968
- Dieterich C, Grossmann S, Tanzer A, Ropcke S, Arndt PF, Stadler PF, Vingron M (2005).
Comparative promoter region analysis powered by CORG.
BMC Genomics 2005; **6**:24
- Dobrev G, Chahrour M, Dautzenberg M, Chirivella L, Kanzler B, Farinas I, Karsenty G, Grosschedl R (2006).
SATB2 is a multifunctional determinant of craniofacial patterning and osteoblast differentiation.
Cell. 2006 Jun 2; **125**(5):971-86.
- Dong H, Osmanova V, Epstein PM, Brocke S (2006).
Phosphodiesterase 8 (PDE8) regulates chemotaxis of activated lymphocytes.
Biochem Biophys Res Commun. 2006; **345**(2):713-9.
- Drissi H, Luc Q, Shakoori R, Chuva De Sousa Lopes S, Choi JY, Terry A, Hu M, Jones S, Neil JC, Lian JB, Stein JL, Van Wijnen AJ, Stein GS (2000).
Transcriptional autoregulation of the bone related CBFA1/RUNX2 gene.
J. Cell. Physiol. 2000 Sep; **184**(3):341-50.

- D'Souza RN, Aberg T, Gaikwad J, Cavender A, Owen M, Karsenty G, Thesleff I (1999).
Cbfa1 is required for epithelial-mesenchymal interactions regulating tooth development in mice.
Development 1999 Jul; **126(13)**:2911-20.
- Duarte WR, Shibata T, Takenaga K, Takahashi E, Kubota K, Ohya K, Ishikawa I, Yamauchi M, Kasugai S (2003).
S100A4: a novel negative regulator of mineralization and osteoblast differentiation.
J. Bone Miner. Res. 2003, **18(3)**:493-501.
- Ducy P, Starbuck M, Priemel M, Shen J, Pinero G, Geoffroy V, Amling M, Karsenty G (1999).
A Cbfa1-dependent genetic pathway controls bone formation beyond embryonic development.
Genes Dev. 1999 Apr 15; **13(8)**:1025-36.
- EL Naaman C, Grum-Schwensen B, Mansouri A, Grigorian M, Santoni-Rugiu E, Hansen T, Kriajevska M, Schafer BW, Heizmann CW, Lukanidin E, Ambartsumian N (2004).
Cancer predisposition in mice deficient for the metastasis-associated Mts1(S100A4) gene.
Oncogene. 2004 Apr 29; **23(20)**:3670-80.
- Epari DR, Schell H, Bail HJ, Duda GN (2006).
Instability prolongs the chondral phase during bone healing in sheep.
Bone 2006 Jun; **38(6)**:864-70. Epub 2005 Dec 15.
- Everett AD, Narron JV, Stoops T, Nakamura H, Tucker A (2004).
Hepatoma-derived growth factor is a pulmonary endothelial cell-expressed angiogenic factor.
Am. J. Physiol. Lung Cell Mol. Physiol. 2004, **286(6)**:L1194-L1201.
- Fen JQ, Zhang J, Dallas SL, Lu Y, Chen S, Tan X, Owen M, Harris SE, MacDougall M (2002).
Dentin matrix protein 1, a target molecule for Cbfa1 in bone, is a unique bone marker gene.
J. Bone Miner. Res. 2002; **17**:1822-1831.
- Filanti C, Dickson GR, Martino DD, Ulivi V, Sanguineti C, Romano P, Palermo C, Manduca P (2000).
The expression of metalloproteinase-2, -9, and -14 and of tissue inhibitors-1 and -2 is developmentally modulated during osteogenesis in vitro, the mature osteoblastic phenotype expressing metalloproteinase-14.
J. Bone Miner. Res. 2000; **15(11)**:2154-2168.
- Fragonas E, Pollesello P, Mlinárik V, Toffanin R, Grando C, Godeas C, Vittur F (1998).
Sensitivity of chondrocytes of growing cartilage to reactive oxygen species.
Biochim Biophys Acta 1998; **1425**:103-111.
- Fransson LA (2003).
Glypicans.
Int. J. Biochem. Cell Biol. 2003; **35**:125-129.
- Friese MA, Hellwage J, Jokiranta TS, Meri S, Muller-Quernheim HJ, Peter HH, Eibel H, Zipfel PF (2000).
Different regulation of factor H and FHL-1/reconectin by inflammatory mediators and expression of the two proteins in rheumatoid arthritis (RA).
Clin Exp Immunol. 2000 Aug; **121(2)**:406-15.
- Fujikawa A, Shirasaka D, Yamamoto S, Ota H, Yahiro K, Fukada M, Shintani T, Wada A, Aoyama N, Hirayama T, Fukamachi H, Noda M (2003).
Mice deficient in protein tyrosine phosphatase receptor type Z are resistant to gastric ulcer induction by VacA of Helicobacter pylori.
Nat Genet. 2003; **33(3)**:375-81.
- Fukazawa N, Ayukawa K, Nishikawa K, Ohashi H, Ichihara N, Hikawa Y, Abe T, Kudo Y, Kiyama H, Wada K, Aoki S (2006).
Identification and functional characterization of mouse SERINC5 as a myelin membrane protein.
Brain Res. 2006; **1070**:1-14.

- Gabison EE, Hoang-Xuan T, Mauviel A, Menashi S (2005).
EMMPRIN/CD147, an MMP modulator in cancer, development and tissue repair.
Biochimie 2005; **87(3-4)**:361-368.
- Garcia-Cao I, Lafuente MJ, Criado LM, Diaz-Meco MT, Serrano M, Moscat J (2003).
Genetic inactivation of Par4 results in hyperactivation of NF-kappaB and impairment of JNK and p38.
EMBO Rep. 2003 Mar; **4(3)**:307-12.
- Gergen JP, Butler BA (1988).
Isolation of the Drosophila segmentation gene runt and analysis of its expression during embryogenesis.
Genes Dev. 1988; **2**:1179-1193.
- Gersdorff N, Müller M, Otto S, Poschadel R, Hübner S, Miosge N (2005).
Basement membrane composition in the early mouse embryo day 7.
Dev. Dyn. 2005; **233(3)**:1140-1148.
- Gerstenfeld LC, Einhorn TA (2003).
Developmental aspects of fracture healing and the use of pharmacological agents to alter healing.
J Musculoskelet Neuronal Interact 2003, **3(4)**:297–303; discussion 320–1.
- Giger RJ, Cloutier JF, Sahay A, Prinjha RK, Levensgood DV, Moore SE, Pickering S, Simmons D, Rastan S, Walsh FS, Kolodkin AL, Ginty DD, Geppert M (2000).
Neuropilin-2 is required in vivo for selective axon guidance responses to secreted semaphorins.
Neuron 2000; **25**:29-41.
- Gilbert, S.F. (1994).
Developmental biology. 4th Ed.
Sinauer Assoc. USA.
- Globus RK, Amblard D, Nishimura Y, Iwaniec UT, Kim JB, Almeida EAC, Damsky CD, Wronski TJ, van der Meulen MCH (2005).
Skeletal phenotype of growing transgenic mice that express a function-perturbing form of beta1 integrin in osteoblasts.
Calcif. Tissue Int. 2005; **76**:39-49.
- Gravallese EM (2003).
Osteopontin: a bridge between bone and the immune system.
J. Clin. Invest. 2003; **112**:147-149.
- Gurlek A, Pittelkow MR, Kumar R (2002).
Modulation of growth factor/cytokine synthesis and signaling by 1alpha,25-dihydroxyvitamin D(3): implications in cell growth and differentiation.
Endocr. Rev. 2002; **23**:763-786.
- Gustafsson E, Aszodi A, Ortega N, Hunziker EB, Denker HW, Werb Z, Fassler R (2003).
Role of collagen type II and perlecan in skeletal development.
Ann. N. Y. Acad. Sci. 2003; **995**:140-150.
- Haas SA, Beissbarth T, Rivals E, Krause A, Vingron M (2000).
GeneNest: automated generation and visualization of gene indices.
Trends Genet 2000; **16(11)**:521-523.
- Hadjiargyrou M, Lombardo F, Zhao S, Ahrens W, Joo J, Ahn H, Jurman M, White DW, Rubin CT (2002).
Transcriptional profiling of bone regeneration. Insight into the molecular complexity of wound repair.
J. Biol. Chem. 2002; **277(33)**:30177-30182.

- Hagood JS, Prabhakaran P, Kumbla P, Salazar L, MacEwen MW, Barker TH, Ortiz LA, Schoeb T, Siegal GP, Alexander CB, Pardo A, Selman M (2005).
Loss of fibroblast Thy-1 expression correlates with lung fibrogenesis.
Am J Pathol. 2005; **167(2)**:365-79.
- Hall BK (1999).
The neural crest in development and evolution.
New York: Springer; 1999
- Hall BK (2005a).
Bones and Cartilage, Developmental and Evolutionary Skeletal Biology.
London: Elsevier Academic Press; 2005.
- Hall BK (2005b).
Consideration of the neural crest and its skeletal derivatives in the context of novelty/innovation.
J. Exp. Zool B Mol. Dev. Evol. 2005; **304**:548-557.
- Harroch S, Furtado GC, Brueck W, Rosenbluth J, Lafaille J, Chao M, Buxbaum JD, Schlessinger J (2002).
A critical role for the protein tyrosine phosphatase receptor type Z in functional recovery from demyelinating lesions.
Nat. Genet. 2002; **32**:411-414.
- Hatori K, Sasano Y, Takahashi I, Kamakura S, Kagayama M, Sasaki K (2004).
Osteoblasts and osteocytes express MMP2 and -8 and TIMP1, -2, and -3 along with extracellular matrix molecules during appositional bone formation.
Anat. Rec. A Discov. Mol. Cell. Evol. Biol. 2004, **277(2)**:262-271.
- Hayman AR, Jones SJ, Boyde A, Foster D, Colledge WH, Carlton MB, Evans MJ, Cox TM (1996).
Mice lacking tartrate-resistant acid phosphatase (Acp 5) have disrupted endochondral ossification and mild osteopetrosis.
Development 1996; **122(10)**:3151-3162.
- Helseth DL, Veis A (1984).
Cathepsin D-mediated processing of procollagen: lysosomal enzyme involvement in secretory processing of procollagen.
Proc. Natl. Acad. Sci. USA 1984; **81(11)**:3302-3306.
- Herzig U, Cadenas C, Sieckmann F, Sierralta W, Thaller C, Visel A, Eichele G (2001).
Development of high-throughput tools to unravel the complexity of gene expression patterns in the mammalian brain.
Novartis Found. Symp. 2001; **239**:129-46; discussion 146-59. Review.
- Hieta N, Impola U, Lóez-Otín C, Saarialho-Kere U, Kähäri VM (2003).
Matrix metalloproteinase-19 expression in dermal wounds and by fibroblasts in culture.
J. Invest. Dermatol. 2003, **121(5)**:997-1004.
- Hong HH, Pischon N, Santana RB, Palamakumbura AH, Chase HB, Gantz D, Guo Y, Uzel MI, Ma D, Trackman PC (2004).
A role for lysyl oxidase regulation in the control of normal collagen deposition in differentiating osteoblast cultures.
J. Cell. Physiol. 2004; **200**:53-62.
- Huelsenbeck JP, Ronquist F (2001).
MRBAYES: Bayesian inference of phylogenetic trees.
Bioinformatics 2001; **17**:754-755.
- Hwang KC, Cui XS, Park SP, Shin MR, Park SY, Kim EY, Kim NH (2004).
Identification of differentially regulated genes in bovine blastocysts using an annealing control primer system.
Mol. Reprod. Dev. 2004; **69**:43-51.

- Inada M, Wang Y, Byrne MH, Rahman MU, Miyaura C, Lopez-Otin C, Krane SM (2004).
Critical roles for collagenase-3 (Mmp13) in development of growth plate cartilage and in endochondral ossification.
Proc. Natl. Acad. Sci. USA 2004; **101(49)**:17192-17197.
- Inoue K, Fei YJ, Zhuang L, Gopal E, Miyauchi S, Ganapathy V (2004).
Functional features and genomic organization of mouse NaCT, a sodium-coupled transporter for tricarboxylic acid cycle intermediates.
Biochem J. 2004; **378(Pt 3)**:949-57.
- Irie S, Hachiya T, Rabizadeh S, Maruyama W, Mukai J, Li Y, Reed JC, Bredesen DE, Sato TA (1999).
Functional interaction of Fas-associated phosphatase-1 (FAP-1) with p75(NTR) and their effect on NF-kappaB activation.
FEBS Lett. 1999; **460**:191-198.
- Ishii M, Koike C, Igarashi A, Yamanaka K, Pan H, Higashi Y, Kawaguchi H, Sugiyama M, Kamata N, Iwata T, Matsubara T, Nakamura K, Kurihara H, Tsuji K, Kato Y (2005).
Molecular markers distinguish bone marrow mesenchymal stem cells from fibroblasts.
Biochem. Biophys. Res. Commun. 2005; **332**:297-303.
- Jakob F, Becker K, Paar E, Ebert-Duemig R, Schutze N (2002).
Expression and regulation of thioredoxin reductases and other selenoproteins in bone.
Methods Enzymol. 2002; **347**:168-179.
- Javed A, Barnes GL, Jasanya BO, Stein JL, Gerstenfeld L, Lian JB, Stein GS (2001).
Runt homology domain transcription factors (Runx, Cbfa, and AML) mediate repression of the bone sialoprotein promoter: evidence for promoter context-dependent activity of Cbfa proteins.
Mol. Cell. Biol. 2001; **21**: 2891-905.
- Jeanmougin F, Thompson JD, Gouy M, Higgins DG, Gibson TJ (1998).
Multiple sequence alignment with Clustal X.
Trends Biochem Sci. 1998 Oct; **23(10)**:403-5.
- Jefferies RPS (1986).
The Ancestry of Vertebrates.
London: British Museum (Natural History); 1986.
- Jeschke M, Brandi ML, Susa M (1998).
Expression of Src family kinases and their putative substrates in the human preosteoclastic cell line FLG 29.1.
J Bone Miner Res. 1998 Dec; **13(12)**:1880-9.
- Jiang XC, Bruce C, Mar J, Lin M, Ji Y, Francone OL, Tall AR (1999).
Targeted mutation of plasma phospholipid transfer protein gene markedly reduces high-density lipoprotein levels.
J. Clin. Invest. 1999; **103**:907-914.
- Jiang XC, Li Z, Liu R, Yang XP, Pan M, Lagrost L, Fisher EA, Williams KJ (2005).
Phospholipid transfer protein deficiency impairs apolipoprotein-B secretion from hepatocytes by stimulating a proteolytic pathway through a relative deficiency of vitamin E and an increase in intracellular oxidants.
J Biol Chem. 2005; **280(18)**:18336-40.
- Jonker L, Arthur HM (2002).
Endoglin expression in early development is associated with vasculogenesis and angiogenesis.
Mech. Dev. 2002; **110(1-2)**:193-196.
- Jukkola T, Sinjushina N, Partanen J (2004).
Apcdd1 expression during mouse embryonic development.
Gene Expr. Patterns 2004; **4**:755-762.

- Kalajzic I, Braut A, Guo D, Jiang X, Kronenberg MS, Mina M, Harris MA, Harris SE, Rowe DW (2004).
Dentin matrix protein 1 expression during osteoblastic differentiation, generation of an osteocyte GFP-transgene.
Bone 2004; **35**:74-82.
- Keene DR, Sakai LY, Burgeson RE (1991).
Human bone contains type III collagen, type VI collagen, and fibrillin: type III collagen is present on specific fibers that may mediate attachment of tendons, ligaments, and periosteum to calcified bone cortex.
J. Histochem. Cytochem. 1991; **39**:59-69.
- Kenner L, Hoebertz A, Beil T, Keon N, Karreth F, Eferl R, Scheuch H, Szremska A, Amling M, Schorpp-Kistner M, Angel P, Wagner EF (2004).
Mice lacking JunB are osteopenic due to cell-autonomous osteoblast and osteoclast defects.
J. Cell. Biol. 2004; **164**(4):613-623.
- Kerstens HMJ, Poddighe PJ, Hanselaar AGJM (1995).
A novel in situ hybridization signal amplification method based on the deposition of biotinylated tyramine.
J. Histochem. Cytochem. 1995; **43**:347-352.
- Kim G, Lee TH, Wetzel P, Geers C, Robinson MA, Myers TG, Owens JW, Wehr NB, Eckhaus MW, Gros G, Wynshaw-Boris A, Levine RL (2004).
Carbonic anhydrase III is not required in the mouse for normal growth, development, and life span.
Mol. Cell. Biol. 2004; **24**:9942-9947.
- Kitching R, Qi S, Li V, Raouf A, Vary CP, Seth A (2002).
Coordinate gene expression patterns during osteoblast maturation and retinoic acid treatment of MC3T3-E1 cells.
J. Bone Miner. Metab. 2002; **20**:269-280.
- Komatsu DE, Hadjiargyrou M (2004).
Activation of the transcription factor HIF-1 and its target genes, VEGF, HO-1, iNOS, during fracture repair.
Bone 2004; **34**(4):680-688.
- Kornak U, Mundlos S (2003).
Genetic disorders of the skeleton: a developmental approach.
Am. J. Hum. Genet. 2003 Sep; **73**(3):447-74. Epub 2003 Jul 31. Review.
- Krause A, Haas SA, Coward E, Vingron M (2002).
SYSTEMS, GeneNest, SpliceNest: exploring sequence space from genome to protein.
Nucleic Acids Res. 2002; **30**:299-300.
- Levanon D, Glusman G, Bangsow T, Ben-Asher E, Male DA, Avidan N, Bangsow C, Hattori M, Taylor TD, Taudien S, Blechschmidt K, Shimizu N, Rosenthal A, Sakaki Y, Lancet D, Groner Y (2001).
Architecture and anatomy of the genomic locus encoding the human leukemia-associated transcription factor RUNX1/AML1.
Gene. 2001 Jan 10; **262**(1-2):23-33.
- Li QL, Ito K, Sakakura C, Fukamachi H, Inoue K, Chi XZ, Lee KY, Nomura S, Lee CW, Han SB, Kim HM, Kim WJ, Yamamoto H, Yamashita N, Yano T, Ikeda T, Itohara S, Inazawa J, Abe T, Hagiwara A, Yamagishi H, Ooe A, Kaneda A, Sugimura T, Ushijima T, Bae SC, Ito Y (2002).
Causal relationship between the loss of RUNX3 expression and gastric cancer.
Cell. 2002 Apr 5; **109**(1):113-24.
- Li X, Quigg RJ, Zhou J, Ryaby JT, Wang H (2005).
Early signals for fracture healing.
J. Cell. Biochem. 2005; **95**:189-205.

- Lin Q, Lu J, Yanagisawa H, Webb R, Lyons GE, Richardson JA, Olson EN (1998).
Requirement of the MADS-box transcription factor MEF2C for vascular development.
Development 1998; **125**:4565-74.
- Liu D, Kang JS, Derynck R (2004).
TGF-beta-activated Smad3 represses MEF2-dependent transcription in myogenic differentiation.
EMBO J. 2004; **23**:1557-1566.
- Liu X, Wu H, Byrne M, Krane S, Jaenisch R (1997).
Type III collagen is crucial for collagen I fibrillogenesis and for normal cardiovascular development.
Proc. Natl. Acad. Sci. USA 1997; **94**(5):1852-1856.
- Liu Y, Shi Z, Silveira A, Liu J, Sawadogo M, Yang H, Feng X (2003).
Involvement of upstream stimulatory factors 1 and 2 in RANKL-induced transcription of tartrate-resistant acid phosphatase gene during osteoclast differentiation.
J. Biol. Chem. 2003; **278**:20603-20611.
- Lockyer PJ, Vanlingen S, Reynolds JS, McNulty TJ, Irvine RF, Parys JB, Cullen PJ (1999).
Tissue-specific expression and endogenous subcellular distribution of the inositol 1,3,4,5-tetrakisphosphatebinding proteins GAP1(IP4BP) and GAP1(m).
Biochem. Biophys. Res. Commun. 1999; **255**:421-426.
- Lowell CA, Soriano P, Varmus HE (1994).
Functional overlap in the src gene family: inactivation of hck and fgr impairs natural immunity.
Genes Dev. 1994; **8**:387-98.
- Lu X, Le Noble F, Yuan L, Jiang Q, De Lafarge B, Sugiyama D, Breant C, Claes F, De Smet F, Thomas JL, Autiero M, Carmeliet P, Tessier-Lavigne M, Eichmann A (2004).
The netrin receptor UNC5B mediates guidance events controlling morphogenesis of the vascular system.
Nature. 2004 Nov 11; **432**(7014):179-86.
- Lu ZH, Books JT, Ley TJ (2005).
YB-1 is important for late-stage embryonic development, optimal cellular stress responses, and the prevention of premature senescence.
Mol. Cell. Biol. 2005; **25**(11):4625-4637.
- Maeda Y, Tsuji K, Nifuji A, Noda M (2004).
Inhibitory helix-loop-helix transcription factors Id1/Id3 promote bone formation in vivo.
J. Cell. Biochem. 2004; **93**(2):337-344.
- Mao J, Qiao X, Luo H, Wu J (2006).
Transgenic drak2 overexpression in mice leads to increased T cell apoptosis and compromised memory T cell development.
J Biol Chem. 2006; **281**(18):12587-95.
- Marie PJ (2002).
Role of N-cadherin in bone formation.
J. Cell. Physiol. 2002; **190**:297-305.
- Mattson MP, Duan W, Chan SL, Camandola S (1999).
Par-4: an emerging pivotal player in neuronal apoptosis and neurodegenerative disorders.
J. Mol. Neurosci. 1999; **13**:17-30.
- May LG, Gay CV (1997).
Multiple G-protein involvement in parathyroid hormone regulation of acid production by osteoclasts.
J. Cell. Biochem. 1997; **64**:161-170.
- McLysaght A, Hokamp K, Wolfe KH (2002).
Extensive genomic duplication during early chordate evolution.
Nat. Genet. 2002; **31**:200-204.

- Mena C, Devlin RD, Reddy SV, Gazitt Y, Choi SJ, Roodman GD (1999).
Annexin II increases osteoclast formation by stimulating the proliferation of osteoclast precursors in human marrow cultures.
J. Clin. Invest. 1999; **103(11)**:1605-1613.
- Michael M, Dieterich C, Vingron M (2005).
SITEBLAST-rapid and sensitive local alignment of genomic sequences employing motif anchors.
Bioinformatics. 2005 May 1; **21(9)**:2093-4. Epub 2004 Dec 14.
- Miller CT, Yelon D, Stainier DY, Kimmel CB (2003).
Two endothelin 1 effectors, hand2 and bapx1, pattern ventral pharyngeal cartilage and the jaw joint.
Development. 2003 Apr; **130(7)**:1353-65.
- Miyazawa T, Ogawa Y, Chusho H, Yasoda A, Tamura N, Komatsu Y, Pfeifer A, Hofmann F, Nakao K (2002).
Cyclic GMP-dependent protein kinase II plays a critical role in C-type natriuretic peptide-mediated endochondral ossification.
Endocrinology 2002; **143**:3604-3610.
- Miyazono K, Maeda S, Imamura T (2004).
Coordinate regulation of cell growth and differentiation by TGF-beta superfamily and Runx proteins.
Oncogene 2004; **23**:4232-4237.
- Miyoshi H, Shimizu K, Kozu T, Maseki N, Kaneko Y, Ohki M (1991).
t(8;21) breakpoints on chromosome 21 in acute myeloid leukemia are clustered within a limited region of a single gene, AML1.
Proc. Natl. Acad. Sci. USA. 1991 Dec 1; **88(23)**:10431-4.
- Mukhopadhyay M, Shtrom S, Rodriguez-Esteban C, Chen L, Tsukui T, Gomer L, Dorward DW, Glinka A, Grinberg A, Huang SP, Niehrs C, Belmonte JC, Westphal H (2001).
Dickkopf1 is required for embryonic head induction and limb morphogenesis in the mouse.
Dev. Cell 2001; **1**:423-34.
- Müller HJ (2001).
PCR - Polymerase-Kettenreaktion
Spektrum Akademischer Verlag, Heidelberg
- Mundlos S, Otto F, Mundlos C, Mulliken JB, Aylsworth AS, Albright S, Lindhout D, Cole WG, Henn W, Knoll JH, Owen MJ, Mertelsmann R, Zabel BU, Olsen BR (1997).
Mutations involving the transcription factor CBF1 cause cleidocranial dysplasia.
Cell 1997; **89(5)**:773-779.
- Myokai F, Oyama M, Nishimura F, Ohira T, Yamamoto T, Arai H, Takashiba S, Murayama Y (2003).
Unique genes induced by mechanical stress in periodontal ligament cells.
J. Periodontal Res. 2003; **38(3)**:255-261.
- Nakashima K, Zhou X, Kunkel G, Zhang Z, Deng JM, Behringer RR, de Crombrughe B (2002).
The novel zinc finger-containing transcription factor osterix is required for osteoblast differentiation and bone formation.
Cell 2002; **108**:17-29.
- Nam S, Jin YH, Li QL, Lee KY, Jeong GB, Ito Y, Lee J, Bae SC (2002).
Expression pattern, regulation, and biological role of runt domain transcription factor, run, in Caenorhabditis elegans.
Mol. Cell Biol. 2002; **22**:547-554.
- Nielsen C (1995).
Animal Evolution: Interrelationships of the Living Phyla.
Oxford: Oxford Univ. Press; 1995.
- Nosten-Bertrand M, Errington ML, Murphy KP, Tokugawa Y, Barboni E, Kozlova E, Michalovich D, Morris RG, Silver J, Stewart CL, Bliss TV, Morris RJ (1996).
Normal spatial learning despite regional inhibition of LTP in mice lacking Thy-1.
Nature 1996; **379**:826-9.

- Ntambi JM, Miyazaki M, Stoehr JP, Lan H, Kendziorowski CM, Yandell BS, Song Y, Cohen P, Friedman JM, Attie AD (2002).
Loss of stearoyl-CoA desaturase-1 function protects mice against adiposity.
Proc. Natl. Acad. Sci. USA 2002; **99**:11482-11486.
- Oda H, Wada H, Tagawa K, Akiyama-Oda Y, Satoh N, Humphreys T, Zhang S, Tsukita S (2002).
A novel amphioxus cadherin that localizes to epithelial adherens junctions has an unusual domain organization with implications for chordate phylogeny.
Evol. Dev. 2002; **4**:426-434.
- Okamoto Y, Kim D, Battaglini R, Sasaki H, Spate U, Stashenko P (2004).
MIP-1 gamma promotes receptor-activator-of-NF-kappa-B ligand-induced osteoclast formation and survival.
J. Immunol. 2004; **173**:2084-2090.
- Otto F, Thornell AP, Crompton T, Denzel A, Gilmour KC, Rosewell IR, Stamp GW, Beddington RS, Mundlos S, Olsen BR, Selby PB, Owen MJ (1997).
Cbfa1, a candidate gene for cleidocranial dysplasia syndrome, is essential for osteoblast differentiation and bone development.
Cell 1997 May 30; **89**(5):765-71.
- Otto F, Kanegane H, Mundlos S (2002).
Mutations in the RUNX2 gene in patients with cleidocranial dysplasia.
Hum. Mutat. 2002; **19**:209-216.
- Ou JH, Yen TS, Wang YF, Kam WK, Rutter WJ (1987).
Cloning and characterization of a human ribosomal protein gene with enhanced expression in fetal and neoplastic cells.
Nucleic Acids Res. 1987; **15**(21):8919-8934.
- Paraoanu LE, Weiss B, Robitzki AA, Layer PG (2005).
Cytochrome-c oxidase is one of several genes elevated in marginal retina of the chick embryo.
Neuroscience 2005; **132**(3):665-672.
- Parsons SJ and Parsons JT (2004).
Src family kinases, key regulators of signal transduction.
Oncogene 2004; **23**:7906-9.
- Peters T, Ausmeier K, Dildrop R, Ruther U (2002).
The mouse Fused toes (Ft) mutation is the result of a 1.6-Mb deletion including the entire Iroquois B gene cluster.
Mamm Genome. 2002; **13**(4):186-8.
- Philippe H, Lartillot N, Brinkmann H (2005).
Multigene analyses of bilaterian animals corroborate the monophyly of Ecdysozoa, Lophotrochozoa, and Protostomia.
Mol. Biol. Evol. 2005; **22**:1246-1253.
- Pichaud F, Delage-Mourroux R, Frenkian M, Frenco JL, Roux S, de Vernejoul MC, Jullienne A (1997).
Type-I 15-hydroxyprostaglandin dehydrogenase: role in macrophage/osteoclast differentiation.
Adv. Exp. Med. Biol. 1997; **433**:399-402.
- Pickering MC, Warren J, Rose KL, Carlucci F, Wang Y, Walport MJ, Cook HT, Botto M (2006).
Prevention of C5 activation ameliorates spontaneous and experimental glomerulonephritis in factor H-deficient mice.
Proc Natl Acad Sci U S A. 2006; **103**(25):9649-54.
- Pineda VV, Athos JI, Wang H, Cerver J, Ippolito D, Boulay G, Birnbaumer L, Storm DR (2004).
Removal of G(alpha1) Constraints on Adenylyl Cyclase in the Hippocampus Enhances LTP and Impairs Memory Formation.
Neuron 2004; **41**:153-63.

- Pinto JP, Conceicao NM, Viegas CS, Leite RB, Hurst LD, Kelsh RN, Cancela ML (2005).
Identification of a new pebp2alphaA2 isoform from zebrafish runx2 capable of inducing osteocalcin gene expression in vitro.
J. Bone Miner. Res. 2005 Aug; **20(8)**:1440-53. Epub 2005 Mar 21.
- Portman MA, Chen SH, Xiao Y, Ning XH (1999).
Maturation changes in gene expression for adenine nucleotide translocator isoforms and betaF1-ATPase in rabbit heart.
Mol. Genet. Metab. 1999; **66(2)**:75-79.
- Qin C, Baba O, Butler WT (2004).
Post-translational modifications of sibling proteins and their roles in osteogenesis and dentinogenesis.
Crit. Rev. Oral. Biol. Med. 2004; **15**:126-136.
- Qin L, Raggatt LJ, Partridge NC (2004b).
Parathyroid hormone: a double-edged sword for bone metabolism.
Trends Endocrinol. Metab. 2004; **15**:60-5.
- Quarles LD (2003).
FGF23, PHEX, and MEPE regulation of phosphate homeostasis and skeletal mineralization.
Am. J. Physiol. Endocrinol. Metab. 2003; **285**:1-9.
- Raouf A, Ganss B, McMahon C, Vary C, Roughley PJ, Seth A (2002).
Lumican is a major proteoglycan component of the bone matrix.
Matrix. Biol. 2002; **21(4)**:361-367.
- Rescher U, Gerke V (2004).
Annexins-unique membrane binding proteins with diverse functions.
J. Cell. Sci. 2004; **117**:2631-2639.
- Robson P, Wright GM, Keeley FW (2000).
Distinct non-collagen based cartilages comprising the endoskeleton of the Atlantic hagfish, Myxine glutinosa.
Anat. Embryol. (Berl.) 2000; **202**:281-290
- Rodriguez de Cordoba S, Esparza-Gordillo J, Goicoechea de Jorge E, Lopez-Trascasa M, Sanchez-Corral P (2004).
The human complement factor H: functional roles, genetic variations and disease associations.
Mol. Immunol. 2004; **41**:355-67.
- Roose J, Huls G, van Beest M, Moerer P, van der Horn K, Goldschmeding R, Logtenberg T, Clevers H (1999).
Synergy between tumor suppressor APC and the beta-catenin-Tcf4 target Tcf1.
Science 1999; **285**:1923-6.
- Rudolph U, Finegold MJ, Rich SS, Harriman GR, Srinivasan Y, Brabet P, Boulay G, Bradley A, Birnbaumer L (1995).
Ulcerative colitis and adenocarcinoma of the colon in G alpha i2-deficient mice.
Nat. Genet. 1995; **10(2)**:143-150.
- Rychel AL, Smith SE, Shimamoto HT, Swalla BJ (2006).
Evolution and Development of the Chordates: Collagen and Pharyngeal Cartilage.
Mol. Biol. Evol. 2006; **23**:541-549
- Saltman LH, Javed A, Ribadeneyra J, Hussain S, Young DW, Osdoby P, Amcheslavsky A, van Wijnen AJ, Stein JL, Stein GS, Lian JB, Bar-Shavit Z (2005).
Organization of transcriptional regulatory machinery in osteoclast nuclei: Compartmentalization of Runx1.
J. Cell. Physiol. 2005 Sep; **204(3)**:871-80
- Sambrook J, Russell DW, Sambrook JE (2001).
Molecular Cloning: A laboratory manual;
Cold Spring Harbor Laboratory Press.

- Sato M, Morii E, Komori T, Kawahata H, Sugimoto M, Terai K, Shimizu H, Yasui T, Ogihara H, Yasui N, Ochi T, Kitamura Y, Ito Y, Nomura S (1998).
Transcriptional regulation of osteopontin gene in vivo by PEBP2alphaA/CBFA1 and ETS1 in the skeletal tissues.
Oncogene 1998; **17**:1517-25.
- Sato M, Yasui N, Nakase T, Kawahata H, Sugimoto M, Hirota S, Kitamura Y, Nomura S, Ochi T (1998b).
Expression of bone matrix proteins mRNA during distraction osteogenesis.
J. Bone Miner. Res. 1998; **13(8)**:1221-1231.
- Satoh N, Jeffery WR (1995).
Chasing tails in ascidians: developmental insights into the origin and evolution of chordates.
Trends Genet. 1995; **11**:354-359.
- Schell H, Epari DR, Kassi JP, Bragulla H, Bail HJ, Duda GN (2005).
The course of bone healing is influenced by the initial shear fixation stability.
J. Orthop. Res. 2005 Sep; **23(5)**:1022-8.
- Schinke T, Karsenty G (1999).
Characterization of Osf1, an osteoblast-specific transcription factor binding to a critical cis-acting element in the mouse Osteocalcin promoters.
J. Biol. Chem. 1999; **274**:30182-30189.
- Schwartzberg PL, Xing L, Hoffmann O, Lowell CA, Garrett L, Boyce BF, Varmus HE (1997).
Rescue of osteoclast function by transgenic expression of kinase-deficient Src in src^{-/-} mutant mice.
Genes Dev. 1997; **11**:2835-44.
- Seemann P (2005).
Zur Bedeutung des Wachstumsfaktors Gdf5 – Funktionelle Analyse und Genotyp-Phänotyp-Korrelation
Dissertation, Freie Universität Berlin, Fachbereich Biologie, Chemie und Pharmazie, Dezember 2005
- Seitz V, Ortiz Garcia S, Liston A (2000).
Alternative coding strategies and the inapplicable data coding problem.
Taxon. 2000; **49**:47-54.
- Selvamurugan N, Pulumati MR, Tyson DR, Partridge NC (2000).
Parathyroid hormone regulation of the rat collagenase-3 promoter by protein kinase A-dependent transactivation of core binding factor alpha1.
J. Biol. Chem. 2000; **275**:5037-5042.
- Shearer J, Fueger PT, Bracy DP, Wasserman DH, Rottman JN (2005).
Partial gene deletion of heart-type fatty acid-binding protein limits the severity of dietary-induced insulin resistance.
Diabetes. 2005; **54(11)**:3133-9.
- Shi GP, Sukhova GK, Kuzuya M, Ye Q, Du J, Zhang Y, Pan JH, Lu ML, Cheng XW, Iguchi A, Perrey S, Lee AME, Chapman HA, Libby P (2003).
Deficiency of the cysteine protease cathepsin S impairs microvessel growth.
Circ. Res. 2003; **92(5)**:493-500.
- Shroff B, Smith T, Norris K, Pileggi R, Sauk JJ (1993).
Hsp 47 is localized to regions of type I collagen production in developing murine femurs and molars.
Connect. Tissue Res. 1993; **29(4)**:273-286.
- Smith AN, Borthwick KJ, Karet FE (2002).
Molecular cloning and characterization of novel tissue-specific isoforms of the human vacuolar H(+)-ATPase C, G and d subunits, and their evaluation in autosomal recessive distal renal tubular acidosis.
Gene 2002; **297**:169-177.

- Steiglitz BM, Keene DR, Greenspan DS (2002).
PCOLCE2 encodes a functional procollagen C-proteinase enhancer (PCPE2) that is a collagen-binding protein differing in distribution of expression and posttranslational modification from the previously described PCPE1.
J. Biol. Chem. 2002; **277**:49820-49830.
- Stewart M, Terry A, Hu M, O'Hara M, Blyth K, Baxter E, Cameron E, Onions DE, Neil JC (1997).
Proviral insertions induce the expression of bone-specific isoforms of PEBP2alphaA (CBFA1): evidence for a new myc collaborating oncogene.
Proc. Natl. Acad. Sci. USA 1997 Aug 5; **94(16)**:8646-51.
- Stickens D, Behonick DJ, Ortega N, Heyer B, Hartenstein B, Yu Y, Fosang AJ, Schorpp-Kistner M, Angel P, Werb Z (2004).
Altered endochondral bone development in matrix metalloproteinase 13-deficient mice.
Development 2004; **131**:5883-95.
- Stipanuk MH (2004).
Role of the liver in regulation of body cysteine and taurine levels: a brief review.
Neurochem. Res. 2004; **29**:105-110.
- Stock M, Schafer H, Fliegauf M, Otto F (2004).
Identification of novel genes of the bonespecific transcription factor Runx2.
J. Bone Miner. Res. 2004; **19**:959-972.
- Stoffel W, Jenke B, Block B, Zumbansen M, Koebke J (2005).
Neutral sphingomyelinase 2 (smpd3) in the control of postnatal growth and development.
Proc. Natl. Acad. Sci. USA 2005; **102**:4554-4559.
- Stricker S, Fundele R, Vortkamp A, Mundlos S (2002).
Role of Runx genes in chondrocyte differentiation.
Dev Biol. 2002 May 1; **245(1)**:95-108.
- Stricker S, Poustka AJ, Wiecha U, Stiege A, Hecht J, Panopoulou G, Vilcinskas A, Mundlos S, Seitz V (2003).
A single amphioxus and sea urchin runt-gene suggests that runt-gene duplications occurred in early chordate evolution.
Dev. Comp. Immunol. 2003 Sep; **27(8)**:673-84.
- Suzuki MM, Satoh N (2000).
Genes expressed in the amphioxus notochord revealed by EST analysis.
Dev. Biol. 2000; **224**:168-177.
- Takahara K, Lyons GE, Greenspan DS (1994).
Bone morphogenetic protein-1 and a mammalian tolloid homologue (mTld) are encoded by alternatively spliced transcripts which are differentially expressed in some tissues.
J. Biol. Chem. 1994; **269(51)**:32572-32578.
- Takashima S, Kitakaze M, Asakura M, Asanuma H, Sanada S, Tashiro F, Niwa H, Miyazaki Ji J, Hirota S, Kitamura Y, Kitsukawa T, Fujisawa H, Klagsbrun M, Hori M (2002).
Targeting of both mouse neuropilin-1 and neuropilin-2 genes severely impairs developmental yolk sac and embryonic angiogenesis.
Proc Natl Acad Sci U S A. 2002; **99(6)**:3657-62.
- Thiele H, Sakano M, Kitagawa H, Sugahara K, Rajab A, Hohne W, Ritter H, Leschik G, Nurnberg P, Mundlos S (2004).
Loss of chondroitin 6-O-sulfotransferase-1 function results in severe human chondrodysplasia with progressive spinal involvement.
Proc. Natl. Acad. Sci. USA 2004 Jul 6; **101(27)**:10155-60. Epub 2004 Jun 23.
- Tintut Y, Parhami F, Le V, Karsenty G, Demer LL (1999).
Inhibition of osteoblast-specific transcription factor Cbfa1 by the cAMP pathway in osteoblastic cells. Ubiquitin/proteasome-dependent regulation.
J. Biol. Chem. 1999; **274**:28875-28879.

- Trivett MK, Potter IC, Power G, Zhou H, Macmillan DL, Martin TJ, Danks JA (2005).
Parathyroid hormone-related protein production in the lamprey *Geotria australis*: developmental and evolutionary perspectives.
Dev. Genes Evol. 2005 Nov; **215(11)**:553-63. Epub 2005 Nov 4.
- Uchida M, Shima M, Chikazu D, Fujieda A, Obara K, Suzuki H, Nagai Y, Yamato H, Kawaguchi H (2001).
Transcriptional induction of matrix metalloproteinase-13 (collagenase-3) by 1 α ,25-dihydroxyvitamin D3 in mouse osteoblastic MC3T3-E1 cells.
J. Bone Miner. Res. 2001; **16**:221-230.
- Uusitalo H, Hiltunen A, Söderström M, Aro HT, Vuorio E (2000).
Expression of cathepsins B, H, K, L, and S and matrix metalloproteinases 9 and 13 during chondrocyte hypertrophy and endochondral ossification in mouse fracture callus.
Calcif. Tissue Int. 2000; **67(5)**:382-390.
- Vaidya A, Pniak A, Lemke G, and Brown A (2003).
EphA3 null mutants do not demonstrate motor axon guidance defects.
Mol. Cell. Biol. 2003; **23**:8092-8.
- Van den Plas D, Merregaert J (2004).
In vitro studies on *Itm2a* reveal its involvement in early stages of the chondrogenic differentiation pathway.
Biol. Cell. 2004 Aug; **96(6)**:463-70.
- van der Meulen T, Kranenbarg S, Schipper H, Samallo J, van Leeuwen JL, Franssen H (2005).
Identification and characterisation of two *runx2* homologues in zebrafish with different expression patterns.
Biochim. Biophys. Acta. 2005 Jun 30; **1729(2)**:105-17. Epub 2005 Apr 26.
- Vortkamp A, Pathi S, Peretti GM, Caruso EM, Zaleske DJ, Tabin CJ (1998).
Recapitulation of signals regulating embryonic bone formation during postnatal growth and in fracture repair.
Mech. Dev. 1998 Feb; **71(1-2)**:65-76.
- Wagner EF (2002) .
Functions of AP1 (Fos/Jun) in bone development.
Ann. Rheum. Dis. 2002; **61 Suppl 2**:ii40-ii42.
- Wang Q, Xie Y, Du QS, Wu XJ, Feng X, Mei L, McDonald JM, Xiong WC (2003) .
Regulation of the formation of osteoclastic actin rings by proline-rich tyrosine kinase 2 interacting with gelsolin.
J. Cell. Biol. 2003; **160**:565-575.
- Wang Y, Middleton F, Horton JA, Reichel L, Farnum CE, Damron TA (2004).
Microarray analysis of proliferative and hypertrophic growth plate zones identifies differentiation markers and signal pathways.
Bone 2004; **35(6)**:1273-1293.
- Watanabe S, Akagi R, Mori M, Tsuchiya T, Sassa S (2004).
Marked developmental changes in heme oxygenase-1 (HO-1) expression in the mouse placenta: correlation between HO-1 expression and placental development.
Placenta 2004, **25(5)**:387-395.
- Wennberg C, Hessle L, Lundberg P, Mauro S, Narisawa S, Lerner UH, Millan JL (2000).
Functional characterization of osteoblasts and osteoclasts from alkaline phosphatase knockout mice.
J. Bone Miner. Res. 2000; **15**:1879-1888.
- Weston AD, Hoffman LM, Underhill TM (2003).
Revisiting the role of retinoid signaling in skeletal development.
Birth Defects Res. C. Embryo Today 2003; **69(2)**:156-173.

- Wiecha U (2003).
Untersuchungen zur Anzahl und Struktur *Runt*-homologer Gene bei den Cyclostomata und *Runt*-Expressionsanalyse bei *Branchiostoma floridae*.
Diplomarbeit, Freie Universität Berlin, Fachbereich Biologie, Chemie und Pharmazie, April 2003
- Winchell CJ, Sullivan J, Cameron CB, Swalla BJ, Mallatt J (2002).
Evaluating hypotheses of deuterostome phylogeny and chordate evolution with new LSU and SSU ribosomal DNA data.
Mol. Biol. Evol. 2002; **19**:762-776.
- Witke W, Li W, Kwiatkowski DJ, Southwick FS (2001).
Comparisons of CapG and gelsolin-null macrophages: demonstration of a unique role for CapG in receptor-mediated ruffling, phagocytosis, and vesicle rocketing.
J. Cell. Biol. 2001; **154**:775-784.
- Wu W, Glinka A, Delius H, Niehrs C (2000).
Mutual antagonism between dickkopf1 and dickkopf2 regulates Wnt/beta-catenin signalling.
Curr. Biol. 2000; **10**:1611-1614.
- Yamaguchi TP, Bradley A, McMahon AP, Jones S (1999).
A Wnt5a pathway underlies outgrowth of multiple structures in the vertebrate embryo.
Development 1999; **126**:1211-23.
- Yang Y, Topol L, Lee H, Wu J (2003).
Wnt5a and Wnt5b exhibit distinct activities in coordinating chondrocyte proliferation and differentiation.
Development. 2003; **130**(5):1003-15.
- Ye L, MacDougall M, Zhang S, Xie Y, Zhang J, Li Z, Lu Y, Mishina Y, Feng JQ (2004).
Deletion of dentin matrix protein-1 leads to a partial failure of maturation of predentin into dentin, hypomineralization, and expanded cavities of pulp and root canal during postnatal tooth development.
J. Biol. Chem. 2004; **279**:19141-8.
- Ying J, Li H, Cui Y, Wong AH, Langford C, Tao Q (2006).
Epigenetic disruption of two proapoptotic genes MAPK10/JNK3 and PTPN13/FAP-1 in multiple lymphomas and carcinomas through hypermethylation of a common bidirectional promoter.
Leukemia. 2006; **20**(6):1173-5.
- Yoshida CA, Furuichi T, Fujita T, Fukuyama R, Kanatani N, Kobayashi S, Satake M, Takada K, Komori T (2002).
Core-binding factor beta interacts with Runx2 and is required for skeletal development.
Nat. Genet. 2002 Dec; **32**(4):633-8. Epub 2002 Nov 18.
- Yoshida CA, Komori T (2005).
Role of Runx proteins in chondrogenesis.
Crit. Rev. Eukaryot. Gene Expr. 2005; **15**(3):243-54. Review.
- Yoshida CA, Yamamoto H, Fujita T, Furuichi T, Ito K, Inoue K, Yamana K, Zanma A, Takada K, Ito Y, Komori T (2004).
Runx2 and Runx3 are essential for chondrocyte maturation, and Runx2 regulates limb growth through induction of Indian hedgehog.
Genes Dev. 2004; **18**: 952-63.
- Yoshitake H, Rittling SR, Denhardt DT, Noda M (1999).
Osteopontin-deficient mice are resistant to ovariectomy-induced bone resorption.
Proc. Natl. Acad. Sci. USA 1999; **96**:8156-60.

- Yoshizawa T, Handa Y, Uematsu Y, Takeda S, Sekine K, Yoshihara Y, Kawakami T, Arioka K, Sato H, Uchiyama Y, Masushige S, Fukamizu A, Matsumoto T, Kato S (1997).
Mice lacking the vitamin D receptor exhibit impaired bone formation, uterine hypoplasia and growth retardation after weaning.
Nat. Genet. 1997; **16**:391-6.
- Zeng W, Yuan W, Wang Y, Jiao W, Zhu Y, Huang C, Li D, Li Y, Zhu C, Wu X, Liu M (2002).
Expression of a novel member of sortingnexin gene family, SNX-L, in human liver development.
Biochem. Biophys. Res. Commun. 2002 ; **299**:542-548.
- Zhang G, Miyamoto MM, Cohn MJ (2006).
Lamprey type II collagen and Sox9 reveal an ancient origin of the vertebrate collagenous skeleton.
Proc. Natl. Acad. Sci. USA 2006; **103**:3180-3185
- Zhao Y, Guan H, Liu SF, Wu RC, Wang Z (2005).
Overexpression of QM induces cell differentiation and mineralization in MC3T3-E1.
Biol. Pharm. Bull. 2005; **28(8)**:1371-1376.
- Zheng Q, Zhou G, Morello R, Chen Y, Garcia-Rojas X, Lee B (2003).
Type X collagen gene regulation by Runx2 contributes directly to its hypertrophic chondrocyte-specific expression in vivo.
J. Cell. Biol. 2003 Sep 1; **162(5)**:833-42.