

7. Literaturverzeichnis

- (1) Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med* 1993; 328(17):1230-1235.
- (2) Strohl KP, Redline S. Recognition of obstructive sleep apnea. *Am J Respir Crit Care Med* 1996; 154(2 Pt 1):279-289.
- (3) Nieto FJ, Young TB, Lind BK, Shahar E, Samet JM, Redline S et al. Association of sleep-disordered breathing, sleep apnea, and hypertension in a large community-based study. Sleep Heart Health Study. *JAMA* 2000; 283(14):1829-1836.
- (4) Peppard PE, Young T, Palta M, Skatrud J. Prospective study of the association between sleep-disordered breathing and hypertension. *N Engl J Med* 2000; 342(19):1378-1384.
- (5) Blazejova K, Sonka K, Skodova Z, Nevsimalova S. [Prevalence of obesity, hypertension and smoking in patients with the sleep apnea syndrome--comparison with the Czech population]. *Cas Lek Cesk* 2000; 139(11):339-342.
- (6) Zhang LQ, Yao WZ, He QY, Wang YZ, Ren B. [Effect of movement arousal and hypoxia on circadian blood pressure in hypertensive patients with obstructive sleep apnea syndrome]. *Zhonghua Yi Xue Za Zhi* 2003; 83(6):475-477.
- (7) Partinen M, Telakivi T. Epidemiology of obstructive sleep apnea syndrome. *Sleep* 1992; 15(6 Suppl):S1-S4.
- (8) Koehler U, Becker HF, Grimm W, Heitmann J, Peter JH, Schafer H. Relations among hypoxemia, sleep stage, and bradyarrhythmia during obstructive sleep apnea. *Am Heart J* 2000; 139(1 Pt 1):142-148.
- (9) Wright J, Johns R, Watt I, Melville A, Sheldon T. Health effects of obstructive sleep apnoea and the effectiveness of continuous positive airways pressure: a systematic review of the research evidence. *BMJ* 1997; 314(7084):851-860.
- (10) Schafer H, Koehler U, Ewig S, Hasper E, Tasci S, Luderitz B. Obstructive sleep apnea as a risk marker in coronary artery disease. *Cardiology* 1999; 92(2):79-84.
- (11) Turkington PM, Allgar V, Bamford J, Wanklyn P, Elliott MW. Effect of upper airway obstruction in acute stroke on functional outcome at 6 months. *Thorax* 2004; 59(5):367-371.
- (12) Davies CW, Crosby JH, Mullins RL, Barbour C, Davies RJ, Stradling JR. Case-control study of 24 hour ambulatory blood pressure in patients with obstructive sleep apnoea and normal matched control subjects. *Thorax* 2000; 55(9):736-740.

- (13) Fichter J, Bauer D, Arampatzis S, Fries R, Heisel A, Sybrecht GW. Sleep-related breathing disorders are associated with ventricular arrhythmias in patients with an implantable cardioverter-defibrillator. *Chest* 2002; 122(2):558-561.
- (14) Fietze I, Rottig J, Quispe-Bravo S, Riedel F, Witte J, Baumann G et al. Sleep apnea syndrome in patients with cardiac pacemaker. *Respiration* 2000; 67(3):268-271.
- (15) Tremel F, Pepin JL, Veale D, Wuyam B, Siche JP, Mallion JM et al. High prevalence and persistence of sleep apnoea in patients referred for acute left ventricular failure and medically treated over 2 months. *Eur Heart J* 1999; 20(16):1201-1209.
- (16) Peker Y, Hedner J, Norum J, Kraiczi H, Carlson J. Increased incidence of cardiovascular disease in middle-aged men with obstructive sleep apnea: a 7-year follow-up. *Am J Respir Crit Care Med* 2002; 166(2):159-165.
- (17) Andreas S, Schulz R, Haro W, Werner GS, Kreuzer H. [Sleep-related breathing disorders in patients with coronary heart disease]. *Dtsch Med Wochenschr* 1995; 120(45):1533-1537.
- (18) Andreas S, Schulz R, Werner GS, Kreuzer H. Prevalence of obstructive sleep apnoea in patients with coronary artery disease. *Coron Artery Dis* 1996; 7(7):541-545.
- (19) Peker Y, Hedner J, Kraiczi H, Loth S. Respiratory disturbance index: an independent predictor of mortality in coronary artery disease. *Am J Respir Crit Care Med* 2000; 162(1):81-86.
- (20) Koehler U, Schafer H. Is obstructive sleep apnea (OSA) a risk factor for myocardial infarction and cardiac arrhythmias in patients with coronary heart disease (CHD)? *Sleep* 1996; 19(4):283-286.
- (21) Kanagala R, Murali NS, Friedman PA, Ammash NM, Gersh BJ, Ballman KV et al. Obstructive sleep apnea and the recurrence of atrial fibrillation. *Circulation* 2003; 107(20):2589-2594.
- (22) Becker HF, Jerrentrup A, Ploch T, Grote L, Penzel T, Sullivan CE et al. Effect of nasal continuous positive airway pressure treatment on blood pressure in patients with obstructive sleep apnea. *Circulation* 2003; 107(1):68-73.
- (23) Pepperell JC, Ramdassingh-Dow S, Crosthwaite N, Mullins R, Jenkinson C, Stradling JR et al. Ambulatory blood pressure after therapeutic and subtherapeutic nasal continuous positive airway pressure for obstructive sleep apnoea: a randomised parallel trial. *Lancet* 2002; 359(9302):204-210.
- (24) Teramoto S, Kume H, Matsuse T. Ambulatory blood pressure after sleep apnoea treatment. *Lancet* 2002; 360(9329):341.

- (25) Malone S, Liu PP, Holloway R, Rutherford R, Xie A, Bradley TD. Obstructive sleep apnoea in patients with dilated cardiomyopathy: effects of continuous positive airway pressure. *Lancet* 1991; 338(8781):1480-1484.
- (26) Sin DD, Logan AG, Fitzgerald FS, Liu PP, Bradley TD. Effects of continuous positive airway pressure on cardiovascular outcomes in heart failure patients with and without Cheyne-Stokes respiration. *Circulation* 2000; 102(1):61-66.
- (27) Marin JM, Carrizo SJ, Vicente E, Agusti AG. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. *Lancet* 2005; 365(9464):1046-1053.
- (28) Ip MS, Lam B, Tang LC, Lauder IJ, Ip TY, Lam WK. A community study of sleep-disordered breathing in middle-aged Chinese women in Hong Kong: prevalence and gender differences. *Chest* 2004; 125(1):127-134.
- (29) Parra O, Garcia-Escasans N, Montserrat JM, Garcia EL, Ruiz J, Lopez JA et al. Should patients with sleep apnoea/hypopnoea syndrome be diagnosed and managed on the basis of home sleep studies? *Eur Respir J* 1997; 10(8):1720-1724.
- (30) Redline S, Tosteson T, Boucher MA, Millman RP. Measurement of sleep-related breathing disturbances in epidemiologic studies. Assessment of the validity and reproducibility of a portable monitoring device. *Chest* 1991; 100(5):1281-1286.
- (31) Kashyap R, Hock LM, Bowman TJ. Higher prevalence of smoking in patients diagnosed as having obstructive sleep apnea. *Sleep Breath* 2001; 5(4):167-172.
- (32) Pack AI. Obstructive sleep apnea. *Adv Intern Med* 1994; 39:517-567.
- (33) Bresnitz EA, Goldberg R, Kosinski RM. Epidemiology of obstructive sleep apnea. *Epidemiol Rev* 1994; 16(2):210-227.
- (34) Davies RJ, Stradling JR. The epidemiology of sleep apnoea. *Thorax* 1996; 51 Suppl 2:S65-S70.
- (35) Ancoli-Israel S, Klauber MR, Kripke DF, Parker L, Cobarrubias M. Sleep apnea in female patients in a nursing home. Increased risk of mortality. *Chest* 1989; 96(5):1054-1058.
- (36) Lloberes P, Levy G, Descals C, Sampol G, Roca A, Sagales T et al. Self-reported sleepiness while driving as a risk factor for traffic accidents in patients with obstructive sleep apnoea syndrome and in non-apnoeic snorers. *Respir Med* 2000; 94(10):971-976.
- (37) Teran-Santos J, Jimenez-Gomez A, Cordero-Guevara J. The association between sleep apnea and the risk of traffic accidents. Cooperative Group Burgos-Santander. *N Engl J Med* 1999; 340(11):847-851.
- (38) Yee B, Campbell A, Beasley R, Neill A. Sleep disorders: a potential role in New Zealand motor vehicle accidents. *Intern Med J* 2002; 32(7):297-304.

- (39) Redline S, Young T. Epidemiology and natural history of obstructive sleep apnea. *Ear Nose Throat J* 1993; 72(1):20-26.
- (40) Dingli K, Coleman EL, Vennelle M, Finch SP, Wraith PK, Mackay TW et al. Evaluation of a portable device for diagnosing the sleep apnoea/hypopnoea syndrome. *Eur Respir J* 2003; 21(2):253-259.
- (41) Cohrs S, Hajak G. Validierungsstudie zum ApnoeScreen I-Gerät bei Patienten mit dem Verdacht auf eine schlafgebundene Atmungsstörung. 1-2-0002.

Ref Type: Generic

- (42) Teschler H, Stampa JW, Farhat AA, Wessendorf TE, Konietzko N. [Comparison of the efficacy of the BiPAP-S/T and the VPAP-S/T ventilators in patients with stable chronic respiratory insufficiency]. *Pneumologie* 1998; 52(6):305-310.
- (43) Brownell LG, West P, Sweatman P, Acres JC, Kryger MH. Protriptyline in obstructive sleep apnea: a double-blind trial. *N Engl J Med* 1982; 307(17):1037-1042.
- (44) Brownell LG, Perez-Padilla R, West P, Kryger MH. The role of protriptyline in obstructive sleep apnea. *Bull Eur Physiopathol Respir* 1983; 19(6):621-624.
- (45) Smith PL, Haponik EF, Allen RP, Bleecker ER. The effects of protriptyline in sleep-disordered breathing. *Am Rev Respir Dis* 1983; 127(1):8-13.
- (46) Hein H, Kirsten D, Jugert C, Magnussen H. [Theophylline in therapy of obstructive sleep apnea?]. *Pneumologie* 1993; 47 Suppl 4:750-753.
- (47) Terra SG, Oberg KC. Medroxyprogesterone acetate in the treatment of obstructive sleep apnea. *Ann Pharmacother* 1997; 31(6):776-778.
- (48) White DP, Zwillich CW, Pickett CK, Douglas NJ, Findley LJ, Weil JV. Central sleep apnea. Improvement with acetazolamide therapy. *Arch Intern Med* 1982; 142(10):1816-1819.
- (49) Winkels W, Klutmann M, Bias R, Wessolowski T, Simon H. [Effectiveness of an anti-snoring prosthesis in obstructive sleep apnea syndrome]. *Pneumologie* 1997; 51 Suppl 3:802-803.
- (50) Henke KG, Frantz DE, Kuna ST. An oral elastic mandibular advancement device for obstructive sleep apnea. *Am J Respir Crit Care Med* 2000; 161(2 Pt 1):420-425.
- (51) Yoshida K. Oral device therapy for the upper airway resistance syndrome patient. *J Prosthet Dent* 2002; 87(4):427-430.
- (52) Sakakibara H, Umemoto M, Kuwahara M, Suetsugu S. [Treatment of obstructive sleep apnea syndrome with a mandibular positioning device and other nonsurgical modalities]. *Nihon Kyobu Shikkan Gakkai Zasshi* 1995; 33 Suppl:76-84.

- (53) Friedman M, Ibrahim H, Lee G, Joseph NJ. Combined uvulopalatopharyngoplasty and radiofrequency tongue base reduction for treatment of obstructive sleep apnea/hypopnea syndrome. *Otolaryngol Head Neck Surg* 2003; 129(6):611-621.
- (54) Osnes T, Rollheim J, Hartmann E. Effect of UPPP with respect to site of pharyngeal obstruction in sleep apnoea: follow-up at 18 months by overnight recording of airway pressure and flow. *Clin Otolaryngol* 2002; 27(1):38-43.
- (55) Sullivan CE, Issa FG, Berthon-Jones M, Eves L. Reversal of obstructive sleep apnoea by continuous positive airway pressure applied through the nares. *Lancet* 1981; 1(8225):862-865.
- (56) Sin DD, Mayers I, Man GC, Pawluk L. Long-term compliance rates to continuous positive airway pressure in obstructive sleep apnea: a population-based study. *Chest* 2002; 121(2):430-435.
- (57) Nunez C, Gallagher D, Visser M, Pi-Sunyer FX, Wang Z, Heymsfield SB. Bioimpedance analysis: evaluation of leg-to-leg system based on pressure contact footpad electrodes. *Med Sci Sports Exerc* 1997; 29(4):524-531.
- (58) Sageman WS, Riffenburgh RH, Spiess BD. Equivalence of bioimpedance and thermodilution in measuring cardiac index after cardiac surgery. *J Cardiothorac Vasc Anesth* 2002; 16(1):8-14.
- (59) Spiess BD, Patel MA, Soltow LO, Wright IH. Comparison of bioimpedance versus thermodilution cardiac output during cardiac surgery: evaluation of a second-generation bioimpedance device. *J Cardiothorac Vasc Anesth* 2001; 15(5):567-573.
- (60) Hahn G, Sipinkova I, Baisch F, Hellige G. Changes in the thoracic impedance distribution under different ventilatory conditions. *Physiol Meas* 1995; 16(3 Suppl A):A161-A173.
- (61) Kunst PW, Vazquez dA, Bohm SH, Faes TJ, Lachmann B, Postmus PE et al. Monitoring of recruitment and derecruitment by electrical impedance tomography in a model of acute lung injury. *Crit Care Med* 2000; 28(12):3891-3895.
- (62) Frerichs I, Hinz J, Herrmann P, Weisser G, Hahn G, Dudykevych T et al. Detection of local lung air content by electrical impedance tomography compared with electron beam CT. *J Appl Physiol* 2002; 93(2):660-666.
- (63) Mond H, Strathmore N, Kertes P, Hunt D, Baker G. Rate responsive pacing using a minute ventilation sensor. *Pacing Clin Electrophysiol* 1988; 11(11 Pt 2):1866-1874.
- (64) Simon R, Ni Q, Willems R, Hartley JW, Daum DR, Lang D et al. Comparison of impedance minute ventilation and direct measured minute ventilation in a rate adaptive pacemaker. *Pacing Clin Electrophysiol* 2003; 26(11):2127-2133.

- (65) Edgar DR, Horwood JM, Woollons DJ, Baker J, Shaw DB. Development and preliminary clinical tests of an impedance sensing VDD recording pacemaker for diagnosis and research. *Med Biol Eng Comput* 1999; 37(4):517-522.
- (66) Ruhsam C, Pfutzner H, Nopp P, Nakesch H, Frais-Kolbl H, Marktl W. Time-frequency spectral analysis of electric field plethysmography signals. *Med Prog Technol* 1995; 21(1):17-28.
- (67) Saarelainen S, Himanen SL, Hasan J, Virkkala J, Koobi T. Whole-body impedance recording--a practical method for the diagnosis of sleep apnoea. *Clin Physiol Funct Imaging* 2003; 23(2):110-113.
- (68) Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep* 1991; 14(6):540-545.
- (69) Johns MW. Daytime sleepiness, snoring, and obstructive sleep apnea. The Epworth Sleepiness Scale. *Chest* 1993; 103(1):30-36.
- (70) Lown B, Wolf M. Approaches to sudden death from coronary heart disease. *Circulation* 1971; 44(1):130-142.
- (71) Nakesch H, Pfutzner H, Ruhsam C, Nopp P, Futschik K. Five-electrode field plethysmography technique for separation of respiration and cardiac signals. *Med Biol Eng Comput* 1994; 32(4 Suppl):S65-S70.
- (72) Pfutzner H, Futschik K, Doblender A, Schenz G, Zwick H. A method for the elimination of artefacts in electric field plethysmography of the lung. *Front Med Biol Eng* 1990; 2(1):53-63.
- (73) Koehler U, Schafer H. Is obstructive sleep apnea (OSA) a risk factor for myocardial infarction and cardiac arrhythmias in patients with coronary heart disease (CHD)? *Sleep* 1996; 19(4):283-286.
- (74) Scharf C, Cho YK, Bloch KE, Brunckhorst C, Duru F, Balaban K et al. Diagnosis of sleep-related breathing disorders by visual analysis of transthoracic impedance signals in pacemakers. *Circulation* 2004; 110(17):2562-2567.
- (75) Defaye P, Pepin JL, Poezevara Y, Mabo P, Murgatroyd F, Levy P et al. Automatic recognition of abnormal respiratory events during sleep by a pacemaker transthoracic impedance sensor. *J Cardiovasc Electrophysiol* 2004; 15(9):1034-1040.
- (76) Yasuda Y, Umezu A, Horihata S, Yamamoto K, Miki R, Koike S. Modified thoracic impedance plethysmography to monitor sleep apnea syndromes. *Sleep Med* 2005; 6(3):215-224.
- (77) Bar A, Pillar G, Dvir I, Sheffy J, Schnall RP, Lavie P. Evaluation of a portable device based on peripheral arterial tone for unattended home sleep studies. *Chest* 2003; 123(3):695-703.
- (78) Stein PK, Duntley SP, Domitrovich PP, Nishith P, Carney RM. A simple method to identify sleep apnea using Holter recordings. *J Cardiovasc Electrophysiol* 2003; 14(5):467-473.

- (79) Suzuki R, Akashiba T, Saito O, Horie T. [Risk factors in development of sleep apnea syndrome--a large population study using a simple monitoring system]. *Nihon Kokyuki Gakkai Zasshi* 2002; 40(8):653-659.
- (80) Zhang Q, He QY, Du QY, Pang GF, Zhao LS, Wu RQ et al. [Epidemiologic study on sleep apnea-hypopnea syndrome by home investigation in Chengde city]. *Zhonghua Jie He He Hu Xi Za Zhi* 2003; 26(5):273-275.
- (81) Taasan VC, Block AJ, Boysen PG, Wynne JW. Alcohol increases sleep apnea and oxygen desaturation in asymptomatic men. *Am J Med* 1981; 71(2):240-245.
- (82) Kohnlein T, Welte T, Tan LB, Elliott MW. Central sleep apnoea syndrome in patients with chronic heart disease: a critical review of the current literature. *Thorax* 2002; 57(6):547-554.
- (83) Mansfield DR, Gollogly NC, Kaye DM, Richardson M, Bergin P, Naughton MT. Controlled trial of continuous positive airway pressure in obstructive sleep apnea and heart failure. *Am J Respir Crit Care Med* 2004; 169(3):361-366.
- (84) Ogawa A, Iwase T, Yamamoto T, Nishiyama S, Narui K, Momomura S. Improvement of cheyne-stokes respiration, central sleep apnea and congestive heart failure by noninvasive bilevel positive pressure and medical treatment. *Circ J* 2004; 68(9):878-882.
- (85) Koehler U, Schafer H. Is obstructive sleep apnea (OSA) a risk factor for myocardial infarction and cardiac arrhythmias in patients with coronary heart disease (CHD)? *Sleep* 1996; 19(4):283-286.
- (86) Bloch KE, Schoch OD, Zhang JN, Russi EW. German version of the Epworth Sleepiness Scale. *Respiration* 1999; 66(5):440-447.
- (87) Hamm M, Krause J, Felsmann M, Barnstorf D, Kothe R, Fabel H. [A computerized processing unit for ambulatory diagnosis of sleep apnea and nocturnal hypoxemia]. *Pneumologie* 1990; 44 Suppl 1:627-628.
- (88) Norman RG, Ahmed MM, Walsleben JA, Rapoport DM. Detection of respiratory events during NPSG: nasal cannula/pressure sensor versus thermistor. *Sleep* 1997; 20(12):1175-1184.