

References

- Abraham, W. C., Mason, S. E., Demmer, J., Williams, J. M., Richardson, C. L., Tate, W. P., et al. (1993). Correlations between immediate early gene induction and the persistence of long-term potentiation. *Neuroscience*, 56(3), 717-727.
- Angel, P., & Karin, M. (1991). The role of Jun, Fos and the AP-1 complex in cell-proliferation and transformation. *Biochim Biophys Acta*, 1072(2-3), 129-157.
- Barroso-Chinea, P., Aymerich, M. S., Castle, M. M., Perez-Manso, M., Tunon, T., Erro, E., et al. (2007). Detection of two different mRNAs in a single section by dual in situ hybridization: a comparison between colorimetric and fluorescent detection. *J Neurosci Methods*, 162(1-2), 119-128.
- Barroso-Chinea, P., Castle, M., Aymerich, M. S., Perez-Manso, M., Erro, E., Tunon, T., et al. (2007). Expression of the mRNAs encoding for the vesicular glutamate transporters 1 and 2 in the rat thalamus. *J Comp Neurol*, 501(5), 703-715.
- Beauchamp, G. K., & Yamazaki, K. (2003). Chemical signalling in mice. *Biochem Soc Trans*, 31(Pt 1), 147-151.
- Belluscio, L., Koentges, G., Axel, R., & Dulac, C. (1999). A map of pheromone receptor activation in the mammalian brain 209-220. *Cell*, 97, 209-220.
- Boehm, T., & Zufall, F. (2006). MHC peptides and the sensory evaluation of genotype. *Trends Neurosci*, 29(2), 100-107.
- Brennan, P., Kaba, H., & Keverne, E. B. (1990). Olfactory recognition: a simple memory system. *Science*, 250(4985), 1223-1226.
- Brennan, P. A., & Keverne, E. B. (1997). Neural mechanisms of mammalian olfactory learning. *Prog Neurobiol*, 51(4), 457-481.

- Cahill, L., & McGaugh, J. L. (1996). Modulation of memory storage. *Curr Opin Neurobiol*, 6(2), 237-242.
- Chiasson, B. J., Dennison, Z., & Robertson, H. A. (1995). Short communication Amygdala kindling and immediate-early genes. *Molecular Brain Research*, 29, 191-199.
- Chowdhury, S., Shepherd, J. D., Okuno, H., Lyford, G., Petralia, R. S., Plath, N., et al. (2006). Arc/Arg3.1 interacts with the endocytic machinery to regulate AMPA receptor trafficking. *Neuron*, 52(3), 445-459.
- Cole, A. J., Saffen, D. W., Baraban, J. M., & Worley, P. F. (1989). Rapid increase of an immediate early gene messenger RNA in hippocampal neurons by synaptic NMDA receptor activation. *Nature*, 340(6233), 474-476.
- Coolen, L. M., & Wood, R. I. (1998). Bidirectional connections of the medial amygdaloid nucleus in the Syrian hamster brain: simultaneous anterograde and retrograde tract tracing. *J Comp Neurol*, 399(2), 189-209.
- Davis, S., Bozon, B., & Laroche, S. (2003). How necessary is the activation of the immediate early gene zif268 in synaptic plasticity and learning? *Behav Brain Res*, 142(1-2), 17-30.
- De Olmos, J. S., Beltramo, C. A., & Alheid., G. F. (2004). Amygdala and extended amygdala of the rat: a cytoarchitectonical, fibroarchitectonical, and chemoarchitectonical survey. In G. Paxinos (Ed.), *The Rat Nervous System* (pp. 509-603). Amsterdam: Elsevier Academic Press.
- de Vries, G. J., & Miller, M. A. (1998). Anatomy and function of extrahypothalamic vasopressin systems in the brain. *Prog Brain Res*, 119, 3-20.
- Dickinson, J. A., Amato, S. F., McManus, B. J., & Chiles, T. C. (1995). Membrane immunoglobulin receptor cross-linking induces fosB mRNA expression in mature B

lymphocytes: assembly of distinct FosB-containing nucleoprotein complexes during B cell stimulation. *Cell Immunol*, 165(1), 92-100.

Dluzen, D. E., Muraoka, S., Engelmann, M., Ebner, K., & Landgraf, R. (2000). Oxytocin induces preservation of social recognition in male rats by activating alpha-adrenoceptors of the olfactory bulb. *Eur J Neurosci*, 12(2), 760-766.

Dluzen, D. E., Muraoka, S., Engelmann, M., & Landgraf, R. (1998). The effects of infusion of arginine vasopressin, oxytocin, or their antagonists into the olfactory bulb upon social recognition responses in male rats. *Peptides*, 19(6), 999-1005.

Dong, H. W., Petrovich, G. D., & Swanson, L. W. (2001). Topography of projections from amygdala to bed nuclei of the stria terminalis. *Brain Res Brain Res Rev*, 38(1-2), 192-246.

Doving, K. B., & Trotier, D. (1998). Structure and function of the vomeronasal organ. *J. Exp. Biol.*, 201((Pt. 21)), 2913- 2925.

Dudley, C. A., & Moss, R. L. (1999). Activation of an anatomical distinct subpopulation of accessory olfactory bulb neurons by chemosensory stimulation. *Neuroscience*, 91(4), 1549–1556.

Dulac, C., & Axel, R. (1995). A novel family of genes encoding putative pheromone receptors in mammals. *Cell*, 83, 195-206.

Engelmann, M., Wotjak, C. T., & Landgraf, R. (1995). Social discrimination procedure: an alternative method to investigate juvenile recognition abilities in rats. *Physiol Behav*, 58(2), 315-321.

Esclapez, M., Tillakaratne, N. J., Kaufman, D. L., Tobin, A. J., & Houser, C. R. (1994). Comparative localization of two forms of glutamic acid decarboxylase and their mRNAs in rat brain supports the concept of functional differences between the forms. *J Neurosci*, 14(3 Pt 2), 1834-1855.

- Esclapez, M., Tillakaratne, N. J., Tobin, A. J., & Houser, C. R. (1993). Comparative localization of mRNAs encoding two forms of glutamic acid decarboxylase with nonradioactive in situ hybridization methods. *J Comp Neurol*, 331(3), 339-362.
- Feldblum, S., Erlander, M. G., & Tobin, A. J. (1993). Different distributions of GAD65 and GAD67 mRNAs suggest that the two glutamate decarboxylases play distinctive functional roles. *J Neurosci Res*, 34(6), 689-706.
- Ferguson, J. N., Aldag, J. M., Insel, T. R., & Young, L. J. (2001). Oxytocin in the medial amygdala is essential for social recognition in the mouse. *J Neurosci*, 21(20), 8278-8285.
- Ferguson, J. N., Young, L. J., Hearn, E. F., Matzuk, M. M., Insel, T. R., & Winslow, J. T. (2000). Social amnesia in mice lacking the oxytocin gene. *Nat Genet*, 25(3), 284-288.
- Finkel, M. P., Biskis, B. O., & Jenkins, P. B. (1966). Virus induction of osteosarcomas in mice. *Science*, 151(711), 698-701.
- Fuller, T. A., & Price, J. L. (1988). Putative glutamatergic and/or aspartatergic cells in the main and accessory olfactory bulbs of the rat. *J Comp Neurol*, 276(2), 209-218.
- Gammie, S. C., & Nelson, R. J. (2001). cFOS and pCREB activation and maternal aggression in mice. *Brain Res*, 898(2), 232-241.
- Gong, L. B., He, L., Liu, Y., Chen, X. Q., & Jiang, B. (2005). Expression of early growth response factor-1 in rats with cerulein-induced acute pancreatitis and its significance. *World J Gastroenterol*, 11(32), 5022-5024.
- Gonzalez, G. A., & Montminy, M. R. (1989). Cyclic AMP stimulates somatostatin gene transcription by phosphorylation of CREB at serine 133. *Cell*, 59(4), 675-680.
- Grimm, R., & Tischmeyer, W. (1997). Complex patterns of immediate early gene induction in rat brain following brightness discrimination training and pseudotraining. *Behav Brain Res*, 84(1-2), 109-116.

- Guillamon, A., & Segovia, S. (1997). Sex differences in the vomeronasal system. *Brain Res Bull*, 44(4), 377-382.
- Halem, H. A., Cherry, J. A., & Baum, M. J. (1999). Vomeronasal neuroepithelium and forebrain Fos responses to male pheromones in male and female mice. *J Neurobiol*, 39(2), 249-263.
- Hall, J., Thomas, K. L., & Everitt, B. J. (2000). Rapid and selective induction of BDNF expression in the hippocampus during contextual learning. *Nat Neurosci*, 3(6), 533-535.
- Halpern, M. (1987). The organization and function of the vomeronasal system. *Annu Rev Neurosci*, 10, 325-362.
- Halpern, M., Shapiro, L. S., & Jia, C. (1995). Differential localization of G proteins in the opossum vomeronasal system. *Brain Res*, 677(1), 157-161.
- Hayashi, Y., Momiyama, A., Takahashi, T., Ohishi, H., Ogawa-Meguro, R., Shigemoto, R., et al. (1993). Role of a metabotropic glutamate receptor in synaptic modulation in the accessory olfactory bulb. *Nature*, 366(6456), 687-690.
- Hermel, E. E., Faccioni-Heuser, M. C., Marcuzzo, S., Rasia-Filho, A. A., & Achaval, M. (2006). Ultrastructural features of neurons and synaptic contacts in the posterodorsal medial amygdala of adult male rats. *J Anat*, 208(5), 565-575.
- Herzog, E., Takamori, S., Jahn, R., Brose, N., & Wojcik, S. M. (2006). Synaptic and vesicular co-localization of the glutamate transporters VGLUT1 and VGLUT2 in the mouse hippocampus. *J Neurochem*, 99(3), 1011-1018.
- Hisano, S., Hoshi, K., Ikeda, Y., Maruyama, D., Kanemoto, M., Ichijo, H., et al. (2000). Regional expression of a gene encoding a neuron-specific Na⁽⁺⁾-dependent inorganic phosphate cotransporter (DNPI) in the rat forebrain. *Brain Res Mol Brain Res*, 83(1-2), 34-43.

- Hollander, E., Bartz, J., Chaplin, W., Phillips, A., Sumner, J., Soorya, L., et al. (2007). Oxytocin increases retention of social cognition in autism. *Biol Psychiatry*, 61(4), 498-503.
- Hunter, T., & Cooper, J. A. (1985). Protein-tyrosine kinases. *Annu Rev Biochem*, 54, 897-930.
- Imamura, K., Mori, K., Fujita, S. C., & Obata, K. (1985). Immunochemical identification of subgroups of vomeronasal nerve fibers and their segregated terminations in the accessory olfactory bulb. *Brain Res*, 328(2), 362-366.
- Inamura, K., Kashiwayanagi, M., & Kurihara, K. (1999). Regionalization of Fos immunostaining in rat accessory olfactory bulb when the vomeronasal organ was exposed to urine. *Eur J Neurosci*, 11(7), 2254-2260.
- Inoue, Y., Udo, H., Inokuchi, K., & Sugiyama, H. (2007). Homer1a regulates the activity-induced remodeling of synaptic structures in cultured hippocampal neurons. *Neuroscience*, 150(4), 841-852.
- Jansen, L. M., Gispen-de Wied, C. C., Wiegant, V. M., Westenberg, H. G., Lahuis, B. E., & van Engeland, H. (2006). Autonomic and neuroendocrine responses to a psychosocial stressor in adults with autistic spectrum disorder. *J Autism Dev Disord*, 36(7), 891-899.
- Jia, C., & Halpern, M. (1996). Subclasses of vomeronasal receptor neurons: differential expression of G proteins (Gi alpha 2 and G(o alpha)) and segregated projections to the accessory olfactory bulb. *Brain Res*, 719(1-2), 117-128.
- Jia, C., & Halpern, M. (1997). Segregated populations of mitral/tufted cells in the accessory olfactory bulb. *NeuroReport*, 8, 1887-1890.
- Johnston, R. E., & Rasmussen, K. (1984). Individual recognition of female hamsters by males: role of chemical cues and of the olfactory and vomeronasal systems. *Physiol Behav*, 33(1), 95-104.

- Kammermeier, P. J., Xiao, B., Tu, J. C., Worley, P. F., & Ikeda, S. R. (2000). Homer proteins regulate coupling of group I metabotropic glutamate receptors to N-type calcium and M-type potassium channels. *J Neurosci*, 20(19), 7238-7245.
- Kaneko, T., & Fujiyama, F. (2002). Complementary distribution of vesicular glutamate transporters in the central nervous system. *Neurosci Res*, 42(4), 243-250.
- Kaufman, D. L., Houser, C. R., & Tobin, A. J. (1991). Two forms of the gamma-aminobutyric acid synthetic enzyme glutamate decarboxylase have distinct intraneuronal distributions and cofactor interactions. *J Neurochem*, 56(2), 720-723.
- Kemble, E. D., Blanchard, D. C., Blanchard, R. J., & Takushi, R. (1984). Taming wild rats following medial amygdaloid lesions. *Physiol. Behav.*, 32, 131-134.
- Kendrick, K. M., Da Costa, A. P., Broad, K. D., Ohkura, S., Guevara, R., Levy, F., et al. (1997). Neural control of maternal behaviour and olfactory recognition of offspring. *Brain Res Bull*, 44(4), 383-395.
- Keverne, E. B. (1999). The vomeronasal organ. *Science*, 286(5440), 716-720.
- Keverne, E. B., & Curley, J. P. (2004). Vasopressin, oxytocin and social behaviour. *Curr Opin Neurobiol*, 14(6), 777-783.
- Kevetter, G. A., & Winans, S. S. (1981a). Connections of the corticomedial amygdala in the golden hamster. I. Efferents of the "vomeronasal amygdala". *J Comp Neurol*, 197(1), 81-98.
- Kevetter, G. A., & Winans, S. S. (1981b). Connections of the corticomedial amygdala in the golden hamster. II. Efferents of the "olfactory amygdala". *J Comp Neurol*, 197(1), 99-111.
- Kosfeld, M., Heinrichs, M., Zak, P. J., Fischbacher, U., & Fehr, E. (2005). Oxytocin increases trust in humans. *Nature*, 435(7042), 673-676.

- Kruijer, W., Schubert, D., & Verma, I. M. (1985). Induction of the proto-oncogene fos by nerve growth factor. *Proc Natl Acad Sci U S A*, 82(21), 7330-7334.
- Kumar, A., Dudley, C. A., & Moss, R. L. (1999). Functional dichotomy within the vomeronasal system: distinct zones of neuronal activity in the accessory olfactory bulb correlate with sex-specific behaviors. *J Neurosci*, 19(20), RC32.
- Lai, W. S., Chen, A., & Johnston, R. E. (2004). Patterns of neural activation associated with exposure to odors from a familiar winner in male golden hamsters. *Horm Behav*, 46(3), 319-329.
- Leinders-Zufall, T., Brennan, P., Widmayer, P., S, P. C., Maul-Pavicic, A., Jager, M., et al. (2004). MHC class I peptides as chemosensory signals in the vomeronasal organ. *Science*, 306(5698), 1033-1037.
- Levai, O., Feistel, T., Breer, H., & Strotmann, J. (2006). Cells in the vomeronasal organ express odorant receptors but project to the accessory olfactory bulb. *J Comp Neurol*, 498(4), 476-490.
- Licht, G., & Meredith, M. (1987). Convergence of main and accessory olfactory pathways onto single neurons in the hamster amygdala. *Exp Brain Res*, 69(1), 7-18.
- Loscher, W., Schirmer, M., Freichel, C., & Gernert, M. (2006). Distribution of GABAergic neurons in the striatum of amygdala-kindled rats: an immunohistochemical and in situ hybridization study. *Brain Res*, 1083(1), 50-60.
- Luo, M., Fee, M. S., & Katz, L. C. (2003). Encoding pheromonal signals in the accessory olfactory bulb of behaving mice. *Science*, 299(5610), 1196-1201.
- Luo, M., & Katz, L. C. (2004). Encoding pheromonal signals in the mammalian vomeronasal system. *Curr Opin Neurobiol*, 14(4), 428-434.
- Lyford, G. L., Yamagata, K., Kaufmann, W. E., Barnes, C. A., Sanders, L. K., Copeland, N. G., et al. (1995). Arc, a growth factor and activity-regulated gene, encodes a novel

cytoskeleton-associated protein that is enriched in neuronal dendrites. *Neuron*, 14(2), 433-445.

Maaswinkel, H., Baars, A. M., Gispen, W. H., & Spruijt, B. M. (1996). Roles of the basolateral amygdala and hippocampus in social recognition in rats. *Physiol Behav*, 60(1), 55-63.

Maki, Y., Bos, T. J., Davis, C., Starbuck, M., & Vogt, P. K. (1987). Avian sarcoma virus 17 carries the jun oncogene. *Proc Natl Acad Sci U S A*, 84(9), 2848-2852.

Martinez-Marcos, A., & Halpern, M. (1999). Differential centrifugal afferents to the anterior and posterior accessory olfactory bulb. *NeuroReport*, 10, 2011-2015.

Matsunami, H., & Buck, L. B. (1997). A multigene family encoding a diverse array of putative pheromone receptors in mammals. *Cell*, 90, 775-784.

Matsuoka, M., Kaba, H., Mori, Y., & Ichikawa, M. (1997). Synaptic plasticity in olfactory memory formation in female mice. *Neuroreport*, 8(11), 2501-2504.

Matsuoka, M., Mori, Y., Hoshino, K., & Ichikawa, M. (1994). Social environment affects synaptic structure in the glomerulus of the accessory olfactory bulb of the hamster. *Neurosci Res*, 19(2), 187-193.

Matsuoka, M., Mori, Y., & Ichikawa, M. (1998). Morphological changes of synapses induced by urinary stimulation in the hamster accessory olfactory bulb. *Synapse*, 28(2), 160-166.

Matsuoka, M., Yamagata, K., Sugiura, H., Yoshida-Matsuoka, J., Norita, M., & Ichikawa, M. (2002). Expression and regulation of the immediate-early gene product Arc in the accessory olfactory bulb after mating in male rat. *Neuroscience*, 111(2), 251-258.

Matsuoka, M., Yoshida-Matsuoka, J., Yamagata, K., Sugiura, H., Ichikawa, M., & Norita, M. (2003). Rapid induction of Arc is observed in the granule cell dendrites in the accessory olfactory bulb after mating. *Brain Res*, 975(1-2), 189-195.

- Meisami, E. (1989). A proposed relationship between increases in the number of olfactory receptor neurons, convergence ratio and sensitivity in the developing rat. *Brain Res Dev Brain Res*, 46(1), 9-19.
- Mercugliano, M., Soghomonian, J. J., Qin, Y., Nguyen, H. Q., Feldblum, S., Erlander, M. G., et al. (1992). Comparative distribution of messenger RNAs encoding glutamic acid decarboxylases (Mr 65,000 and Mr 67,000) in the basal ganglia of the rat. *J Comp Neurol*, 318(3), 245-254.
- Meredith, M. (1998). Vomeronasal, olfactory, hormonal convergence in the brain. Cooperation or coincidence? *Ann NY Acad Sci*, 855, 349-361.
- Meredith, M., & Westberry, J. M. (2004). Distinctive responses in the medial amygdala to same-species and different-species pheromones. *J Neurosci*, 24(25), 5719-5725.
- Modahl, C., Green, L., Fein, D., Morris, M., Waterhouse, L., Feinstein, C., et al. (1998). Plasma oxytocin levels in autistic children. *Biol Psychiatry*, 43(4), 270-277.
- Mohedano-Moriano, A., Pro-Sistiaga, P., Ubeda-Banon, I., Crespo, C., Insausti, R., & Martinez-Marcos, A. (2007). Segregated pathways to the vomeronasal amygdala: differential projections from the anterior and posterior divisions of the accessory olfactory bulb. *Eur J Neurosci*, 25(7), 2065-2080.
- Mombaerts, P. (1999). Molecular biology of odorant receptors in vertebrates. *Annu Rev Neurosci*, 22, 487-509.
- Moore, A. J., Brodie, E. D., & Wolf, J. B. (1997). Interacting phenotypes and the evolutionary process .1. Direct and indirect genetic effects of social interactions. *Evolution*, 51(5), 1352-1362.
- Mucignat-Caretta, C., Colivicchi, M. A., Fattori, M., Ballini, C., Bianchi, L., Gabai, G., et al. (2006). Species-specific chemosignals evoke delayed excitation of the vomeronasal amygdala in freely-moving female rats. *J Neurochem*, 99(3), 881-891.

- Mumby, D. G., & Pinel, J. P. (1994). Rhinal cortex lesions and object recognition in rats. *Behav. Neurosci.*, 108, 11-18.
- Newman, S. W. (1999). The medial extended amygdala in male reproductive behavior. A node in the mammalian social behavior network. *Ann N Y Acad Sci*, 877, 242-257.
- Ngai, J., Chess, A., Dowling, M. M., Necles, N., Macagno, E. R., & Axel, R. (1993). Coding of olfactory information: topography of odorant receptor expression in the catfish olfactory epithelium. *Cell*, 72(5), 667-680.
- Nikolaev, E., Werka, T., & Kaczmarek, L. (1992). C-fos protooncogene expression in rat brain after long-term training of two-way active avoidance reaction. *Behav Brain Res*, 48(1), 91-94.
- Nishi, A., Snyder, G. L., & Greengard, P. (1997). Bidirectional regulation of DARPP-32 phosphorylation by dopamine. *J Neurosci*, 17(21), 8147-8155.
- Nitecka, L., & Ben-Ari, Y. (1987). Distribution of GABA-like immunoreactivity in the rat amygdaloid complex. *J Comp Neurol*, 266(1), 45-55.
- Norlin, E. M., Gussing, F., & Berghard, A. (2003). Vomeronasal phenotype and behavioral alterations in G alpha i2 mutant mice. *Curr Biol*, 13(14), 1214-1219.
- Panka, D. J., Mudgett-Hunter, M., Parks, D. R., Peterson, L. L., Herzenberg, L. A., Haber, E., et al. (1988). Variable region framework differences result in decreased or increased affinity of variant anti-digoxin antibodies. *Proc Natl Acad Sci U S A*, 85(9), 3080-3084.
- Pankevich, D. E., Cherry, J. A., & Baum, M. J. (2006a). Accessory olfactory neural Fos responses to a conditioned environment are blocked in male mice by vomeronasal organ removal. *Physiol Behav*, 87(4), 781-788.

- Pankevich, D. E., Cherry, J. A., & Baum, M. J. (2006b). Effect of vomeronasal organ removal from male mice on their preference for and neural Fos responses to female urinary odors. *Behav Neurosci*, 120(4), 925-936.
- Paxinos, G., & Watson, C. (1998). *Rat Brain Atlas (in stereotaxic coordination)* (Fourth ed.): Academic Press.
- Paxinos, G., & watson, C. (2004). *The Rat Nervous System*: Academic Press.
- Pearse-Pratt, R., Schellinck, H., Brown, R., Singh, P. B., & Roser, B. (1998). Soluble MHC antigens and olfactory recognition of genetic individuality: the mechanism. *Genetica*, 104(3), 223-230.
- Pedersen, C. A. (2004). Biological aspects of social bonding and the roots of human violence. *Ann N Y Acad Sci*, 1036, 106-127.
- Petrovich, G. D., Canteras, N. S., & Swanson, L. W. (2001). Combinatorial amygdala inputs to hippocampal domains and hypothalamic behavior systems. *Brain Res Brain Res Rev*, 38(1-2), 247-289.
- Pittman, J. Q., & Spencer, S. j. (2005). Neurohypophysial peptides: gatekeepers in the amygdala. *TRENDS in Endocrinology and Metabolism*, 16(8), 343-344.
- Polston, E. K., Heitz, M., Barnes, W., Cardamone, K., & Erskine, M. S. (2001). NMDA-Mediated Activation of the Medial Amygdala Initiates a Downstream Neuroendocrine Memory Responsible for Pseudopregnancy in the Female Rat. *The Journal of Neuroscience*, 21(11), 4104–4110.
- Popik, P., & van Ree, J. M. (1991). Oxytocin but not vasopressin facilitates social recognition following injection into the medial preoptic area of the rat brain. *Eur Neuropsychopharmacol*, 1(4), 555-560.

- Pro-Sistiaga, P., Mohedano-Moriano, A., Ubeda-Banon, I., Del Mar Arroyo-Jimenez, M., Marcos, P., Artacho-Perula, E., et al. (2007). Convergence of olfactory and vomeronasal projections in the rat basal telencephalon. *J Comp Neurol*, 504(4), 346-362.
- Restrepo, D., Arellano, J., Oliva, A. M., Schaefer, M. L., & Lin, W. (2004). Emerging views on the distinct but related roles of the main and accessory olfactory systems in responsiveness to chemosensory signals in mice. *Horm Behav*, 46(3), 247-256.
- Reti, I. M., Miskimon, M., Dickson, M., Petralia, R. S., Takamiya, K., Bland, R., et al. (2008). Activity-dependent secretion of neuronal activity regulated pentraxin from vasopressin neurons into the systemic circulation. *Neuroscience*, 151(2), 352-360.
- Richter, K., Wolf, G., & Engelmann, M. (2005). Social recognition memory requires two stages of protein synthesis in mice. *Learn Mem*, 12(4), 407-413.
- Rigby, P. W., Dieckmann, M., Rhodes, C., & Berg, P. (1977). Labeling deoxyribonucleic acid to high specific activity in vitro by nick translation with DNA polymerase I. *J Mol Biol*, 113(1), 237-251.
- Rodriguez, I. (2004). Pheromone receptors in mammals. *Hormones and Behavior*, 46, 219-230.
- Royer, J., C, S., F, J., & R, G. (1988). Morphometric study of the glomerular population in the mouse olfactory bulb: numerical density and size distribution along the rostrocaudal axis. *J Comp Neurology*, 270, 559-568.
- Scalia, F., & Winans, S. S. (1975). The differential projections of the olfactory bulb and accessory olfactory bulb in mammals. *J Comp Neurol*, 161(1), 31-55.
- Schaefer, M. L., Yamazaki, K., Osada, K., Restrepo, D., & Beauchamp, G. K. (2002). Olfactory fingerprints for major histocompatibility complex-determined body odors II: relationship among odor maps, genetics, odor composition, and behavior. *J. Neurosci*, 22, 9513-9521.

- Schaefer, M. L., Young, D. A., & Restrepo, D. (2001). Olfactory fingerprints for major histocompatibility complex-determined body odors. *J. Neurosci*, 21, 2481-2487.
- Shinohara, H., Kato, K., & Asano, T. (1992). Differential localization of G proteins, Gi and Go, in the accessory olfactory bulb of the rat. *J. Neuroscience*, 12, 1275-1279.
- Speel, E. J., Hopman, A. H., & Komminoth, P. (1999). Amplification methods to increase the sensitivity of in situ hybridization: play card(s). *J. Histochem. Cytochem.*, 47(3), 281-288.
- Steward, O., Wallace, C. S., Lyford, G. L., & Worley, P. F. (1998). Synaptic activation causes the mRNA for the IEG Arc to localize selectively near activated postsynaptic sites on dendrites. *Neuron*, 21(4), 741-751.
- Sugai, T., Sugitani, M., & Onoda, N. (1997). Subdivisions of the guinea-pig accessory olfactory bulb revealed by the combined method with immunohistochemistry, electrophysiological, and optical recordings. *Neuroscience*, 79(3), 871-885.
- Sugai, T., Sugitani, M., & Onoda, N. (2000). Novel subdivisions of the rat accessory olfactory bulb revealed by the combined method with lectin histochemistry, electrophysiological and optical recordings. *Neuroscience*, 95(1), 23-32.
- Swanson, L. W., & Petrovich, G. D. (1998). What is the amygdala? *Trends Neurosci*, 21(8), 323-331.
- Takami, S., & Graziadei, P. P. (1990). Morphological complexity of the glomerulus in the rat accessory olfactory bulb--a Golgi study. *Brain Res*, 510(2), 339-342.
- Takami, S., & Graziadei, P. P. (1992). The morphology of GABA-immunoreactive neurons in the accessory olfactory bulb of rats. *Brain Res*, 588, 317-323.
- Taziaux, M., Cornil, C. A., Dejace, C., Arckens, L., Ball, G. F., & Balthazart, J. (2006). Neuroanatomical specificity in the expression of the immediate early gene c-fos following expression of appetitive and consummatory male sexual behaviour in Japanese quail. *Eur J Neurosci*, 23(7), 1869-1887.

- Thor, D. H., & Holloway, W. R., Jr. (1982). Social Memory of the Male laboratory Rat. *J Comp Psychol*, 96(6), 1000-1006.
- Tischmeyer, W., & Grimm, R. (1999). Activation of immediate early genes and memory formation. *Cell Mol Life Sci*, 55(4), 564-574.
- Trombley, P. Q., & Westbrook, G. L. (1990). Excitatory synaptic transmission in cultures of rat olfactory bulb. *J Neurophysiol*, 64(2), 598-606.
- Tzingounis, A. V., & Nicoll, R. A. (2006). Arc/Arg3.1: linking gene expression to synaptic plasticity and memory. *Neuron*, 52(3), 403-407.
- Varoqui, H., Schafer, M. K., Zhu, H., Weihe, E., & Erickson, J. D. (2002). Identification of the differentiation-associated Na⁺/Pi transporter as a novel vesicular glutamate transporter expressed in a distinct set of glutamatergic synapses. *J Neurosci*, 22(1), 142-155.
- Veinante, P., & Freund-Mercier, M. J. (1997). Distribution of oxytocin- and vasopressin-binding sites in the rat extended amygdala: a histoautoradiographic study. *J Comp Neurol*, 383(3), 305-325.
- Vochteloo, J. D., & Koolhaas, J. M. (1987). Medial amygdala lesions in male rats reduce aggressive behavior: Interference with experience. *Physiol. Behav. Neurosci*, 41, 99-102.
- Wahle, P., Sanides-Kohlrausch, C., Meyer, G., & Lubke, J. (1990). Substance P- and opioid-immunoreactive structures in olfactory centers of the cat: adult pattern and postnatal development. *J Comp Neurol*, 302(2), 349-369.
- Williams, J. M., Beckmann, A. M., Mason-Parker, S. E., Abraham, W. C., Wilce, P. A., & Tate, W. P. (2000). Sequential increase in Egr-1 and AP-1 DNA binding activity in the dentate gyrus following the induction of long-term potentiation. *Brain Res Mol Brain Res*, 77(2), 258-266.

- Winslow, J. T., & Insel, T. R. (2004). Neuroendocrine basis of social recognition. *Curr Opin Neurobiol*, 14(2), 248-253.
- Yan, Z., Hsieh-Wilson, L., Feng, J., Tomizawa, K., Allen, P. B., Fienberg, A. A., et al. (1999). Protein phosphatase 1 modulation of neostriatal AMPA channels: regulation by DARPP-32 and spinophilin. *Nat Neurosci*, 2(1), 13-17.
- Zeman, W., & Innes, J. R. M. (1963). *Craigie's Neuroanatomy of the Rat*. New York: ACADEMIC PRESS.
- Zhang, X., & Firestein, S. (2002). The olfactory receptor gene superfamily of the mouse. *Nat Neurosci*, 5(2), 124-133.