

9 Literaturverzeichnis

- Adler M., Gervais F., Siedel U. (2000) Phytoplankton species composition in the chemocline of mesotrophic lakes. *Arch.Hydrobiol.Spec.Issues Advanc.Limnol.* 55:513–530
- Adrian, R. Untersuchungen zur herbivoren und carnivoren Ernährungsweise von *Cyclops kolen-sis* und *C. vicinus* (Crustacea, Copepoda). 1988. Freie Universität Berlin. Ref Type: Thesis/Dissertation
- Adrian R., Frost T. M. (1992) Comparative feeding ecology of *Tropocyclops prasinus mexicanus* (Copepoda, Cyclopoida). *Journal of Plankton Research* 14:1369–1382
- Adrian R., Frost T. M. (1993) Omnivory in cyclopoid copepods: comparisions of algae and invertebrates as food for three, differently sized species. *Journal of Plankton Research* 15:643–658
- Adrian R., Schneider-Olt B. (1999) Top down effects of crustacean zooplankton on pelagic microorganisms in a mesotrophic lake. *Journal of Plankton Research* 21:2175–2190
- Adrian R., Wickham S. A., Butler N. M. (2001) Trophic interactions between zooplankton and the microbial community in contrasting food webs: the epilimnion and deep chlorophyll maximum of a mesotrophic lake. *Aquat.Microb.Ecol.* 24:83–97
- Adrian R., Schipolowski T. (2003) Bacterial and protozoan mass accumulation in the deep chlorophyll maximum of a mesotrophic lake. *Arch.Hydrobiol.* 157:27–46
- Ahlgren G., Lundstedt L., Brett M., Forsberg C. (1990) Lipid composition and food quality of some freshwater phytoplankton for cladoceran zooplankters. *Journal of Plankton Research* 12:809–818
- Anraku M. (1964) Some technical problems in quantitative studies of grazing and predation by marine planktonic copepods. *Oceanogr.Soc.Jpn.* 20:19–29
- Azam F., Fenchel T., Field J. G., Gray J. S., Meyer-Reil L. A., Thingstad F. (1983) The ecological role of water-column microbes in the sea. *Mar.Ecol.Prog.Ser.* 10:257–263
- Ban S. (1994) Effect of temperature and food concentration on post-embryonic development, egg production and body size of calanoid copepod *Eurytemora affinis*. *Journal of Plankton Research* 16:721–735
- Bernardi R., Guissani G. (1990) Are blue-green algae suitable as food for zooplankton? An overview. *Hydrobiologia* 200/201:29–41
- Boëchat I.G. (2005) Biochemical composition of protists: Dependency on diet and trophic mode and consequences for their nutritional quality. Humboldt-Universität zu Berlin. Thesis/Dissertation
- Boëchat I. G., Adrian R. (2005) Biochemical composition of algivorous freshwater ciliates: You are not what you eat. *FEMS Microbiology Ecology* 53:393–400
- Boëchat I. G., Schuran S., Adrian R. (2005) Supplementation of the protist *Chilomonas parame-cium* with a highly unsaturated fatty acid enhances its nutritional quality for the rotifer *Keratella quadrata*. *Journal of Plankton Research* 27:663–670
- Boëchat I. G., Krüger A., Adrian R. (2006) Sterol composition of four freshwater heterotrophic protists and their diet. *Microb.Ecol.* in press:
- Børshem K.Y., Bratbak G. (1987) Cell volume to cell carbon conversion factors for a bacterivo-rous *Monas* sp. enriched seawater. *Mar.Ecol.Prog.Ser.* 20:171–175
- Brandl Z. (1973) Laboratory culture of cyclopoid copepods in a defined food. *Vest.Cesk.Spol.Zool.* 73:81–88
- Brandl Z., Fernando C. H. (1978) Prey selection by the cyclopoid copepods *Mesocyclops edax* and *Cyclops vicinus*. *Verh.Internat.Verein.Limnol.* 20:2505–2510

- Brandl Z., Fernando C. H. (1986) Feeding and food consumption of *Mesocyclops edax*. Syllogeus 58:254–258
- Brandl Z. (1998) Feeding strategies of planktonic cyclopoids in lacustrine ecosystems 9. J.Mar.Systems 15:87–95
- Brett M. T., Wiackowski K., Lubnow F. S., Mueller-Solger A., Elser J. J., Goldman C. R. (1994) Species-dependent effects of zooplankton on planktonic ecosystem processes in Castle Lake, California. Ecology 75:2243–2254
- Brett M. T., Müller-Navarra D. C. (1997) The role of highly unsaturated fatty acids in aquatic foodweb processes. Freshwater Biology 38:483–499
- Burns C. W., Xu Z. (1990) Utilization of colonial cyanobacteria and algae by freshwater calanoid copepods: survivorship and reproduction of adult *Boeckella* spp. Arch.Hydrobiol. 117:257–270
- Burns C. W., Gilbert J. J. (1993) Predation on ciliates by freshwater calanoid copepods: rates of predation and relative vulnerabilities of prey. Freshwater Biology 30:377–393
- Burns C. W., Schallenberg M. (1996) Relative impacts of copepods, cladocerans and nutrients on the microbial food web of a mesotrophic lake. Journal of Plankton Research 18:683–714
- Burns C. W., Schallenberg M. (1998) Impacts of nutrients and zooplankton on the microbial food web of an ultra-oligotrophic lake. Journal of Plankton Research 20:1501–1525
- Burns C. W., Schallenberg M. (2001) Calanoid copepods versus cladocerans: Consumer effects on protozoa in lakes of different trophic status. Limnol.Oceanogr. 46:1558–1565
- Carrick H. J., Fahnstiel G. L., Stoermer E. F., Wetzel R. G. (1991) The importance of zooplankton-protozoan trophic coupling in Lake Michigan. Limnol.Oceanogr. 36:1335–1345
- Clarke N. V. (1978) The food of adult copepods from Lake Kainji, Nigeria. Freshwater Biology 8:321–326
- Cognard C. (1973) Influence de la température sur les développements embryonnaire et post-embryonnaire du copépode diaptomide *Eudiaptomus gracilis* (Sars, 1863). Crustaceana 24:323–331
- Corkett C. J., McLaren I. A. (1969) Egg production and oil storage by the copepod *Pseudocalanus* in the laboratory. J.Exp.Mar.Biol.Ecol. 3:90–105
- Culver D. A., Brunskill G. J. (1969) Fayetteville Green Lake, New York. V. Studies of primary production and zooplankton in a meromictic marl lake. Limnol.Oceanogr. 14:862–873
- DeBiase A. D., Sanders R. W., Porter K. G. (1990) Relative nutritional value of ciliate protozoa and algae as food for *Daphnia*. Microb.Ecol. 19:199–210
- Demott W. R. (1986) The role of taste in food selection by freshwater zooplankton. Oecologia 69:334–340
- Demott W. R. (1989) Optimal foraging theory as a predictor of chemically mediated food selection by suspension-feeding copepods. Limnol.Oceanogr. 34:140–154
- Demott W. R. (1995) Food selection by calanoid copepods in response to between-lake variation in food abundance. Freshwater Biology 33:171–180
- Desviletes C. H., Bourdier G., Amblard C. H., Barth B. (1997) Use of fatty acids for the assessment of zooplankton grazing on bacteria, protozoans and microalgae. Freshwater Biology 38:629–637
- Eckstein H. (1964) Untersuchungen über den Einfluß des Rheinwassers auf die Limnologie des Schluchsees. IV. Untersuchungen über die Einbürgerung des *Eudiaptomus gracilis* Sars in den Schluchsee. Arch.Hydrobiol./Suppl. 28:119–182
- Ederington M. C., McManus G. B., Harvey H. R. (1995) Trophic transfer of fatty acids, sterols, and a triterpenoid alcohol between bacteria, a ciliate, and the copepod *Acartia tonsa*. Limnol.Oceanogr. 40:860–867

- Ehret, H. Experimentelle Untersuchungen zum Nahrungsspektrum von *Eudiaptomus gracilis*.
2000. Albert-Ludwigs-Universität Freiburg. Diplom-Thesis
- Einsle U. (1993) Crustacea, Copepoda; Calanoida und Cyclopoida. Gustav Fischer Verlag, Stuttgart, Jena, New York
- Elgmork K. (1958) On the phenology of Mesocyclops oithonoides (G.O. Sars).
Verh. Internat. Verein. Limnol. 13:778–784
- Fee E. J. (1976) The vertical and seasonal distribution of chlorophyll in lakes of the experimental lakes area, northwestern Ontario: Implications for primary production estimates. Limnol. Oceanogr. 21:767–783
- Fessenden L., Cowles T. J. (1994) Copepod predation on phagotrophic ciliates in Oregon coastal waters. Mar. Ecol. Prog. Ser. 107:103–111
- Fogg G. E., Nalewajko C., Watt W. D. (1965) Extracellular products of phytoplankton photosynthesis. 1. Proc. Roy. Soc. London B 162:517–534
- Foissner W., Berger H., Schaumburg J. (1999) Identification and ecology of limnetic plankton ciliates.
- Frost B. W. (1972) Effects of size and concentration of food particles on the feeding behavior of the marine planktonic copepod *Calanus pacificus*. Limnol. Oceanogr. 17:805–815
- Fryer G. (1957) The food of some freshwater cyclopoid copepods. J. Anim. Ecol. 26:263–286
- Fussmann G., Weithoff G., Yoshida T. (2005) A direct, experimental test of resource vs. consumer dependence. Ecology 86:2924–2930
- Gasol J. M., Guerrero R., Pedrós-Alio C. (1992) Spatial and temporal dynamics of a metalimnetic *Cryptomonas* peak. Journal of Plankton Research 14:1565–1579
- Gervais F., Siedel U., Heilmann B., Weithoff G., Heisig-Gunkel G., Nicklisch A. (2003) Small scale vertical distribution of phytoplankton, nutrients and sulphide below the oxicline of a mesotrophic lake. Journal of Plankton Research 25:273–278
- Gophen M. (1976) Temperature effect of lifespan, metabolism, and development time of *Mesocyclops leuckarti* (Claus). Oecologia 25:271–277
- Gras R., Saint-Jean L. (1981) Durée du développement juvénile de quelques Copepodes planctinique du lac Tchad. Revue Hydrobiol. trop. 14:39–51
- Guillard R. R. L., Lorenzen C. J. (1972) Yellow-green algae with chlorophyllide c. J. Phycol. 8:10–14
- Hamburger K., Boëtius F. (1987) Ontogeny of growth, respiration and feeding rate of the freshwater calanoid copepode *Eudiaptomus graciloides*. Journal of Plankton Research 9:589–606
- Hansen A., Jeppesen E. (1992) Life cycle of Cyclops vicinus in relation to food availability, predation, diapause and temperature. Journal of Plankton Research 14:591–605
- Hansen A., Santer B. (1995) The influence of food resources on the development, survival and reproduction of the two cyclopoid copepods: *Cyclops vicinus* and *Mesocyclops leuckarti*. Journal of Plankton Research 17:631–646
- Hansen A., Hairston Jr. N. G. (1998) Food limitation in a wild cyclopoid copepod population: direct and indirect life history response. Oecologia 115:320–330
- Hansen A. (2000) Response of ciliates and Cryptomonas to the spring cohort of a cyclopoid copepod in a shallow hypereutrophic lake. Journal of Plankton Research 22:185–203
- Hart R. C. (1990) Copepod post-embryonic durations: pattern, conformity, and predictability. The realities of isochronal and equiproportional development, and trends in the copepodid-naupliar duration ratio. Hydrobiologia 206:175–206

- Hart R. C. (1996) Naupliar and copepodite growth and survival of two freshwater calanoids at various food levels: Demographic contrasts, similarities, and food needs. Limnol.Oceanogr. 41:648–658
- Harvey H. R., Eglinton G., O'Hara S. C. M., Corner E. D. S. (1987) Biotransformation and assimilation of dietary lipids by *Calanus* feeding on a dinoflagellate. Geochimica et Cosmochimica Acta 51:3031–3040
- Harvey H. R., Ederington M .C., McManus G. B. (1997) Lipid composition of the marine ciliates *Pleuronema* sp. and *Fabrea salina*: Shifts in response to changes in diet. J.Euk.Microbiol. 44:189–193
- Heinbokel J. F. (1978) Studies of the functional role of tintinnids in the southern California Bight I. Grazing and growth rates in laboratory cultures. Mar. Biol. 47: 177–189
- Herzig A. (1983) The ecological significance of the relationship between temperature and duration of embryonic development in planktonic freshwater copepods. Hydrobiologia 100:65–91
- Hessen, D. O. Eksperimentelle innhegningsforsøk i Gjersjøen med spesiell vekt på zooplanktonets relasjon til fytoplankton, bakterier og planktivor fisk. 1982. Univ. Oslo. Ref Type: Thesis/Dissertation
- Hessen D. O., Nilssen JP, Eriksen TO (1986) Food size spectra and species replacement within herbivorous zooplankton. Int.Revue ges.Hydrobiol. 71:1–10
- Hofmann W. (1979) Characteristics of syntopic populations of *Eudiaptomus gracilis* (SARS) and *E. graciloides* (LILLJEBORG) in three lakes of different trophic levels. Arch.Hydrobiol. 86:1–12
- Holling C. S. (1959) The components of predation as revealed by a study of small mammal predation of the European Pine Sawfly. Can.Entomol. 91:293–320
- Hopp U., Maier G., Bleher R. (1997) Reproduction and adult longevity of five species of planktonic cyclopoid copepods reared on different diets: a comparative study. Freshwater Biology 38:289–300
- Jack J. D., Gilbert J. J. (1993) Susceptibilities of different-sized ciliates to direct suppression by small and large cladocerans. Freshwater Biology 29:19–29
- Jack J. D., Gilbert J. J. (1997) Effects of Metazoan Predators on Ciliates in Freshwater Plankton Communities. J.Euk.Microbiol. 44:194–199
- Jacobs J. (1974) Quantitative measurement of food selection. A modification of the ratio and Ivlev's electivity index. Oecologia 14:413–417
- Jamieson C. D. (1980) The predatory feeding of copepodid stages III to adult *Mesocyclops leuckarti* (Claus). In: Kerfoot WC (ed) Evolution and ecology of zooplankton communities. University Press, London, p 518–536
- Jamieson C. D., Burns C. W. (1988) The effects of temperature and food on copepodite development, growth and reproduction in three species of *Boeckella* (Copepoda; Calanoida). Hydrobiologia 164:235–257
- Jeschke J. M., Kopp M., Tollrian R. (2002) Predator functional responses: discriminating between handling and digesting prey. Ecological Monographs 72:95–112
- Jiménez-Melero R., Santer B., Guerrero F. (2005) Embryonic and naupliar development of *Eudiaptomus gracilis* and *Eudiaptomus graciloides* at different temperatures: comments on individual variability. Journal of Plankton Research 27:1175–1187
- Jürgens K., Wickham S. A., Rothaupt K. O., Santer B. (1996) Feeding rates of macro- and microzooplankton on heterotrophic nanoflagellates. Limnol.Oceanogr. 41:1833–1839

- Jürgens K., Skibbe O., Jeppesen E. (1999) Impact of metazooplankton on the composition and population dynamics of planktonic ciliates in a shallow, hypertrophic lake. *Aquat.Microb.Ecol.* 17:61–75
- Kasprzak P., Gervais F., Adrian R., Weiler W., Radke R., Jäger I., Riest S., Siedel U., Schneider-Olt B., Böhme M., Eckmann R., Walz N. (2000) Trophic characteristics and pelagic food web structure of two mesotrophic lakes in Brandenburg (Germany). *Internat.Rev.Hydrobiol.* 85:167–189
- Kessler, K. (2003) Vertical distribution of Daphnia in a trade-off between food and temperature. Christian-Albrechts-Universität zu Kiel. Thesis/Dissertation
- Kirchman D. L. (1993) Leucine incorporation as a measure of biomass production by heterotrophic bacteria. In: Kemp PF, Sherr BF, Sherr EB, Cole JJ (eds) *Handbook of methods in aquatic microbial ecology*. Lewis Publishers, Boca Raton, p 509–512
- Knisly K., Geller W. (1986) Selective feeding of four zooplankton species on natural lake phytoplankton. *Oecologia* 69:86–94
- Korniyenko G. S. (1976) Contribution of infusoria to the nutrition of *Acanthocyclops vernalis* and *Cyclops vicinus*. *Hydrobiol.Journal* 12:62–65
- Lampert W. (1977) Studies on the carbon balance of *Daphnia pulex* de Geer as related to environmental conditions. II. The dependence of carbon assimilation on animal size, temperature, food concentration, and diet species 4. *Arch.Hydrobiol.Supp.* 48:310–335
- Lampert W., Fleckner H., Rai H., Taylor B. E. (1986) Phytoplankton control by grazing zooplankton: a study on the spring clear water phase 6. *Limnol.Oceanogr.* 31:478–490
- Lampert W., Sommer U. (1993) *Limnoökologie*. Georg Thieme Verlag, Stuttgart
- Lancelot C. (1979) Gross excretion rates of natural marine phytoplankton and heterotrophic uptake of excreted products in the southern North Sea as determined by short-term kinetics 2. *Mar.Ecol.Prog.Ser.* 1:179–186
- Landry M. R. (1983) The development of marine calanoid copepods with comment on the iso-chronal rule. *Limnol.Oceanogr.* 28:614–624
- Larsson U., Hagström Å. (1982) Fractionated phytoplankton primary production, exudate release and bacterial production in a Baltic eutrophication gradient 3. *Mar.Biol.* 67:57–70
- Lindholm T. (1992) Ecological role of depth maxima of phytoplankton. *Arch.Hydrobiol.Beih.Ergebn.Limnol.* 35:33–45
- Llewellyn C. A., Fishwick J. R., Blackford J. C. (2005) Phytoplankton community assemblage in the English Channel: a comparision using chlorophyll a derived from HPLC-CHEMTAX and carbon derived from microscopy cell counts. *Journal of Plankton Research* 27:103–119
- Maier G. (1989a) The seasonal cycle of *Thermocyclops crassus* (Fischer, 1853) (Copepoda: Cyclopoida) in a shallow, eutrophic lake. *Hydrobiologia* 178:43–58
- Maier G. (1989b) The effect of temperature on the development times of eggs, naupliar and copepodite stages of five species of cyclopoid copepods. *Hydrobiologia* 184:79–88
- Maier G. (1989) Variable life cycles in freshwater copepod *Cyclops vicinus* (Uljanin 1875): Support for the predator avoidance hypothesis? *Arch.Hydrobiol.* 115:203–219
- Maier G. (1990) Reproduction and life history of a Central European, planktonic *Diacyclops bicuspidatus* population. *Arch.Hydrobiol.* 117:485–495
- Maier G. (1992) The reproductive biology of *Cyclops vicinus*. *Journal of Plankton Research* 14:127–135

- Maier G. (1994) Patterns of life history among cyclopoid copepods of central Europe. Freshwater Biology 31:77–86
- Maier G. (1995) Mating frequency and interspecific matings in some freshwater cyclopoid copepods. Oecologia 101:245–250
- Maier G. (1996) Copepod communities in lakes of varying trophic degree. Arch.Hydrobiol. 136:455–465
- Marshall S. M., Orr A. P. (1972) The biology of a marine copepod. Springer,
- Massana R., Gasol J. M., Jürgens K., Pedrós-Alio C. (1994) Impact of *Daphnia pulex* on a metalimnetic microbial community. Journal of Plankton Research 16:1379–1399
- McLaren I. A. (1965) Some relationships between temperatures and egg size, body size, development rate, and fecundity. Limnol.Oceanogr. 10:528–538
- Mohr S., Adrian R. (2000) Functional responses of rotifers *Brachionus calyciflorus* and *Brachionus rubens* feeding on armored and unarmored ciliates. Limnol.Oceanogr. 45:1175–1180
- Mohr S., Adrian R. (2002) Effects of *Brachionus calyciflorus* and *Brachionus rubens* on a manipulated freshwater microbial community. Journal of Plankton Research 24:25–31
- Mohr S., Adrian R. (2002) Reproductive success of the rotifer *Brachionus calyciflorus* feeding on ciliates and flagellates of different trophic modes. Freshwater Biology 47:1832–1839
- Mohr S., Gerten D., Adrian R. (2002) Minimal algal food requirements in the presence of protozoan prey for the rotifer *Brachionus calyciflorus*. Journal of Plankton Research 24:723–728
- Montagnes D. J. S., Berges J. A., Harrison P. J., Taylor F. J. R. (1994) Estimating carbon, nitrogen, protein, and chlorophyll a from volume in marine phytoplankton. Limnol.Oceanogr. 39:1044–1066
- Montagnes D. J. S., Berger J. D., Taylor F. J. R. (1996) Growth rate of the marine planktonic ciliate *Strombidinopsis cheshiri* Snyder and Ohman as a function of food concentration and interclonal variability. J.Exp.Mar.Biol.Ecol. 206:121–132
- Moriarty D. J. W., Darlington J. P. E. C., Dunn I. G., Moriarty M., Tevlin M. P. (1973) Feeding and grazing in Lake George, Uganda. Proc.R.Soc.Lond.B. 184:299–319
- Muck P., Lampert W. (1980) Feeding of freshwater filter-feeders at very low food concentrations: Poor evidence for "threshold feeding" and "optimal foraging" in *Daphnia longispina* and *Eudiaptomus gracilis*. Journal of Plankton Research 2:367–379
- Muck P., Lampert W. (1984) An experimental study on the importance of food conditions for the relative abundance of calanoid copepods and cladocerans. Arch.Hydrobiol./Suppl. 66:157–179
- Munro I. G. (1974) The effect of Temperature on the development of egg, naupliar and copepodite stages of two species of copepods, *Cyclops vicinus* Uljanin and *Eudiaptomus gracilis* Sars. Oecologia 16:355–367
- Nilssen J. P., Elgmork K. (1977) *Cyclops abyssorum* - life cycle dynamics and habitat selection. Mem.Ist.Ital.Idrobiol. 34:197–238
- Nilssen J. P. (1980) When and how to reproduce: a dilemma for limnetic cyclopoid copepods. In: Kerfoot WC (ed) Evolution and Ecology of Zooplankton Communities. University Press of New England, Hanover, p 418–426
- Nilssen J. P., Wærvågen SB (2000) Superficial ecosystem similarities vs autecological stripping: the "twin species" *Mesocyclops leuckarti* (Claus) and *Thermocyclops oithonoides* (Sars) - seasonal habitat utilisation and life history traits. J.Limnol. 59:79–102
- Northcote T. G., Halsey T. G. (1969) Seasonal changes in the limnology of some meromictic lakes in southern British Columbia. J.Fish.Res.Board Can. 26:1763–1787

- Nyberg, J. Zooplankton samfunnet i Borrevannet (1981), Vestfold, med spesiell vekt på livshistorene til Cyclops strenuus, Mesocyclops leuckarti og Eurytemora velox. Univ. Oslo. Thesis/Dissertation
- Oksum-Eriksen, T. (1982) Eksperimentelle innhengningsforsøk i Gjersjøen: beiteeffekt og interaksjoner i zooplankton samfunnet analysert i småinnhegninger. Univ. Oslo. Thesis/Dissertation
- Opitz, M. (2005) Studien zur Ökologie des Phytoplanktons in zwei mesotrophen Seen, unter Berücksichtigung der Ökophysiologie der Spezies im Bereich der Chemokline. Humboldt-Universität zu Berlin. Thesis/Dissertation
- Papinska K. (1985) Carnivorous and detritivorous feeding of *Mesocyclops leuckarti* Claus (Cyclopoida, Copepoda). *Hydrobiologia* 120:249–257
- Pasternak A. F., Huntingford F. A., Crompton D. W. T. (1995) Changes in metabolism and behaviour of freshwater copepod *Cyclops strenuus abyssorum* infected with *Diphyllobothrium* spp. *Parasitology* 110:395–399
- Pasternak A. F., Arashkevich E. G., Semenova T. N. (1996) Summer delay in the development of *Eudiaptomus graciloides*: is this a diapause? *Russian Journal of Aquatic Ecology* 5:39–47
- Pasternak A. F., Arashkevich E. G. (1999) Resting stages in the life cycle of *Eudiaptomus graciloides* (Lill.) (Copepoda: Calanoida) in Lake Glubokoe. *Journal of Plankton Research* 21:309–325
- Pedrós-Alió C., Gasol J. M., Guerrero R. (1987) On the ecology of a *Cryptomonas phaseolus* population forming a metalimnetic bloom in Lake Cisó, Spain: Annual distribution and loss factors. *Limnol.Oceanogr.* 32:285–298
- Peters R. H. (1984) Methods for the study of feeding, grazing and assimilation by zooplankton. In: Downing J. A., Rigler F. H. (eds) *A Manual on Methods for the Assessment of Secondary Productivity in Fresh Waters*. Blackwell Scientific Publications, Oxford, p 336–412
- Pomeroy L. R. (1984) Significance of Microrganisms in Carbon and Energy Flow in Marine Ecosystems. In: Klug MJ, Reddy CA (eds) *Current Perspectives in Microbial Ecology*. American Society for Microbiology, Washington,D.C., p 405–411
- Porter K. G., Feig Y. S. (1980) The use of DAPI for identifying and counting aquatic microflora. *Limnol.Oceanogr.* 25:141–153
- Porter K. G., Sherr E. B., Sherr B. F., Pace M., Sanders R. W. (1985) Protozoa in planktonic food webs. *J.Plotozool.* 32:409–415
- Porter K. G. (1996) Integrating the microbial loop and the classic food chain into a realistic planktonic food web. In: Polis G. A., Winemiller K. O. (eds) *Food Webs – Integration of Patterns & Dynamics*. Chapman & Hall, New York, p 51–59
- Putt M., Stoecker D. K. (1989) An experimentally determined carbon : volume ratio for marine "oligotrichous" ciliates from estuarine and coastal waters. *Limnol.Oceanogr.* 34:1097–1103
- Richman S., Dodson S. I. (1983) The effect of food quality on feeding and respiration by *Daphnia* and *Diaptomus* 5. *Limnol.Oceanogr.* 28:948–956
- Sanders R. W., Wickham S. A. (1993) Planktonic protozoa and metazoa: predation, food quality and population control. *Marine Microbial Food Webs* 7:197–223
- Sanders R. W., Williamson C. E., Stutzman P. L., Moeller R. E., Goulden C. E., Aoki-Goldsmith R. (1996) Reproductive success of "herbivorous" zooplankton fed algal and nonalgal food resources. *Limnol.Oceanogr.* 41:1295–1305
- Santer B. (1993) Potential importance of algae in the diet of adult *Cyclops vicinus*. *Freshwater Biology* 30:269–278

- Santer B., Bosch F.vd (1994) Herbivorous nutrition of *Cyclops vicinus*: the effect of a pure algal diet on feeding, development, reproduction and life cycle. Journal of Plankton Research 16:171–195
- Santer B. (1994) Influence of food type and concentration on the development of *Eudiaptomus gracilis* and implications for interactions between calanoid and cyclopoid copepods. Arch.Hydrobiol. 131:141–159
- Santer B., Lampert W. (1995) Summer diapause in cyclopoid copepods: adaptative response to a food bottleneck? Journal of Animal Ecology 64:600–613
- Santer B. (1996) Nutritional suitability of the dinoflagellate *Ceratium furcoides* for four copepod species. Journal of Plankton Research 18:323–333
- Santer B., Blohm-Sievers E., Cáceres C. E., Hairston Jr. N. G. (2000) Life-history variation in the coexisting freshwater copepods *Eudiaptomus gracilis* and *Eudiaptomus graciloides*. Arch.Hydrobiol. 149:441–458
- Schönberg N. (1961) Über die Biologie einiger pelagischer Kopepodenarten im See Võrtsjärv. Hüdrobioloogilised Uurimused. Academy of sciences of the Estonian S.S.R., p 159–193
- Schulze P. C., Folt C. L. (1989) Effects of conspecifics and phytoplankton on predation rates of the omnivorous copepods *Epischura lacustris* and *Epischura nordenskioldi*. Limnol.Oceanogr. 34:444–450
- Sherr E. B., Sherr B. F. (1988) Role of microbes in pelagic food webs: A revised concept. Limnol.Oceanogr. 33:1225–1227
- Simek, K., Bobková, J., Macek, M., Nedoma, J., and Psenner, R. Ciliate grazing on picoplankton in a eutrophic reservoir during the summer phytoplankton maximum: A study at the species and community level. Limnol.Oceanogr. 40(6), 1077–1090. 1995.
Ref Type: Generic
- Simek K., Macek M., Pernthaler J., Straskraba V., Psenner R. (1996) Can freshwater planktonic ciliates survive on a diet of picoplankton? Journal of Plankton Research 18:597–613
- Smyly W. J. P. (1970) Observations on rate of development, longevity and fecundity of *Acanthocyclops viridis* (Jurine) (Copepoda, Cyclopoida) in relation to type of prey. Crustaceana 18:21–36
- Stabell T. (1996) Ciliate bacterivory in epilimnetic waters. Aquat.Microb.Ecol. 10:265–272
- Steiner, S. (2002) Dynamik und Energietransfer einer planktischen Crustaceengemeinschaft in Abhängigkeit von der Nahrungsgrundlage und den Planktivoren. Institut für Hydrobiologie, Technische Universität Dresden. Thesis/Dissertation
- Stemberger R. S. (1985) Prey selection by copepod *Diacyclops thomasi*. Oecologia 65:492–497
- Stockner J. G., Porter K. G. (1988) Microbial Food Webs in Freshwater Planktonic Ecosystems. In: Carpenter S. R. (ed) Complex Interactions in Lake Communities. Springer, New York, p 69–84
- Stoecker D. K., Evans G. T. (1985) Effects of protozoan herbivory and carnivory in a microplankton food web. Mar.Ecol.Prog.Ser. 25:159–167
- Stoecker D. K., Egloff D. A. (1987) Predation by *Acartia tonsa* Dana on planktonic ciliates and rotifers. J.Exp.Mar.Biol.Ecol. 110:53–68
- Stoecker D. K., Capuzzo J. M. (1990) Predation on Protozoa: its importance to zooplankton. Journal of Plankton Research 12:891–908
- Tóth L. G., Zánkai N. P. (1985) Feeding of *Cyclops vicinus* (Uljanin)(Copepoda: Cyclopoida) in Lake Balaton on the basis of gut content analyses. Hydrobiologia 122:251–260
- Tóth L. G., Zánkai N. P., Messner O. M. (1987) Alga consumption of four dominant planktonic crustaceans in Lake Balaton (Hungary). Hydrobiologia 145:323–332

- Twombly S., Burns C. W. (1996) Exuvium analysis: A nondestructive method of analyzing copepod growth and development. Limnol.Oceanogr. 41:1324–1329
- Twombly S., Burns C. W. (1996) Effects of food quality on individual growth and development in the freshwater copepod *Boeckella triarticulata*. Journal of Plankton Research 18:2179–2196
- Utermöhl H. (1958) Zur Vervollkommnung der quantitativen Phytoplankton-Methodik. Internat.Verein.Theor.Angew.Limnol. 9:1–38
- van den Bosch F., Santer B. (1993) Cannibalism in *Cyclops abyssorum*. OIKOS 67:19–28
- Vanderploeg H. A., Pfaffenröder G. (1985) Modes of algal capture by the freshwater copepod *Diaptomus sicilis* and their relation to food-size selection. Limnol.Oceanogr. 30:871–885
- Vijverberg J. (1980) Effect of temperature in laboratory studies on development and growth of Cladocera and Copepoda from Tjeukemeer, The Netherlands. Freshwater Biology 10:317–340
- Vijverberg J., Richter A. F. (1982) Population dynamics and production of *Acanthocyclops robustus* (Sars) and *Mesocyclops leuckarti* (Claus) in Tjeukemeer. Hydrobiologia 95:261–274
- von Elert E., Stampfli P. (2000) Food quality for *Eudiaptomus gracilis*: the importance of particular highly unsaturated fatty acids. Freshwater Biology 45:189–200
- Walker K. F. (1975) The seasonal phytoplankton cycles of two saline lakes in central Washington. Limnol.Oceanogr. 20:40–53
- Wærvågen, S. B. En limnologisk studie av Gjerstadvann i Aust-Agder, med spesiell vekt på zooplanktonsamfunnets livshistorier og populasjonsdynamikk. 0–177. 1985. Univ. Oslo. Ref Type: Thesis/Dissertation
- Weisse T., Müller H., Pinto-Coelho R. M., Springmann D. (1990) Response of the microbial loop to the phytoplankton spring bloom in a large prealpine lake. Limnol.Oceanogr. 35:781–794
- Weisse T., Frahm A. (2001) Species specific interactions between small planktonic ciliates (*Urotricha* spp.) and rotifers (*Keratella* spp.). Journal of Plankton Research 23:1329–1338
- Weisse T., Karstens N., Meyer V. C. L., Janke L., Lettner S., Teichgräber K. (2001) Niche separation in common prostome freshwater ciliates: the effect of food and temperature. Aquat.Microb.Ecol. 26:167–179
- Weisse T., Frahm A. (2002) Direct and indirect impact of two common rotifer species (*Keratella* spp.) on two abundant ciliate species (*Urotricha furcata*, *Balanion plancticum*). Freshwater Biology 47:53–64
- Whitehouse J. W., Lewis B. G. (1973) The effect of diet and density on development, size and egg production in *Cyclops abyssorum* Sars, 1863 (Copepoda, Cyclopoida). Crustaceana 25:225–236
- Wiackowski K., Brett M. T., Goldman C. R. (1994) Differential effects of zooplankton species on ciliate community structure. Limnol.Oceanogr. 39:486–492
- Wickham S. A. (1995a) *Cyclops* predation on ciliates: species-specific differences and functional responses. Journal of Plankton Research 17:1633–1646
- Wickham S. A. (1995b) Trophic relations between cyclopoid copepods and ciliated protists: Complex interactions link the microbial and classical food webs. Limnol.Oceanogr. 40:1173–1181
- Williamson C. E. (1980) The predatory behavior of *Mesocyclops edax*: Predator preferences, prey defenses, and starvation-induced changes. Limnol.Oceanogr. 25:903–909
- Williamson C. E., Gilbert J. J. (1980) Variation among zooplankton predators: The potential of *Asplanchna*, *Mesocyclops*, and *Cyclops* to attack, capture, and eat various rotifer prey. In:

- Kerfoot WC (ed) Evolution and Ecology of Zooplankton Communities. University Press of New England, Hanover, p 509–517
- Williamson C. E., Sanders R. W., Moeller R. E., Stutzman P. L. (1996) Utilization of subsurface food resources for zooplankton reproduction: Implications for diel vertical migration theory. Limnol.Oceanogr. 41:224-233
- Woitke P., Martin C. D., Nicklisch S., Kohl J. G. (1994) HPLC-determination of lipophilic photosynthesis pigments in algal cultures and lake water samples using a non-endcapped C 18-RP column. Fres.J.Anal.Chem. 348:762–768
- Xu Z., Burns C. W. (1991) Development, growth and survivorship of juvenile calanoid copepods on diets of cyanobacteria and algae. Internat.Rev.Hydrobiol. 76:73–87
- Zánkai N. P., Ponyi J. E. (1974) On the feeding of *Eudiaptomus gracilis* in Lake Balaton. Annal.Biol.Tihany 41:363–371
- Zánkai N. P. (1978) The duration of development of *Eudiaptomus gracilis* (G. O. Sars) Copepoda in lake Balaton. Acta Biol.Debrecina 15:193–198
- Zánkai N. P. (1984) Predation of *Cyclops vicinus* (Copepoda: Cyclopoida) on small zooplankton animals in Lake Balaton (Hungary). Arch.Hydrobiol. 99:360–378
- Zánkai N. P. (1987) Post-embryonic development of cyclopoid copepods in various seasons at Lake Balaton (Hungary). Journal of Plankton Research 9:1057–1068
- Zánkai N. P. (1994) Feeding of copepodite and adult stages of *Eudiaptomus gracilis* (G. O. Sars, 1863) (Copepoda, Calanoida) on mixed plastic beads. Crustaceana 66:90–109