

## 7 Literaturverzeichnis

**Bader,A., Al-Dubai,H., and Weitzer,G.** (2000). Leukemia inhibitory factor modulates cardiogenesis in embryoid bodies in opposite fashions. *Circ. Res* 2000. Apr. 14;86. (7):787. - 94. **86**, 787-794.

**Bagutti,C., Wobus,A.M., Fassler,R., and Watt,F.M.** (1996). Differentiation of embryonal stem cells into keratinocytes: comparison of wild-type and beta 1 integrin-deficient cells. *Dev Biol* **179**, 184-196.

**Baharvand,H. and Matthaei,K.I.** (2004). Culture condition difference for establishment of new embryonic stem cell lines from the C57BL/6 and BALB/c mouse strains. *In Vitro Cell Dev Biol Anim* **40**, 76-81.

**Bain,G., Ray,W.J., Yao,M., and Gottlieb,D.I.** (1996). Retinoic acid promotes neural and represses mesodermal gene expression in mouse embryonic stem cells in culture. *Biochem Biophys Res Commun* **223**, 691-694.

**Berstine,E.G., Hooper,M.L., Grandchamp,S., and Ephrussi,B.** (1973). Alkaline phosphatase activity in mouse teratoma. *Proc Natl Acad Sci U S A* **70**, 3899-3903.

**Bohets,H.H., Nouwen,E.J., de Broe,M.E., and Dierickx,P.J.** (1994). Effects of foetal calf serum on cell viability, cytotoxicity and detoxification in the two kidney-derived cell lines LLC-PK1 and MDCK. *Toxic. in Vitro* **8**, 559-561.

**Bradley,A., Evans,M., Kaufman,M.H., and Robertson,E.** (1984). Formation of germ-line chimaeras from embryo-derived teratocarcinoma cell lines. *Nature* **309**, 255-256.

**Buesen,R., Visan,A., Genschow,E., Slawik,B., Spielmann,H., and Seiler,A.** (2004). Trends in improving the embryonic stem cell test (EST): an overview. *ALTEX*. 2004. ;21. (1):15. -22. **21**, 15-22.

**Chambers,I., Colby,D., Robertson,M., Nichols,J., Lee,S., Tweedie,S., and Smith,A.** (2003). Functional expression cloning of Nanog, a pluripotency sustaining factor in embryonic stem cells. *Cell* 2003. May. 30. ;113. (5. ):643. -55. **113**, 643-655.

- Chambers,I. and Smith,A.** (2004). Self-renewal of teratocarcinoma and embryonic stem cells. *Oncogene 2004 Sep 20;23(43. ):7150. -60.* **23**, 7150-7160.
- Conover,J.C., Ip,N.Y., Poueymirou,W.T., Bates,B., Goldfarb,M.P., DeChiara,T.M., and Yancopoulos,G.D.** (1993). Ciliary neurotrophic factor maintains the pluripotentiality of embryonic stem cells. *Development* **119**, 559-565.
- Cui,L., Johkura,K., Yue,F., Ogiwara,N., Okouchi,Y., Asanuma,K., and Sasaki,K.** (2004). Spatial Distribution and Initial Changes of SSEA-1 and Other Cell Adhesion-related Molecules on Mouse Embryonic Stem Cells Before and During Differentiation. *J Histochem Cytochem 2004. Nov. ;52. (11. ):1447. -57.* **52**, 1447-1457.
- Dani,C., Chambers,I., Johnstone,S., Robertson,M., Ebrahimi,B., Saito,M., Taga,T., Li,M., Burdon,T., Nichols,J. et al.** (1998). Paracrine induction of stem cell renewal by LIF-deficient cells: a new ES cell regulatory pathway. *Dev Biol* **203**, 149-162.
- Doetschman,T.C., Eistetter,H., Katz,M., Schmidt,W., and Kemler,R.** (1985). The in vitro development of blastocyst-derived embryonic stem cell lines: formation of visceral yolk sac, blood islands and myocardium. *J Embryol Exp Morphol* **87:27-45.**, 27-45.
- Drab,M., Haller,H., Bychkov,R., Erdmann,B., Lindschau,C., Haase,H., Morano,I., Luft,F.C., and Wobus,A.M.** (1997). From totipotent embryonic stem cells to spontaneously contracting smooth muscle cells: a retinoic acid and db-cAMP in vitro differentiation model. *FASEB J* **11**, 905-915.
- Evans,M.J. and Kaufman,M.H.** (1981). Establishment in culture of pluripotential cells from mouse embryos. *Nature* **292**, 154-156.
- Fairchild,P.J., Brook,F.A., Gardner,R.L., Graca,L., Strong,V., Tone,Y., Tone,M., Nolan,K.F., and Waldmann,H.** (2000). Directed differentiation of dendritic cells from mouse embryonic stem cells. *Curr Biol 2000. Nov. 30. ;10. (23):1515. -8.* **10**, 1515-1518.
- Fijnvandraat,A.C., van Ginneken,A.C., de Boer,P.A., Ruijter,J.M., Christoffels,V.M., Moorman,A.F., and R.H.** (2003). Cardiomyocytes derived from embryonic stem cells resemble cardiomyocytes of the embryonic heart tube. *Cardiovasc Res JID - 0077427* **58**, 399-409.

- Gajovic,S., Chowdhury,K., and Gruss,P.** (1998). Genes expressed after retinoic acid-mediated differentiation of embryoid bodies are likely to be expressed during embryo development. *Exp Cell Res* **242**, 138-143.
- Gardner,R.L. and Brook,F.A.** (1997). Reflections on the biology of embryonic stem (ES) cells. *Int J Dev Biol* **41**, 235-243.
- Gearing,D.P., Thut,C.J., VandeBos,T., Gimpel,S.D., Delaney,P.B., King,J., Price,V., Cosman,D., and Beckmann,M.P.** (1991). Leukemia inhibitory factor receptor is structurally related to the IL-6 signal transducer, gp130. *EMBO J* **10**, 2839-2848.
- Genschow,E., Spielmann,H., Scholz,G., Pohl,I., Seiler,A., Clemann,N., Bremer,S., and Becker,K.** (2004). Validation of the embryonic stem cell test in the international ECVAM validation study on three in vitro embryotoxicity tests. *Altern. Lab Anim* **32**, 209-244.
- Genschow,E., Spielmann,H., Scholz,G., Seiler,A., Brown,N., Piersma,A., Brady,M., Clemann,N., Huuskonen,H., Paillard,F. et al.** (2002). The ECVAM international validation study on in vitro embryotoxicity tests: results of the definitive phase and evaluation of prediction models. European Centre for the Validation of Alternative Methods. *Altern. Lab Anim.* 2002 Mar. -Apr. ;30(2):151. -76. **30**, 151-176.
- Giard,D.J.** (1987). Routine heat inactivation of serum reduces its capacity to promote cell attachment. *In Vitro Cell Dev Biol* **23**, 691-697.
- Gossler,A., Doetschman,T., Korn,R., Serfling,E., and Kemler,R.** (1986). Transgenesis by means of blastocyst-derived embryonic stem cell lines. *Proc Natl Acad Sci U S A* **83**, 9065-9069.
- Gómez-Lechón,M.J.** (1999). Oncostatin M: signal transduction and biological activity. *Life Sci* **65**, 2019-2030.
- Guan,K., Chang,H., Rolletschek,A., and Wobus,A.M.** (2001). Embryonic stem cell-derived neurogenesis. Retinoic acid induction and lineage selection of neuronal cells. *Cell Tissue Res* 2001. Aug. ;305(2):171. -6. **305**, 171-176.

- Hahnel,A.C., Rappolee,D.A., Millan,J.L., Manes,T., Ziomek,C.A., Theodosiou,N.G., Werb,Z., Pedersen,R.A., and Schultz,G.A.** (1990). Two alkaline phosphatase genes are expressed during early development in the mouse embryo. *Development* **110**, 555-564.
- Hibi,M., Murakami,M., Saito,M., Hirano,T., Taga,T., and Kishimoto,T.** (1990). Molecular cloning and expression of an IL-6 signal transducer, gp130. *Cell* **63**, 1149-1157.
- Ichihara,M., Hara,T., Kim,H., Murate,T., and Miyajima,A.** (1997). Oncostatin M and leukemia inhibitory factor do not use the same functional receptor in mice. *Blood* **90**, 165-173.
- Jakel,H.P., Wobus,A.M., Bloch,C., Jakel,N., and Schoneich,J.** (1983). A micromethod for the determination of alkaline phosphatase in mammalian cells. *Biomed Biochim Acta* **42**, 1123-1128.
- Johnson,L.V., Calarco,P.G., and Siebert,M.L.** (1977). Alkaline phosphatase activity in the preimplantation mouse embryo. *J Embryol Exp Morphol* **40**, 83-89.
- Kanno,S., Kim,P.K., Sallam,K., Lei,J., Billiar,T.R., and Shears,L.L.** (2004). Nitric oxide facilitates cardiomyogenesis in mouse embryonic stem cells. *Proc Natl Acad Sci U S A* 2004 Aug 17. ;101(33. ):12277. -81. Epub. 2004 Aug 10. **101**, 12277-12281.
- Labosky,P.A., Barlow,D.P., and Hogan,B.L.** (1994). Mouse embryonic germ (EG) cell lines: transmission through the germline and differences in the methylation imprint of insulin-like growth factor 2 receptor (Igf2r) gene compared with embryonic stem (ES) cell lines. *Development* **120**, 3197-3204.
- Leahy,A., Xiong,J.W., Kuhnert,F., and Stuhlmann,H.** (1999). Use of developmental marker genes to define temporal and spatial patterns of differentiation during embryoid body formation. *J Exp Zool* **284**, 67-81.
- Ling,V. and Neben,S.** (1997). In vitro differentiation of embryonic stem cells: immunophenotypic analysis of cultured embryoid bodies. *J Cell Physiol* **171**, 104-115.

- Maltsev,V.A., Ji,G.J., Wobus,A.M., Fleischmann,B.K., and Hescheler,J.** (1999). Establishment of beta-adrenergic modulation of L-type Ca<sup>2+</sup> current in the early stages of cardiomyocyte development. *Circ. Res* **84**, 136-145.
- Maltsev,V.A., Rohwedel,J., Hescheler,J., and Wobus,A.M.** (1993). Embryonic stem cells differentiate in vitro into cardiomyocytes representing sinusnodal, atrial and ventricular cell types. *Mech. Dev* **44**, 41-50.
- Maltsev,V.A., Wobus,A.M., Rohwedel,J., Bader,M., and Hescheler,J.** (1994). Cardiomyocytes differentiated in vitro from embryonic stem cells developmentally express cardiac-specific genes and ionic currents. *Circ. Res* **75**, 233-244.
- Martin,G.R.** (1980). Teratocarcinomas and mammalian embryogenesis. *Science* **209**, 768-776.
- Martin,G.R.** (1981). Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proc Natl Acad Sci U. S A* **78**, 7634-7638.
- Martin,G.R. and Evans,M.J.** (1975a). Differentiation of clonal lines of teratocarcinoma cells: formation of embryoid bodies in vitro. *Proc Natl Acad Sci U. S A* **72**, 1441-1445.
- Martin,G.R. and Evans,M.J.** (1975b). Multiple Differentiation of Clonal Teratocarcinoma Stem Cells Following Embryoid Body Formation in Vitro. *Cell* **6**, 467-474.
- Matsuda,T., Nakamura,T., Nakao,K., Arai,T., Katsuki,M., Heike,T., and Yokota,T.** (1999). STAT3 activation is sufficient to maintain an undifferentiated state of mouse embryonic stem cells. *EMBO J* **18**, 4261-4269.
- Matsui,Y., Zsebo,K., and Hogan,B.L.** (1992). Derivation of pluripotential embryonic stem cells from murine primordial germ cells in culture. *Cell* **70**, 841-847.
- Meszaros,K., Aberle,S., White,M., and Parent,J.B.** (1995). Immunoreactivity and bioactivity of lipopolysaccharide-binding protein in normal and heat-inactivated sera. *Infect Immun.* **63**, 363-365.

- Metcalf,D.** (2003). The unsolved enigmas of leukemia inhibitory factor. *Stem. Cells* 2003. ;21. (1):5. -14. **21**, 5-14.
- Mitsui,K., Tokuzawa,Y., Itoh,H., Segawa,K., Murakami,M., Takahashi,K., Maruyama,M., Maeda,M., and Yamanaka,S.** (2003). The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells. *Cell* 2003. May. 30. ;113. (5. ):631. -42. **113**, 631-642.
- Moreau,J.F., Donaldson,D.D., Bennett,F., Witek-Giannotti,J., Clark,S.C., and Wong,G.G.** (1988). Leukaemia inhibitory factor is identical to the myeloid growth factor human interleukin for DA cells. *Nature* **336**, 690-692.
- Murray,P. and Edgar,D.** (2001). The regulation of embryonic stem cell differentiation by leukaemia inhibitory factor (LIF). *Differentiation* 2001. Oct. ;68. (4-5. ):227-34. **68**, 227-234.
- Nichols,J., Evans,E.P., and Smith,A.G.** (1990). Establishment of germ-line-competent embryonic stem (ES) cells using differentiation inhibiting activity. *Development* **110**, 1341-1348.
- Nicolas,J.F., Avner,P., Gaillard,J., Guenet,J.L., Jakob,H., and Jacob,F.** (1976). Cell lines derived from teratocarcinomas. *Cancer Res* **36**, 4224-4231.
- Niwa,H., Burdon,T., Chambers,I., and Smith,A.** (1998). Self-renewal of pluripotent embryonic stem cells is mediated via activation of STAT3. *Genes Dev* **12**, 2048-2060.
- Niwa,H., Miyazaki,J., and Smith,A.G.** (2000). Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nat. Genet* 2000. Apr. ;24. (4):372. -6. **24**, 372-376.
- Oka,M., Tagoku,K., Russell,T.L., Nakano,Y., Hamazaki,T., Meyer,E.M., Yokota,T., and Terada,N.** (2002). CD9 Is Associated with Leukemia Inhibitory Factor-mediated Maintenance of Embryonic Stem Cells. *Mol Biol Cell* 2002. Apr. ;13(4):1274. -81. **13**, 1274-1281.

**Okamoto,K., Okazawa,H., Okuda,A., Sakai,M., Muramatsu,M., and Hamada,H.** (1990). A novel octamer binding transcription factor is differentially expressed in mouse embryonic cells. *Cell* **60**, 461-472.

**Oyamada,Y., Komatsu,K., Kimura,H., Mori,M., and Oyamada,M.** (1996). Differential regulation of gap junction protein (connexin) genes during cardiomyocytic differentiation of mouse embryonic stem cells in vitro. *Exp Cell Res* **229**, 318-326.

**Pasumarthi,K.B. and Field,L.J.** (2002). Cardiomyocyte enrichment in differentiating ES cell cultures: strategies and applications. *Methods Mol Biol* 2002. ;185. :157. -68. **185:157-68.**, 157-168.

**Pease,S., Braghetta,P., Gearing,D., Grail,D., and Williams,R.L.** (1990). Isolation of embryonic stem (ES) cells in media supplemented with recombinant leukemia inhibitory factor (LIF). *Dev Biol* **141**, 344-352.

**Pennica,D., Shaw,K.J., Swanson,T.A., Moore,M.W., Shelton,D.L., Zioncheck,K.A., Rosenthal,A., Taga,T., Paoni,N.F., and Wood,W.I.** (1995). Cardiotrophin-1. Biological activities and binding to the leukemia inhibitory factor receptor/gp130 signaling complex. *J Biol Chem* **270**, 10915-10922.

**Piquet-Pellorce,C., Grey,L., Mereau,A., and Heath,J.K.** (1994). Are LIF and related cytokines functionally equivalent? *Exp Cell Res* **213**, 340-347.

**Qi,X., Li,T.G., Hao,J., Hu,J., Wang,J., Simmons,H., Miura,S., Mishina,Y., and Zhao,G.Q.** (2004). BMP4 supports self-renewal of embryonic stem cells by inhibiting mitogen-activated protein kinase pathways. *Proc Natl Acad Sci U S A* %20;**101**, 6027-6032.

**Rathjen,J., Lake,J.A., Bettess,M.D., Washington,J.M., Chapman,G., and Rathjen,P.D.** (1999). Formation of a primitive ectoderm like cell population, EPL cells, from ES cells in response to biologically derived factors. *J Cell Sci* **112**, 601-612.

**Rathjen,P.D., Nichols,J., Toth,S., Edwards,D.R., Heath,J.K., and Smith,A.G.** (1990a). Developmentally programmed induction of differentiation inhibiting activity and the control of stem cell populations. *Genes Dev* **4**, 2308-2318.

- Rathjen,P.D., Toth,S., Willis,A., Heath,J.K., and Smith,A.G.** (1990b). Differentiation inhibiting activity is produced in matrix-associated and diffusible forms that are generated by alternate promoter usage. *Cell* **62**, 1105-1114.
- Raz,R., Lee,C.K., Cannizzaro,L.A., d'Eustachio,P., and Levy,D.E.** (1999). Essential role of STAT3 for embryonic stem cell pluripotency. *Proc Natl Acad Sci U. S A* **96**, 2846-2851.
- Resnick,J.L., Bixler,L.S., Cheng,L., and Donovan,P.J.** (1992). Long-term proliferation of mouse primordial germ cells in culture. *Nature* **359**, 550-551.
- Roach,M.L. and McNeish,J.D.** (2002). Methods for the isolation and maintenance of murine embryonic stem cells. *Methods Mol Biol* 2002. ;185. :1-16. **185:1-16.**, 1-16.
- Robson,P., Stein,P., Zhou,B., Schultz,R.M., and Baldwin,H.S.** (2001). Inner cell mass-specific expression of a cell adhesion molecule (PECAM-1/CD31) in the mouse blastocyst. *Dev Biol* 2001. Jun. 15. ;234. (2):317. -29. **234**, 317-329.
- Rohwedel,J., Guan,K., and Wobus,A.M.** (1999). Induction of cellular differentiation by retinoic acid in vitro. *Cells Tissues Organs* **165**, 190-202.
- Rohwedel,J., Kleppisch,T., Pich,U., Guan,K., Jin,S., Zuschratter,W., Hopf,C., Hoch,W., Hescheler,J., Witzemann,V. et al.** (1998). Formation of postsynaptic-like membranes during differentiation of embryonic stem cells in vitro. *Exp Cell Res* **239**, 214-225.
- Rolletschek,A., Blyszczuk,P., and Wobus,A.M.** (2004). Embryonic stem cell-derived cardiac, neuronal and pancreatic cells as model systems to study toxicological effects. *Toxicol Lett* 2004. Apr. 1;149. (1-3. ):361. -9. **149**, 361-369.
- Rose,T.M., Weiford,D.M., Gunderson,N.L., and Bruce,A.G.** (1994). Oncostatin M (OSM) inhibits the differentiation of pluripotent embryonic stem cells in vitro. *Cytokine* **6**, 48-54.
- Rosner,M.H., Vigano,M.A., Ozato,K., Timmons,P.M., Poirier,F., Rigby,P.W., and Staudt,L.M.** (1990). A POU-domain transcription factor in early stem cells and germ cells of the mammalian embryo. *Nature* **345**, 686-692.



- Rutledge,J.C.** (1997). Developmental toxicity induced during early stages of mammalian embryogenesis. *Mutat Res* **396**, 113-127.
- Sabapathy,K., Klemm,M., Jaenisch,R., and Wagner,E.F.** (1997). Regulation of ES cell differentiation by functional and conformational modulation of p53. *EMBO J* **16**, 6217-6229.
- Schmidt,M.M., Guan,K., and Wobus,A.M.** (2001). Lithium influences differentiation and tissue-specific gene expression of mouse embryonic stem (ES) cells in vitro. *Int J Dev Biol* 2001. Apr. ;45. (2):421. -9. **45**, 421-429.
- Scholer,H.R., Dressler,G.R., Balling,R., Rohdewohld,H., and Gruss,P.** (1990a). Oct-4: a germline-specific transcription factor mapping to the mouse t-complex. *EMBO J* **9**, 2185-2195.
- Scholer,H.R., Ruppert,S., Suzuki,N., Chowdhury,K., and Gruss,P.** (1990b). New type of POU domain in germ line-specific protein Oct-4. *Nature* **344**, 435-439.
- Scholz,G., Genschow,E., Brown,N.A., Piersma,A., Brady,M., Clemann,N. et al.** Final evaluation of the ECVAM Project "Prevalidation and validation of three in vitro embryotoxicity tests". Archives of Pharmakology Supplement to Volume 361[Number 4], R 162. 2000.
- Scholz,G., Pohl,I., Genschow,E., Klemm,M., and Spielmann,H.** (1999). Embryotoxicity screening using embryonic stem cells in vitro: correlation to in vivo teratogenicity. *Cells Tissues Organs* **165**, 203-211.
- Seiler,A., Visan,A., Pohl,I., Genschow,E., Buesen,R., and Spielmann,H.** (2002). [Improving the embryonic stem cell test (EST) by establishing molecular endpoints of tissue specific development using murine embryonic stem cells (D3 cells)]. *ALTEX*. 2002. ;19. Suppl 1:55. -63. **19 Suppl 1:55-63.**, 55-63.
- Shen,M.M. and Leder,P.** (1992). Leukemia inhibitory factor is expressed by the preimplantation uterus and selectively blocks primitive ectoderm formation in vitro. *Proc Natl Acad Sci U S A* **89**, 8240-8244.

- Smith,A.G.** (1992). Mouse embryo stem cells: their identification, propagation and manipulation. *Semin Cell Biol* **3**, 385-399.
- Smith,A.G., Heath,J.K., Donaldson,D.D., Wong,G.G., Moreau,J., Stahl,M., and Rogers,D.** (1988). Inhibition of pluripotential embryonic stem cell differentiation by purified polypeptides. *Nature* **336**, 688-690.
- Smith,A.G. and Hooper,M.L.** (1987). Buffalo rat liver cells produce a diffusible activity which inhibits the differentiation of murine embryonal carcinoma and embryonic stem cells. *Dev Biol* **121**, 1-9.
- Smith,T.A. and Hooper,M.L.** (1983). Medium conditioned by feeder cells inhibits the differentiation of embryonal carcinoma cultures. *Exp Cell Res* **145**, 458-462.
- Solter,D. and Knowles,B.B.** (1978). Monoclonal antibody defining a stage-specific mouse embryonic antigen (SSEA-1). *Proc Natl Acad Sci U. S A* **75**, 5565-5569.
- Solter,D. and Knowles,B.B.** (1979). Developmental stage-specific antigens during mouse embryogenesis. *Curr Top. Dev Biol* **13 Pt 1:139-65.**, 139-165.
- Spielmann,H.** (1998). Reproduction and development. *Environ Health Perspect* **106 Suppl 2:571-6.**, 571-576.
- Spielmann,H. and Scholz,G.** (1999) Invitox Protokoll 113 - Embryonic stem cell test (EST) <http://ecvam-sis.jrc.it>. 28-1-1999.
- Stewart,C.L., Gadi,I., and Bhatt,H.** (1994). Stem cells from primordial germ cells can reenter the germ line. *Dev Biol* **161**, 626-628.
- Stewart,C.L., Kaspar,P., Brunet,L.J., Bhatt,H., Gadi,I., Kontgen,F., and Abbondanzo,S.J.** (1992). Blastocyst implantation depends on maternal expression of leukaemia inhibitory factor. *Nature* **359**, 76-79.
- Strubing,C., Ahnert-Hilger,G., Shan,J., Wiedenmann,B., Hescheler,J., and Wobus,A.M.** (1995a). Differentiation of pluripotent embryonic stem cells into the neuronal lineage in vitro gives rise to mature inhibitory and excitatory neurons. *Mech. Dev* **53**, 275-287.

- Strubing,C., Wobus,A.M., and Hescheler,J.** (1995b). [Establishment of an in vitro model system for the differentiation of synaptically coupled neurons from mouse embryonic stem cells]. *ALTEX*. **12**, 129-137.
- Suda,Y., Suzuki,M., Ikawa,Y., and Aizawa,S.** (1987). Mouse embryonic stem cells exhibit indefinite proliferative potential. *J Cell Physiol* **133**, 197-201.
- Tessarollo,L.** (2001). Manipulating mouse embryonic stem cells. *Methods Mol Biol* 2001. ;158:47. -63. **158:47-63.**, 47-63.
- Toumadje,A., Kusumoto,K.I., Parton,A., Mericko,P., Dowell,L., Ma,G., Chen,L., Barnes,D.W., and Sato,J.D.** (2003). PLURIPOTENT DIFFERENTIATION IN VITRO OF MURINE ES-D3 EMBRYONIC STEM CELLS. *In Vitro Cell Dev Biol Anim.* 2003. Nov. ;39(10. ):449. -453. **39**, 449-453.
- Triglia,R.P. and Linscott,W.D.** (1980). Titers of nine complement components, conglutinin and C3b-inactivator in adult and fetal bovine sera. *Mol Immunol* **17**, 741-748.
- Viswanathan,S., Benatar,T., Rose-John,S., Lauffenburger,D.A., and Zandstra,P.W.** (2002). Ligand/receptor signaling threshold (LIST) model accounts for gp130-mediated embryonic stem cell self-renewal responses to LIF and HIL-6. *Stem. Cells* 2002. ;20(2):119. -38. **20**, 119-138.
- Ward,C.M., Barrow,K.M., and Stern,P.L.** (2004). Significant variations in differentiation properties between independent mouse ES cell lines cultured under defined conditions. *Exp Cell Res* **293**, 229-238.
- Ward,R.L.** (1979). Destruction of bacterial viruses in serum by heat and radiation under conditions that sustain the ability of serum to support growth of cells in suspended culture. *J Clin Microbiol* **10**, 650-656.
- Weiss,M.J. and Orkin,S.H.** (1996). In vitro differentiation of murine embryonic stem cells. New approaches to old problems. *J Clin Invest* **97**, 591-595.

**Williams,R.L., Hilton,D.J., Pease,S., Willson,T.A., Stewart,C.L., Gearing,D.P., Wagner,E.F., Metcalf,D., Nicola,N.A., and Gough,N.M.** (1988). Myeloid leukaemia inhibitory factor maintains the developmental potential of embryonic stem cells. *Nature* **336**, 684-687.

**Wobus,A.M.** (2001). Potential of embryonic stem cells. *Mol Aspects Med* 2001. Jun. ;22. (3. ):149. -64. **22**, 149-164.

**Wobus,A.M., Guan,K., and Pich,U.** (2001). In vitro differentiation of embryonic stem cells and analysis of cellular phenotypes. *Methods Mol Biol* 2001. ;158:263. -86. **158:263-86.**, 263-286.

**Wobus,A.M., Guan,K., Yang,H.T., and Boheler,K.R.** (2002). Embryonic stem cells as a model to study cardiac, skeletal muscle, and vascular smooth muscle cell differentiation. *Methods Mol Biol* 2002. ;185. :127. -56. **185:127-56.**, 127-156.

**Wobus,A.M., Holzhausen,H., Jakel,P., and Schoneich,J.** (1984). Characterization of a pluripotent stem cell line derived from a mouse embryo. *Exp Cell Res* **152**, 212-219.

**Wobus,A.M. and Kaomei Guan** (1998). Embryonic Stem Cell-Derived Cardiac Differentiation: Modulation of Differentiation and "Loss-of-Function" Analysis In Vitro. *Trends Cardiovasc Med* **8**, 64-74.

**Wobus,A.M., Kaomei,G., Shan,J., Wellner,M.C., Rohwedel,J., Ji,G., Fleischmann,B., Katus,H.A., Hescheler,J., and Franz,W.M.** (1997). Retinoic acid accelerates embryonic stem cell-derived cardiac differentiation and enhances development of ventricular cardiomyocytes. *J Mol Cell Cardiol* **29**, 1525-1539.

**Wobus,A.M., Rohwedel,J., Maltsev,V., and Hescheler,J.** (1995). Development of cardiomyocytes expressing cardiac-specific genes, action potentials, and ionic channels during embryonic stem cell-derived cardiogenesis. *Ann N Y Acad Sci* **752:460-9.**, 460-469.

**Wobus,A.M., Wallukat,G., and Hescheler,J.** (1991). Pluripotent mouse embryonic stem cells are able to differentiate into cardiomyocytes expressing chronotropic responses to adrenergic and cholinergic agents and Ca<sup>2+</sup> channel blockers. *Differentiation* **48**, 173-182.

- Yamane,T., Dylla,S.J., Muijtjens,M., and Weissman,I.L.** (2005). Enforced Bcl-2 expression overrides serum and feeder cell requirements for mouse embryonic stem cell self-renewal. *Proc Natl Acad Sci U S A* **102**, 3312-3317.
- Ying,Q.L., Nichols,J., Chambers,I., and Smith,A.** (2003). BMP induction of Id proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. *Cell* 2003. Oct. 31. ;115. (3. ):281. -92. **115**, 281-292.
- Yoshida,K., Chambers,I., Nichols,J., Smith,A., Saito,M., Yasukawa,K., Shoyab,M., Taga,T., and Kishimoto,T.** (1994). Maintenance of the pluripotential phenotype of embryonic stem cells through direct activation of gp130 signalling pathways. *Mech. Dev* **45**, 163-171.
- Zandstra,P.W., Le,H.V., Daley,G.Q., Griffith,L.G., and Lauffenburger,D.A.** (2000). Leukemia inhibitory factor (LIF) concentration modulates embryonic stem cell self-renewal and differentiation independently of proliferation. *Biotechnol. Bioeng.* 2000. Sep. 20;69. (6. ):607. -17. **69**, 607-617.
- zur Nieden,N.I., Kempka,G., and Ahr,H.J.** (2003). In vitro differentiation of embryonic stem cells into mineralized osteoblasts. *Differentiation* 2003. Jan. ;71. (1):18-27. **71**, 18-27.
- zur Nieden,N.I., Ruf,L.J., Kempka,G., Hildebrand,H., and Ahr,H.J.** (2001). Molecular markers in embryonic stem cells. *Toxicol In Vitro* 2001. Aug. -Oct. ;15. (4-5. ):455. -61. **15**, 455-461.