

6. Literatur

Accili, D., N. Perrotti, R. Rees-Jones und S.I. Taylor:

Tissue distribution and subcellular localization of an endogenous substrate (pp 120) for the insulin receptor-associated tyrosine kinase.
Endocrinology, 119 (1986), 1274-1280.

Albelda, S.M. und C.A. Buck:

Integrins and other cell adhesion molecules.
FASEB J., 4 (1990), 2868-2880.

Alberts, B., D. Bray, J. Lewis, M. Raff, K. Roberts und J.D. Watson:

Zell-Zell-Adhäsion, Zellverbindungen und die extrazelluläre Matrix.
In: Molekularbiologie der Zelle, Alberts, B. and e. al. (eds.), VCH, Weinheim (1995), 943-996.

Alonso-Lebrero, J.L., J.M. Serrador, C. Dominguez-Jimenez, O. Barreiro, A. Luque, M.A. del Pozo, K. Snapp, G. Kansas, R. Schwartz-Albiez, H. Furthmayr, F. Lozano und F. Sanchez-Madrid:

Polarization and interaction of adhesion molecules P-selectin glycoprotein ligand 1 and intercellular adhesion molecule 3 with moesin and ezrin in myeloid cells.
Blood, 95 (2000), 2413-2419.

Aplin, A.E., A. Howe, S.K. Alahari und R.L. Juliano:

Signal transduction and signal modulation by cell adhesion receptors: the role of integrins, cadherins, immunoglobulin-cell adhesion molecules, and selectins.
Pharmacol Rev, 50 (1998), 197-263.

Barnett, T.R., L. Drake und W.d. Pickle:

Human biliary glycoprotein gene: characterization of a family of novel alternatively spliced RNAs and their expressed proteins.
Mol. Cell. Biol., 13 (1993), 1273-1282.

Baum, O., W. Reutter, D. Flanagan, H. Callanan, Y.P. Lim, S.H. Lin und D.C. Hixson:

Anti-peptide sera against cell-CAM 105 determine high molecular-mass variants of the long isoform in rat hepatocytes.
Eur. J. Biochem., 228 (1995), 316-322.

Baum, O., S. Troll und D.C. Hixson:

The long and the short isoform of cell-CAM 105 show variant-specific modifications in adult rat organs.

Biochem. Biophys. Res. Commun., 227 (1996), 775-781.

Beauchemin, N., P. Draber, G. Dveksler, P. Gold, S. Gray-Owen, F. Grunert, S.

Hammarstrom, K.V. Holmes, A. Karlsson, M. Kuroki, S.H. Lin, L. Lucka, S.

Najjar, M. Neumaier, B. Obrink, J.E. Shively, K.M. Skubitz, C.P. Stanners, P.

Thomas, J.A. Thompson, M. Virji, S. v.Kleist, C. Wagener, S. Watt und W.

Zimmermann:

Redefined Nomenclature for Members of the Carcineembryonic Antigen Family.

Exp. Cell Res., 252 (1999), 243-249.

Beauchemin, N., T. Kunath, J. Robitaille, B. Chow, C. Turbide, E. Daniels und A.

Veillette:

Association of biliary glycoprotein with protein tyrosine phosphatase SHP-1 in malignant colon epithelial cells.

Oncogene, 14 (1997), 783-790.

Becker, A.:

Vergleichende Charakterisierung des Membranglycoproteins gp110 in der Leber und anderen Organen.

Dissertation, Freie Universität Berlin, Fachbereich Biochemie (1989).

Becker, A., R. Gossrau, C. Hoffmann und W. Reutter:

Localization of a putative cell adhesion molecule (gp110) in Wistar and Fischer rat tissues.

Histochemistry, 93 (1989), 55-61.

Becker, A., L. Lucka, C. Kilian, C. Kannicht und W. Reutter:

Characterisation of the ATP-dependent taurocholate-carrier protein (gp110) of the hepatocyte canalicular membrane.

Eur. J. Biochem., 214 (1993), 539-548.

Becker, A., R. Neumeier, C.S. Park, R. Gossrau und W. Reutter:

Identification of a transformation-sensitive 110-kDa plasma membrane glycoprotein of rat hepatocytes.

Eur. J. Cell Biol., 39 (1985), 417-423.

Berndorff, D., R. Gessner, B. Kreft, N. Schnoy, A.M. Lajous-Petter, N. Loch, W.

Reutter, M. Hortsch und R. Tauber:

Liver-intestine cadherin: molecular cloning and characterization of a novel Ca(2+)-dependent cell adhesion molecule expressed in liver and intestine.

J. Cell Biol., 125 (1994), 1353-1369.

Bertolotti, R., U. Rutishauser und G.M. Edelman:

A cell surface molecule involved in aggregation of embryonic liver cells.

Proc. Natl. Acad. Sci. U S A, 77 (1980), 4831-4835.

Birnboim, H.C. und T. Doly:

A rapid alkaline procedure for screening recombinant plasmid DNA.

Nucl. Acids Res., 7 (1979), 1513-1523.

Blomberg, L.A., S.M. Wu, G. Dirami, M. Dym, J.Y. Chou und W.Y. Chan:

Characterization and cellular localization of PSG in rat testis.

Mol. Cell. Biochem., 177 (1997), 229-237.

Blume-Jensen, P. und T. Hunter:

Oncogenic kinase signalling.

Nature, 411 (2001), 355-365.

Boehm, M.K., M.O. Mayans, J.D. Thornton, R.H. Begent, P.A. Keep und S.J. Perkins:

Extended glycoprotein structure of the seven domains in human carcinoembryonic antigen by X-ray and neutron solution scattering and an automated curve fitting procedure: implications for cellular adhesion.

J. Mol. Biol., 259 (1996), 718-736.

Bork, P., L. Holm und C. Sander:

The immunoglobulin fold. Structural classification, sequence patterns and common core.

J. Mol. Biol., 242 (1994), 309-320.

Boulton, I.C. und S.D. Gray-Owen:

Neisserial binding to CEACAM1 arrests the activation and proliferation of CD4+ T lymphocytes.

Nat. Immunol., 19 (2002), 19.

Brandley, B.K., S.J. Swiedler und P.W. Robbins:

Carbohydrate ligands of the LEC cell adhesion molecules.

Cell, 63 (1990), 861-863.

Brenner, S.L. und E.D. Korn:

On the mechanism of actin monomer-polymer subunit exchange at steady state.
J. Biol. Chem., 258 (1983), 5013-5020.

Brooks, J.E.:

Properties and uses of restriction endonucleases.

Methods of enzymology, Vol.152. Guide to molecular cloning technique, 113 (1987),

Brummer, J., A. Ebrahimnejad, R. Flayeh, U. Schumacher, T. Loning, A.M. Bamberger und C. Wagener:

cis Interaction of the cell adhesion molecule CEACAM1 with integrin beta(3).

Am. J. Pathol., 159 (2001), 537-546.

Brummer, J., M. Neumaier, C. Gopfert und C. Wagener:

Association of pp60c-src with biliary glycoprotein (CD66a), an adhesion molecule of the carcinoembryonic antigen family downregulated in colorectal carcinomas.

Oncogene, 11 (1995), 1649-1655.

Budt, M., B. Michely, M.M. Müller, W. Reutter und L. Lucka:

Secreted CEACAM1 Splice Variants in Rat Cell Lines and *in Vivo* in Rat Serum.
Biochem. Biophys. Res. Comm., *im Druck* (2002)

Burridge, K. und M. Chrzanowska-Wodnicka:

Focal adhesions, contractility, and signaling.

Ann. Rev. Cell. Dev. Biol., 12 (1996), 463-518.

Burshyn, D.N., W. Yang, T. Yi und E.O. Long:

A novel phosphotyrosine motif with a critical amino acid at position -2 for the SH2 domain-mediated activation of the tyrosine phosphatase SHP-1.
J. Biol. Chem., 272 (1997), 13066-13072.

Carey, D.J.:

Syndecans: multifunctional cell-surface co-receptors.

Biochem. J., 327 (1997), 1-16.

Carriquiry, L.A. und A. Pineyro:

Should carcinoembryonic antigen be used in the management of patients with colorectal cancer?

Dis. Colon Rectum, 42 (1999), 921-929.

Chen, C.J., T.T. Lin und J.E. Shively:

Role of interferon regulatory factor-1 in the induction of biliary glycoprotein (cell CAM-1) by interferon-gamma.
J. Biol. Chem., 271 (1996), 28181-28188.

Chen, T., W. Zimmermann, J. Parker, I. Chen, A. Maeda und S. Bolland:

Biliary glycoprotein (BGPa, CD66a, CEACAM1) mediates inhibitory signals.
J. Leukoc. Biol., 70 (2001), 335-340.

Cheung, P.H., O. Culic, Y. Qiu, K. Earley, N. Thompson, D.C. Hixson und S.H. Lin:

The cytoplasmic domain of C-CAM is required for C-CAM-mediated adhesion function: studies of a C-CAM transcript containing an unspliced intron.
Biochem. J., 295 (1993a), 427-435.

Cheung, P.H., W. Luo, Y. Qiu, X. Zhang, K. Earley, P. Millirons und S.H. Lin:

Structure and function of C-CAM1. The first immunoglobulin domain is required for intercellular adhesion.
J. Biol. Chem., 268 (1993b), 24303-24310.

Cheung, P.H., N.L. Thompson, K. Earley, O. Culic, D. Hixson und S.H. Lin:

Cell-CAM105 isoforms with different adhesion functions are coexpressed in adult rat tissues and during liver development.
J. Biol. Chem., 268 (1993c), 6139-6146.

Chiba, R., N. Nakagawa, K. Kurasawa, Y. Tanaka, Y. Saito und I. Iwamoto:

Ligation of CD31 (PECAM-1) on endothelial cells increases adhesive function of alphavbeta3 integrin and enhances beta1 integrin-mediated adhesion of eosinophils to endothelial cells.
Blood, 94 (1999), 1319-1329.

Clark, E.A. und J.S. Brugge:

Integrins and signal transduction pathways: the road taken.
Science, 268 (1995), 233-239.

Cleveland, D.W.:

Tubulin and associated proteins.
In: Guidebook to the Cytoskeletal and Motor Proteins, Kreis, T. and R. Vale (eds.), Oxford University Press, Oxford (1993), 101-105.

Coggeshall, K.M.:

Inhibitory signaling by B cell Fc gamma RIIb.
Curr. Opin. Immunol., 10 (1998), 306-312.

Coligan, J.E., J.T. Lautenschleger, M.L. Egan und C.W. Todd:

Isolation and characterization of carcinoembryonic antigen.

Immunochemistry, 9 (1972), 377-386.

Colucci-Guyon, E., M.M. Portier, I. Dunia, D. Paulin, S. Pournin und C. Babine:

Mice lacking vimentin develop and reproduce without an obvious phenotype.

Cell, 79 (1994), 679-694.

Cooper, G.M.:

The cytoskeleton and cell movement.

In: The Cell. A Molecular Approach, Cooper, G. M. (eds.), ASM Press, Washington (1997), 423-431.

Coulombe, P.A., O. Bousquet, L. Ma, S. Yamada und D. Wirtz:

The 'ins' and 'outs' of intermediate filament organization.

Trends Cell Biol., 10 (2000), 420-428.

Crossin, K.L. und L.A. Krushel:

Cellular signaling by neural cell adhesion molecules of the immunoglobulin superfamily.

Dev. Dyn., 218 (2000), 260-279.

Da Silva-Azevedo, L. und W. Reutter:

The Long Isoform of the Cell Adhesion Molecule C-CAM Binds to Actin.

Biochem. Biophys. Res. Commun., 256 (1999), 404-408.

Dabora, S.L. und M.P. Sheetz:

Cultured cell extracts support organelle movement on microtubules in vitro.

Cell. Motil. Cytoskeleton, 10 (1988), 482-495.

Daeron, M., S. Latour, O. Malbec, E. Espinosa, P. Pina, S. Pasmans und W.H. Fridman:

The same tyrosine-based inhibition motif, in the intracytoplasmic domain of Fc gamma

RIIB, regulates negatively BCR-, TCR-, and FcR-dependent cell activation.

Immunity, 3 (1995), 635-646.

Dagert, M. und S.D. Ehrlich:

Prolonged incubation in calcium chlorid improves the competence of Escherichia coli cells.

Gene, 6 (1979), 23-28.

Danielpour, D.:

Transdifferentiation of NRP-152 rat prostatic basal epithelial cells toward a luminal phenotype: regulation by glucocorticoid, insulin-like growth factor-I and transforming growth factor-&bgr; [In Process Citation].
J. Cell. Sci., 112 (1999), 169-179.

den Hartigh, J.C., P.M. van Bergen en Henegouwen, A.J. Verkleij und J. Boonstra:

The EGF receptor is an actin-binding protein.
J. Cell Biol., 119 (1992), 349-355.

Diamond, M.S., D.E. Staunton, S.D. Marlin und T.A. Springer:

Binding of the integrin Mac-1 (CD11b/CD18) to the third immunoglobulin-like domain of ICAM-1 (CD54) and its regulation by glycosylation.
Cell, 65 (1991), 961-971.

Donda, A., L. Mori, A. Shamshiev, I. Carena, C. Mottet, M.H. Heim, C. Beglinger, F.**Grunert, C. Rochlitz, L. Terracciano, P. Jantschkeff und G. De Libero:**

Locally inducible CD66a (CEACAM1) as an amplifier of the human intestinal T cell response.
Eur. J. Immunol., 30 (2000), 2593-2603.

Draberova, L., H. Cerna, H. Brodska, M. Boubelik, S.M. Watt, C.P. Stanners und P.**Draber:**

Soluble isoforms of CEACAM1 containing the A2 domain: increased serum levels in patients with obstructive jaundice and differences in 3-fucosyl-N-acetyl-lactosamine moiety.
Immunology, 101 (2000), 279-287.

Dveksler, G.S., M.N. Pensiero, C.B. Cardellichio, R.K. Williams, G.S. Jiang, K.V.**Holmes und C.W. Dieffenbach:**

Cloning of the mouse hepatitis virus (MHV) receptor: expression in human and hamster cell lines confers susceptibility to MHV.
J. Virol., 65 (1991), 6881-6891.

Earley, K., W. Luo, Y. Qiu, N.L. Thompson, J. Chou, D.C. Hixson und S.H. Lin:

Identification of a new isoform of cell-cell adhesion molecule 105 (C-CAM), C-CAM4: a secretory protein with only one Ig domain.
Biochem. J., 315 (1996), 799-806.

Ebrahimnejad, A., R. Flayeh, G. Unteregger, C. Wagener und J. Brummer:

Cell adhesion molecule CEACAM1 associates with paxillin in granulocytes and epithelial and endothelial cells.
Exp. Cell Res., 260 (2000), 365-373.

Edelman, E.M.:

Morphoregulatory molecules.
Biochem., 27 (1988), 3533-3543.

Edelman, G.M.:

CAMs and IgGs: cell adhesion and the evolutionary origins of immunity.
Immunol. Rev., 100 (1987), 11-45.

Edlund, M., H. Gaardsvoll, E. Bock und B. Obrink:

Different isoforms and stock-specific variants of the cell adhesion molecule C-CAM (cell-CAM 105) in rat liver.
Eur. J. Biochem., 213 (1993), 1109-1116.

Edlund, M., K. Wikstrom, R. Toomik, P. Ek und B. Obrink:

Characterization of protein kinase C-mediated phosphorylation of the short cytoplasmic domain isoform of C-CAM.
FEBS Lett., 425 (1998), 166-170.

Ergun, S., N. Kilik, G. Ziegeler, A. Hansen, P. Nollau, J. Gotze, J.H. Wurmbach, A.**Horst, J. Weil, M. Fernando und C. Wagener:**

CEA-related cell adhesion molecule 1: a potent angiogenic factor and a major effector of vascular endothelial growth factor.
Mol. Cell, 5 (2000), 311-320.

Estrera, V.T., D.T. Chen, W. Luo, D.C. Hixson und S.H. Lin:

Signal transduction by the CEACAM1 tumor suppressor. Phosphorylation of serine 503 is required for growth-inhibitory activity.
J. Biol. Chem., 276 (2001), 15547-15553.

Folkman, J. und P.A. D'Amore:

Blood vessel formation: what is its molecular basis?
Cell, 87 (1996), 1153-1155.

Fournes, B., S. Sadekova, C. Turbide, S. Letourneau und N. Beauchemin:

The CEACAM1-L Ser503 residue is crucial for inhibition of colon cancer cell tumorigenicity.
Oncogene, 20 (2001), 219-230.

Fuchs, E. und K. Weber:

Intermediate filaments: structure, dynamics, function, and disease.
Annu. Rev. Biochem., 63 (1994), 345-382.

Godfraind, C., S.G. Langreth, C.B. Cardellichio, R. Knobler, J.P. Coutelier, M. Dubois-Dalcq und K.V. Holmes:

Tissue and cellular distribution of an adhesion molecule in the carcinoembryonic antigen family that serves as a receptor for mouse hepatitis virus.
Lab. Invest., 73 (1995), 615-627.

Gold, P. und S.O. Freedman:

Specific carcinoembryonic antigens of the human digestive system.
J. Exp. Med., 122 (1965), 467-481.

Graham, R.A., S. Wang, P.J. Catalano und D.G. Haller:

Postsurgical surveillance of colon cancer: preliminary cost analysis of physician examination, carcinoembryonic antigen testing, chest x-ray, and colonoscopy.
Ann. Surg., 228 (1998), 59-63.

Gray-Owen, S.D., D.R. Lorenzen, A. Haude, T.F. Meyer und C. Dehio:

Differential Opa specificities for CD66 receptors influence tissue interactions and cellular response to *Neisseria gonorrhoeae*.
Mol. Microbiol., 26 (1997), 971-980.

Greene, L.A. und A.S. Tischler:

Establishment of a noradrenergic clonal line of rat adrenal pheochromocytoma cells which respond to nerve growth factor.
Proc. Natl. Acad. Sci. U S A, 73 (1976), 2424-2428.

Grunert, F., M. Kuroki und S.C. Stocks:

CEA Family Members Expressed on Hematopoietic Cells and their Possible Role in Cell Adhesion and Signaling.
In: Cell Adhesion and Communication Mediated by the CEA Family, Stanners, C. P. (eds.), OPA, Amsterdam (1998), 99-120.

Hammarström, S., W.N. Khan, S. Teglund, M.L. Hammarström, T. Ramos, V. Baranov, M.M.W. Yeung und L. Frängsmyr:

The Carcinoembryonic antigen family.
In: 2, Structure of Antigens, Van Regenmortel, M. H. V. (eds.), CRC Press, Boca Raton (1993), 341-376.

Hauck, C.R., H. Grassme, J. Bock, V. Jendrossek, K. Ferlinz, T.F. Meyer und E. Gulbins:

Acid sphingomyelinase is involved in CEACAM receptor-mediated phagocytosis of *Neisseria gonorrhoeae*.
FEBS Lett., 478 (2000), 260-266.

Hauck, C.R., T.F. Meyer, F. Lang und E. Gulbins:

CD66-mediated phagocytosis of Opa52 *Neisseria gonorrhoeae* requires a Src-like tyrosine kinase- and Rac1-dependent signalling pathway.
EMBO J., 17 (1998), 443-454.

Hauck, W., P. Nedellec, C. Turbide, C.P. Stanners, T.R. Barnett und N. Beauchemin:

Transcriptional control of the human biliary glycoprotein gene, a CEA gene family member down-regulated in colorectal carcinomas.
Eur. J. Biochem., 223 (1994), 529-541.

Henke, R.C., O. Tolhurst, J.W. Sentry, P. Gunning und P.L. Jeffrey:

Expression of actin and myosin genes during PC12 cell differentiation.
Neurochem. Res., 16 (1991), 675-679.

Higgs, H.N. und T.D. Pollard:

Regulation of actin polymerization by Arp2/3 complex and WASP/Scar proteins.
J. Biol. Chem., 274 (1999), 32531-32534.

Hill, D.J., M.A. Toleman, D.J. Evans, S. Villullas, L. Van Alphen und M. Virji:

The variable P5 proteins of typeable and non-typeable *Haemophilus influenzae* target human CEACAM1.
Mol. Microbiol., 39 (2001), 850-862.

Hitt, A.L., J.H. Hartwig und E.J. Luna:

Ponticulin is the major high affinity link between the plasma membrane and the cortical actin network in *Dictyostelium*.
J. Cell Biol., 126 (1994), 1433-1444.

Hixson, D.C., K.D. McEntire und B. Obrink:

Alterations in the expression of a hepatocyte cell adhesion molecule by transplantable rat hepatocellular carcinomas.
Cancer Res., 45 (1985), 3742-3749.

Hof, P., S. Pluskey, S. Dhe-Paganon, M.J. Eck und S.E. Shoelson:

Crystal structure of the tyrosine phosphatase SHP-2.
Cell, 92 (1998), 441-450.

Horwitz, A., K. Duggan, C. Buck, M.C. Beckerle und K. Burridge:

Interaction of plasma membrane fibronectin receptor with talin--a transmembrane linkage.

Nature, 320 (1986), 531-533.

Howe, A., A.E. Aplin, S.K. Alahari und R.L. Juliano:

Integrin signaling and cell growth control.

Curr. Opin. Cell Biol., 10 (1998), 220-231.

Hsieh, J.T., W. Luo, W. Song, Y. Wang, D.I. Kleinerman, N.T. Van und S.H. Lin:

Tumor suppressive role of an androgen-regulated epithelial cell adhesion molecule (C-CAM) in prostate carcinoma cell revealed by sense and antisense approaches.

Cancer Res., 55 (1995), 190-197.

Huang, J., J.D. Hardy, Y. Sun und J.E. Shively:

Essential role of biliary glycoprotein (CD66a) in morphogenesis of the human mammary epithelial cell line MCF10F.

J. Cell. Sci., 112 (1999), 4193-4205.

Hubbard, A.K. und R. Rothlein:

Intercellular adhesion molecule-1 (ICAM-1) expression and cell signaling cascades.

Free Radic. Biol. Med., 28 (2000), 1379-1386.

Huber, M., L. Izzi, P. Grondin, C. Houde, T. Kunath, A. Veillette und N. Beauchemin:

The carboxyl-terminal region of biliary glycoprotein controls its tyrosine phosphorylation and association with protein-tyrosine phosphatases SHP-1 and SHP-2 in epithelial cells.

J. Biol. Chem., 274 (1999), 335-344.

Hunter, I., H. Sawa, M. Edlund und B. Obrink:

Evidence for regulated dimerization of cell-cell adhesion molecule (C- CAM) in epithelial cells.

Biochem. J., 320 (1996), 847-853.

Hunter, T.:

Signaling - 2000 and beyond.

Cell, 100 (2000), 113-127.

Ilan, N., A. Mohsenin, L. Cheung und J.A. Madri:

PECAM-1 shedding during apoptosis generates a membrane-anchored truncated molecule with unique signaling characteristics.

FASEB J., 15 (2001), 362-372.

Iwabuchi, K., S. Yamamura, A. Prinetti, K. Handa und S. Hakomori:

GM3-enriched microdomain involved in cell adhesion and signal transduction through carbohydrate-carbohydrate interaction in mouse melanoma B16 cells.
J. Biol. Chem., 273 (1998), 9130-9138.

Izzi, L., C. Turbide, C. Houde, T. Kunath und N. Beauchemin:

cis-Determinants in the cytoplasmic domain of CEACAM1 responsible for its tumor inhibitory function.
Oncogene, 18 (1999), 5563-5572.

Jeng, R.L. und M.D. Welch:

Cytoskeleton: actin and endocytosis - no longer the weakest link.
Curr. Biol., 11 (2001), R691-694.

Jirsa, M., L. Muchova, L. Draberova, P. Draber, F. Smid, M. Kuroki, Z. Marecek und A.K. Groen:

Carcinoembryonic antigen-related cell adhesion molecule 1 is the 85-kilodalton pronase-resistant biliary glycoprotein in the cholesterol crystallization promoting low density protein-lipid complex.
Hepatology, 34 (2001), 1075-1082.

Kammerer, R., S. Hahn, B.B. Singer, J.S. Luo und S. von Kleist:

Biliary glycoprotein (CD66a), a cell adhesion molecule of the immunoglobulin superfamily, on human lymphocytes: structure, expression and involvement in T cell activation.
Eur. J. Immunol., 28 (1998), 3664-3674.

Kammerer, R., D. Stober, B.B. Singer, B. Obrink und J. Reimann:

Carcinoembryonic antigen-related cell adhesion molecule 1 on murine dendritic cells is a potent regulator of T cell stimulation.
J. Immunol., 166 (2001), 6537-6544.

Kannicht, C., L. Lucka, R. Nuck, W. Reutter und M. Gohlke:

N-Glycosylation of the carcinoembryonic antigen related cell adhesion molecule, C-CAM, from rat liver: detection of oversialylated bi- and triantennary structures.
Glycobiology, 9 (1999), 897-906.

Kataoka, K., Y. Takata, A. Nakajima, S. Saito und N. Huh:

A carcinoembryonic antigen family cDNA from mouse placenta encoding a protein with a rare domain composition.
Placenta, 21 (2000), 610-614.

Katz, B.Z., S. Levenberg, K.M. Yamada und B. Geiger:

Modulation of cell-cell adherens junctions by surface clustering of the N-cadherin cytoplasmic tail.

Exp. Cell Res., 243 (1998), 415-424.

Keck, U., P. Nedellec, N. Beauchemin, J. Thompson und W. Zimmermann:

The cea10 gene encodes a secreted member of the murine carcinoembryonic antigen family and is expressed in the placenta, gastrointestinal tract and bone marrow.

Eur. J. Biochem., 229 (1995), 455-464.

Kleinerman, D.I., C.P. Dinney, W.W. Zhang, S.H. Lin, N.T. Van und J.T. Hsieh:

Suppression of human bladder cancer growth by increased expression of C-CAM1 gene in an orthotopic model.

Cancer Res., 56 (1996), 3431-3435.

Kleinerman, D.I., P. Troncoso, S.H. Lin, L.L. Pisters, E.R. Sherwood, T. Brooks, A.C. von Eschenbach und J.T. Hsieh:

Consistent expression of an epithelial cell adhesion molecule (C-CAM) during human prostate development and loss of expression in prostate cancer: implication as a tumor suppressor.

Cancer Res., 55 (1995), 1215-1220.

Koch, P.J. und W.W. Franke:

Desmosomal cadherins: another growing multigene family of adhesion molecules.

Curr. Opin. Cell Biol., 6 (1994), 682-687.

Kodelja, V., K. Lucas, S. Barnert, S. von Kleist, J.A. Thompson und W. Zimmermann:

Identification of a carcinoembryonic antigen gene family in the rat. Analysis of the N-terminal domains reveals immunoglobulin-like, hypervariable regions.

J. Biol. Chem., 264 (1989), 6906-6912.

Krebs, E.G. und J.A. Beavo:

Phosphorylation-dephosphorylation of enzymes.

Annu. Rev. Biochem., 48 (1979), 923-959.

Kuijpers, T.W., M. Hoogerwerf, L.J. van der Laan, G. Nagel, C.E. van der Schoot, F. Grunert und D. Roos:

CD66 nonspecific cross-reacting antigens are involved in neutrophil adherence to cytokine-activated endothelial cells.

J. Cell Biol., 118 (1992), 457-466.

Kuroki, M., F. Arakawa, Y. Matsuo, S. Oikawa, H. Nakazato und Y. Matsuoka:

Three novel molecular forms of biliary glycoprotein deduced from cDNA clones from a human leukocyte library.

Biochim. Biophys. Res. Commun., 176 (1991), 578-585.

Laemmli, U.K.:

Cleavage of structural proteins during the assembly of the head of bacteriophage T4.

Nature, 227 (1970), 680-685.

Lasky, L.A.:

Selectin-carbohydrate interactions and the initiation of the inflammatory response.

Annu. Rev. Biochem., 64 (1995), 113-139.

Lee, C., M. Ferguson und L.B. Chen:

Construction of the endoplasmic reticulum.

J. Cell Biol., 109 (1989), 2045-2055.

Leung-Hagesteijn, C.Y., K. Milankov, M. Michalak, J. Wilkins und S. Dedhar:

Cell attachment to extracellular matrix substrates is inhibited upon downregulation of expression of calreticulin, an intracellular integrin alpha-subunit-binding protein.

J. Cell. Sci., 107 (1994), 589-600.

Leventhal, P.S. und E.L. Feldman:

Tyrosine phosphorylation and enhanced expression of paxillin during neuronal differentiation in vitro.

J. Biol. Chem., 271 (1996), 5957-5960.

Li, W., R. Nishimura, A. Kashishian, A.G. Batzer, W.J. Kim, J.A. Cooper und J.**Schlessinger:**

A new function for a phosphotyrosine phosphatase: linking GRB2-Sos to a receptor tyrosine kinase.

Mol. Cell. Biol., 14 (1994), 509-517.

Lin, S.H. und G. Guidotti:

Cloning and expression of a cDNA coding for a rat liver plasma membrane ecto-ATPase.

The primary structure of the ecto-ATPase is similar to that of the human biliary glycoprotein I.

J. Biol. Chem., 264 (1989), 14408-14414.

Lucka, L., M. Budt, I. Cichocka, K. Danker, R. Horstkorte und W. Reutter:

C-CAM-mediated adhesion leads to an outside-in dephosphorylation signal.

Eur. J. Biochem., 262 (1999), 541-546.

Lucka, L., I. Cichocka, K. Bäumler, K. Bechler und W. Reutter:

A short isoform of carcinoembryonic-antigen-related rat liver cell-cell adhesion molecule (C-CAM/gp110) mediates intercellular adhesion. Sequencing and recombinant functional analysis.

Eur. J. Biochem., 234 (1995), 527-535.

Lucka, L., S. Sel, K. Danker, R. Horstkorte und W. Reutter:

Carcinoembryonic antigen-related cell-cell adhesion molecule C-CAM is greatly increased in serum and urine of rats with liver diseases.

FEBS Lett., 438 (1998), 37-40.

Luna, E.J. und A.L. Hitt:

Cytoskeleton--plasma membrane interactions.

Science, 258 (1992), 955-964.

Mandel, M. und A. Higa:

Calcium dependent bacteriophage DNA infection.

J. Biol. Chem., 53 (1970), 154.

Margolis, R.N., M.J. Schell, S.I. Taylor und A.L. Hubbard:

Hepatocyte plasma membrane ECTO-ATPase (pp120/HA4) is a substrate for tyrosine kinase activity of the insulin receptor.

Biochem. Biophys. Res. Commun., 166 (1990), 562-566.

Michely, B.:

Struktur, Expression und Funktion des Zell-Adhäsionsmoleküls CEACAM1 in Rattenbrust-Adenocarcinom-Zelllinien.

Dissertation, Freie Universität Berlin, Fachbereich Humanmedizin (2000).

Mitchison, T. und M. Kirschner:

Dynamic instability of microtubule growth.

Nature, 312 (1984), 237-242.

Moller, M.J., R. Kammerer, F. Grunert und S. von Kleist:

Biliary glycoprotein (BGP) expression on T cells and on a natural-killer-cell subpopulation.

Int. J. Cancer, 65 (1996), 740-745.

Morales, V.M., A. Christ, S.M. Watt, H.S. Kim, K.W. Johnson, N. Utku, A.M. Texeira, A. Mizoguchi, E. Mizoguchi, G.J. Russell, S.E. Russell, A.K. Bhan, G.J. Freeman und R.S. Blumberg:

Regulation of human intestinal intraepithelial lymphocyte cytolytic function by biliary glycoprotein (CD66a).
J. Immunol., 163 (1999), 1363-1370.

Morris, H.P. und L.J. Slaughter:

Historical development of transplantable hepatomas.
Adv. Exp. Med. Biol., 92 (1977), 1-19.

Mowery, J. und D.C. Hixson:

Detection of cell-CAM 105 in the pericanalicular domain of the rat hepatocyte plasma membrane.
Hepatology, 13 (1991), 47-56.

Muenzner, P., C. Dehio, T. Fujiwara, M. Achtman, T.F. Meyer und S.D. Gray-Owen:

Carcinoembryonic antigen family receptor specificity of *Neisseria meningitidis* Opa variants influences adherence to and invasion of proinflammatory cytokine-activated endothelial cells.
Infect. Immun., 68 (2000), 3601-3607.

Muenzner, P., M. Naumann, T.F. Meyer und S.D. Gray-Owen:

Pathogenic *Neisseria* trigger expression of their carcinoembryonic antigen-related cellular adhesion molecule 1 (CEACAM1; previously CD66a) receptor on primary endothelial cells by activating the immediate early response transcription factor, nuclear factor-kappaB.
J. Biol. Chem., 276 (2001), 24331-24340.

Mullins, R.D. und T.D. Pollard:

Structure and function of the Arp2/3 complex.
Curr. Opin. Struct. Biol., 9 (1999), 244-249.

Mullis, K.B. und F.A. Faloona:

Specific synthesis of DNA in vitro via a polymerase-catalysed chain reaction.
Methods. in Enzymol., 155 (1987), 335-350.

Najjar, S.M., D. Accili, N. Philippe, J. Jernberg, R. Margolis und S.I. Taylor:

pp120/ecto-ATPase, an endogenous substrate of the insulin receptor tyrosine kinase, is expressed as two variably spliced isoforms.
J. Biol. Chem., 268 (1993), 1201-1206.

Najjar, S.M., N. Philippe, Y. Suzuki, G.A. Ignacio, P. Formisano, D. Accili und S.I.

Taylor:

Insulin-stimulated phosphorylation of recombinant pp120/HA4, an endogenous substrate of the insulin receptor tyrosine kinase.

Biochemistry, 34 (1995), 9341-9349.

Nakajima, A., H. Iijima, M.F. Neurath, T. Nagaishi, E.E. Nieuwenhuis, R.

Raychowdhury, J. Glickman, D.M. Blau, S. Russell, K.V. Holmes und R.S.

Blumberg:

Activation-induced expression of carcinoembryonic antigen-cell adhesion molecule 1 regulates mouse T lymphocyte function.

J. Immunol., 168 (2002), 1028-1035.

Nap, M., M.L. Hammarstrom, O. Bormer, S. Hammarstrom, C. Wagener, S. Handt, M.

Schreyer, J.P. Mach, F. Buchegger, S. von Kleist und et al.:

Specificity and affinity of monoclonal antibodies against carcinoembryonic antigen.

Cancer Res., 52 (1992), 2329-2339.

Nedellec, P., G.S. Dveksler, E. Daniels, C. Turbide, B. Chow, A.A. Basile, K.V. Holmes

und N. Beauchemin:

Bgp2, a new member of the carcinoembryonic antigen-related gene family, encodes an alternative receptor for mouse hepatitis viruses.

J. Virol., 68 (1994), 4525-4537.

Noguchi, T., T. Matozaki, K. Horita, Y. Fujioka und M. Kasuga:

Role of SH-PTP2, a protein-tyrosine phosphatase with Src homology 2 domains, in insulin-stimulated Ras activation.

Mol. Cell. Biol., 14 (1994), 6674-6682.

Obrink, B.:

CEA adhesion molecules: multifunctional proteins with signal-regulatory properties.

Curr. Opin. Cell Biol., 9 (1997), 616-626.

Ocklind, C., K. Rubin und B. Obrink:

Different cell surface glycoproteins are involved in cell-cell and cell-collagen adhesion of rat hepatocytes.

FEBS Lett., 121 (1980), 47-50.

Odin, P. und B. Obrink:

Dynamic expression of the cell adhesion molecule cell-CAM 105 in fetal and regenerating rat liver.

Exp. Cell Res., 164 (1986), 103-114.

Odin, P. und B. Obrink:

Quantitative determination of the organ distribution of the cell adhesion molecule cell-CAM 105 by radioimmunoassay.
Exp. Cell Res., 171 (1987), 1-15.

Odin, P. und B. Obrink:

The cell-surface expression of the cell adhesion molecule cellCAM 105 in rat fetal tissues and regenerating liver.
Exp. Cell Res., 179 (1988), 89-103.

Ohmori, T., Y. Yatomi, Y. Wu, M. Osada, K. Satoh und Y. Ozaki:

Wheat germ agglutinin-induced platelet activation via platelet endothelial cell adhesion molecule-1: involvement of rapid phospholipase C gamma 2 activation by Src family kinases.

Biochemistry, 40 (2001), 12992-13001.

Ohwada, A., H. Takahashi, I. Nagaoka und S. Kira:

Biliary glycoprotein mRNA expression is increased in primary lung cancer, especially in squamous cell carcinoma.

Am. J. Respir. Cell. Mol. Biol., 11 (1994), 214-220.

Oikawa, S., M. Kuroki, Y. Matsuoka, G. Kosaki und H. Nakazato:

Homotypic and heterotypic Ca(++)-independent cell adhesion activities of biliary glycoprotein, a member of carcinoembryonic antigen family, expressed on CHO cell surface.

Biochem. Biophys. Res. Commun., 186 (1992), 881-887.

Olsen, A., S. Teglund, D. Nelson, L. Gordon, A. Copeland, A. Georgescu, A. Carrano und S. Hammarstrom:

Gene organization of the pregnancy-specific glycoprotein region on human chromosome 19: assembly and analysis of a 700-kb cosmid contig spanning the region.
Genomics, 23 (1994), 659-668.

Ong, S.H., G.R. Guy, Y.R. Hadari, S. Laks, N. Gotoh, J. Schlessinger und I. Lax:

FRS2 proteins recruit intracellular signaling pathways by binding to diverse targets on fibroblast growth factor and nerve growth factor receptors.
Mol. Cell. Biol., 20 (2000), 979-989.

Otey, C.A., F.M. Pavalko und K. Burridge:

An interaction between alpha-actinin and the beta 1 integrin subunit in vitro.
J. Cell Biol., 111 (1990), 721-729.

Owens, G.C., G.M. Edelman und B.A. Cunningham:

Organization of the neural cell adhesion molecule (N-CAM) gene: alternative exon usage as the basis for different membrane-associated domains.
Proc. Natl. Acad. Sci. U S A, 84 (1987), 294-298.

Pavalko, F.M., D.M. Walker, L. Graham, M. Goheen, C.M. Doerschuk und G.S.**Kansas:**

The cytoplasmic domain of L-selectin interacts with cytoskeletal proteins via alpha-actinin: receptor positioning in microvilli does not require interaction with alpha-actinin.
J. Cell Biol., 129 (1995), 1155-1164.

Paves, H., T. Neuman, M. Metsis und M. Saarma:

Nerve growth factor induces rapid redistribution of F-actin in PC12 cells.
FEBS Lett., 235 (1988), 141-143.

Pawson, T. und G.D. Gish:

SH2 and SH3 domains: from structure to function.
Cell, 71 (1992), 359-362.

Perrotti, N., D. Accili, B. Marcus-Samuels, R.W. Rees-Jones und S.I. Taylor:

Insulin stimulates phosphorylation of a 120-kDa glycoprotein substrate (pp120) for the receptor-associated protein kinase in intact H-35 hepatoma cells.
Proc. Natl. Acad. Sci. USA, 84 (1987), 3137-3140.

Petit, V. und J.P. Thiery:

Focal adhesions: structure and dynamics.
Biol. Cell, 92 (2000), 477-494.

Phan, D., X. Sui, D.T. Chen, S.M. Najjar, G. Jenster und S.H. Lin:

Androgen regulation of the cell-cell adhesion molecule-1 (Ceacam1) gene.
Mol. Cell. Endocrinol., 184 (2001), 115-123.

Pollard, T.D., S. Almo, S. Quirk, V. Vinson und E.E. Lattman:

Structure of actin binding proteins: insights about function at atomic resolution.
Annu. Rev. Cell Biol., 10 (1994), 207-249.

Pollard, T.D. und J.A. Cooper:

Actin and actin-binding proteins. A critical evaluation of mechanisms and functions.
Annu. Rev. Biochem., 55 (1986), 987-1035.

Pollard, T.D. und M.S. Mooseker:

Direct measurement of actin polymerization rate constants by electron microscopy of actin filaments nucleated by isolated microvillus cores.

J. Cell Biol., 88 (1981), 654-659.

Poy, M.N., R.J. Ruch, M.A. Fernstrom, Y. Okabayashi und S.M. Najjar:

Shc and CEACAM1 interact to regulate the mitogenic action of insulin.

J. Biol. Chem., 277 (2002a), 1076-1084.

Poy, M.N., Y. Yang, K. Rezaei, M.A. Fernstrom, A.D. Lee, Y. Kido, S.K. Erickson und**S.M. Najjar:**

CEACAM1 regulates insulin clearance in liver.

Nat. Genet., 19 (2002b), 19-26.

Prall, F., P. Nollau, M. Neumaier, H.D. Haubeck, Z. Drzeniek, U. Helmchen, T. Loning**und C. Wagener:**

CD66a (BGP), an adhesion molecule of the carcinoembryonic antigen family, is expressed in epithelium, endothelium, and myeloid cells in a wide range of normal human tissues.

J. Histochem. Cytochem., 44 (1996), 35-41.

Rebstock, S., K. Lucas, J.A. Thompson und W. Zimmermann:

cDNA and gene analyses imply a novel structure for a rat carcinoembryonic antigen-related protein.

J. Biol. Chem., 265 (1990), 7872-7879.

Rivera, D.L., S.M. Oilater, X. Liu, J.H. Thompson, X.J. Zhang, K. Pennline, R. Azuero,**D.A. Clark und M.J. Miller:**

Interleukin-10 attenuates experimental fetal growth restriction and demise.

FASEB J., 12 (1998), 189-197.

Robinson, D.R., Y.M. Wu und S.F. Lin:

The protein tyrosine kinase family of the human genome.

Oncogene, 19 (2000), 5548-5557.

Rosen, S.D. und C.R. Bertozzi:

The selectins and their ligands.

Curr. Opin. Cell Biol., 6 (1994), 663-673.

Ruoslahti, E. und B. Obrink:

Common principles in cell adhesion.

Exp. Cell Res., 227 (1996), 1-11.

Ruoslahti, E. und M.D. Pierschbacher:

New perspectives in cell adhesion: RGD and integrins.
Science, 238 (1987), 491-497.

Russ, W.P. und D.M. Engelman:

The G_{xxx}G motif: a framework for transmembrane helix-helix association.
J. Mol. Biol., 296 (2000), 911-919.

Rybakova, I.N., K.J. Amann und J.M. Ervasti:

A new model for the interaction of dystrophin with F-actin.
J. Cell Biol., 135 (1996), 661-672.

Sadekova, S., N. Lamarche-Vane, X. Li und N. Beauchemin:

The CEACAM1-L glycoprotein associates with the actin cytoskeleton and localizes to cell-cell contact through activation of Rho-like GTPases.
Mol. Biol. Cell, 11 (2000), 65-77.

Saiki, R.K., S. Scharf, F. Saloona, K.B. Mullis, G.T. Horn, H.A. Erlich und N. Arnheim:

Enzymatic amplification of β-globin genomic sequence and restriction site analysis for diagnosis of sickle cell anemia.
Science, 230 (1985), 1350-1354.

Sanger, F., S. Nicklen und A.R. Coulson:

DNA sequencing with chain-terminating inhibitors.
Proc. Natl. Acad. Sci. USA, 74 (1977), 5463-5467.

Sano, M. und M. Iwanaga:

Requirement for specific protein kinase activities during the rapid redistribution of F-actin that precedes the outgrowth of neurites in PC12D cells.
Cell. Struct. Funct., 17 (1992), 341-350.

Sawa, H., K. Kamada, H. Sato, S. Sendo, A. Kondo, I. Saito, M. Edlund und B. Obrink:

C-CAM expression in the developing rat central nervous system.
Brain. Res. Dev. Brain. Res., 78 (1994), 35-43.

Schmucker, D., J.C. Clemens, H. Shu, C.A. Worby, J. Xiao, M. Muda, J.E. Dixon und S.L. Zipursky:

Drosophila Dscam is an axon guidance receptor exhibiting extraordinary molecular diversity.
Cell, 101 (2000), 671-684.

Schumann, D., C.J. Chen, B. Kaplan und J.E. Shively:

Carcinoembryonic antigen cell adhesion molecule 1 directly associates with cytoskeleton proteins actin and tropomyosin.

J. Biol. Chem., 276 (2001), 47421-47433.

Senes, A., M. Gerstein und D.M. Engelman:

Statistical analysis of amino acid patterns in transmembrane helices: the GxxxG motif occurs frequently and in association with beta-branched residues at neighboring positions.

J. Mol. Biol., 296 (2000), 921-936.

Serrador, J.M., J.L. Alonso-Lebrero, M.A. del Pozo, H. Furthmayr, R. Schwartz-Albiez, J. Calvo, F. Lozano und F. Sanchez-Madrid:

Moesin interacts with the cytoplasmic region of intercellular adhesion molecule-3 and is redistributed to the uropod of T lymphocytes during cell polarization.

J. Cell Biol., 138 (1997), 1409-1423.

Serrador, J.M., M. Vicente-Manzanares, J. Calvo, O. Barreiro, M.C. Montoya, R. Schwartz-Albiez, H. Furthmayr, F. Lozano und F. Sanchez-Madrid:

A novel serine-rich motif in the intercellular adhesion molecule 3 is critical for its ERM-directed subcellular targeting.

J. Biol. Chem., 9 (2002), 9.

Shimoyama, Y., S. Hirohashi, S. Hirano, M. Noguchi, Y. Shimosato, M. Takeichi und O. Abe:

Cadherin cell-adhesion molecules in human epithelial tissues and carcinomas.

Cancer Res., 49 (1989), 2128-2133.

Sicheri, F., I. Moarefi und J. Kuriyan:

Crystal structure of the Src family tyrosine kinase Hck.

Nature, 385 (1997), 602-609.

Sikorski, M.A., D.E. Staunton und J.W. Mier:

L-Selectin crosslinking induces integrin-dependent adhesion: evidence for a signaling pathway involving PTK but not PKC.

Cell Adhes. Commun., 4 (1996), 355-367.

Sinclair, N.R.:

Why so many coinhibitory receptors?

Scand. J. Immunol., 50 (1999), 10-13.

Singer, B.B., I. Scheffrahn und B. Obrink:

The tumor growth-inhibiting cell adhesion molecule CEACAM1 (C-CAM) is differently expressed in proliferating and quiescent epithelial cells and regulates cell proliferation.
Cancer Res., 60 (2000), 1236-1244.

Skubitz, K.M., K.D. Campbell, K. Ahmed und A.P. Skubitz:

CD66 family members are associated with tyrosine kinase activity in human neutrophils.
J. Immunol., 155 (1995), 5382-5390.

Skubitz, K.M., K.D. Campbell und A.P. Skubitz:

CD66a, CD66b, CD66c, and CD66d each independently stimulate neutrophils.
J. Leukoc. Biol., 60 (1996), 106-117.

Springer, T.A.:

Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm.
Cell, 76 (1994), 301-314.

Stein-Gerlach, M., C. Wallasch und A. Ullrich:

SHP-2, SH2-containing protein tyrosine phosphatase-2.
Int. J. Biochem. Cell Biol., 30 (1998), 559-566.

Sundberg, U. und B. Obrink:

CEACAM1 isoforms with different cytoplasmic domains show different localization, organization and adhesive properties in polarized epithelial cells.
J. Cell. Sci., 115 (2002), 1273-1284.

Svenberg, T.:

Carcinoembryonic antigen-like substances of human bile, isolation and partial characterisation.
Int. J. Cancer, 17 (1976), 588-596.

Svenberg, T., B. Wahren und S. Hammarstrom:

Elevated serum levels of a biliary glycoprotein (BGP I) in patients with liver or biliary tract disease.
Clin. Exp. Immunol., 36 (1979), 317-325.

Taguchi, F. und S. Matsuyama:

Soluble receptor potentiates receptor-independent infection by murine coronavirus.
J. Virol., 76 (2002), 950-958.

Takahashi, H., Y. Okai, R.J. Paxton, L.J. Hefta und J.E. Shively:

Differential regulation of carcinoembryonic antigen and biliary glycoprotein by gamma-interferon.

Cancer Res., 53 (1993), 1612-1619.

Takeichi, M.:

Cadherins in cancer: implications for invasion and metastasis.

Curr. Opin. Cell Biol., 5 (1993), 806-811.

Takeichi, M.:

Morphogenetic roles of classic cadherins.

Curr. Opin. Cell Biol., 7 (1995), 619-627.

Tapley, P., A. Horwitz, C. Buck, K. Duggan und L. Rohrschneider:

Integrins isolated from Rous sarcoma virus-transformed chicken embryo fibroblasts.

Oncogene, 4 (1989), 325-333.

Tedder, T.F., D.A. Steeber, A. Chen und P. Engel:

The selectins: vascular adhesion molecules.

FASEB J., 9 (1995), 866-873.

Teglund, S., A. Olsen, W.N. Khan, L. Frangsmyr und S. Hammarstrom:

The pregnancy-specific glycoprotein (PSG) gene cluster on human chromosome 19: fine structure of the 11 PSG genes and identification of 6 new genes forming a third subgroup within the carcinoembryonic antigen (CEA) family.

Genomics, 23 (1994), 669-684.

Towbin, H., T. Staehlin und J. Gordon:

Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications.

Proc. Natl. Acad. Sci. USA, 76 (1979), 4530-4354.

Tsutsumi, Y., N. Onoda, M. Misawa, M. Kuroki und Y. Matsuoka:

Immunohistochemical demonstration of nonspecific cross-reacting antigen in normal and neoplastic human tissues using a monoclonal antibody. Comparison with carcinoembryonic antigen localization.

Acta Pathol. Jpn., 40 (1990), 85-97.

Tuma, P.L., C.M. Finnegan, J.H. Yi und A.L. Hubbard:

Evidence for apical endocytosis in polarized hepatic cells: phosphoinositide 3-kinase inhibitors lead to the lysosomal accumulation of resident apical plasma membrane proteins.

J. Cell Biol., 145 (1999), 1089-1102.

Turbide, C., T. Kunath, E. Daniels und N. Beauchemin:

Optimal ratios of biliary glycoprotein isoforms required for inhibition of colonic tumor cell growth.

Cancer Res., 57 (1997), 2781-2788.

van der Geer, P. und T. Pawson:

The PTB domain: a new protein module implicated in signal transduction.

Trends Biochem. Sci., 20 (1995), 277-280.

Vandekerckhove, J. und K. Vancompernolle:

Structural relationships of actin-binding proteins.

Curr. Opin. Cell Biol., 4 (1992), 36-42.

Venetianer, A., D.L. Schiller, T. Magin und W.W. Franke:

Cessation of cytokeratin expression in a rat hepatoma cell line lacking differentiated functions.

Nature, 305 (1983), 730-733.

Virji, M., D. Evans, J. Griffith, D. Hill, L. Serino, A. Hadfield und S.M. Watt:

Carcinoembryonic antigens are targeted by diverse strains of typable and non-typable Haemophilus influenzae.

Mol. Microbiol., 36 (2000), 784-795.

Virji, M., K. Makepeace, D.J. Ferguson und S.M. Watt:

Carcinoembryonic antigens (CD66) on epithelial cells and neutrophils are receptors for Opa proteins of pathogenic neisseriae.

Mol. Microbiol., 22 (1996), 941-950.

Wang, J., S.D. Gray-Owen, A. Knorre, T.F. Meyer und C. Dehio:

Opa binding to cellular CD66 receptors mediates the transcellular traversal of Neisseria gonorrhoeae across polarized T84 epithelial cell monolayers.

Mol. Microbiol., 30 (1998), 657-671.

Watt, S.M., A.M. Teixeira, G.Q. Zhou, R. Doyonnas, Y. Zhang, F. Grunert, R.S.

Blumberg, M. Kuroki, K.M. Skubitz und P.A. Bates:

Homophilic adhesion of human CEACAM1 involves N-terminal domain interactions:
structural analysis of the binding site.

Blood, 98 (2001), 1469-1479.

Weiss, B., S. Jacquenmin, T.R. Live, G.C. Fareed und C.C. Richardson:

Enzymatic breakage and joining of deoxyribonucleic acid.

J. Biol. Chem., 243 (1968), 4543.

Wessells, J., D. Wessner, R. Parsells, K. White, D. Finkenzeller, W. Zimmermann und G. Dveksler:

Pregnancy specific glycoprotein 18 induces IL-10 expression in murine macrophages.
Eur. J. Immunol., 30 (2000), 1830-1840.

Williams, A.F. und A.N. Barclay:

The immunglobuline superfamily-domains for cell surface recognition.

Ann. Rev. Immunol., 6 (1988), 381-405.

Williams, J.C., A. Weijland, S. Gonfloni, A. Thompson, S.A. Courtneidge, G. Superti-Furga und R.K. Wierenga:

The 2.35 Å crystal structure of the inactivated form of chicken Src: a dynamic molecule with multiple regulatory interactions.

J. Mol. Biol., 274 (1997), 757-775.

Wright, J.H., P. Drueckes, J. Bartoe, Z. Zhao, S.H. Shen und E.G. Krebs:

A role for the SHP-2 tyrosine phosphatase in nerve growth-induced PC12 cell differentiation.

Mol. Biol. Cell, 8 (1997), 1575-1585.

Yaffe, M.B. und L.C. Cantley:

Signal transduction. Grabbing phosphoproteins.

Nature, 402 (1999), 30-31.

Yonemura, S., M. Hirao, Y. Doi, N. Takahashi, T. Kondo und S. Tsukita:

Ezrin/radixin/moesin (ERM) proteins bind to a positively charged amino acid cluster in the juxta-membrane cytoplasmic domain of CD44, CD43, and ICAM-2.

J. Cell Biol., 140 (1998), 885-895.

Zelus, B.D., D.R. Wessner, G.S. Dveksler und K.V. Holmes:

Neutralization of MHV-A59 by soluble recombinant receptor glycoproteins.

Adv. Exp. Med. Biol., 440 (1998), 3-9.

Zhao, T. und P.J. Newman:

Integrin activation by regulated dimerization and oligomerization of platelet endothelial cell adhesion molecule (PECAM)-1 from within the cell.

J. Cell Biol., 152 (2001), 65-73.

Zhou, G.Q. und S. Hammarstrom:

Pregnancy-specific glycoprotein (PSG) in baboon (*Papio hamadryas*): family size, domain structure, and prediction of a functional region in primate PSGs.

Biol. Reprod., 64 (2001), 90-99.

Zhou, G.Q., Y. Zhang und S. Hammarstrom:

The carcinoembryonic antigen (CEA) gene family in non-human primates.

Gene, 264 (2001), 105-112.