

Probability that  $n$  random points are in convex  
position<sup>◇</sup>

Pavel Valtr\*

B 94-01  
January 1994

**Abstract**

We show that  $n$  random points chosen independently and uniformly from a parallelogram are in convex position with probability

$$\left( \frac{\binom{2n-2}{n-1}}{n!} \right)^2.$$

<sup>◇</sup>The work on this paper was supported by the “Deutsche Forschungsgemeinschaft”, grant We 1265/2-1.

\*Department of Applied Mathematics, Charles University, Malostranské nám. 25, 118 00 Praha 1, Czech Republic and

Graduierertenkolleg “Algorithmische Diskrete Mathematik”, Fachbereich Mathematik und Informatik, Freie Universität Berlin, Takustrasse 9, D-14195 Berlin, Germany

# 1 The result

A finite set of points in the plane is called *convex* if its points are vertices of a convex polygon. In this paper we show the following result.

**Theorem 1** *The set  $A$  of  $n$  random points chosen independently and uniformly from a parallelogram  $S$  is convex with probability*

$$\left( \frac{\binom{2n-2}{n-1}}{n!} \right)^2.$$

A large part of studies in stochastic geometry deals with the convex hull  $C$  of a set of  $n$  points placed independently and uniformly in a fixed convex body  $K$  in  $\mathbb{R}^d$ . Typical questions are: How many vertices does  $C$  have? What is the volume of  $C$ ? What is the surface area of  $C$ ? See [WW] for a survey. In this paper we settle one very special case – the probability that  $C$  has  $n$  vertices in the case  $K$  is a parallelogram. It is interesting that our approach is purely combinatorial, with no use of integration. We think that our method based on an approximation of the uniform distribution in a square by a large grid might have other applications. However, it is already not clear how to apply our method for  $K$  a triangle or in three dimensions.

In this section we prove Theorem 1, and in the next section we mention some applications of Theorem 1.

*Proof of Theorem 1.* Let  $n > 2$  be a fixed integer. Since a proper affine transformation transfers the uniform distribution on  $S$  onto the uniform distribution on a square, we may and shall assume that  $S$  is a square. We shall approximate the square  $S$  by a grid whose size tends to infinity.

Let  $m$  be a positive integer (denoting the size of the grid). Partition the (axis-parallel) square  $S$  by  $m - 1$  horizontal and by  $m - 1$  vertical lines into  $m^2$  squares  $S_1, \dots, S_{m^2}$  of equal size. The centers of the squares  $S_1, \dots, S_{m^2}$  form a square grid  $m \times m$ . Every point of  $A$  lies in each of the squares  $S_1, \dots, S_{m^2}$  with the same probability  $1/m^2$ . Move every point of  $A$  to the center of the square  $S_i$  in which it lies, and denote the obtained multiset by  $A(m)$ . It is not difficult to see that

$$\text{Prob}(A \text{ is convex}) = \lim_{m \rightarrow \infty} \text{Prob}(A(m) \text{ is convex}).$$

Thus,

$$\text{Prob}(A \text{ is convex}) = \lim_{m \rightarrow \infty} \text{Prob}(R_m \text{ is convex}),$$

where, for every  $m \geq 1$ ,  $R_m$  is a multiset of  $n$  points chosen randomly and independently from the square grid  $G_m = \{(i, j) : i, j = 1, 2, \dots, m\}$  (each point of  $G_m$  is always taken with the same probability  $1/m^2$ ).

Let  $\mathcal{M}(G_m)$  be the set of all multisets of size  $n$  with elements from  $G_m$ , and let  $\mathcal{C}(G_m)$  be the set of all convex  $n$ -element subsets of  $G_m$ . It is easy to see that

$$\text{Prob}(A \text{ is convex}) = \lim_{m \rightarrow \infty} \text{Prob}(R_m \text{ is convex}) = \lim_{m \rightarrow \infty} \frac{|\mathcal{C}(G_m)|}{|\mathcal{M}(G_m)|} = \lim_{m \rightarrow \infty} \frac{|\mathcal{C}(G_m)|}{\binom{m^2}{n}}.$$

In the sequel we shall estimate the size of  $\mathcal{C}(G_m)$ .

Every convex set  $R \in \mathcal{C}(G_m)$  is uniquely defined by the smallest axis-parallel rectangle  $Q(R)$  containing  $R$  and by the set  $V(R)$  of the  $n$  integer vectors forming the boundary of the convex hull of  $R$  oriented in counterclockwise order.

Let  $X(R)$  and  $Y(R)$  be the multisets of the first and of the second coordinates of vectors in  $V(R)$ , respectively. Formally,

$$X(R) = \bigcup_{(x,y) \in V(R)} \{x\}, \quad Y(R) = \bigcup_{(x,y) \in V(R)} \{y\}.$$

Let  $\mathcal{C}'(G_m)$  be the set of all convex sets  $R \in \mathcal{C}(G_m)$  such that  $0 \notin X(R) \cup Y(R)$  and that the directions of the  $n^2$  vectors  $(x, y)$  formed by all the  $n^2$  pairs  $x \in X(R), y \in Y(R)$  are distinct. Thus, in particular, the multisets  $X(R)$  and  $Y(R)$  are sets for any  $R \in \mathcal{C}'(G_m)$ . It is not difficult to see that

$$\lim_{m \rightarrow \infty} \frac{|\mathcal{C}'(G_m)|}{|\mathcal{C}(G_m)|} = 1.$$

Therefore,

$$\text{Prob}(A \text{ is convex}) = \lim_{m \rightarrow \infty} \frac{|\mathcal{C}(G_m)|}{\binom{m^2}{n}} = \lim_{m \rightarrow \infty} \frac{|\mathcal{C}'(G_m)|}{\binom{m^2}{n}}.$$

In the estimation of the size of  $\mathcal{C}'(G_m)$  we use an auxiliary set  $\mathcal{S}$  defined by

$$\mathcal{S} = \{(X(R), Y(R), Q(R)) : R \in \mathcal{C}'(G_m)\}.$$

The following construction shows that, for every  $(X, Y, Q) \in \mathcal{S}$ , there are exactly  $n!$  sets  $R \in \mathcal{C}'(G_m)$  with  $(X(R), Y(R), Q(R)) = (X, Y, Q)$ :

Take any of the  $n!$  one-to-one correspondences  $f : X \rightarrow Y$  between  $X$  and  $Y$ , and define a set  $V$  of  $n$  vectors by  $V = \{(x, f(x)) : x \in X\}$ . Due to the definitions of  $\mathcal{C}'(G_m)$  and  $\mathcal{S}$ , vectors in  $V$  have distinct directions and, consequently, form the (counterclockwise oriented) boundary of the convex hull of a unique set  $R \in \mathcal{C}'(G_m)$  fitting into the rectangle  $Q$ .

Thus,

$$|\mathcal{C}'(G_m)| = n! \cdot |\mathcal{S}|$$

and

$$\text{Prob}(A \text{ is convex}) = \lim_{m \rightarrow \infty} \frac{|\mathcal{C}'(G_m)|}{\binom{m^2}{n}} = \lim_{m \rightarrow \infty} \frac{n! \cdot |\mathcal{S}|}{\binom{m^2}{n}}.$$

It remains to estimate the size of the set  $\mathcal{S}$  which is done in the sequel technical part of the proof.

For  $(X, Y, Q) \in \mathcal{S}$ , partition each of the two sets  $X$  and  $Y$  into two subsets containing elements with the same sign:

$$X^+ = \{x \in X : x > 0\}, \quad X^- = \{x \in X : x < 0\},$$

$$Y^+ = \{y \in Y : y > 0\}, \quad Y^- = \{y \in Y : y < 0\}.$$

Suppose that each of the sets  $X^+, X^-, Y^+, Y^-$  is ordered in an arbitrary way. Denote  $s = |X^+|$  and  $t = |Y^+|$ . Thus,

$$X^+ = \{x_1, \dots, x_s\}, \quad X^- = \{x_{s+1}, \dots, x_n\},$$

$$Y^+ = \{y_1, \dots, y_t\}, \quad Y^- = \{y_{t+1}, \dots, y_n\}.$$

For every  $(X, Y, Q) \in \mathcal{S}$ , where  $Q = \{(x, y) : a_1 \leq x \leq a_2, b_1 \leq y \leq b_2\}$ , the orders on the sets  $X^+, X^-, Y^+, Y^-$  uniquely determine four sets  $D^-, E^-, D^+, E^+$  of integers from the set  $\{1, 2, \dots, m\}$  in the following way:

$$D^+ = \{a_1 + \sum_{i=1}^k x_i : k = 0, 1, \dots, s\}, \quad D^- = \{a_2 + \sum_{i=s+1}^k x_i : k = s, s+1, \dots, n\},$$

$$E^+ = \{b_1 + \sum_{i=1}^k y_i : k = 0, 1, \dots, t\}, \quad E^- = \{b_2 + \sum_{i=t+1}^k y_i : k = t, t+1, \dots, n\}.$$

Note that the sets  $D^-, E^-, D^+, E^+$  satisfy the following conditions:

$$|D^+| + |D^-| = n + 2, \quad a_1 = \min D^+ = \min D^-, \quad a_2 = \max D^+ = \max D^-, \quad (1)$$

$$|E^+| + |E^-| = n + 2, \quad b_1 = \min E^+ = \min E^-, \quad b_2 = \max E^+ = \max E^-. \quad (2)$$

For any  $(X, Y, Q) \in \mathcal{S}$ , we obtain  $|X^+|!|X^-|!|Y^+|!|Y^-|!$  different 4-tuples of sets  $D^-, E^-, D^+, E^+$  corresponding to different orders on the sets  $X^+, X^-, Y^+, Y^-$ . Denote the set of all these 4-tuples  $(D^-, E^-, D^+, E^+)$  by  $\mathcal{F}(X, Y, Q)$ . Thus,

$$\begin{aligned} |\mathcal{F}(X, Y, Q)| &= |X^+|!|X^-|!|Y^+|!|Y^-|! \\ &= (|D^+| - 1)!(|D^-| - 1)!(|E^+| - 1)!(|E^-| - 1)!, \end{aligned}$$

where  $(D^-, E^-, D^+, E^+)$  is an arbitrary 4-tuple in  $\mathcal{F}(X, Y, Q)$ . For  $0 \leq i \leq n - 2$  and  $0 \leq j \leq n - 2$ , we say that a 4-tuple  $(D^-, E^-, D^+, E^+)$  of sets of integers has property  $\mathcal{P}_{i,j}$  if

$$\mathcal{P}_{i,j}: |D^+| = i + 2, |E^+| = j + 2, \text{ and the sets } D^-, E^-, D^+, E^+ \text{ satisfy (1) and (2) for some } 1 \leq a_1 < a_2 \leq m \text{ and } 1 \leq b_1 < b_2 \leq m.$$

There are  $\binom{n-2}{i} \binom{m}{n} \cdot \binom{n-2}{j} \binom{m}{n}$  4-tuples  $(D^-, E^-, D^+, E^+)$  with  $\mathcal{P}_{i,j}$  and  $|D^+ \cap D^-| = |E^+ \cap E^-| = 2$ . It follows that there are  $(1 + o(1)) \cdot \binom{n-2}{i} \binom{m}{n} \cdot \binom{n-2}{j} \binom{m}{n}$  4-tuples  $(D^-, E^-, D^+, E^+)$  with  $\mathcal{P}_{i,j}$ . (Throughout the proof,  $o(1)$  denotes functions of  $m$

which tend to 0 as  $m$  tends to infinity.) Most of them (i.e., a  $(1 - o(1))$ -fraction of them) lie in the disjoint union

$$\bigcup_{(X,Y,Q) \in \mathcal{S}} \mathcal{F}(X,Y,Q).$$

Thus,

$$\begin{aligned} |\mathcal{S}| &= \sum_{(X,Y,Q) \in \mathcal{S}} 1 = \sum_{(X,Y,Q) \in \mathcal{S}} \frac{|\mathcal{F}(X,Y,Q)|}{|X^+|!|X^-|!|Y^+|!|Y^-|!} = \\ &= \sum_{i=0}^{n-2} \sum_{j=0}^{n-2} \frac{(1 - o(1)) \cdot (1 + o(1)) \binom{n-2}{i} \binom{m}{n} \binom{n-2}{j} \binom{m}{n}}{(i+1)!(n-i-1)!(j+1)!(n-j-1)!} = \\ &= (1 + o(1)) \binom{m}{n}^2 \cdot \sum_{i=0}^{n-2} \sum_{j=0}^{n-2} \frac{\binom{n}{n-i-1} \cdot \binom{n}{n-j-1}}{(n!)^2} \binom{n-2}{i} \binom{n-2}{j} = \\ &= (1 + o(1)) \binom{m}{n}^2 \frac{1}{(n!)^2} \left( \sum_{i=0}^{n-2} \binom{n}{n-i-1} \binom{n-2}{i} \right) \left( \sum_{j=0}^{n-2} \binom{n}{n-j-1} \binom{n-2}{j} \right) = \\ &= (1 + o(1)) \binom{m}{n}^2 \frac{1}{(n!)^2} \binom{2n-2}{n-1}^2. \end{aligned}$$

Hence,

$$\begin{aligned} \text{Prob}(A \text{ is convex}) &= \lim_{m \rightarrow \infty} \frac{n! \cdot |\mathcal{S}|}{\binom{m^2}{n}} = \lim_{m \rightarrow \infty} \frac{(1 + o(1)) \binom{m}{n}^2 \frac{1}{n!} \binom{2n-2}{n-1}^2}{\binom{m^2}{n}} = \\ &= \left( \frac{\binom{2n-2}{n-1}}{n!} \right)^2. \end{aligned}$$

□

## 2 Applications and related results

In this section we sketch some applications of our result.

1. *Replacing the parallelogram by a convex body.* It is known that, for every bounded convex body  $K$  in the plane, there are two parallelograms, one containing  $K$  and one contained in  $K$ , whose areas differ from the area of  $K$  at most by a constant factor (e.g., see [Ba] for analogous results). Using this result and Theorem 1, it is not difficult to show that there are two positive constants  $c_1$  and  $c_2$  such that the set of  $n$  points chosen independently and uniformly from an arbitrary convex body is convex with probability at least  $\left(\frac{c_1}{n}\right)^n$  and at most  $\left(\frac{c_2}{n}\right)^n$ .

2. *The expected area of a random triangle.* It is not difficult to show that

$$\text{Prob}(A \text{ is convex}) + 4 \cdot E[\text{Area of } T] = 1,$$

where  $A$  is a set of four random points selected independently and uniformly from a convex body  $S$  of area 1, and  $T$  is a triangle with random vertices selected also independently and uniformly from  $S$ . If  $S$  is a parallelogram, Theorem 1 yields that the expected area of  $T$  is

$$\frac{1 - (5/6)^2}{4} = \frac{11}{144},$$

which was also shown in [He] by a different method.

3. *Convex subsets of a random set.* The author originally considered Theorem 1 in connection with the following result.

**Theorem 2** *Let  $A$  be a set of  $n$  random points chosen independently and uniformly from a parallelogram. Let  $c(A)$  be the largest convex subset of  $A$ . Set  $h = 2^{4/3}e \approx 6.85$ . Then  $c(A) \geq \lambda n^{1/3}$  with probability smaller than  $(\frac{h}{\lambda})^{3\lambda n^{1/3}}$ , for any  $\lambda \geq h$ .*

*Proof.* Let  $\lambda \geq h$ . For simplicity, assume that  $\lambda n^{1/3}$  is an integer. The set  $A$  contains  $\binom{n}{\lambda n^{1/3}}$  subsets of size  $\lambda n^{1/3}$ . According to Theorem 1, each of them is convex with probability

$$\left( \frac{\binom{2\lambda n^{1/3}-2}{\lambda n^{1/3}-1}}{(\lambda n^{1/3})!} \right)^2.$$

It follows that the expected number of convex independent subsets of  $A$  of size  $\lambda n^{1/3}$  is at most

$$\begin{aligned} \binom{n}{\lambda n^{1/3}} \cdot \left( \frac{\binom{2\lambda n^{1/3}-2}{\lambda n^{1/3}-1}}{(\lambda n^{1/3})!} \right)^2 &< \frac{n^{\lambda n^{1/3}}}{\left(\frac{\lambda n^{1/3}}{e}\right)^{\lambda n^{1/3}}} \cdot \left( \frac{4^{\lambda n^{1/3}}}{\left(\frac{\lambda n^{1/3}}{e}\right)^{\lambda n^{1/3}}} \right)^2 = \\ &= \left( \frac{4^2 e^3}{\lambda^3} \right)^{\lambda n^{1/3}} = \left( \frac{h}{\lambda} \right)^{3\lambda n^{1/3}}. \end{aligned}$$

Consequently,  $A$  contains a convex independent subset of size  $\geq \lambda n^{1/3}$  with probability smaller than  $(\frac{h}{\lambda})^{3\lambda n^{1/3}}$ .  $\square$

One application of Theorem 2 on so-called dense sets may be found in the author's PhD. thesis [Va].

By a more careful handling with the result of Theorem 1, one can prove that, for any  $\varepsilon > 0$  and any sufficiently large  $n \geq n(\varepsilon)$ ,

$$(h/2 - \varepsilon)n^{1/3} \leq c(A) \leq hn^{1/3}$$

holds with a high probability.

4. *Construction of random convex sets.* Emo Welzl pointed out that the above proof of Theorem 1 yields a fast way how to construct a random convex set of size  $n$  in a square. Let  $M_n$  be the set of all  $n$ -element subsets of a square  $S$ , and let  $\mu$  be the probabilistic measure on  $M_n$  corresponding to a choice of  $n$  points selected independently and uniformly from the square  $S$ . Let  $C_n$  be the set of all convex  $n$ -element subsets of  $S$ . Theorem 1 gives  $\mu(C_n) = \left(\frac{\binom{2n-2}{n-1}}{n!}\right)^2$ . The measure  $\mu' = \mu/(\mu(C_n))|_{C_n}$  is a probabilistic measure on  $C_n$ . With respect to  $\mu'$ , a random convex set  $A \in C_n$  can be constructed in a straightforward way by repeated choosings of an  $n$ -point random subset  $A$  of  $S$  with respect to  $\mu$ , until the set  $A$  is convex. However, this procedure has the expected running time at least  $\Omega(R \cdot (n!/ \binom{2n-2}{n-1})^2) = \Omega(R \cdot (n/4e)^{2n+1})$ , where  $R$  is the time required for finding a random real number uniformly distributed in the interval  $[0, 1]$ . The above proof of Theorem 1 yields a procedure which constructs a random convex set with respect to  $\mu'$  essentially faster, in time  $\mathcal{O}(n \log n + n \cdot R + P(n))$ , where  $R$  is as above and  $P(n)$  is the time required for constructing a random permutation of the set  $\{1, 2, \dots, n\}$ . Of course, the argument also applies for any parallelogram.

5. *The limit shape of a random convex set.* Scale and shift the square grid  $n \times n$  so that it fits into the square  $S = \{(x, y) : -1 \leq x, y \leq 1\}$ , and consider the set  $K(n)$  of all its convex subsets. Bárány [Bá] proved that for every  $\varepsilon > 0$  there exists  $n_0$  such that for every  $n \geq n_0$  the following holds: if we randomly choose an element  $A$  of  $K(n)$ , each with the same probability, then the Hausdorff distance between the boundary of the convex hull of  $A$  and the curve  $\{(x, y) : \sqrt{1 - |x|} + \sqrt{1 - |y|} = 1\}$  is smaller than  $\varepsilon$  with a high probability. (Hausdorff distance between two sets is the maximum distance of a point in any of the two sets to the other set.)

It is interesting that random convex sets have the same limit shape. Consider the square  $S = \{(x, y) : -1 \leq x, y \leq 1\}$  again, and define  $C_n$  and  $\mu'$  as in the above paragraph “*Construction of random convex sets*”. With a help of the above proof of Theorem 1, it can be shown that for every  $\varepsilon > 0$  there exists  $n_0$  such that for every  $n \geq n_0$  the following holds: if we randomly choose an element  $A$  of  $C_n$  with respect to the measure  $\mu'$ , then the Hausdorff distance between the boundary of the convex hull of  $A$  and the curve  $\{(x, y) : \sqrt{1 - |x|} + \sqrt{1 - |y|} = 1\}$  is smaller than  $\varepsilon$  with a high probability.

Let us note that the limit shape curve of the boundary of the convex hull of  $n$  random points chosen independently and uniformly inside any planar convex body  $K$  is (obviously) the perimeter of  $K$ .

**Acknowledgment.** I would like to thank Imre Bárány for interesting discussions which lead to this paper, and Stephan Brandt for valuable comments which helped me to improve the style of this paper.

## References

- [Ba] K. Ball, Volume ratios and a reverse isoperimetric inequality, *J. London Math. Soc.* (2) 44 (1991), 351-359.
- [Bá] I. Bárány, The limit shape theorem for convex lattice polygons, to appear.
- [He] N. Henze, Random triangles in convex regions, *J. Appl. Prob.* 20 (1983), 111-125.
- [Va] P. Valtr, Planar point sets with bounded ratios of distances, PhD. thesis, Free Univ. Berlin (1994).
- [WW] W. Weil and J.A. Wieacker, Stochastic geometry, Chapter 5.2 in: P.M. Gruber and J.M. Wills (eds.), *Handbook of Convex Geometry, II*, North-Holland (1993), pp. 1393-1438.