

Generalized Davenport-Schinzel Sequences

Martin Klazar*
Pavel Valtr**

B 94-04
January 1994

Abstract

The extremal function $Ex(u, n)$ (introduced in the theory of Davenport-Schinzel sequences in other notation) denotes for a fixed finite alternating sequence $u = ababa \dots$ the maximum length of a finite sequence v over n symbols with no immediate repetition which does not contain u . Here (following the idea of J. Nešetřil) we generalize this concept for arbitrary sequence u . We summarize the already known properties of $Ex(u, n)$ and we present also two new theorems which give good upper bounds on $Ex(u, n)$ for u consisting of (two) smaller subsequences u_i provided we have good upper bounds on $Ex(u_i, n)$. We use these theorems to describe a wide class of sequences u ("linear sequences") for which $Ex(u, n) = O(n)$. Both theorems are used for obtaining new superlinear upper bounds as well. We partially characterize linear sequences over three symbols. We also present several problems about $Ex(u, n)$.

Key words: Davenport-Schinzel sequence, extremal problem, maximum length

*Department of Applied Mathematics, Charles University, Malostranské nám. 25, 118 00 Praha 1, Czech Republic, E-mail: klazar@cspguk11.bitnet.

**Department of Applied Mathematics, Charles University, Malostranské nám. 25, 118 00 Praha 1, Czech Republic, and Graduiertenkolleg "Algorithmische Discrete Mathematik", Fachbereich Mathematik, Freie Universität Berlin, Takustrasse 9, 14195 Berlin, Germany, E-mail: valtr@inf.fu-berlin.de. Supported by "Deutsche Forschungsgemeinschaft", grant We 1265/2-1.

1 Introduction

In this paper we shall investigate the maximum length $Ex(u, n)$ of finite sequences over n symbols not containing a fixed sequence u . We search for sequences u for which there is a linear upper bound on $Ex(u, n)$. We call them *linear* sequences.

First we give a brief informal overview of results concerning extremal problems of this type (belonging to a branch which could be called "Extremal theory of sequences"). After that all necessary definitions will be introduced. The first section concludes by the formulation of our main result: we present two operations which enable us to derive upper bounds on $Ex(u, n)$ from upper bounds for shorter u . In the second section we summarize the properties of $Ex(u, n)$ which are useful in the proofs. In the third section we show four applications of our operations: we prove the linearity of certain relatively complicated sequences, we show on examples how to derive nonlinear upper bounds on $Ex(u, n)$, we discuss the linearity of sequences over three symbols and we describe which linear sequences we are able to obtain at present. The remaining two sections are devoted to the proofs of the main theorems. In the second and in the third sections we list some problems which might stimulate further research in this area.

1.1 History

Davenport-Schinzel sequences are finite sequences over n symbols with no immediate repetition of the same symbol which contain no five-term alternating subsequence (or, more generally, no alternating subsequence of the length s). Davenport and Schinzel posed [4] the problem to estimate the maximum length of such sequences. They proved in [4] $Ex(ababa, n) = O(n \cdot \log n)$ ($Ex(ababa, n) = O(n \cdot \log n / \log \log n)$ in [5]) and $Ex(ababab \dots, n) = O(n \cdot \exp(\sqrt{n}))$ (s fixed) which was improved by Szemerédi [14] to $Ex(ababab \dots, n) = O(n \cdot \log^* n)$. As usual $\log^* n$ denotes the minimum number of iterations of the power function 2^m (starting with $m = 1$) which are needed to get a number bigger or equal to n . However, the problem whether $Ex(ababa, n) = O(n)$ remained open until 1986 when it was answered [7] by Hart and Sharir in the negative: they proved $Ex(ababa, n) = \Theta(n \cdot \alpha(n))$ where $\alpha(n)$ is the inverse to the Ackerman function and goes to infinity but extremely slowly. Later simpler constructions proving $Ex(ababa, n) = \Omega(n \cdot \alpha(n))$ were found ([11], [15]). M. Sharir [12] derived the upper bounds ($ababab \dots$ is of the length s)

$$Ex(ababab \dots, n) = O(n \cdot \alpha(n)^{O(\alpha(n)^{s-5})}).$$

Agarwal, Sharir and Shor [3] found almost tight upper and lower bounds:

$$Ex(ababab \dots, n) \leq n \cdot 2^{(\alpha(n))^{\frac{s-5}{2}} \log_2 \alpha(n) + C_s(n)} \quad \text{for } s \geq 5 \text{ odd}$$

$$Ex(ababab \dots, n) \leq n \cdot 2^{(\alpha(n))^{\frac{s-4}{2}} + C_s(n)} \quad \text{for } s \geq 6 \text{ even}$$

$$Ex(ababab \dots, n) = \Omega(n \cdot 2^{K_s \cdot (\alpha(n))^{\frac{s-4}{2}} + Q_s(n)}) \quad \text{for } s \geq 6 \text{ even}$$

where $K_s = \frac{1}{(\frac{s-4}{2})!}$ and the functions $C_s(n)$ and $Q_s(n)$ are asymptotically smaller than the main terms in the exponent. For $s = 6$ they found a stronger estimate $Ex(ababab, n) = \Theta(n \cdot 2^{\alpha(n)})$.

Füredi and Hajnal [6] investigated a similar problem what is the maximum number of 1's in a 0-1 matrix of the size $n \times n$ if some configurations are forbidden.

The primary motivation of Davenport-Schinzel sequences was geometrical and now they play an important role in computational geometry. See the books [2] and [13] for more information and references.

The function $Ex(u, n)$ extending functions $Ex(ababa, n)$ resp. $Ex(ababab \dots, n)$ was defined in [1]. Note here that this definition which follows in the next subsection was suggested by J. Nešetřil. Some other results from [1] will be mentioned in the second section. In [8] it was proved that

$$Ex(u, n) = n \cdot 2^{O(\alpha(n)^{|u|-4})}$$

($|u|$ denotes the length of u) for any fixed finite sequence u . Thus, from the practical point of view, $Ex(u, n)$ has a "linear" upper bound for any u . In this paper we study the question for which u actually $Ex(u, n) = O(n)$ and we give a partial answer to it. It is easy to prove that it holds for $u = abab$ but it is not so easy to prove the same thing for $u = aabbaabb$. The more general sequence

$$x_1 \dots x_1 x_2 \dots x_2 \dots x_k \dots x_k x_1 \dots x_1 x_2 \dots x_2 \dots x_k \dots x_k$$

is linear as well ([9]) but here we prove stronger results.

1.2 Definitions

For any finite sequence u we denote by $|u|$ the *length* of u , by $S(u)$ the set of all symbols occurring in u and by $\|u\|$ the size of $S(u)$. Thus $\|u\| \leq |u|$ for all u . The sequences for which equality is achieved are called *chains*. Hence in chains no symbol repeats.

Two sequences $u = a_1 a_2 \dots a_n$ and $v = b_1 b_2 \dots b_n$ of the same length are *equivalent* if there exists a bijection $f : S(u) \rightarrow S(v)$ such that $f(a_i) = b_i$ for all $i = 1, 2, \dots, n$. Thus u and v coincide after a renaming of symbols. The notation $u \subseteq v$ means that u is a *subsequence* of v . We say that v *contains* u , formally $u \prec v$, if u is equivalent to some t , $t \subseteq v$.

We shall refer to the occurrences of a symbol $a \in S(u)$ in the sequence u as to *a-occurrences*. The sequence $u = aa \dots a$, $|u| = i$ is denoted shortly by a^i .

The mirror image $a_n \dots a_2 a_1$ of the sequence $u = a_1 a_2 \dots a_n$ is denoted by \bar{u} . The sequence $u = a_1 a_2 \dots a_m$ is called *k-regular* if $a_i = a_j, i \neq j$ implies $|i - j| \geq k$. Thus any at most k consecutive elements in u are different to each other. Davenport-Schinzel sequences are 2-regular.

Definition 1.2.1

$$Ex(u, n) = \max\{|v| \mid u \not\prec v, v \text{ is } \|u\|\text{-regular, } \|v\| \leq n\}$$

where u is a fixed sequence, v is arbitrary sequence and $n \geq 1$ is an integer.

We investigate the behaviour of this function for u fixed and n growing to infinity. Sometimes a more general definition will be useful:

Definition 1.2.2

$$Ex(u, n, l) = \max\{|v| \mid u \not\prec v, v \text{ is } l\text{-regular}, \|v\| \leq n\}$$

where $l \geq \|u\|$ is an integer.

Obviously $Ex(u, n) = Ex(u, n, l) = \|u\| - 1$ for any chain u and $Ex(u, n, l) \geq n$ for any nonchain u . Also $Ex(a^i, n) = (i - 1)n$ and $Ex(abab, n) = 2n - 1$. Our interest is focused on the set

Definition 1.2.3

$$Lin = \{u \mid Ex(u, n) = O(n)\}.$$

We call its elements *linear* sequences. Other sequences, for instance $ababa$, are called *nonlinear*.

1.3 Results

The following two theorems are the main result of this paper.

Theorem A Suppose that $u = u_1a^2u_2$ and v are two sequences such that $S(u) \cap S(v) = \emptyset$ and a is a symbol. If v is not a chain then $Ex(u_1avau_2, n) = O(Ex(v, 2Ex(u, n)))$. In the case v is a chain $Ex(u_1avau_2, n) \leq Ex(u, n)$ holds.

Theorem B Suppose a, b are two symbols and $u = u_1a^2u_2a$ is a sequence such that $b \notin S(u)$. Then $Ex(u_1ab^iau_2ab^i, n) = \Theta(Ex(u, n))$ for any $i \geq 1$.

We are interested in linear sequences. For them it follows immediately

Consequence A Let $u, v \in Lin$ be as in Theorem A. Then $u_1avau_2 \in Lin$.

Consequence B Let $u \in Lin$ and b be as in Theorem B. Then $u_1ab^iau_2ab^i \in Lin$ for all $i \geq 1$.

We use both consequences in the obvious manner: by repeated applications of both transformations $u, v \rightarrow u_1avau_2$ and $u \rightarrow u_1ab^iau_2ab^i$ we can generate a wide class of linear sequences.

2 Properties of $\text{Ex}(u, n)$

Fact 2.1 *Suppose two sequences u and v and two integers $l > k \geq \|u\|$ are given. Then for any $n \geq 1$*

1. $\text{Ex}(u, n, k)$ is finite and $\text{Ex}(u, n, k) = O(\|u\| \cdot \|u\| \cdot n^{\|u\|})$.
2. $\text{Ex}(u, n, l) \leq \text{Ex}(u, n, k) \leq (\text{Ex}(u, l-1, k) + 1)\text{Ex}(u, n, l)$.
3. If $v \prec u$ then $\text{Ex}(v, n) = O(\text{Ex}(u, n))$. Especially if u is linear then v is linear too.
4. $\text{Ex}(u, n) = \text{Ex}(\bar{u}, n)$.

Proof: The first claim follows from Pigeon-Hole argumentation (we mentioned in the Introduction that a far stronger estimate holds ([8])). The second inequality in the second claim is obtained by deleting occurrences by a greedy algorithm. For both proofs we refer to [1]. The first inequality as well as the fourth claim are obvious. We prove the third claim. Suppose w is a $\|u\|$ -regular sequence not containing v . Hence it cannot contain u and therefore $\text{Ex}(v, n, \|u\|) \leq \text{Ex}(u, n, \|u\|) = \text{Ex}(u, n)$. The second inequality in 2. yields the estimate

$$\text{Ex}(v, n) = \text{Ex}(v, n, \|v\|) = O(\text{Ex}(v, n, \|u\|) = O(\text{Ex}(u, n))).$$

□

The reason why we exclude from the following considerations chains is that the extremal function of any chain is constant and hence for these singular sequences the nice estimates are not valid.

Fact 2.2 *Suppose u, v and w are three sequences, $i \geq 2, j \geq 2$ are two integers and a is a symbol such that au is not a chain. Then for $n \geq 1$*

1. $\text{Ex}(a^i u, n) = \Theta(\text{Ex}(au, n)), \text{Ex}(ua^i, n) = \Theta(\text{Ex}(ua, n))$.
2. $\text{Ex}(wa^j v, n) = \Theta(\text{Ex}(wa^2 v, n))$.

Proof: See in [1].

□

Let us reformulate Fact 2.2. Suppose the sequence u is written in the "exponential form" $u = a_1^{i_1} a_2^{i_2} \dots a_r^{i_r}$, $a_i \neq a_{i+1}, i_j \geq 1$ (note that $i_1 = i_2 = \dots = i_r = 1$ iff u is 2-regular). The reduced sequence is defined by $\text{red}(u) = a_1^{j_1} a_2^{j_2} \dots a_r^{j_r}$ where $j_1 = j_r = 1$ and $j_k = \min\{2, i_k\}$ for $1 < k < r$. The fully reduced sequence is defined by $\text{fred}(u) = a_1 a_2 \dots a_r$.

Fact 2.3 $\text{Ex}(u, n) = \Theta(\text{Ex}(\text{red}(u), n))$ for any sequence u and any integer $n \geq 1$ provided that $\text{red}(u)$ is not a chain.

Proof: Follows from the previous Fact 2.2. \square

There is the question whether $Ex(u, n) = \Theta(Ex(fred(u), n))$ (suppose $fred(u)$ is not a chain) which is equivalent to the following problem.

Problem 2.1 Does $Ex(ua^2v, n) = O(Ex(uav, n))$ hold for any symbol a and all sequences u, v (provided uav is not a chain)?

Fact 2.4 Suppose $i \geq 1$ is an integer, u is a sequence over two symbols ($\|u\| \leq 2$) and a, b are two symbols. Then

1. $a^i b^i a^i b^i$ is linear.
2. $ababa$ is nonlinear.
3. u is nonlinear iff $ababa \prec u$.

Proof: For the proof of the first statement we refer to [1]. Or apply Consequence B to the sequence a^{2i} . The second claim is proved in [7].

We deduce the last statement from the first two. If $ababa \prec u$ then clearly u is nonlinear. If the sequence u is over two symbols and does not contain $ababa$ then clearly $u = a^{m_1} b^{m_2} a^{m_3} b^{m_4}$ where $m_i \geq 0$. Thus $u \prec a^m b^m a^m b^m$ for $m = \max(m_i \mid i = 1, 2, 3, 4)$ and u is linear according to 1. and according to 3. of Fact 2.1. \square

3 Applications of Theorems A and B

3.1 Linearity of $abcdcbabcd$

The sequence $w(k, i) = a_1^i a_2^i \dots a_k^i a_1^i a_2^i \dots a_k^i$ which we mentioned in Introduction is a k -symbol analog to $a^i b^i a^i b^i$. It was proved in [9] by methods different from those presented here that $w(k, i)$ is linear. Now we prove a stronger result.

Theorem 3.1.1 The sequence $z(k, i) = a_1^i a_2^i \dots a_{k-1}^i a_k^i a_{k-1}^i \dots a_2^i a_1^i a_2^i \dots a_{k-1}^i a_k^i$, where the symbols a_1, a_2, \dots, a_k are mutually distinct, is linear for all $i, k \geq 1$.

Proof: By induction on k . For $k = 1$ the sequence $z(1, i) = a^{2i}$ is linear. Suppose that the assertion holds for $k > 1$. Thus $a_1^i a_2^i \dots a_{k-1}^i a_k^{2i} a_{k-1}^i \dots a_2^i a_1^i a_2^i \dots a_{k-1}^i a_k^i \prec z(k, 2i)$ is linear because $z(k, 2i)$ is a linear sequence. Applying Consequence B we conclude that $z(k+1, i)$ is linear as well. \square

3.2 Superlinear bounds

Despite the fact that we stress the linear case here we cannot resist the temptation to present an application of our theorems to nonlinear sequences. The difficulty is that the strong superlinear bounds of [7] and [3] were derived only for the 2-regular case and we need a bit more here. However, it may be checked that the method of [7] giving $Ex(ababa, n) = \Theta(n\alpha(n))$ works for $a^i b^i a^i b^i a^i$ as well and so $Ex(a^i b^i a^i b^i a^i, n) = \Theta(n\alpha(n))$. We present two examples of new strong superlinear upper bounds. It is possible to derive other similar results.

Theorem 3.2.1

$$\Omega(n\alpha(n)) = Ex(abacdcdcab, n) = O(n\alpha^2(n))$$

Proof: The lower bound follows trivially from $ababa \prec abacdcdcab$. The upper bound follows from Theorem A:

$$\begin{aligned} Ex(abacdcdcab, n) &= O(Ex(ababa, 2Ex(aba^2ba, n))) = \\ &= O(O(n\alpha(n)) \cdot \alpha(O(n\alpha(n)))) = O(n\alpha^2(n)) \end{aligned}$$

where $\alpha(O(n\alpha(n))) = O(\alpha(n))$ follows from the extreme slow growth of $\alpha(n)$. \square

Theorem 3.2.2

$$Ex(abacdcbacd, n) = \Theta(n\alpha(n))$$

Proof: This result follows from Theorem B:

$$Ex(abacdcbacd, n) = \Theta(Ex(abac^2abac, n)) = \Theta(Ex(aba^2ba, n)) = \Theta(n\alpha(n)).$$

\square

3.3 Linear sequences over three symbols

The third point of Fact 2.4 gives a complete characterization of linear sequences over two symbols. We would like to obtain a similar characterization for the whole set Lin of linear sequences but this seems to be a hard task. Though we are unable to decide linearity even in the slightly more general case of sequences over three symbols, we give for these sequences a characterization theorem which puts away only few sequences as undecided.

Theorem 3.3.1 *Suppose u is a sequence over three symbols ($\|u\| \leq 3$) and neither u nor \bar{u} contains any of the three sequences*

$$ababa \quad cababcb \quad acbabcb.$$

Then u is a linear sequence.

Proof: Suppose that $ababa \not\prec u$. It is not difficult to show that any 2-regular sequence u , $ababa \not\prec u$, $\|u\| \leq 3$ is contained in one of three sequences $u_1 = ababcba$, $u_2 = abcabac$ and $u_3 = acababc$. Thus $ababa \not\prec u$, $\|u\| \leq 3$ implies $u \prec u_1(i)$ or $u \prec u_2(i)$ or $u \prec u_3(i)$ (for large i) where $u_1(i) = a^i b^i a^i b^i c^i b^i c^i$, similarly for $u_2(i)$ and $u_3(i)$. But, as may be deduced from Consequence B and Fact 2.4.1., $u_1(i)$ and $u_2(i)$ are linear (actually $u_2(i) = z(3, i)$). Hence u is linear or $u \prec u_3(i)$.

Using Consequences A and B it may be proved that any subsequence of $u_3(i)$ which does not contain any of the four sequences $s_1 = cababcba$, $s_2 = acbabcb$, $s_3 = acabacb$ and $s_4 = acababc$ is linear. Thus u is linear or $u \succ s_j$ for some $j \in \{1, 2, 3, 4\}$. But s_1 is equivalent to \bar{s}_4 and s_2 is equivalent to \bar{s}_3 . We are finished. \square

In the opposite direction we are able to say only that $ababa \prec u$ implies the nonlinearity of u . The linearity of three sequences over three symbols $u_3 = acababc$, $s_1 = cababcba$ and $s_2 = acbabcb$ remains open.

Problem 3.3.1 Are u_3, s_1 and s_2 resp. $u_3(i), s_1(i)$ and $s_2(i)$ linear?

3.4 How to get many linear sequences

We conclude this section by a compact description of the widest class of sequences M such that $M \subseteq Lin$ may be proved from above.

Let M be the minimal (to inclusion) set of sequences satisfying the following rules

1. $a^i \in M$ for any symbol a and any integer $i, i \geq 1$.
2. If $u = u_1 a^2 u_2$ and v (a is a symbol) are two sequences of M such that $S(u) \cap S(v) = \emptyset$ then $u_1 a v a u_2$ lies in M as well.
3. If $u = u_1 a^2 u_2 a$ (a is a symbol) is a sequence of M and $b, b \notin S(u)$ is a new symbol then $u_1 a b^i a u_2 a b^i$ lies in M for any $i \geq 1$ as well.
4. If u is a sequence of M and $v, v \prec u$ is another sequence then v lies in M as well.
5. If u is a sequence of M then \bar{u} lies in M as well.

Then

Theorem 3.4.1 $M \subseteq Lin$

Proof: We see immediately that the first and the last two rules preserve linearity and hence it suffices to prove only that the set Lin is closed on the second rule as on the third one. But this is exactly the statement of Consequence A and B. \square

We present, as a concrete example of an application of the previous theorem, three linear sequences. The reader will be surely able to establish how the previous five

rules were used to derive these sequences as well as (s)he will be able to obtain many others.

Example

The following three sequences belong to M and therefore they all are linear.

$ababcbedcdededefefgfg$ $ccaaccaabbdefedefbabb$ $ccaaccabgggbdefedefbbbaabggg$

It is worth noting that $u \in M$ iff $fred(u) \in M$. Hence the answer to Problem 2.1 is affirmative if we restrict ourselves to M . This supports the conjecture that the change of any exponent in u does not influence (except the trivial case of a chain) the growth rate of $Ex(u, n)$.

Problem 3.4.1 *Characterize the set Lin . Are there 2-regular linear sequences over n symbols which are longer than $3n - 2$?*

This bound is achieved for instance by $z(n, 1)$ or by the sequences which are constructed as the first sequence in the example above.

Problem 3.4.2 *Suppose B is the set of all minimal (to \prec) nonlinear sequences. What can be said about the elements of B ? Is B finite?*

The set Lin is closed to \prec ($u \prec v \in Lin \Rightarrow u \in Lin$) and thus B characterizes Lin : $u \notin Lin$ iff there is a sequence v that $v \in B, v \prec u$. The set B serves for Lin as a collection of "forbidden pictures". An immediate observation is that $ababa \in B$. Another observation, nontrivial, is that the set B must contain at least two elements: the simple and nice construction of [15] proving $Ex(ababa, n) = \Omega(n\alpha(n))$ proves also implicitly $Ex(w, n) = \Omega(n\alpha(n))$ where $w = abcbadadabcd$ but $ababa \not\prec w$. Thus there is a sequence $u^*, u^* \neq ababa, u^* \prec w, u^* \in B$. For details we refer to [10].

4 Proof of Theorem A

Suppose a is a symbol and $u = u_1 a^2 u_2$ and v are two sequences such that $S(u) \cap S(v) = \emptyset$. We denote by t the sequence $u_1 a v a u_2$.

We start with the simpler case of Theorem A when v is a chain. Suppose w is a $\|t\|$ -regular sequence not containing t . We show that w cannot even contain u . Let $s \subseteq w$ be equivalent to u . The $\|t\|$ -regularity of w implies that there occur $\|v\|$ distinct symbols between the two "a's" in s which are not elements of $S(s)$. Thus $t \prec w$ which is a contradiction. We get an inequality $Ex(t, n, \|t\|) \leq Ex(u, n, \|t\|)$. Hence $Ex(t, n) = Ex(t, n, \|t\|) \leq Ex(u, n, \|t\|) \leq Ex(u, n, \|u\|) = Ex(u, n)$.

Before proving the first part of Theorem A we give an auxiliary lemma. We say that a nondecreasing integral function $f : \{1, 2, \dots\} \rightarrow \{1, 2, \dots\}$ is *big* if $\sum_i f(n_i) \leq c f(\sum_i n_i)$ for some fixed constant c for any sum of integers $\sum_i n_i$.

Lemma 4.1 *The function $Ex(u, n)$ is big for any nonchain u .*

Proof: We call in this proof a sequence u *irreducible* if there does not exist a nontrivial decomposition $u = u_1u_2$ such that $S(u_1) \cap S(u_2) = \emptyset$. Otherwise we call it *reducible*. It is easy to see that $Ex(u, n)$ is big with constant $c = 1$ for any irreducible u . Indeed, if $n = n_1 + n_2 + \dots + n_m$ then by concatenation of m $\|u\|$ -regular sequences $v_i, u \not\prec v_i, \|v_i\| \leq n_i, |v_i| = Ex(u, n_i), S(v_i) \cap S(v_j) = \emptyset$ we get a $\|u\|$ -regular sequence $v = v_1v_2 \dots v_m, \|v\| \leq n$ not containing u which proves the desired inequality. To manage the case of reducible sequences we need the following claim.

Claim $Ex(u_1u_2, n) = O(Ex(u_1, n) + Ex(u_2, n))$ for any two sequences $u_1, u_2, S(u_1) \cap S(u_2) = \emptyset$ except when $|u_1| = |u_2| = 1$.

Proof: Suppose v is $\|u_1\| + \|u_2\|$ -regular, $|v| \geq c(Ex(u_1, \|v\|) + Ex(u_2, \|v\|))$ for some large fixed constant c . We show that this implies $u_1u_2 \prec v$. We split $v = v_1v_2, |v_1| \doteq |v_2| \doteq \frac{1}{2}c(Ex(u_1, \|v\|) + Ex(u_2, \|v\|))$. There is a sequence $t, t \subseteq v_1$ equivalent to u_1 . After deleting all a -occurrences, $a \in S(t)$ from v_2 we get a $\|u_2\|$ -regular subsequence of v_2 long enough to contain inevitably u_2 . Thus $u_1u_2 \prec v$ which proves the claim.

For a given reducible nonchain sequence u we decompose $u = u_1u_2 \dots u_m$ where u_i are irreducible and $S(u_i)$ are mutually disjoint. The previous claim yields $Ex(u, n) = O(Ex(u_1, n) + Ex(u_2, n) + \dots + Ex(u_m, n))$. Any term of this sum is a big function and $Ex(u_i, n) = O(Ex(u, n))$ (3. of Fact 2.1). Hence the lemma follows. \square

Now we are able to continue in the proof. Let u, v, t be as above, v is a nonchain and w is a $\|t\| = \|u\| + \|v\|$ -regular sequence not containing t . Let $f : \{1, 2, \dots\} \rightarrow \{1, 2, \dots\}$ be a nondecreasing big function that will be specified later. We take the leftmost occurrence in w and add occurrence after occurrence maintaining in every step the condition $|s| \leq f(\|s\|)$ for currently constructed contiguous subsequence s of w . In case we get equality we finish s and start a new s by the next occurrence. On the end we get the decomposition

$$w = w_1w_2 \dots w_hw_0, |w_i| = f(\|w_i\|), i > 0, |w_0| \leq f(\|w_0\|).$$

We need $f(n)$ be sufficiently large but not too much, we need namely:

$$(*) \quad f(1) > Ex(t, 2\|u\| - 2).$$

$$(**) \quad f(n) > \frac{1}{\|v\|}(\|u\| + \|v\|)(Ex(v, n) + \|u\|) \text{ for any } n \geq 1.$$

$$(***) \quad f(n) = O(Ex(v, n)) \text{ for } n \geq 1.$$

Thus the choice $f(n) = dEx(v, n)$ for a large constant d clearly meets all conditions $(*), (**)$ and $(***)$. Obviously $f(n)$ is nondecreasing and big because $Ex(v, n)$ is. For any $i = 1, 2, \dots, h$ let $w_i^*, w_i^* \subseteq w_i$ be such that $\|w_i^*\| = |w_i^*| = \|w_i\|$ (we take for any $x \in S(w_i)$ exactly one x -occurrences). Thus $|w_i^*| \geq 2\|u\| - 1$ according to $(*)$.

Then we choose $w_i^{**}, w_i^{**} \subseteq w_i^* (i > 0)$ such that $|w_i^{**}| \geq |w_i^*| - (\|u\| - 1) > \frac{1}{2}|w_i^*|$ and that the two sequences

$$w_{odd} = w_1^{**} w_3^{**} \dots w_{h-1}^{**}, w_{even} = w_2^{**} w_4^{**} \dots w_h^{**}$$

(suppose for simplicity h is even) are $\|u\|$ -regular. This is achieved by deleting at most $\|u\| - 1$ occurrences from w_i^* which equal to one of the last $\|u\| - 1$ occurrences of w_{i-2}^* . It suffices to show $w_{odd} \not\prec u, w_{even} \not\prec u$. Then conclude:

$$\begin{aligned} |w| &= \sum_{i=1}^h |w_i| + |w_0| \leq \sum_{i=1}^h f(|w_i^*|) + f(\|w_0\|) \leq \sum_{i=1}^h f(2|w_i^{**}|) + f(\|w_0\|) = \\ &= \sum_{i \text{ odd}}^h f(2|w_i^{**}|) + \sum_{i \text{ even}}^h f(2|w_i^{**}|) + f(\|w_0\|) \leq \end{aligned}$$

(bigness of f)

$$\leq cf(2|w_{odd}|) + cf(2|w_{even}|) + f(\|w\|) \leq 2cf(2Ex(u, \|w\|)) + f(\|w\|) =$$

(property (***))

$$= 2cdEx(v, 2Ex(u, \|w\|)) + dEx(v, \|w\|) = O(Ex(v, 2Ex(u, \|w\|))).$$

Suppose now on the contrary that, say, w_{odd} contains u (the sequence w_{even} is treated similarly). Thus $u^* = u_1^*(a^*)^2 u_2^* \subseteq w_{odd}$ is equivalent to u . The two a^* -occurrences must lie in two distinct segments $w_{2i+1}^{**}, w_{2j+1}^{**}, i < j$ for there is no repetition in any w_i^{**} and hence $u_1^* a^* w_{2j} a^* u_2^* \subseteq w$. Now we proceed as in the proof of the claim above, we delete all x -occurrences, $x \in S(u^*)$, from w_{2j} and obtain a $\|v\|$ -regular subsequence $w'_{2j}, w'_{2j} \subseteq w_{2j}$. Clearly

$$|w'_{2j}| \geq \frac{\|v\|}{\|u\| + \|v\|} |w_{2j}| - \|u\|$$

which follows from the fact that the deletion of all occurrences of a symbol from a k -regular sequence s yields a $k - 1$ -regular subsequence s' of s of the length at least $\frac{k-1}{k}(|s| - 1)$. But then, according to (**),

$$\begin{aligned} |w'_{2j}| &> \frac{\|v\|}{\|u\| + \|v\|} \cdot \frac{\|u\| + \|v\|}{\|v\|} (Ex(v, \|w_{2j}\|) + \|u\|) - \|u\| = \\ &= Ex(v, \|w_{2j}\|) \geq Ex(v, \|w'_{2j}\|) \end{aligned}$$

and $v \prec w'_{2j}$. Consequently $t = u_1 a v a u_2 \prec w$ which is a contradiction.

5 Proof of Theorem B

This theorem is in some sense stronger than Theorem A because in Theorem B the sequence $v = b^{2^i}$ is split into two parts which are inserted in two places of u , in the middle and on the end. In theorem A we put simply the whole v in the middle of u . Therefore, one can expect a more complicated proof. We start with three preliminary lemmas.

Lemma 5.1 *Let $l \geq 2$ be an integer and let $\varepsilon, 0 < \varepsilon < 1$ be a constant. Suppose v is a $\lceil \frac{l}{\varepsilon} \rceil$ -regular sequence. Then any $v_1, v_1 \subseteq v, |v_1| \geq \varepsilon|v|$ contains a subsequence $v_2, v_2 \subseteq v_1$ such that*

1. v_2 is l -regular
2. $|v_2| \geq (\frac{\varepsilon}{l})^2|v|$

Proof: Let v and v_1 be as described. We split $v = w_1 w_2 \dots w_h w$ where $|w_i| = \lceil \frac{l}{\varepsilon} \rceil, i = 1, 2, \dots, h, |w| \leq \lceil \frac{l}{\varepsilon} \rceil$. Let

$$A = \{i \mid |v_1 \cap w_i| \leq l - 1\}, a = |A|, B = \{i \mid |v_1 \cap w_i| \geq l\}, b = |B|.$$

Clearly

$$a + b = h = \lfloor \frac{|v|}{\lceil \frac{l}{\varepsilon} \rceil} \rfloor \leq \frac{\varepsilon}{l}|v|.$$

Further $a(l - 1) + b\lceil \frac{l}{\varepsilon} \rceil \geq |v_1| \geq \varepsilon|v|$. Hence $\varepsilon|v| \leq (\frac{\varepsilon}{l}|v| - b)(l - 1) + b\lceil \frac{l}{\varepsilon} \rceil$ and

$$b \geq \frac{\frac{\varepsilon}{l}|v|}{\lceil \frac{l}{\varepsilon} \rceil - l + 1} \geq (\frac{\varepsilon}{l})^2|v|.$$

By appropriately taking one occurrences from any $v_1 \cap w_i, i \in B$ we create an l -regular subsequence $v_2, v_2 \subseteq v_1, |v_2| = b$. \square

Suppose $u = a_1 a_2 \dots a_m$ is a sequence. An *interval* $I = \langle a_j, a_{j+k} \rangle$ in u is any contiguous subsequence $a_j a_{j+1} \dots a_{j+k}, k \geq 1$ of length at least 2. In the case a_j and a_{j+k} are both a -occurrences we call I an *a -interval*. An *ordered sequence* $(u, <)$ is a sequence enriched by a linear order $(S(u), <)$.

Let $(u, <), u = a_1 a_2 \dots a_m$ be an ordered sequence, a_i be an a -occurrences in u ($a \in S(u)$) and $c, d \geq 1$ be two integers. We say that a_i is *(c, d) -covered* (in u) if there is an interval I in u such that

1. $a_i \in I$
2. there are at most c a -occurrences in I
3. there are $2d$ occurrences of d (not necessarily distinct) symbols $x_j \in S(u), j = 1 \dots d, x_j < a$ in I , each of these symbols occurs at least twice in I .

Lemma 5.2 *For any 2-regular ordered sequence $(u, <)$ and any integer $r \geq 2$ either $|u| \leq 720r||u||$ or there are at least $\frac{1}{10}|u|$ occurrences in u which are $(8r, r - 1)$ -covered.*

Proof: We can suppose that $S(u) = \{1, 2, \dots, n\}$ and that j coincides with the standard order of integers. We will define by induction sets U_0, U_1, \dots, U_n of disjoint intervals in u . For any $j = 1, 2, \dots, n$ the set U_j will contain some k -intervals, $k = 1, 2, \dots, j$. First put $U_0 = \emptyset$. Fix j and suppose that the set U_{j-1} have been defined. We split all j -occurrences in u in m $8r$ -tuples T_1, T_2, \dots, T_m and a residual tuple T of the size at most $8r - 1$ so that T_1 consists of the $8r$ leftmost j -occurrences, T_2 consists of the next $8r$ j -occurrences etc. Define

$$\mathcal{S}_j = \{i \mid \text{at least one } j\text{-occurrences of } T_i \text{ is not } (8r, r - 1)\text{-covered} \}.$$

The elements of $T_i, i \in \mathcal{S}_j$ group in $4r$ pairs (x, x') of consecutive elements generating j -intervals $\langle x, x' \rangle$. The set U_j consists of all those intervals $\langle x, x' \rangle$ and of all members of U_{j-1} not intersecting them. Now, crucially, $4r$ intervals $\langle x, x' \rangle$ corresponding to one $T_i, i \in \mathcal{S}_j$ intersect all together at most $(r - 2) + 2 = r$ intervals of U_{j-1} . This holds by the definition of \mathcal{S}_j . Thus

$$|U_j| \geq 4r|\mathcal{S}_j| + |U_{j-1}| - r|\mathcal{S}_j| = 3r|\mathcal{S}_j| + |U_{j-1}|$$

and

$$|U_n| \geq 3r \sum_{j=1}^n |\mathcal{S}_j|.$$

Since u is 2-regular and U_n consists of disjoint k -intervals,

$$|u| \geq 3|U_n| \geq 9r \sum_{j=1}^n |\mathcal{S}_j|.$$

The number of occurrences in u which are $(8r, r - 1)$ -covered is therefore at least (suppose $|u| \geq 720rn$)

$$|u| - \sum_{j=1}^n 8r|\mathcal{S}_j| - (8r - 1)n \geq |u| - \frac{8r}{9r}|u| - \frac{1}{90}|u| = \frac{1}{10}|u|.$$

□

In the remaining lemma we force the symbols x_j to be distinct. Suppose $(u, <)$, $u = a_1 a_2 \dots a_m$ is an ordered sequence, $v \subseteq u$ is a subsequence, $y \in v$ is an a -occurrences ($a \in S(u)$) and $c, d \geq 1$ are two integers. We say that y is strongly (v, c, d) -covered if there is an interval I in u such that

1. $y \in I$

2. there are at most c a -occurrences in $I \cap v$
3. there are $2d$ occurrences of d distinct symbols $x_j \in S(u), j = 1 \dots d, x_j < a$ in I , any of these symbols occurs at least twice in I .

Lemma 5.3 *Suppose two integers $k \geq 2, d \geq 1$ are given. Then there exist two integers $l \geq 2, c \geq 1$ and two positive constants $\varepsilon, \Delta > 0$ such that for any l -regular ordered sequence $(u, <)$ one of the following statements holds.*

1. $|u| \leq \Delta \|u\|$
2. there is a k -regular subsequence $v, v \subseteq u, |v| \geq \varepsilon |u|$ such that any occurrence in v is strongly (v, c, d) -covered.

Proof: We proceed by induction on d . To prove the statement for $d = 1$ we take a $10k$ -regular ordered sequence $(u, <)$. Then, according to the previous lemma ($r = 2$), either $|u| \leq 1440 \|u\|$ or there is a subsequence $v \subseteq u, |v| \geq \frac{1}{10} |u|$ whose each element is in u $(16, 1)$ -covered. We apply Lemma 5.1 and choose a k -regular subsequence $v' \subseteq v, |v'| \geq (\frac{1}{10k})^2 |u|$. Any element of this sequence is still $(16, 1)$ -covered. We see that for $k \geq 2, d = 1$ we can put $c = 16, l = 10k, \Delta = 1440$ and $\varepsilon = (\frac{1}{10k})^2$. Suppose the statement has been proved for $d > 1$ and any $k \geq 2$. Our task is to derive the existence of the numbers $c(k, d + 1), l(k, d + 1), \Delta(k, d + 1)$ and $\varepsilon(k, d + 1)$ corresponding to a given $d + 1, k$. We show that it is possible to take

$$l(k, d + 1) = l(10k, d), c(k, d + 1) = 8r, \varepsilon(k, d + 1) = (\frac{1}{10k})^2 \varepsilon(10k, d)$$

and

$$\Delta(k, d + 1) = \max(\Delta(10k, d), \frac{720r}{\varepsilon(10k, d)})$$

where

$$r = d \cdot c(10k, d) + 2.$$

Suppose $(u, <)$ is an ordered $l(k, d + 1)$ -regular sequence. Then, according to the induction hypothesis, either $|u| \leq \Delta(10k, d) \|u\|$ or there is a $10k$ -regular subsequence $v_1, v_1 \subseteq u, |v_1| \geq \varepsilon(10k, d) |u|$ whose each element is strongly $(v_1, c(10k, d), d)$ -covered. Suppose the latter possibility holds.

Now, according to Lemma 5.2, either $|v_1| \leq 720r \|v_1\|$ or there is a subsequence $v_2, v_2 \subseteq v_1, |v_2| \geq \frac{1}{10} |v_1|$ any element of which is $(8r, r - 1)$ -covered in v_1 . Suppose the latter possibility holds.

Finally, we choose according to Lemma 5.1 a k -regular subsequence $v_3, v_3 \subseteq v_2$ where $|v_3| \geq (\frac{1}{10k})^2 |v_1|$. We prove that any element of v_3 is strongly $(v_3, c(k, d + 1), d + 1)$ -covered. Let $x \in v_3$ be an a -occurrences. According to the choice of v_2 there is an interval I in v_1 that $(8r, r - 1)$ -covers x . This implies that there are $r - 1$ symbols $a_j \in S(u), j = 1, 2, \dots, r - 1, a_j < a$ any of which occurs twice in I and that there are at most $8r$ a -occurrences in I .

We define I^u as the interval in u spanned by I and show that I^u strongly $(v_3, c(k, d + 1), d + 1)$ -covers x . Property 1. is obvious, property 2. requires that $I^u \cap v_3$ contains at most $c(k, d + 1)$ a -occurrences which is true already for $I = I^u \cap v_1$. It remains to show that 3. holds.

In case at least $d + 1$ symbols a_j are distinct we are finished. Otherwise (see the definition of r) some symbol, say a_1 , has at least $2c(10k, d) + 1$ occurrences in I^u . Denote the subsequence consisting of these occurrences as p . Now we make use of the choice of v_1 from u . We denote by J the interval in u which strongly $(v_1, c(10k, d), d)$ -covers the middle element of p . Clearly $J \subseteq I^u$. We see that there are again $d + 1$ distinct symbols satisfying 3. for I^u . Namely a_1 and those d symbols less than a_1 any of which occurs twice in J . In the former possibilities it is easy to check that $|u| \leq \Delta(k, d + 1)$. \square

Now we are able to prove Theorem B. Suppose $u = u_1 a^2 u_2 a$ is a sequence, $b \notin S(u)$ is a new symbol and $i \geq 1$ is an integer. Our task is to prove $Ex(u_1 a b^i a u_2 a b^i, n) = O(Ex(u, n))$. The lower bound $Ex(u_1 a b^i a u_2 a b^i, n) = \Omega(Ex(u, n))$ follows from $u \prec u_1 a b^i a u_2 a b^i$. By Fact 2.2 it suffices to prove $Ex(t, n) = O(Ex(u, n))$ where $t = u_1 a b^2 a u_2 a b$. Put $k = d = \|u\|$ in the previous lemma and let c, l, ε and Δ be the corresponding constants guaranteed by this lemma.

Suppose the sequence w is l -regular and does not contain t . We define the linear order $(S(w), <)$ by $a < b$ iff the rightmost a -occurrences lies to the right of the rightmost b -occurrences. In the first case of the previous lemma $|w| \leq \Delta \|w\|$. Otherwise there is a $\|u\|$ -regular subsequence $v, v \subseteq w, |v| \geq \varepsilon |w|$ whose each element is strongly $(v, c, \|u\|)$ -covered. We show that v does not contain the sequence $u' = u_1 a^{2c+1} u_2 a$. Suppose on the contrary that $s = u_1^* (a^*)^{2c+1} u_2^* a^*, s \subseteq v$ is equivalent to u' . We denote by p the subsequence $(a^*)^{2c+1}$ of s . The middle a^* -occurrences in the subsequence p must be strongly $(v, c, \|u\|)$ -covered by an interval I in w . Let J be interval in w spanned by the first and by the last a^* -occurrences in p . Clearly $I \subseteq J$. There are $\|u\|$ distinct symbols $x_i, x_i < a^*, i = 1, 2, \dots, \|u\|$ and each of them occurs at least twice in I . By the definition of \jmath each of these symbols occurs to the right of s . At least one of them is not an element of $S(s)$ and we get $t \prec v$ which is a contradiction. Thus $u' \not\prec v$ and $|v| \leq Ex(u', \|v\|) \leq g Ex(u, \|v\|)$ for some constant g according to Fact 2.2. Hence

$$|w| \leq \frac{1}{\varepsilon} |v| \leq \frac{g}{\varepsilon} Ex(u, \|v\|) \leq \frac{g}{\varepsilon} Ex(u, \|w\|).$$

Together

$$|w| \leq \max\{\Delta \|w\|, \frac{g}{\varepsilon} Ex(u, \|w\|)\} = O(Ex(u, \|w\|)).$$

We have proved that

$$Ex(t, n, l) = O(Ex(u, n)).$$

Finally, according to 2. of Fact 2.1, the estimate

$$Ex(t, n) = Ex(t, n, \|t\|) = O(Ex(t, n, l)) = O(Ex(u, n))$$

holds. The Theorem is proved.

Acknowledgement

The authors are grateful to Jaroslav Nešetřil and to Jiří Matoušek for their fruitful comments.

(This paper is to appear in *Combinatorica*.)

References

- [1] R. Adamec, M. Klazar and P. Valtr, Generalized Davenport-Schinzel sequences with linear upper bound, . **Discrete Math.** (108), 1992219–229
- [2] P. K. Agarwal, *Intersection and decomposition algorithms for planar arrangements*, Cambridge University Press 1991.
- [3] P. K. Agarwal, M. Sharir and P. Shor, Sharp upper and lower bounds on the length of general Davenport-Schinzel sequences, *J. Comb. Theory A* **52** (1989), 228–274.
- [4] H. Davenport and A. Schinzel, A combinatorial problem connected with differential equations, *Amer. J. Math.* **87** (1965), 684–689.
- [5] H. Davenport, A combinatorial problem connected with differential equations II, *Acta Arith.* **17** (1971), 363–372.
- [6] Z. Füredi and P. Hajnal, Davenport-Schinzel theory of matrices, *Discrete Math.* **103** (1992), 233–251.
- [7] S. Hart and M. Sharir, Nonlinearity of Davenport-Schinzel sequences and of generalized path compression schemes, *Combinatorica* **6** (1986), 151–177.
- [8] M. Klazar, A general upper bound in Extremal theory of sequences, *Comment. Math. Un. Carolinae* **33** (1992), 737–746.
- [9] M. Klazar, A linear upper bound in Extremal theory of sequences (to appear in *J. of Comb. Th. A*).
- [10] M. Klazar, Two results on a partial ordering of finite sequences (to appear in *Comment. Math. Un. Carolinae*).
- [11] P. Komjath, A simplified construction of nonlinear Davenport-Schinzel sequences, *J. Comb. Th. A* **49** (1988), 262–267.
- [12] M. Sharir, Almost linear upper bounds on the length of general Davenport-Schinzel sequences, *Combinatorica* **7** (1987), 131–143.

- [13] M. Sharir and P. K. Agarwal, *Davenport-Schinzel sequences and their geometric applications*, manuscript 1993.
- [14] E. SzmerÉDI, On a problem by Davenport and Schinzel, *Acta Arith.* **25** (1974), 213–224.
- [15] A. Wiernik and M. Sharir, Planar realization of nonlinear Davenport-Schinzel sequences by segments, *Discrete Comput. Geometry* **3** (1988), 15–47