

## 6. LITERATURVERZEICHNIS

1. Sorensen OH, Nielsen SP, Charles P, et al. Consensus development statement on osteoporosis [letter]. *Osteoporos Int* 1997; 7:589.
2. Kanis JA. Assessment of fracture risk and its application to screening for postmenopausal osteoporosis: synopsis of a WHO report. WHO Study Group. *Osteoporos Int* 1994; 4:368-81.
3. DAGO. Epidemiologie und Gesundheitsökonomie, Hrsg. Osteoporose Leitlinien. Die Empfehlung der Deutschen Arbeitsgemeinschaft Osteoporose (DAGO). Deutsche Grünes Kreuz. 1. Aufl. Marburg: Kilian, 1996:13-20.
4. Eastell R, Cedel SL, Wahner HW, Riggs BL, Melton LJ. Classification of vertebral fractures. *J Bone Miner Res* 1991; 6:207-215.
5. McCloskey EV, Spector TD, Eyres KS, et al. The assessment of vertebral deformity: a method for use in population studies and clinical trials [see comments]. *Osteoporos Int* 1993; 3:138-147.
6. Minne HW, Leidig G, Wuster C, et al. A newly developed spine deformity index (SDI) to quantitate vertebral crush fractures in patients with osteoporosis. *Bone Miner* 1988; 3:335-349.
7. Felsenberg D, Kalender W. Computer-assisted morphometry of vertebral fractures. In: Genant HK JM, v. Kuijk C, ed. Vertebral fractures in osteoporosis, 1995:309-317.
8. Beard CM, Lane AW, O'Fallon WM, Riggs BL, Melton LJ, 3rd. Comparison of respondents and nonrespondents in an osteoporosis study. *Ann Epidemiol* 1994; 4:398-403.
9. Heilbrun LK, Ross PD, Wasnich RD, Yano K, Vogel JM. Characteristics of respondents and nonrespondents in a prospective study of osteoporosis. *J Clin Epidemiol* 1991; 44:233-9.

10. Matthis C, Raspe A, Holzmann M, Raspe H. Die Epidemiologie der vertebraalen Osteoporose in Europa: Prävalenz, Risikofaktoren und Krankheitslast in Deutschland (EVOS);Schlußbericht. Lübeck, 1998:24.
11. van Bodegom JW, Kuiper JW, van Rijn RR, van Kuijk C, Zwamborn AW, Grashuis JL. Vertebral dimensions: influence of X-ray technique and patient size on measurements. *Calcif Tissue Int* 1998; 62:214-8.
12. Kellgren H, Lawrence J. Osteoarthritis and disk degeneration in an urban population. *Ann. Rheum. Dis* 1958; 17:388-397.
13. Smith-Bindman R, Cummings SR, Steiger P, Genant HK. A comparison of morphometric definitions of vertebral fracture. *J Bone Miner Res* 1991; 6:25-34.
14. Raymakers JA, Kapelle JW, van Beresteijn EC, Duursma SA. Assessment of osteoporotic spine deformity. A new method. *Skeletal Radiol* 1990; 19:91-97.
15. Ross PD. Clinical consequences of vertebral fractures. *Am J Med* 1997; 103:30S-42S; discussion 42S-43S.
16. Albright F, Smith P, Richardson A. Postmenopausal Osteoporosis. *JAMA* 1941; 116:2465-2474.
17. Melton LJd, Kan SH, Frye MA, Wahner HW, O'Fallon WM, Riggs BL. Epidemiology of vertebral fractures in women. *Am J Epidemiol* 1989; 129:1000-11.
18. Bengner U, Johnell O, Redlund-Johnell I. Changes in incidence and prevalence of vertebral fractures during 30 years. *Calcif Tissue Int* 1988; 42:293-6.
19. Felsenberg D, Kalender WA. Computerassisted morphometry of vertebral bodies. In: Genant HK, Jergas M, eds. *Vertebral fractures in osteoporosis*. San Francisco: UCSF-Print, 1996.

20. O'Neill TW, Varlow J, Felsenberg D, et al. Variation in vertebral height ratios in population studies. European Vertebral Osteoporosis Study Group. *J Bone Miner Res* 1994; 9:1895-907.
21. Black D, Palermo L. Use of vertebral reference values to define prevalent vertebral deformities. In: Genant HK JM, v. Kuijk C, ed. *Vertebral fractures in osteoporosis*. Radiology Research and Education Foundation, San Francisco, 1995:189-204.
22. Black DM, Cummings SR, Stone K, Hudes E, Palermo L, Steiger P. A new approach to defining normal vertebral dimensions. *J Bone Miner Res* 1991; 6:883-92.
23. Leidig-Bruckner G, Limberg B, Felsenberg D, et al. Sex difference in the validity of vertebral deformities as an index of prevalent vertebral osteoporotic fractures: a population survey of older men and women. *Osteoporos Int* 2000; 11:102-19.
24. Gowin W, Müller C, Alenfeld F, Diessel E, Felsenberg D. Differential diagnosis of osteoporotic vertebral fractures/deformities. *J Bone Miner Res* 1996; 11(Suppl):355.
25. O'Neill TW, Felsenberg D, Varlow J, Cooper C, Kanis JA, Silman AJ. The prevalence of vertebral deformity in european men and women: the European Vertebral Osteoporosis Study. *J Bone Miner Res* 1996; 11:1010-8.
26. Black DM, Arden NK, Palermo L, Pearson J, Cummings SR. Prevalent vertebral deformities predict hip fractures and new vertebral deformities but not wrist fractures. Study of Osteoporotic Fractures Research Group. *J Bone Miner Res* 1999; 14:821-8.
27. Klotzbuecher CM, Ross PD, Landsman PB, Abbott TA, 3rd, Berger M. Patients with prior fractures have an increased risk of future fractures: a summary of the literature and statistical synthesis. *J Bone Miner Res* 2000; 15:721-39.

28. Ross PD, Genant HK, Davis JW, Miller PD, Wasnich RD. Predicting vertebral fracture incidence from prevalent fractures and bone density among non-black, osteoporotic women. *Osteoporos Int* 1993; 3:120-6.
29. Nevitt MC, Ross PD, Palermo L, Musliner T, Genant HK, Thompson DE. Association of prevalent vertebral fractures, bone density, and alendronate treatment with incident vertebral fractures: effect of number and spinal location of fractures. The Fracture Intervention Trial Research Group. *Bone* 1999; 25:613-9.
30. Genant HK, Jergas M, Palermo L, et al. Comparison of semiquantitative visual and quantitative morphometric assessment of prevalent and incident vertebral fractures in osteoporosis The Study of Osteoporotic Fractures Research Group. *J Bone Miner Res* 1996; 11:984-96.
31. Cooper C, Atkinson EJ, O'Fallon WM, Melton LJd. Incidence of clinically diagnosed vertebral fractures: a population- based study in Rochester, Minnesota, 1985-1989. *J Bone Miner Res* 1992; 7:221-7.
32. Genant HK, Wu CY, van Kuijk C, Nevitt MC. Vertebral fracture assessment using a semiquantitative technique. *J Bone Miner Res* 1993; 8:1137-1148.
33. Hedlund LR, Gallagher JC, Meeger C, Stoner S. Change in vertebral shape in spinal osteoporosis. *Calcif Tissue Int* 1989; 44:168-72.
34. Melton LJd, Lane AW, Cooper C, Eastell R, O'Fallon WM, Riggs BL. Prevalence and incidence of vertebral deformities. *Osteoporos Int* 1993; 3:113-9.
35. Spector TD, McCloskey EV, Doyle DV, Kanis JA. Prevalence of vertebral fracture in women and the relationship with bone density and symptoms: the Chingford Study. *J Bone Miner Res* 1993; 8:817-22.
36. Wu CY, Li J, Jergas M, Genant HK. Comparison of semiquantitative and quantitative techniques for the assessment of prevalent and incident vertebral fractures. *Osteoporos Int* 1995; 5:354-70.

37. Ross PD, Wasnich RD, Davis JW, Vogel JM. Vertebral dimension differences between Caucasian populations, and between Caucasians and Japanese. *Bone* 1991; 12:107-12.
38. Ismail AA, Cooper C, Felsenberg D, et al. Number and type of vertebral deformities: epidemiological characteristics and relation to back pain and height loss. European Vertebral Osteoporosis Study Group. *Osteoporos Int* 1999; 9:206-213.
39. Black DM, Palermo L, Nevitt MC, Genant HK, Christensen L, Cummings SR. Defining incident vertebral deformity: A prospective comparison of several approaches. *J Bone Miner Res* 1999; 1:90-101.