

5. Literaturverzeichnis

- [1] Agarwal R.
Strategies and feasibility of hypertension control in a prevalent hemodialysis cohort.
Clin Nephrol. 2000 May;53(5):344-53.
- [2] Andreassen AK, Hartmann A, Offstad J, Geiran O, Kvernebo K, Simonsen S.
Hypertension prophylaxis with omega-3 fatty acids in heart transplant recipients.
J Am Coll Cardiol. 1997 May;29(6):1324-31.
- [3] Artz MA, Boots JM, Ligtenberg G, Roodnat JI, Christiaans MH, Vos PF, Blom HJ, Sweep FC, Demacker PN, Hilbrands LB.
Improved cardiovascular risk profile and renal function in renal transplant patients after randomized conversion from cyclosporine to tacrolimus.
J Am Soc Nephrol. 2003 Jul;14(7):1880-8.
- [4] Bahous SA, Stephan A, Blacher J, Safar ME.
Aortic stiffness, living donors, and renal transplantation.
Hypertension. 2006 Feb;47(2):216-21.
- [5] Barenbrock M, Hausberg M, Kosch M, Kisters K, Hoeks AP, Rahn KH.
A longitudinal study of vessel wall properties in normotensive and hypertensive renal transplant recipients.
J Hum Hypertens. 1998 Oct;12(10):707-11.
- [6] Baumgart P, Reinbach R, Akbulut T, Walger P, Thiel M, von Eiff M, Gerke M, Rahn KH.
Sprechstundenblutdruck, Heimblutdruck, Ergometer-Blutdruck und 24-Stunden-Blutdruck.
Dtsch Med Wochenschr. 1990 Apr 27;115(17):643-7.

- [7] Baumgart P, Walger P, Gemen S, von Eiff M, Raidt H, Rahn KH.
Blood pressure elevation during the night in chronic renal failure, hemodialysis
and after renal transplantation.
Nephron. 1991;57(3):293-8.
- [8] Bechstein WO, Wullstein C.
Transplantation and outcome quality.
Chirurg. 2002 Jun;73(6):576-81.
- [9] Beige J, Offermann G, Zidek W.
Pulsdruck, Transplantatfunktion und antihypertensive Medikation bei
Nierentransplantation. Ergebnisse der Neoral® Most Multicenterstudie.
[unveröffentl. Manuskript]
- [10] Benedetti E, Troppmann C, Gillingham K, Sutherland DE, Payne WD, Dunn DL,
Matas AJ, Najarian JS, Grussner RW.
Short- and long-term outcomes of kidney transplants with multiple renal arteries.
Ann Surg. 1995 Apr;221(4):406-14.
- [11] Benetos A, Waeber B, Izzo J, Mitchell G, Resnick L, Asmar R, Safar M.
Influence of age, risk factors, and cardiovascular and renal disease on arterial
stiffness: clinical applications.
Am J Hypertens. 2002 Dec;15(12):1101-8.
- [12] Bock HA.
Chronic rejection and hypertension: a chicken-and-egg problem.
Nephrol Dial Transplant. 1995;10(7):1126-8.
- [13] Boom H, Mallat MJ, de Fijter JW, Zwinderman AH, Paul LC.
Delayed graft function influences renal function, but not survival.
Kidney Int. 2000 Aug;58(2):859-66.

- [14] Braun WE, Protiva DA.
Emerging profiles in 105 recipients of renal allografts functioning for 20 to 35 years: the "watershed" effect.
Transplant Proc. 2001 Feb-Mar;33(1-2):1131-3.
- [15] Budde K, Waiser J, Fritsche L, Zitzmann J, Schreiber M, Kunz R, Neumayer HH.
Hypertension in patients after renal transplantation.
Transplant Proc. 1997 Feb-Mar;29(1-2):209-11.
- [16] Carvalho MF, Soares V.
Factors associated with arterial hypertension after renal transplantation.
Transplant Proc. 1998 Sep;30(6):2872-3.
- [17] Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, Jones DW, Materson BJ, Oparil S, Wright JT Jr, Roccella EJ; Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. National Heart, Lung, and Blood Institute; National High Blood Pressure Education Program Coordinating Committee.
Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure.
Hypertension. 2003 Dec;42(6):1206-52.
- [18] Cockcroft DW, Gault MH.
Prediction of creatinine clearance from serum creatinine.
Nephron. 1976;16(1):31-41.
- [19] Cosio FG, Dillon JJ, Falkenhain ME, Tesi RJ, Henry ML, Elkhammas EA, Davies EA, Bumgardner GL, Ferguson RM.
Racial differences in renal allograft survival: the role of systemic hypertension.
Kidney Int. 1995 Apr;47(4):1136-41.

- [20] Cosio FG, Pelletier RP, Pesavento TE, Henry ML, Ferguson RM, Mitchell L, Lemeshow S.
Elevated blood pressure predicts the risk of acute rejection in renal allograft recipients.
Kidney Int. 2001 Mar;59(3):1158-64.
- [21] Cosio FG, Pelletier RP, Sedmak DD, Pesavento TE, Henry ML, Ferguson RM.
Renal allograft survival following acute rejection correlates with blood pressure levels and histopathology.
Kidney Int. 1999 Nov;56(5):1912-9.
- [22] Daberkow U.
Einflussfaktoren auf die Funktion transplantierte Leichennieren unter Berücksichtigung der organerhaltenden Intensivtherapie und der Organentnahme.
Dissertationsarbeit 1996 (Charité Berlin)
- [23] Di Paolo S, Stallone G, Schena A, Infante B, Gesualdo L, Paolo Schena F.
Hypertension is an independent predictor of delayed graft function and worse renal function only in kidneys with chronic pathological lesions.
Transplantation. 2002 Feb 27;73(4):623-7.
- [24] Dragun D, Brasen JH, Schonemann C, Fritsche L, Budde K, Neumayer HH, Luft FC, Wallukat G.
Patients with steroid refractory acute vascular rejection develop agonistic antibodies targeting angiotensin II type 1 receptor.
Transplant Proc. 2003 Sep;35(6):2104-5.
- [25] Dragun D, Müller DN, Bräsen JH, Fritsche L, Nieminen-Kelhä M, Dechend R, Kintschner U, Rudolph B, Hoebeke J, Eckert D, Mazak I, Plehm R, Schönemann C, Unger T, Budde K, Neumayer HH, Luft FC, Wallukat G.
Angiotensin II type 1-receptor activating antibodies in renal-allograft rejection.
N Engl J Med. 2005 Feb 10;352(6):558-69.

- [26] Fellstrom B, Holdaas H, Jardine AG, Nyberg G, Gronhagen-Riska C, Madsen S, Neumayer HH, Cole E, Maes B, Ambuhl P, Olsson AG, Staffler B, Pedersen TR; Assessment of Lescol in Renal Transplantation Study Investigators.
Risk factors for reaching renal endpoints in the assessment of Lescol in renal transplantation (ALERT) trial.
Transplantation. 2005 Jan 27;79(2):205-12.
- [27] Fernández-Fresnedo G, Escallada R, Martín de Francisco AL, Ruiz JC, Rodrigo E, Sanz de Castro S, González Cotorruelo J, Arias M.
Association between pulse pressure and cardiovascular disease in renal transplant patients.
Am J Transplant. 2005 Feb;5(2):394-8.
- [28] Ferreira SR, Moisés VA, Tavares A, Pacheco-Silva A.
Cardiovascular effects of successful renal transplantation: a 1-year sequential study of left ventricular morphology and function, and 24-hour blood pressure profile.
Transplantation. 2002 Dec 15;74(11):1580-7.
- [29] First MR, Peddi VR, Weiskittel P, Woodle ES.
Control of risk factors for cardiovascular disease in long-term renal transplant recipients.
Transplant Proc. 2001 Nov-Dec;33(7-8):3674-5.
- [30] Fleiner F, Fritsche L, Glander P, Neumayer HH, Budde K.
Reporting of rejection after renal transplantation in large immunosuppressive trials: biopsy-proven, clinical, presumed, or treated rejection?
Transplantation. 2006 Mar 15;81(5):655-9.
- [31] Frank H, Schmieder RE, Vogt-Ladner G, Schobel HP, Neumayer HH.
Determinants of left ventricular structure after kidney transplantation.
Transplant Proc. 2000 Dec;32(8):2801-6.

- [32] Frei U, Schindler R, Wieters D, Grouven U, Brunkhorst R, Koch KM.
Pre-transplant hypertension: a major risk factor for chronic progressive renal allograft dysfunction?
Nephrol Dial Transplant. 1995;10(7):1206-11.
- [33] Frei U.
Stand der Nierentransplantation.
ZFA Stuttgart. 1995; 71(11): 897-903.
- [34] Frei, U., Schober-Halstenberg, H.-J.:
Nierenersatztherapie in Deutschland.
QuaSi-Niere Jahresbericht 2005/2006, Berlin, Deutschland.
- [35] Foley RN, Parfrey PS, Kent GM, Harnett JD, Murray DC, Barre PE.
Serial change in echocardiographic parameters and cardiac failure in end-stage renal disease.
J Am Soc Nephrol. 2000 May;11(5):912-6.
- [36] Gatzka CD, Schobel HP, Klingbeil AU, Neumayer HH, Schmieder RE.
Normalization of circadian blood pressure profiles after renal transplantation.
Transplantation. 1995 May 15;59(9):1270-4.
- [37] Gatzka CD, Schobel HP, Vogt-Ladner G, Neumayer HH, Schmieder RE.
Blutdruckabfall im Schlaf und linksventrikuläre Hypertrophie bei nierentransplantierten Patienten.
Z Kardiol. 1992;81 Suppl 2:17-9.
- [38] Guidi E, Cozzi MG, Minetti E, Bianchi G.
Donor and recipient family histories of hypertension influence renal impairment and blood pressure during acute rejections.
J Am Soc Nephrol. 1998 Nov;9(11):2102-7.

- [39] Hariharan S, Johnson CP, Bresnahan BA, Taranto SE, McIntosh MJ, Stablein D.
Improved graft survival after renal transplantation in the United States, 1988 to 1996.
N Engl J Med. 2000 Mar 2;342(9):605-12.
- [40] Held PJ, Kahan BD, Hunsicker LG, Liska D, Wolfe RA, Port FK, Gaylin DS, Garcia JR, Agodoa LY, Krakauer H.
The impact of HLA mismatches on the survival of first cadaveric kidney transplants.
N Engl J Med. 1994 Sep 22;331(12):765-70.
- [41] Hjeltnesaeth J, Hartmann A, Leivestad T, Holdaas H, Sagedal S, Olstad M, Jenssen T.
The impact of early-diagnosed new-onset post-transplantation diabetes mellitus on survival and major cardiac events.
Kidney Int. 2006 Feb;69(3):588-95.
- [42] Hohage H, Bruckner D, Arlt M, Buchholz B, Zidek W, Spieker C.
Influence of cyclosporine A and FK506 on 24 h blood pressure monitoring in kidney transplant recipients.
Clin Nephrol. 1996 May;45(5):342-4.
- [43] Ishida K, Ishida H, Narita M, Sairenchi T, Saito Y, Fukutomi H, Takahashi H, Yamagata K, Koyama A.
Factors affecting renal function in 119 985 adults over three years.
QJM. 2001 Oct;94(10):541-50.
- [44] Jacobi J, Rockstroh J, John S, Schreiber M, Schlaich MP, Neumayer HH, Schmieder RE.
Prospective analysis of the value of 24-hour ambulatory blood pressure on renal function after kidney transplantation.
Transplantation. 2000 Sep 15;70(5):819-27.

- [45] Jelakovic B, Mayer G.
A renocentric view of essential hypertension: lessons to be learnt from kidney transplantation.
Nephrol Dial Transplant. 1995;10(9):1510-2.
- [46] Jindal RM, Hariharan S.
Chronic rejection in kidney transplants. An in-depth review.
Nephron. 1999 Sep;83(1):13-24.
- [47] Kain HK, Hinman AT, Sokolow M.
Arterial blood pressure measurements with a portable recorder in hypertensive patients. I. Variability and correlation with "casual" pressure
Circulation. 1964 Dec;30:882-92.
- [48] Kannel WB.
Vital epidemiologic clues in heart failure.
J Clin Epidemiol. 2000 Mar 1;53(3):229-35.
- [49] Kasiske BL, Chakkera HA, Roel J.
Explained and unexplained ischemic heart disease risk after renal transplantation.
J Am Soc Nephrol. 2000 Sep;11(9):1735-43.
- [50] Kew CE 2nd, Curtis JJ.
The best way to manage hypertension after renal transplantation.
J Ren Nutr. 2000 Jan;10(1):3-6.
- [51] Kim HC, Kwon JK, Park SB, Cho WH, Park CH.
Hypertension in kidney transplantation recipients: effect on long-term renal allograft survival.
Transplant Proc. 1998 Nov;30(7):3906-7.

- [52] Klassen DK.
How aggressively should blood pressure be treated in renal transplant recipients?
Curr Hypertens Rep. 2000 Oct;2(5):473-7.
- [53] Koch K-M.
Klinische Nephrologie.
Urban & Fischer München Jena 2000.
- [54] Ligtenberg G, Hené RJ, Blankestijn PJ, Koomans HA.
Cardiovascular risk factors in renal transplant patients: cyclosporin A versus tacrolimus.
J Am Soc Nephrol. 2001 Feb;12(2):368-73.
- [55] Lipkin GW, Tucker B, Giles M, Raine AE.
Ambulatory blood pressure and left ventricular mass in cyclosporin- and non-cyclosporin-treated renal transplant recipients.
J Hypertens. 1993 Apr;11(4):439-42.
- [56] London GM, Pannier B, Guerin AP, Blacher J, Marchais SJ, Darne B, Metivier F, Adda H, Safar ME.
Alterations of left ventricular hypertrophy in and survival of patients receiving hemodialysis: follow-up of an interventional study.
J Am Soc Nephrol. 2001 Dec;12(12):2759-67.
- [57] Magee C, Pascual M.
The growing problem of chronic renal failure after transplantation of a nonrenal organ.
N Engl J Med. 2003 Sep 4;349(10):994-6.
- [58] Mancia G.
Methods for assessing blood pressure values in humans.
Hypertension. 1983 Sep-Oct;5(5 Pt 2):III5-13.

- [59] Mange KC, Cherikh WS, Maghirang J, Bloom RD.
A comparison of the survival of shipped and locally transplanted cadaveric renal allografts.
N Engl J Med. 2001 Oct 25;345(17):1237-42.
- [60] Mange KC, Cizman B, Joffe M, Feldman HI.
Arterial hypertension and renal allograft survival.
JAMA. 2000 Feb 2;283(5):633-8.
- [61] Mange KC, Feldman HI, Joffe MM, Fa K, Bloom RD.
Blood pressure and the survival of renal allografts from living donors.
J Am Soc Nephrol. 2004 Jan;15(1):187-93.
- [62] Martinez-Castelao A, Hueso M, Sanz V, Rejas J, Alsina J, Grinyo JM.
Treatment of hypertension after renal transplantation: long-term efficacy of verapamil, enalapril, and doxazosin.
Kidney Int Suppl. 1998 Dec;68:S130-4.
- [63] Matsumae T, Matsumae M, Hasegawa Y, Tanaka T, Yoshitake K, Noda R, Ogahara S, Murata T, Kaneoka H.
Relationship between fluctuation pattern of blood pressure during hemodialysis treatment and cardiovascular morphology: an autopsy study of 53 cases.
Nephron. 2001 Jun;88(2):113-9.
- [64] Matteucci MC, Giordano U, Calzolari A, Turchetta A, Santilli A, Rizzoni G.
Left ventricular hypertrophy, treadmill tests, and 24-hour blood pressure in pediatric transplant patients.
Kidney Int. 1999 Oct;56(4):1566-70.
- [65] Mehrens T, Thiele S, Suwelack B, Kempkes M, Hohage H.
The beneficial effects of calcium channel blockers on long-term kidney transplant survival are independent of blood-pressure reduction.
Clin Transplant. 2000 Jun;14(3):257-61.

- [66] Meier-Kriesche HU, Arndorfer JA, Kaplan B.
The impact of body mass index on renal transplant outcomes: a significant independent risk factor for graft failure and patient death.
Transplantation. 2002 Jan 15;73(1):70-4.
- [67] Midtvedt K, Hartmann A, Foss A, Fauchald P, Nordal KP, Rootwelt K, Holdaas H.
Sustained improvement of renal graft function for two years in hypertensive renal transplant recipients treated with nifedipine as compared to lisinopril.
Transplantation. 2001 Dec 15;72(11):1787-92.
- [68] Midtvedt K, Hartmann A, Holdaas H, Fauchald P.
Efficacy of nifedipine or lisinopril in the treatment of hypertension after renal transplantation: a double-blind randomised comparative trial.
Clin Transplant. 2001 Dec;15(6):426-31.
- [69] Midtvedt K, Neumayer HH.
Management strategies for posttransplant hypertension.
Transplantation. 2000 Dec 15;70(11 Suppl):SS64-9.
- [70] Neumayer HH, Luft FC; Transplantationsgruppe Mittelfranken.
Twenty-five years of renal transplantation from a single center: a risk factor analysis for short- and long-term outcomes.
Clin Investig. 1993 May;71(5):341-50.
- [71] Neumayer HH, Frei U, Wagner K.
Vaskuläre Mechanismen der Ciclosporin-Nephrotoxizität.
Nieren-Hochdruckkr. 1993; 22(Suppl1):SS62-70.
- [72] Ojo AO, Hanson JA, Wolfe RA, Leichtman AB, Agodoa LY, Port FK.
Long-term survival in renal transplant recipients with graft function.
Kidney Int. 2000 Jan;57(1):307-13.

- [73] Ojo AO, Held PJ, Port FK, Wolfe RA, Leichtman AB, Young EW, Arndorfer J, Christensen L, Merion RM.
Chronic renal failure after transplantation of a nonrenal organ.
N Engl J Med. 2003 Sep 4;349(10):931-40.
- [74] Opelz G; Collaborative Transplant Study.
Investigation of the relationship between maintenance dose of cyclosporine and nephrotoxicity or hypertension.
Transplant Proc. 2001 Nov-Dec;33(7-8):3351-4.
- [75] Opelz G, Dohler B; Collaborative Transplant Study.
Improved long-term outcomes after renal transplantation associated with blood pressure control.
Am J Transplant. 2005 Nov;5(11):2725-31.
- [76] Opelz G, Wujciak T, Ritz E.
Association of chronic kidney graft failure with recipient blood pressure.
Collaborative Transplant Study.
Kidney Int. 1998 Jan;53(1):217-22.
- [77] Pascual M, Theruvath T, Kawai T, Tolckoff-Rubin N, Cosimi AB.
Strategies to improve long-term outcomes after renal transplantation.
N Engl J Med. 2002 Feb 21;346(8):580-90.
- [78] Paul LC.
Treatment of posttransplant hypertension: too little, too late?
Transplantation. 2003 Dec 15;76(11):1645-6.
- [79] Perez Fontan M, Rodriguez-Carmona A, Garcia Falcon T, Fernandez Rivera C, Valdes F.
Early immunologic and nonimmunologic predictors of arterial hypertension after renal transplantation.
Am J Kidney Dis. 1999 Jan;33(1):21-8.

- [80] Pichlmayr R, Bunzendahl H, Luachart W.
Der Verlauf nach Nierentransplantation.
In: Pichlmayr (Hrsg). Allgemeine und spezielle Operationslehre,
Transplantationschirurgie. Springer Berlin Bd. 3 (1981): 595-686
- [81] Pickering TG, James GD, Boddie C, Harshfield GA, Blank S, Laragh JH.
How common is white coat hypertension?
JAMA. 1988 Jan 8;259(2):225-8.
- [82] Pschyrembel – Klinisches Wörterbuch.
257. Auflage Walter de Gruyter Berlin New York 1994.
- [83] Puyoo O, Rostaing L, Benard E, Cisterne JM, Modesto A, Durand D.
Risk factors of chronic rejection after renal transplantation: retrospective study of
201 patients.
Transplant Proc. 1998 Sep;30(6):2804-6.
- [84] Raine AE.
Does antihypertensive therapy modify chronic allograft failure?
Kidney Int Suppl. 1995 Dec;52:S107-11.
- [85] Roodnat JI, Zietse R, Mulder PG, Rischen-Vos J, van Gelder T, IJzermans JN,
Weimar W.
The vanishing importance of age in renal transplantation.
Transplantation. 1999 Feb 27;67(4):576-80.
- [86] Ruggenenti P, Perna A, Lesti M, Pisoni R, Mosconi L, Arnoldi F, Ciocca I,
Gaspari F, Remuzzi G.
Pretreatment blood pressure reliably predicts progression of chronic
nephropathies. GISEN Group.
Kidney Int. 2000 Nov;58(5):2093-101.

- [87] Schachinger H, Schmieder RE.
Bedeutung der 24-Stunden-Blutdruckmessung zur Erkennung
hypertoniebedingter Endorganschädigungen.
Z Kardiol. 1991;80 Suppl 1:41-7.
- [88] Schindler R, Tanriver Y, Frei U.
Hypertension and allograft nephropathy--cause, consequence, or both?
Nephrol Dial Transplant. 2000 Jan;15(1):8-10.
- [89] Schwenger V, Zeier M, Ritz E.
Hypertension after renal transplantation.
Curr Hypertens Rep. 2001 Oct;3(5):434-9.
- [90] Srivastava T.
Nondiabetic consequences of obesity on kidney.
Pediatr Nephrol. 2006 Apr;21(4):463-70.
- [91] Stobbe, Baumann.
Innere Medizin; Grundlagen und Klinik innerer Krankheiten.
Ullstein Mosby, Berlin Wiesbaden 1996.
- [92] Takemoto SK, Terasaki PI, Gjertson DW, Cecka JM.
Twelve years' experience with national sharing of HLA-matched cadaveric
kidneys for transplantation.
N Engl J Med. 2000 Oct 12;343(15):1078-84.
- [93] Thiel H.
Excuses of Nephrologists Not to Transplant, in Land, W.; Dossetor, J.B. (Hrsg.):
Organ Replacement Therapy: Ethics, Justice and commerce,
Springer Berlin (1991): 353-363.

- [94] Tutone VK, Mark PB, Stewart GA, Tan CC, Rodger RS, Geddes CC, Jardine AG. Hypertension, antihypertensive agents and outcomes following renal transplantation.
Clin Transplant: 2005 Apr;19(2):181-92.
- [95] Wellek, S.
Statistische Methoden zum Nachweis von Äquivalenz.
Gustav Fischer, Jena 1994
- [96] Wesslau C, Scholz D, May G, Ullrich W.
Procurement of organ donors--initial results of the Berlin-Potsdam-Frankfurt model
Z Urol Nephrol. 1990 Apr;83(4):171-4.
- [97] White WB.
Assessment of patients with office hypertension by 24-hour noninvasive ambulatory blood pressure monitoring.
Arch Intern Med. 1986 Nov;146(11):2196-9.
- [98] Wolfe RA, Ashby VB, Milford EL, Ojo AO, Ettenger RE, Agodoa LY, Held PJ, Port FK.
Comparison of mortality in all patients on dialysis, patients on dialysis awaiting transplantation, and recipients of a first cadaveric transplant.
New Engl J Med. 1999 Dec 2;341(23):1725-30.
- [99] Zeier M, Mandelbaum A, Ritz E.
Hypertension in the transplanted patient.
Nephron. 1998 Nov;80(3):257-68.