

5 Abkürzungen

1D	Eindimensional
2D	Zweidimensional
aa	<i>Amino acid</i>
Abb.	Abbildung
Agp	Agrobakterienphytochrom
ALA	5-Aminolävulinsäure
AmS	Ammoniumsulfat
APC	Allophycocyanin
AtBphP	<i>Agrobacterium tumefaciens bacteriophytochrome photoreceptor</i>
AUZ	Analytische Ultrazentifugation
BBO	<i>Broad band optimized</i>
BphP	<i>Bacterial phytochrome photoreceptor</i>
BrBphP	<i>Bradyrhizobium bacteriophytochrome photoreceptor</i>
BV	Biliverdin
CD	Circular dichroismus
CHES	N-cyclohexyl-2-aminoethansulfonsäure
Chl a	Chlorophyll a
cikA	<i>Circadian input kinase A</i> , synonym: hik24
Cph	Cyanobakterielles Phytochrom
Da	Dalton
DB	Doppelbindung
d. h.	Das heißt

DTT	Dithiothreitol
E. coli	Escherichia coli
EDTA	Ethyldiaminessigsäure
EPR-ENDOR	<i>Electron paramagnetic resonance electron-nuclear double resonance</i>
ESI-MS	<i>Electrospray ionisation mass spectroscopy</i>
evtl.	Eventuell
FR	Dunkelrot, ca. 730 nm (<i>farred</i>)
FT-IR	Fourier-Transform Infrarot
FT-RR	Fourier-Transform Resonanzraman
griech.	Griechisch
Hik	Histidinkinase
HIR	<i>High irradiance response</i>
HMQC	<i>Heteronuclear multiple quantum coherence</i>
HO	Hämoxygenase
HOOP	<i>Hydrogen out of plane</i>
HPLC	<i>High pressure liquid pressure chromatography</i>
HY	(long) Hypocotyl
i. d. R.	In der Regel
ip	<i>In plane</i>
IPTG	Isopropyl-β-D-thiogalactosid
IR	Infrarot
ital.	Italienisch
kDa	Kilodalton

lat.	Lateinisch
L _{CM}	<i>Core-membrane linker biliprotein</i>
LED	<i>Light emitting diode</i>
LFR	<i>Low fluency response</i>
LHC	<i>Light-harvesting complex</i>
m	Masse
Ni-NTA	Nickel-Nitriloacetic acid
NMR	<i>Nuclear magnetic resonance</i>
NOE	<i>Nuclear Overhauser effect</i>
OD	Optische Dichte
oop	<i>Out of plane</i>
ORF	<i>Open reading frame</i>
PaBphP	<i>Pseudomonas aeruginosa bacteriophytocrome photoreceptor</i>
PAS	Acronym aus den drei Proteinen, in denen diese Domäne zum ersten Mal beschrieben wurde: Per- <i>period circadian protein</i> , Arnt- <i>aryl hydrocarbon receptor nuclear translocator</i> , Sim- <i>single-minded protein</i>
PBS	Phycobilisom
PΦB	Phytochromobilin
PC	Phycocyanin
PCB	Phycocyanobilin
PCC	<i>Pasteur culture collection</i>
PcyA	Phycocyanobilin:Ferredoxinoxidoreduktase
PE	Phycoerythrin
PEB	Phycoerythrobilin

PebA	15,16-Dihydrobiliverdin:Ferredoxinoxidoreduktase
PebB	3Z-PEB:Ferredoxinoxidoreduktase
PEC	Phycoerythrocyanin
Phy	Phytochrom
Pfr	Phytochrom dunkelrot absorbierend (<i>phytochrome farred-absorbing</i>)
ppm	<i>Parts per million</i>
Pr	Phytochrom rot absorbierend (<i>phytochrome red-absorbing</i>)
PTFE	Polytetrafluorethylen
PUB	Phycourobilin
PVB	Phycoviolobilin
R	Rot, ca. 660 nm
RB	<i>Rich broth</i>
Rcp	<i>Response regulator for phytochrome</i>
r.m.s.d	<i>Root mean squared deviation</i>
RpBphB	<i>Rhodopseudomonas palustris bacteriophytochrome photoreceptor</i>
RR	Resonanz-Raman
RT	Raumtemperatur
s.	Siehe
SAR	<i>Specific absorbance ratio</i>
SEC	<i>Size-exclusion chromatography</i> , Größenausschlußchromatographie
sp.	<i>Species</i>
span.	Spanisch
TCI	<i>Triple carbon inverse</i>

Tris	Tris(hydroxymethyl)aminomethan
U	Umdrehung
u. a.	Unter anderem
UV	Ultraviolet
\bar{v}	Partielles spezifisches Volumen
vgl.	Vergleiche
vis	<i>Visible</i>
Vit.	Vitamin
VLFR	<i>Very low fluency response</i>
Vol.	Volumen
z	Ladung
z. B.	Zum Beispiel

6 Abbildungen

Abb. 1	Strukturen freier Biline	9
Abb. 2	Übersicht über die Biosynthese von Tetrapyrrolen und Bilinchromophoren	11
Abb. 3	Schematischer Aufbau eines hemidiscoidalen PBS	15
Abb. 4	Schematische Darstellung der Funktion von PBS	16
Abb. 5	Schematischer Aufbau von Phytochrom	22
Abb. 6	Schwingungsspektroskopische Ergebnisse zur Chromophorstruktur	28
Abb. 7	Strukturhinweise durch chemisch modifizierte Chromophore	30
Abb. 8	Schematische Darstellung des dimerisierten Phytochrommoleküls	31
Abb. 9	Kulturgefäße für unmarkierte und markierte Zellen	36
Abb. 10	ESI-MS-Spektren von PCB	39
Abb. 11	HPLC-Chromatogramm von PCB	40
Abb. 12	Größenausschlußchromatogramm von Cph1Δ2 in der Pr- und der Pfr-Form	41
Abb. 13	Dimerisierung der Pr-Form	42
Abb. 14	Variation des Assoziationskoeffizienten der Pr-Form	43
Abb. 15	Partialkonzentration von $(\text{Cph1}\Delta 2)_2\text{-Pr}$	44
Abb. 16	Massenänderung bei Phototransformation	45
Abb. 17	Pfr-Probe, mit der das in Abb. 18. gezeigte Experiment durchgeführt wurde	47
Abb. 18	Dimerisierung der Pfr-Form	47
Abb. 19	Simultane Bestimmung von $K_{A, \text{Pr}}$ und K_{A, Pfr^*}	49
Abb. 20	RR-Spektrum der Pr-Form, Übersicht	53
Abb. 21	RR-Spektrum der Pfr-Form, Übersicht	54
Abb. 22	Bandenanpassung an die Pfr-Form	57
Abb. 23	RR-Spektrum der Pr-Form von 1800-1400 cm ⁻¹	58
Abb. 24	RR-Spektrum der Pr-Form von 1400-900 cm ⁻¹	61
Abb. 25	RR-Spektrum der Pr-Form von 900-600 cm ⁻¹	64
Abb. 26	RR-Spektrum der Pfr-Form von 1800-1400 cm ⁻¹	67

Abb. 27 RR-Spektrum der Pfr-Form von 1400-900 cm ⁻¹	70
Abb. 28 RR-Spektrum der Pfr-Form von 900-600 cm ⁻¹	72
Abb. 29 Berechnete und experimentelle RR-Spektren der Pr- und Pfr-Formen	75
Abb. 30 Vergleich für die Pr-Form im Bereich 1800-1400 cm ⁻¹	78
Abb. 31 Vergleich für die Pr-Form im Bereich 1400-900 cm ⁻¹	79
Abb. 32 Vergleich für die Pr-Form im Bereich 900-600 cm ⁻¹	80
Abb. 33 Vergleich für die Pfr-Form im Bereich 1800-1400 cm ⁻¹	81
Abb. 34 Vergleich für die Pfr-Form im Bereich 1400-900 cm ⁻¹	82
Abb. 35 Vergleich für die Pfr-Form im Bereich 900-600 cm ⁻¹	83
Abb. 36 Eindimensionale ¹ H-Spektren der Pr-Form, deuteriertes Protein	88
Abb. 37 NOE-Spektren Pr-Form, deuteriertes Protein	89
Abb. 38 ¹ H- ¹⁵ N-HMQC nach 130 Stunden Meßzeit, Pr- und Pfr-Form	90
Abb. 39 1D- ¹⁵ N-Spektren der Pr- und Pfr-Form	91
Abb. 40 ¹ H- ¹³ C-HMQC nach 45 Minuten Meßzeit	93
Abb. 41 ¹ H- ¹³ C-HMQC der Pr-Form nach 70 Stunden Meßzeit	94
Abb. 42 Struktur von 3 ¹ -Cys-PCB aus der Kristallstruktur von Phycocyanin	95
Abb. 43 Berechnete und gemessene ¹⁵ N-chemische Verschiebungen von PCB	98

7 Tabellen

Tab. 1	Modellfreie Analyse der globalen Massenmittel (Mw_{av})	46
Tab. 2	Vergleich der durch Kurvenanpassung bestimmten Parameter für Pr und Pfr	48
Tab. 3	Durch Bandenanpassung erhaltene Maxima der Pr-Form im Bereich 1800-1400 cm ⁻¹	59
Tab. 4	Durch Bandenanpassung erhaltene Maxima der Pr-Form im Bereich 1400-900 cm ⁻¹	62
Tab. 5	Durch Bandenanpassung erhaltene Maxima der Pr-Form im Bereich 900-600 cm ⁻¹	65
Tab. 6	Durch Bandenanpassung erhaltene Maxima der Pfr-Form im Bereich 1800-1400 cm ⁻¹	68
Tab. 7	Durch Bandenanpassung erhaltene Maxima der Pfr-Form im Bereich 1400-900 cm ⁻¹	71
Tab. 8	Durch Bandenanpassung erhaltene Maxima der Pfr-Form im Bereich 900-600 cm ⁻¹	73
Tab. 9	Mengenangaben für je 1L BG11-Medium	112
Tab. 10	Modenzusammensetzung für Pr	142
Tab. 11	Modenzusammensetzung für Pfr	143

8 Literatur

- Ahmad M., Jarillo J.A., Smirnova O. & Cashmore A.R. (1998) The CRY1 blue light photoreceptor of Arabidopsis interacts with phytochrome A in vitro. *Mol. Cell* 1: 939-948.
- Alberte R.S., Wood A.M., Kursar T.A. & Guillard R.L. (1984) Novel phycoerythrins in marine *Synechococcus* sp. Characterization and evolutionary and ecological implications. *Plant Physiol.* 75: 732-739.
- Andel F., Lagarias D.M. & Mathies R.A. (1996) Resonance Raman analysis of Chromophore Structure in the lumir-R-photoproduct of phytochrome. *Biochemistry* 35: 15997-16008.
- Andel F., Murphy J.T., Haas J.A., McDowell M.T., van der Hoef I., Lugtenburg J., Lagarias J.L. & Mathies R.A. (2000) Probing the Photoreaction Mechanism of Phytochrome through Analysis of Resonance Raman Vibrational Spectra of Recombinant Analogues. *Biochemistry* 39: 2667-2676.
- Arciero D.M., Bryant D.A. & Glazer A.N. (1988a) In vitro attachment of bilins to apophycocyanin: I. Specific covalent adduct formation at cysteinyl residues involved in phycocyanobilin binding in C-phycocyanin. *J. Biol. Chem.* 263: 18343-18349.
- Arciero D.M., Dallas J.L. & Glazer A.N. (1988b) In vitro attachment of bilins to apophycocyanin: II. Determination of the structures of tryptic bilin peptides derived from the phycocyanobilin adduct. *J. Biol. Chem.* 263: 18350-18357.
- Bassi R., Rigoni F. & Giacometti G.M. (1990) Chlorophyll binding proteins with antenna function in higher plants and green algae. *Photochem. Photobiol.* 52: 1187-1206.
- Bax A., Griffey R.H. & Hawkins B.L. (1983) Sensitivity-enhanced correlation of ¹⁵N and ¹H chemical shifts in natural abundance samples via multiple quantum coherence. *J. Am. Chem. Soc.* 105: 7188-7190.
- Beale S. (1993) Biosynthesis of phycobilins. *Am. Chem. Rev.* 93: 785-802.
- Becke A.D.J. J. (1993) Density-functional thermochemistry. III. The role of exact exchange. *J. Chem. Phys.* 98: 5648-5652.
- Berzelius J.J. (1808) Forelasningar Djurkemine 2: 243.
- Beuhler R.J., Pierce R.C., Friedman L. & Siegelman H.W. (1976) Cleavage of phycocyanobilin from C-phycocyanin. Separation and mass spectral identification of the products. *J. Biol. Chem.* 251: 2405-2411.
- Beynon R.J. & Easterby J.S. (1996) Buffer solutions: The basics. BIOS Scientific publishers, Oxford, GB.
- Bhoo S.H., Davis S.J., Walker J., Karniol B. & Vierstra R.D. (2001) Bacteriophytochromes are photochromic histidine kinases using a biliverdin chromophore. *Nature* 414: 776-779.
- Bhoo S.H., Hirano T., Jeong H.Y., Lee J.G., Furuya M. & Song P.S. (1997) Phytochrome photochromism probed by site-directed mutations and chromophore esterification. *J. Am. Chem. Soc.* 119: 11717-11718.
- Bishop J.E., Lagarias J.C., Nagy J.O., Schoenleber R.W. & Rapoport H. (1986) Phycobiliprotein-bilin linkage diversity. I. Structural studies on A- and D-ring-linked phycocyanobilins. *J. Biol. Chem.* 261: 6790-6796.
- Bishop J.E., Rapoport H., Klotz A.V., Chan C.F., Glazer A.N., Füglister P. & Zuber H. (1987) Chromopeptides from phycoerythrocyanin: Structure and linkage of the three bilin groups. *J. Am. Chem. Soc.* 110: 875-881.
- Borthwick H.A., Hendricks S.B., Parker M.W., Toole E.H. & Toole V.K. (1952a) A reversible photoreaction controlling seed germination. *Proc. Natl. Acad. Sci. USA* 38: 662-666.

- Borthwick H.A., Hendricks S.B. & Parker M.W. (1952b) The reaction controlling floral initiation. Proc. Natl. Acad. Sci. USA 38: 929-934.
- Borucki B., Otto H., Rottwinkel G., Hughes J., Heyn M.P. & Lamparter T. (2003) Mechanism of Cph1 phytochrome assembly from stopped-flow kinetics and circular dichroism. Biochemistry 42: 13684-13697.
- Borucki B., von Stetten D., Seibeck S., Lamparter T., Michael N., Mroginski M.A., Otto H., Murgida D.H., Heyn M.P. & Hildebrandt P. (2005) Light-induced proton release of phytochrome is coupled to the transient deprotonation of the tetrapyrrole chromophore. J. Biol. Chem. 280: 34358-34364.
- Brandlmeier T., Scheer H. & Rüdiger W. (1981) Chomophore content and molar absorptivity of phytochrome in the Pr-form. Z. Naturforsch. 36: 431-439.
- Butler W.L., Hendricks S.B. & Siegelman H.A. (1960) Plant Physiol. Suppl. 35, xxxii.
- Butler W.L., Norris, K.H., Siegelman H.W. & Hendricks S.B. (1959) Detection, assay and preliminary purification of the pigment controlling photoresponsive development of plants. Proc. Natl. Acad. Sci. USA 45: 1703-1708.
- Canton F.R. & Quail P.H. (1999) Both phyA and phyB mediate light-imposed repression of PHYA gene expression in Arabidopsis. Plant Physiol. 121: 1207-1216.
- Castenholz R.W. (1988) Culturing methods for cyanobacteria. Meth. Enzymol. 167: 68-93.
- Chen M., Chory J. & Fankhauser C. (2004) Light signal transduction in higher plants. Annu. Rev. Genet. 38: 87-117.
- Choi J.S., Chung Y.H., Moon Y.J., Kim C., Watanabe M., Song P.S., Joe C.O., Bogorad L. & Park Y.M. (1999) Photomovement of the gliding cyanobacterium *Synechocystis* sp. PCC6803. Photochem. Photobiol. 70: 95-102.
- Clough R.C., Jordan-Beebe E.T., Lohman K.N., Marita J.M., Walker J.M., Gatz C. & Vierstra R.D. (1999) Sequences within both the N- and C-terminal domains of phytochrome A are required for PFR ubiquitination and degradation. Plant J. 17: 155-167.
- Cole W.J., Chapman D.J. & Siegelman H.W. (1967) Structure of phycocyanobilin. J. Am. Chem. Soc. 89: 3643-3645.
- Cornah J.E., Terry M.J. & Smith A.G. (2003) Green or red: what stops the traffic in the tetrapyrrole pathway? Trends Plant Sci. 8: 224-230.
- Davis S.J., Kurepa J. & Vierstra R.D. (1999a) The *Arabidopsis thaliana* HY1 locus, required for phytochrome-chromophore biosynthesis, encodes a protein related to heme oxygenases. Proc. Natl. Acad. Sci. USA 96: 6541-6654.
- Davis S.J., Vener A.V. & Vierstra R.D. (1999b) Bacteriophytochromes: Phytochrome-like photoreceptors from nonphotosynthetic eubacteria. Science 286: 2517-2520.
- Dürring M., Huber R., Bode W., Rümbeli R. & Zuber H. (1990) Refined three-dimensional structure of phycoerythrocyanin from the cyanobacterium *Mastigocladius laminosus* at 2.7 Å. J. Mol. Biol. 211: 633-644.
- Dürring M., Schmidt G.B. & Huber R. (1991) Isolation, crystallization, crystal structure analysis and refinement of constitutive C-phycocyanin from the chromatically adapting cyanobacterium *Fremyella diplosiphon* at 1.66 Å resolution. J. Mol. Biol. 217: 577-592.

- Dux P., Rubinstenn G., Vuister G.W., Boelens R., Mulder F.A., Hard K., Hoff W.D., Kroon A.R., Crielaard W., Hellingwerf K.J. & Kaptein R. (1998) Solution structure and backbone dynamics of the photoactive yellow protein. *Biochemistry* 37: 12689-12699.
- Eisenmesser E.Z., Millet O., Labeikovsky W., Korzhnev D.M., Wolf-Watz M., Bosco D.A., Skalicky J.J., Kay L.E. & Kern D. (2005) Intrinsic dynamics of an enzyme underlies catalysis. *Nature* 438: 117-121.
- Esenbeck N. (1836) Über einen blau-rothen Farbstoff, der sich bei der Zersetzung von Oscillatoren bildet. *Liebigs Ann. Chem.* 17: 75-82.
- Esteban B., Carrascal M., Abian J. & Lamparter T. (2005) Light-induced conformational changes of cyanobacterial phytochrome Cph1 probed by limited proteolysis and autophosphorylation. *Biochemistry* 44: 450-461.
- Fairchild C.D., Zhao J., Zhou J., Colson S.E., Bryant D.A., Fisher W.R. & Glazer A.N. (1992) Phycocyanin α -subunit phycocyanobilin lyase. *Proc. Natl. Acad. Sci. USA* 89: 7017-7021.
- Falk H. & Müller N. (1985) On the chemistry of pyrrole pigments. *Magn. Reson. Chem.* 23: 353-357.
- Falk H. (1989) The chemistry of linear oligopyrrols and bile pigments. Springer Verlag, Wien.
- Fankhauser C. & Chory J. (1997) Light control of plant development. *Annu. Rev. Cell Dev. Biol.* 13: 203-229.
- Fankhauser C., Yeh K.C., Lagarias J.C., Zhang H., Elich T.D. & Chory J. (1999) Pks1, a substrate phosphorylated by phytochrome that modulates light signalling in *Arabidopsis*. *Science* 284: 1539-1541.
- Fiedler B., Broc D., Schubert H., Rediger A., Börner T. & Wilde A. (2004) Involvement of cyanobacterial phytochromes in growth under different light qualities and quantities. *Photochem. Photobiol.* 79: 551-555.
- Fischer H. & Plieninger H. (1942) Synthese des Biliverdins (Uteroverdins) und Bilirubins, der Biliverdine XIII α und III α , sowie der Vinylneoxanthosäure. *Hoppe-Seylers Z. Physiol. Chem.* 274: 231.
- Flint L.M. & McAlister E.D. (1937) Wavelengths of radiation in the visible spectrum promoting the germination of light sensitive lettuce seed. *Smithson. Misc. Collection*, 96: 1-11.
- Fodor S.P.A., Lagarias J.C. & Mathies R.A. (1988) Resonance Raman spectra of the Pr-form of phytochrome. *Photochem. Photobiol.* 48: 129-136.
- Fodor, S.P.A., Lagarias, J.C. & Mathies, R.A. (1990) Resonance Raman Analysis of the Pr and Pfr Forms of Phytochrome, *Biochemistry* 29: 11141-11146.
- Förstendorf H., Mummert E., Schäfer E., Scheer H. & Siebert F. (1996) Fourier-transform infrared spectroscopy of phytochrome: Difference spectra of the intermediates of the photoreactions. *Biochemistry* 35: 10793-10799.
- Förstendorf H., Benda C., Gärtner W., Storf M., Scheer H. & Siebert F. (2001) FTIR studies of phytochrome photoreactions reveal the C=O bands of the chromophore: consequences for its protonation states, conformation, and protein interaction. *Biochemistry* 40: 14952-14959.
- Förster T. (1967) Mechanism of energy transfer. *Comprehensive Biochemistry* 22: 61-80.
- Frankenberg N., Mukougawa K., Kohchi T., & Lagarias J.C. (2001) Functional genomic analysis of the HY2 family of ferredoxin-dependent bilin reductases from oxygenic photosynthetic organisms. *Plant Cell* 13: 965-978.
- Frauenfelder H., Parak F. & Young R.D. (1988) Conformational substates in proteins. *Annu. Rev. Biophys. Biophys. Chem.* 17: 451-79.
- Gambetta G.A. & Lagarias J.C. (2001) Genetic engineering of phytochrome biosynthesis in bacteria. *Proc. Natl. Acad. Sci. USA* 98: 10566-10571.

- Gantt E. & Conti S.F. (1965) The ultrastructure of Porphyridium cruentum. *J. Cell Biol.* 26: 365-381.
- Gantt E. & Conti S.F. (1966a) Phycobiliprotein localization in algae. *Brookhaven Symp. Biol.* 19: 393-405.
- Gantt E. & Conti S.F. (1966b) Granule associate with the chloroplast lamellae of Porphyridium cruentum. *J. Cell Biol.* 29: 423-434.
- Garcia-Dominguez M., Muro-Pastor M.I., Reyes J.C. & Florencio F.J. (2000) Light-dependent regulation of cyanobacterial phytochrome expression. *J. Bacteriol.* 182: 38-44.
- Gardner G., Pike C.S., Rice H.V. & Briggs W.R. (1971) "Disaggregation" of Phytochrome in Vitro—A Consequence of Proteolysis. *Plant Physiol.* 48: 686-693.
- Gardner K.H. & Kay L.E. (1998) The use of 2H, 13C, 15N multidimensional NMR to study the structure and dynamics of proteins. *Annu. Rev. Biophys. Biomol. Struct.* 27: 357-406.
- Garner W.W. & Allard H.A. (1920) Effect of the relative length of day and night and other factors of the environment on growth and reproduction in plants. *J. Agric. Res.* 18: 553-606.
- Giraud E., Fardoux J., Fourrier N., Hannibal L., Genty B., Bouyer P., Dreyfus B. & Vermeglio A. (2002) Bacteriophytochrome controls photosystem synthesis in anoxygenic bacteria. *Nature* 417: 202-205.
- Giraud E., Zappa S., Vuillet L., Adriano J.M., Hannibal L., Fardoux J., Berthomieu C., Bouyer P., Pignol D. & Vermeglio A. (2005) A new type of bacteriophytochrome acts in tandem with a classical bacteriophytochrome to control the antennae synthesis in Rhodopseudomonas palustris. *J. Biol. Chem.* 280: 32389-32397.
- Glazer A.N. & Fang S. (1973) Chromophore content of blue-green algal phycobiliproteins. *J. Biol. Chem.* 248: 659-662.
- Glazer, A.N. & Hixson C.S. (1975) Characterization of R-phycocyanin. Chromophore content of R-phycocyanin and C-phycocerythrin. *J. Biol. Chem.* 250: 5487-5495.
- Glazer A.N. (1985) Light harvesting by phycobilisomes. *Annu. Rev. Biophys. Biophys. Chem.* 14: 47-77.
- Glazer A.N. (1988) Phycobiliproteins. *Meth. Enzymol.* 167: 291-303.
- Gossauer A. & Hirsch W. (1974) Bile pigments. IV. Total synthesis of racemic phycocyanobilin (phycobiliverdin) and of homophycobiliverdin. *Justus Liebigs Ann. Chem.* 9: 1496-1513.
- Grubmayer K. & Stanek M. (1998) Deprotonated 2,3-Dihydrobilindiones: Models for the chromophore of the far-red absorbing form of phytochrome. *Chem. – Eur. J.* 4: 1660-1666.
- Hanzawa H., Inomata K., Kinoshita H., Kakiuchi T., Jayasundera K.P., Sawamoto D., Ohta A., Uchida K., Wada K. & Furuya M. (2001) In vitro assembly of phytochrome B apoprotein with synthetic analogs of the phytochrome chromophore. *Proc. Natl. Acad. Sci. USA* 98: 3612-3617.
- Creeth M.J. & Harding S.E. (1982) Some observations on a new type of point average molecular weight. *J. Biochem. Biophys. Methods* 7: 25-34.
- Heintz W. (1851) Über den in den Gallensteinen enthaltenen Farbstoff. *Poggendorfs Ann.* 84: 106.
- Hendricks S.B., Borthwick H.A. & Downs R.J. (1956) Pigment conversion in the formative responses of plants to radiation. *Proc. Natl. Acad. Sci. USA* 42: 19-26.
- Herdman M., Coursin T., Rippka R., Houmard J. & Tandeau de Marsac N. (2000) A new appraisal of the prokaryotic origin of eukaryotic phytochromes. *J. Mol. Evol.* 51: 205-213.

- Hershey H.P., Barker R.F., Idler K.B., Lissemore J.L. & Quail P.H. (1985) Analysis of cloned cDNA and genomic sequences for phytochrome: complete amino acid sequences for two gene products expressed in etiolated *Avena*. *Nucleic Acids Res.* 13: 8543-8559.
- Hershey H.P., Barker R.F., Idler K.B., Murray M.G. & Quail P.H. (1987) Nucleotide sequence and characterization of a gene encoding the phytochrome polypeptide from *Avena*. *Gene* 61: 339-348.
- Heyne K. (2001) Ultraschnelle Dynamik Protein-gesteuerter Reaktionen in Bakteriorhodopsin und bakteriellem Phytochrom. Dissertation, FU Berlin.
- Hoiland, H. (1986) Partial molecular volumes of biochemical model compounds in aqueous solutions. In: *Thermodynamic Data for Biochemistry and Biotechnology* (Hinz, H.J., Hrsg.), pp. 17-44, Springer Verlag, Berlin.
- Hübschmann T., Börner T., Hartmann E. & Lamparter T. (2001) Characterisation of the Cph1 holo-phytochrome from *Synechocystis* sp. PCC6803. *Eur. J. Biochem.* 268: 2055-2063.
- Hübschmann T., Yamamoto H., Gieler T., Murata N. & Börner T. (2005) Red and far-red light alter the transcript profile in the cyanobacterium *Synechocystis* sp. PCC6803: Impact of cyanobacterial phytochromes. *FEBS Lett.* 579: 1613-1618.
- Hughes J., Lamparter T., Mittmann F., Hartmann E., Gärtner W., Wilde A. & Börner T. (1997) A prokaryotic phytochrome. *Nature* 386: 663.
- Hunt R.E. & Pratt L.H. (1980) Partial characterization of undegraded oat phytochrome. *Biochemistry* 19: 390-394.
- Huq E., Al-Sady B. & Quail P.H. (2003) Nuclear translocation of the photoreceptor phytochrome B is necessary for its biological function in seedling photomorphogenesis. *Plant J.* 35: 660-664.
- Inomata K., Hammam M.A., Kinoshita H., Murata Y., Khawn H., Noack S., Michael N. & Lamparter T. (2005) Sterically locked synthetic bilin derivatives and phytochrome Agp1 from *Agrobacterium tumefaciens* form photoinsensitive Pr- and Pfr-like adducts. *J. Biol. Chem.* 280: 24491-24497.
- Jacobi P.A. & Pippin D. (2001) Enantioselective syntheses of (13)C-labeled (2R)- and (2S)-phytochromobilin dimethyl ester. *Org. Lett.* 3:827-830.
- Jaenicke R. (1999) Stability and folding of domain proteins. *Prog. Biophys. Mol. Biol.* 71: 155-241.
- Jones A.M. & Erickson H.P. (1989) Domain structure of phytochrome from *Avena sativa* visualized by electron microscopy. *Photochem. Photobiol.* 49: 479-483.
- Jones A.M. & Quail P.H. (1986) Quaternary structure of 124-kilodalton phytochrome from *Avena sativa* L. *Biochemistry* 25: 2987-2995.
- Jorissen H.J., Quest B., Remberg A., Coursin T., Braslavsky S.E., Schaffner K., Tandeau de Marsac, N. & Gärtner W. (2002) Two independent, light-sensing two-component systems in a filamentous cyanobacterium. *Eur. J. Biochem.* 269: 2662-2671.
- Karniol B. & Vierstra R.D. (2003) The pair of bacteriophytochromes from *Agrobacterium tumefaciens* are histidine kinases with opposing photobiological properties. *Proc. Natl. Acad. Sci. USA* 100: 2807-2812.
- Katz J.J. & Crespi H.L. (1966) Deuterated organisms: cultivation and uses. *Science* 151: 1187-1194.
- Kehoe D.M. & Grossman A.R. (1996) Similarity of a chromatic adaption sensor to phytochrome and ethylene receptors. *Science* 273: 1409-1412.

- Klein G., Grombein S. & Rüdiger W. (1977) On the linkages between chromophore and protein in biliproteins, VI[1]. Structure and protein linkage of the phytochrome chromophore. Hoppe Seylers Z. Physiol. Chem. 358: 1077-1079.
- Kneip C., Hildebrandt P., Schlamann W., Braslavsky S.E., Mark F. & Schaffner K. (1999) Protonation state and structural changes of the tetrapyrrole chromophore during the Pr → Pfr phototransformation of phytochrome: A resonance Raman spectroscopic study. Biochemistry 38: 15185-15192.
- Kneip C., Mozley D., Hildebrandt P., Gärtner W., Braslavsky S.E. & Schaffner K. (1997) Effect of chromophore exchange on the resonance Raman spectra of recombinant phytochrome. FEBS Lett. 414: 23-26.
- Kneip, C. (1998) FT-NIR-Resonanz-Raman-spektroskopische Untersuchungen an Phytochrom und Tetrapyrrolpigmenten aus Cyanobakterien. Dissertation, Gerhard-Mercator-Universität-Gesamthochschule Duisburg.
- Knipp B., Kneip C., Matysik J., Gärtner W., Hildebrandt P., Braslavsky S.E. & Schaffner K. (1997) Regioselective deuteration and resonance Raman spectroscopic characterization of biliverdin and phycocyanobilin. Chem. – Eur. J. 3: 363-367.
- Kohchi T., Mukougawa K., Frankenberg N., Masuda M., Yokota A. & Lagarias J.C. (2001) The Arabidopsis hy2 gene encodes phytochromobilin synthase, a ferredoxin-dependent biliverdin reductase. Plant Cell 13: 425-436.
- Koorneef M., Rolff E. & Spruit C.J.P. (1980) Genetic control of light-inhibited hypocotyl elongation in *Arabidopsis thaliana* L. Heynh. Z. Pflanzenphysiologie 100: 147-160.
- Kützing, F.T. (1843) *Phycologia generalis, oder Anatomie, Physiologie und Systemkunde der Tange*. FA Brockhaus, Leipzig.
- Kylin H. (1910) Über Phykoerythrin und Phykocyan bei *Ceramium rubrum* (Huds.). Hoppe Seylers Z. Physiol. Chem. 69: 169-239.
- Kylin H. (1912) Über die roten und blauen Farbstoffe der Algen. Hoppe Seylers Z. Physiol. Chem. 76: 396-425.
- Lagarias J.C. & Rapoport H. (1980) Chromopeptides from phytochrome. The structure and linkage of the Pr form of the phytochrome chromophore. J. Am. Chem. Soc. 102: 4821-4828.
- Lagarias J.C. & Lagarias D.M. (1989) Self-assembly of synthetic phytochrome holoprotein in vitro. Proc. Natl. Acad. Sci. USA 86: 5778-5780.
- Lamarter T., Mittmann F., Gärtner W., Börner T., Hartmann E. & Hughes J. (1997) Characterization of recombinant phytochrome from the cyanobacterium *Synechocystis*. Proc. Natl. Acad. Sci. USA 94: 11792-11797.
- Lamarter T., Esteban B. & Hughes J. (2001) Phytochrome Cph1 from the cyanobacterium *Synechocystis PCC6803*. Purification, assembly, and quaternary structure. Eur. J. Biochem. 268: 4720-4730.
- Lamarter T., Michael N., Mittmann F. & Esteban B. (2002) Phytochrome from *Agrobacterium tumefaciens* has unusual spectral properties and reveals an N-terminal chromophore attachment site. Proc. Natl. Acad. Sci. USA 99: 11628-11633.
- Lamarter T., Michael N., Caspani O., Miyata T., Shirai K. & Inomata K. (2003) Biliverdin binds covalently to *Agrobacterium* phytochrome 1 via its ring A vinyl side chain. J. Biol. Chem. 278: 33786-33792.

- Lamparter T., Carrascal M., Michael N., Martinez E., Rottwinkel G. & Abian J. (2004a) The Biliverdin chromophore binds covalently to a conserved cysteine residue in the N-terminus of Agrobacterium phytochrome Agp1. *Biochemistry* 43: 3659-3669.
- Lamparter T. (2004b) Evolution of cyanobacterial and plant phytochromes. *FEBS Lett.* 573: 1-5.
- Landgraf F.T., Forreiter C., Hurtado Pico A., Lamparter T. & Hughes J. (2001) Recombinant holophytochrome in *Escherichia coli*. *FEBS Lett.* 508: 459-462.
- Lemberg R. (1930) Chromoproteide der Rotalgen. II. Spaltung mit Pepsin und Säuren. Isolierung eines Pyrrolfarbstoffs. *Liebigs Ann. Chem.* 477: 195-245.
- Lemberg R. & Bader G. (1933) Die Phycobilin der Rotalgen. Überführung in Mesobilirubin und Dehydromesobiliverdin. *Liebigs Ann. Chem.* 505: 151-177.
- Limbach H.H., Hennig J., Kendrick R. & Yannoni C.S. (1984) Proton-transfer kinetics in solids: Tautomerism in free base porphines by ^{15}N CPMAS NMR. *J. Am. Chem. Soc.* 106: 4059-4060.
- Lindner I., Braslavsky S.E., Schaffner K. & Gärtner W. (2000) Model studies of phytochrome photochromism: Protein-mediated photoisomerization of a linear tetrapyrrole in the absence of covalent bonding. *Angew. Chem. Int. Ed.* 39: 3269-3271.
- Lindner I., Knipp B., Braslavsky S.E., Gärtner W. & Schaffner K. (1998) Ein neuartiger Chromophor verändert selektiv die spektralen Eigenschaften eines der beiden stabilen Zustände des pflanzlichen Photorezeptors Phytochrom. *Angew. Chem.* 110: 1943-1946.
- Magdó I., Németh K., Mark F., Hildebrandt P. & Schaffner K. (1999) Calculation of Vibrational Spectra of Linear Tetrapyrroles. 1. Global Sets of Scaling Factors for Force Fields Derived by ab Initio and Density Functional Theory Methods. *J. Phys. Chem. A* 103: 289-303.
- Mas P., Devlin P.F., Panda S. & Kay S.A. (2000) Functional interaction of phytochrome B and Cryptochrome 2. *Nature* 408: 207-211.
- Matsushita T., Mochizuki N. & Nagatani A. (2003) Dimers of the N-terminal domain of phytochrome B are functional in the nucleus. *Nature* 424: 571-573.
- Matysik J., Hildebrandt P., Schlamann W., Braslavsky S.E. & Schaffner K. (1995) Fourier-Transform resonance Raman spectroscopy of intermediates of the phytochrome photocycle. *Biochemistry* 34: 10497-10507.
- Mayer G., Anderka O., Ludwig B. & Schubert D. (2002) The state of association of the cytochrome bc1 complex from *Paracoccus denitrificans* in solution of dodecyl maltoside. *Progr. Colloid Polym. Sci.* 119: 77-83.
- Micura R. & Grubmayr K. (1995) Ein Seryliminoester des Phycocyanobilins als neues Modell für die Chromophor-Wechselwirkungen des Phytochroms. *Angew. Chem.* 107: 1896-1898.
- Micura R. & Grubmayr K. (1994) Long-wavelength absorbing derivatives of phycocyanobilin: New structural aspects on phytochrome. *Bioorg. Med. Chem. Lett.* 4: 2517-2522.
- Mittmann F., Brucker G., Zeidler M., Repp A., Abts T., Hartmann E. & Hughes J. (2004) Targeted knockout in *Phycomitrella* reveals direct actions of phytochrome in the cytoplasm. *Proc. Natl. Acad. Sci. USA* 101: 13939-13944.

- Mizutani Y., Tokutomi S. & Kitagawa T. (1994) Resonance Raman Spectra of the Intermediates in Phototransformation of Large Phytochrome: Deprotonation of the Chromophore in the bleached Intermediate. *Biochemistry* 33: 153-158.
- Mizutani Y., Tokutomi S., Aoyagi K., Horitsu K. & Kitagawa T. (1991) Resonance Raman Study on Intact Pea Phytochrome and Its Model Compounds: Evidence for Proton Migration during the Phototransformation. *Biochemistry* 30: 10693-10700.
- Montgomery B.L. & Lagarias J.C. (2002) Phytochrome ancestry: sensors of bilins and light. *Trends Plant Sci.* 7: 357-366.
- Morgan W.D., Kragt A. & Feeney J. (2000) Expression of deuterium-isotope-labelled protein in the yeast *pichia pastoris* for NMR studies. *J. Biomol. NMR.* 17: 337-347.
- Morgan M.S. & Morrison M. (1999) (Hrsg.) Models as Mediators. Perspectives on Natural and Social Science. Cambridge University Press, Cambridge, GB.
- Moss G.P. (1988) Nomenclature of tetrapyrroles. Recommendations 1986 IUPAC-IUB Joint Commission on Biochemical Nomenclature (JCBN). *Eur. J. Biochem.* 178: 277-328.
- Mroginski M.A., Németh K., Magdó I., Müller M., Robben U., Della Védova C., Hildebrandt P. & Mark F. (2000) Calculation of the Vibrational Spectra of Linear Tetrapyrroles. 2. Resonance Raman Spectra of Hexamethylpyrromethene Monomers. *J. Phys. Chem. B* 104: 10885-10899.
- Mroginski, M.A., Murgida, D.H., von Stetten, D., Kneip, C., Mark, F. & Hildebrandt, P. (2004) Determination of the chromophore structures in the photoinduced reaction cycle of phytochrome. *J. Am. Chem. Soc.* 126: 16734-16735.
- Mumford F.E. & Jenner E.L. (1966) Purification and characterization of phytochrome from oat seedlings. *Biochemistry* 5: 3657-3662.
- Nagy F. & Schäfer E. (2000) Nuclear and cytosolic events of light-induced, phytochrome-regulated signalling in higher plants. *EMBO J.* 19: 157-163.
- Nakasako M., Iwata T., Inoue K. & Tokutomi S. (2005) Light-induced global structural changes in phytochrome A regulating photomorphogenesis in plants. *FEBS J.* 272: 603-612.
- Ollerenshaw J.E., Tugarinov V. & Kay L.E. (2003) Methyl TROSY: Explanation and experimental verification. *Magn. Reson. Chem.* 41: 843-852.
- Otto H., Lamparter T., Borucki B., Hughes J. & Heyn M.P. (2003) Dimerization and inter-chromophore distance of Cph1 phytochrome from *Synechocystis*, as monitored by fluorescence homo- and hetero-energy transfer. *Biochemistry* 42: 5885-5895.
- Park C.M., Shim J.Y., Yang S.S., Kang J.G., Kim J.I., Luka Z. & Song P.S. (2000a) Chromophore-apoprotein interactions in *Synechocystis* sp. PCC6803 phytochrome Cph1. *Biochemistry* 39: 6349-6356.
- Park C.M., Kim J.I., Yang S.S., Kang J.G., Kang J.H., Shim J.Y., Chung Y.H., Park Y.M. & Song P.S. (2000b) A second photochromic bacteriophytochrome from *Synechocystis* sp. PCC6803: Spectral analysis and down-regulation by light. *Biochemistry* 39: 10840-10847.
- Parker M.W., Hendricks S.B., Borthwick H.A. & Scully N.J. (1945) Action spectrum for the photoperiodic control of floral initiation in biloxi soybean. *Science* 102: 152-155.

- Patzelt H., Simon B., ter Laak A., Kessler B., Kühne R., Schmieder P., Oesterhelt D. & Oschkinat H. (2002) The structure of the active center in dark-adapted bacteriorhodopsin by solution-state NMR spectroscopy. Proc. Natl. Acad. Sci. USA 99: 9765-9770.
- Pratt L.H. (1978) Molecular properties of phytochrome. Photochem. Photobiol. 27: 81-105.
- Prytulla S. (1994) Rekonstitution, Reinigung und NMR-spektroskopische Untersuchung eines Detergens-solubilisierten Membranproteins: Der Chlorophyll a/b-bindende Lichtsammlerkomplex LHCII. Dissertation, Technische Universität München.
- Quail P.H. (1991) Phytochrome: a light-activated molecular switch that regulates plant gene expression. Annu. Rev. Genet. 25: 389-409.
- Quail P.H. (1997) An emerging molecular map of the phytochromes. Plant Cell Environ. 20: 657-665.
- Quail P.H. (2002) Phytochrome photosensory signalling networks. Nat. Rev. Mol. Cell Biol. 3: 85-93.
- Quest B. & Gärtner W. (2004) Chromophore selectivity in bacterial phytochromes: dissecting the process of chromophore attachment. Eur. J. Biochem. 271: 1117-1126.
- Quest, Benjamin (2003) Biochemische und Spektroskopische Charakterisierung zweier cyanobakterieller Phytochrome aus Calothrix PCC7601. Dissertation, Ludwig-Maximilian Universität, München.
- Remberg A., Lindner I., Lamarter T., Hughes J., Kneip K., Hildebrandt P., Braslavsky S.E., Gärtner W. & Schaffner K. (1997) Raman spectroscopic and light-induced kinetic characterization of a recombinant phytochrome of the cyanobacterium Synechocystis. Biochemistry 36: 13389-13395.
- Richmond A. (1986) (Hrsg.) CRC Handbook of microalgal mass culture. CRC, Boca Raton, Florida, USA.
- Rice H.V. & Briggs W.R. (1973) Partial Characterization of Oat and Rye Phytochrome. Plant Physiol. 51: 927-938.
- Robben U. (2000) Darstellung neuartiger Tetrapyrrolverbindungen und Charakterisierung ihrer Eigenschaften als Chromophore in rekombinantem Phytochrom. Dissertation, Universität Duisburg.
- Romero L.C. & Lam E. (1993) Guanine nucleotide binding protein involvement in early steps of phytochrome-regulated gene expression. Proc. Natl. Acad. Sci. USA 90: 1465-1469.
- Rubinstenn G., Vuister G.W., Mulder F.A., Dux P.E., Boelens R., Hellingwerf K.J. & Kaptein R. (1998) Structural and dynamic changes of photoactive yellow protein during its photocycle in solution. Nat. Struct. Biol. 5: 568-570.
- Rüdiger W. & Correl D.L. (1969) Über die Struktur des Phytochrom-Chromophors und seine Proteinbindung. Liebigs Ann. Chem. 723: 208-212.
- Rüdiger W., Thümmler F., Cmiel E. & Schneider S. (1983) Chromophore structure of the physiologically active form (Pfr) of phytochrome. Proc. Natl. Acad. Sci. USA 80: 6244-6248.
- Ryu J.S., Kim J.I., Kunkel T., Kim B.C., Cho D.S., Hong S.H., Kim S.H., Fernandez A.P., Kim Y., Alonso J.M., Ecker J.R., Nagy F., Lim P.O., Song P.S., Schäfer E. & Nam H.G. (2005) Phytochrome-specific type 5 phosphatase controls light signal flux by enhancing phytochrome stability and affinity for a signal transducer. Cell 120: 395-406.
- Sambrook J., Fritsch E.F. & Maniatis M. (1989) Molecular cloning: A laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, Connecticut, USA.
- Scheer H. (1981) Biliproteine. Angew. Chem. 93: 230-250.

- Schmitz O., Katayama M., Williams S.B., Kondo T. & Golden S.S. (2000) CikA, a bacteriophytocrome that resets the cyanobacterial circadian clock. *Science* 289: 765-768.
- Schneider-Poetsch H.A.W. (1992) Signal transduction by phytochrome: phytochromes have a module related to the transmitter modules of bacterial sensor proteins. *Photochem. Photobiol.* 56: 839-846.
- Sidler W. (1994) Phycobilisomes and phycobiliprotein structures. In: *Advances in photosynthesis: The molecular biology of cyanobacteria* (Bryant D.A., Hrsg.), pp. 139-216, Kluwer Academic Publisher, Dordrecht, Niederlande.
- Siebert F., Grimm R., Rüdiger W., Schmidt G. & Scheer H. (1990) Infrared spectroscopy of phytochrome and model pigments. *Eur. J. Biochem.* 194: 921-928.
- Siebzehnrübel S., Fischer R., Kufer W. & Scheer H. (1989) Photochemistry of phycobiliproteins: reciprocity of reversible photochemistry and aggregation in phycoerythrocyanin from *Mastigocladus laminosus*. *Photochem. Photobiol.* 49: 753-761.
- Siegelman H.W. & Firer E.M. (1964) Purification of phytochrome from oat seedlings. *Biochemistry* 3: 418-423.
- Sineshchekov V., Koppel' L., Esteban B., Hughes J. & Lamparter T. (2002) Fluorescence investigation of the recombinant cyanobacterial phytochrome (Cph1) and its C-terminally truncated monomeric species (Cph1Δ2): implication for holoprotein assembly, chromophore-apoprotein interaction and photochemistry. *J. Photochem. Photobiol.* 67: 39-50.
- Sineshchekov V.A. (1995) Photobiophysics and photobiochemistry of the heterogeneous phytochrome system. *Biochim. Biophys. Acta* 1228: 125-164.
- Sineshchekov V.A., Hughes J., Hartmann E. & Lamparter T. (1998) Fluorescence and photochemistry of recombinant phytochrome from the cyanobacterium *Synechocystis*. *Photochem. Photobiol.* 67: 263-267.
- Singh A.K. & Sherman L.A. (2005) Pleiotropic effect of a histidine kinase on carbohydrate metabolism in *Synechocystis* sp. strain PCC6803 and its requirement for heterotrophic growth. *J. Bacteriol.* 187: 2368-2376.
- Sklenal V. & Bax A. (1987) Spin-echo water suppression for the generation of pure-phase two-dimensional NMR spectra. *J. Magn. Res.* 74: 469-479.
- Smith H. (2000) Phytochromes and light signal perception by plants – an emerging synthesis. *Nature* 407: 585-591.
- Somers D.E. & Quail P.H. (1995) Temporal and spatial expression patterns of PHYA and PHYB genes in *Arabidopsis*. *Plant J.* 7: 413-427.
- Song P.S., Sarkar H.K., Tabba H. & Smith K.M. (1982) The phototransformation of phytochrome probed by 360 MHz proton NMR spectra. *Biochem. Biophys. Res. Commun.* 105: 179-287.
- Städeler G. (1864) Über die Farbstoffe der Galle. *Liebigs Ann. Chem.* 132: 323.
- Stock A.M., Robinson V.L. & Goudreau P.N. (2000) Two-component signal transduction. *Annu. Rev. Biochem.* 69: 183-215.
- Storf M. (2003) Chromophorbindung und Photochemie der α-Untereinheit des Phycoerythrocyanins aus *Mastigocladus laminosus*. Dissertation, Ludwig-Maximilian Universität, München.
- Storf M., Parbel A., Meyer M., Strohmann B., Scheer H., Deng M.G., Zheng M., Zhou M. & Zhao K.H. (2001) Chromophore attachment to biliproteins: Specificity of PecE/PecF, a lyase-isomerase for the photoactive 3¹-Cys-α84-phycoviolobilin chromophore of phycoerythrocyanin. *Biochemistry* 40: 12444-12456.

- Straume M. & Johnson M.L. (1992) Monte Carlo method for determining complete confidence probability distributions of estimated model parameters. *Methods Enzymol.* 210: 117-129.
- Svedberg, T. & Pedersen, K.O. (1940) Die Ultrazentrifuge. Theodor Steinkopff, Dresden.
- Swanson R.V., Zhou J., Leary J.A., Williams T., de Lorimier R., Bryant D.A. & Glazer A.N. (1992) Characterization of phycocyanin produced by cpcE and cpcF mutants and identification of an intergenic suppressor of the defect in bilin attachment. *J. Biol. Chem.* 267: 16146-16154.
- Sweere U., Eichenberg K., Lohrmann J., Mira-Rodado V., Baurle I., Kudla J., Nagy F., Schäfer E. & Harter K. (2001) Interaction of the response regulator ARR4 with phytochrome B in modulating red light signalling. *Science* 294: 1108-1111.
- Tasler R., Moises T. & Frankenberg-Dinkel N. (2005) Biochemical and spectroscopic characterization of the bacterial phytochrome of *Pseudomonas aeruginosa*. *FEBS J.* 272: 1927-1936.
- Terauchi K. & Ohmori M. (2004) Blue light stimulates cyanobacterial motility via a cAMP signal transduction system. *Mol. Microbiol.* 52: 303-309.
- Thenard L.J. (1807) *Mem. Phys. Chim. Arc.* 1: 23.
- Thümmel F. & Rüdiger W. (1983a) Models for the photoreversibility of phytochrome. Z,E-isomerization of chromopeptides from phycocyanin and phytochrome. *Tetrahedron* 39: 1943-1951.
- Thümmel F., Rüdiger W., Cmiel E. & Schneider S. (1983b) Chromopeptides from phytochrome and phycocyanin. NMR studies of the Pfr and Pr chromophore of phytochrome and E, Z isomeric chromophores of phycocyanin. *Z. Naturforsch.* 38c: 359-368.
- Tooley A.J., Cai Y.A. & Glazer A.N. (2001) Biosynthesis of a fluorescent cyanobacterial C-phycocyanin holo- α subunit in a heterologous host. *Proc. Natl. Acad. Sci. USA* 98: 10560-10565.
- Tugarinov V., Hwang P.M., Ollerenshaw J.E. & Kay L.E. (2003) Cross-correlated relaxation enhanced ^1H - ^{13}C NMR spectroscopy of methyl groups in very high molecular weight-proteins and protein complexes. *J. Am. Chem. Soc.* 125: 10420-10428.
- van Thor J.J., Borucki B., Crielaard W., Otto H., Lamparter T., Hughes J., Hellingwerf K.J. & Heyn M.P. (2001) Light-induced proton release and proton uptake reactions in the cyanobacterial phytochrome Cph1. *Biochemistry* 40: 11460-11471.
- Vierstra R.D. & Quail P. (1982) Native phytochrome: Inhibition of proteolysis yields a homogeneous monomer of 124 kilodaltons from *Avena*. *Proc. Natl. Acad. Sci. USA* 79: 5272-5276.
- Virchow R. (1847) Die pathologischen Pigmente. *Arch. Pathol. Anatom. Physiol. Klin. Med.* 1: 379-486.
- Volkman B.F., Lipson D., Wemmer D.E. & Kern D. (2001) Two-state allosteric behavior in a single-domain signaling protein. *Science*, 291: 2429-2433.
- von Stetten, D. (2005) Resonanz-Raman-spektroskopische Untersuchungen an *Agrobacterium*-Phytochrom. Diplomarbeit, Technische Universität Berlin.
- Wagner D., Fairchild C.D., Kuhn R.M. & Quail P.H. (1996) Chromophore-bearing NH₂-terminal domains of phytochromes A and B determine their photosensory specificity and differential light lability. *Proc. Natl. Acad. Sci. USA* 93: 4011-4015.

- Wang H. & Deng X.W. (2004) Phytochrome signaling mechanism. The Arabidopsis Book, <http://www.bioone.org/bioone/?request=get-toc&issn=1543-8120> (November 2005).
- Wang Y. & Poirier R.A. (1997) Factors that influence the C=N stretching frequency in Imines. *J. Phys. Chem.* 101: 907-912.
- Wegele R., Tasler R., Zeng Y., Rivera M. & Frankenberg-Dinkel N. (2004) The heme oxygenase(s)-phytochrome system of *Pseudomonas aeruginosa*. *J. Biol. Chem.* 279: 45791-45802.
- Weller J.-P. & Gossauer A. (1980) Synthese und Photoisomerisierung des racem. Phytochromobilin-dimethylesters. *Chem. Ber.* 113: 1603-1611.
- Wilde A., Fiedler B. & Börner T. (2002) The cyanobacterial phytochrome Cph2 inhibits phototaxis towards blue light. *Mol Microbiol.* 44: 981-988.
- Wu S.H. & Lagarias J.C. (2000) Defining the bilin lyase domain: lessons from the extended phytochrome superfamily. *Biochemistry* 39: 13487-13495.
- Yeh K.C. & Lagarias J.C. (1998) Eukaryotic phytochromes: Light-regulated serine/threonine protein kinases with histidine kinase ancestry. *Proc. Natl. Acad. Sci. USA* 95: 13976-13981.
- Yeh K.C., Wu S.H., Murphy J.T. and Lagarias J.C. (1997) A cyanobacterial phytochrome two-component light sensory system. *Science* 277: 1505-1508.
- Yoshihara S. & Ikeuchi M. (2004a) Phototactic motility in the unicellular cyanobacterium *Synechocystis* sp. PCC6803. *Photochem. Photobiol. Sci.* 3: 512-518.
- Yoshihara S., Katayama M., Geng X. & Ikeuchi M. (2004b) Cyanobacterial phytochrome-like PixJ1 holoprotein shows novel reversible photoconversion between blue- and green-absorbing forms. *Plant Cell Physiol.* 45: 1729-1737.
- Yphantis, D.A. (1964) Equilibrium Ultracentrifugation of Dilute Solutions. *Biochemistry* 3: 297-317.
- Zambre M., Terryn N., de Clercq J., de Buck S., Dillen W., van Montagu M., van der Straeten D. & Angenon G. (2003) Light strongly promotes gene transfer from *Agrobacterium tumefaciens* to plant cells. *Planta* 216: 580-586.
- Zhao K.H. & Scheer H. (1995) Type I and type II reversible photochemistry of phycoerythrocyanin α -subunit from *Mastigocladus laminsosus* both involve Z-E-isomerization of phycobilin chromophore and are controlled by sulfhydryls in apoprotein. *Biochim. Biophys. Acta* 1228: 244-253.