

## 8 Literaturverzeichnis

1. **Abe, A., M. de Grado, R. A. Pfuetzner, C. Sanchez-Sanmartin, R. Devinney, J. L. Puente, N. C. Strynadka, and B. B. Finlay.** 1999. Enteropathogenic *Escherichia coli* translocated intimin receptor, Tir, requires a specific chaperone for stable secretion. *Mol. Microbiol.* **33**(6):1162-75.
2. **Adhikari, P., G. Allison, B. Whittle, and N. K. Verma.** 1999. Serotype 1a O-antigen modification: molecular characterization of the genes involved and their novel organization in the *Shigella flexneri* chromosome. *J. Bacteriol.* **181**(15):4711-8.
3. **Adu-Bobie, J., G. Frankel, C. Bain, A. G. Goncalves, L. R. Trabulsi, G. Douce, S. Knutton, and G. Dougan.** 1998. Detection of intimins alpha, beta, gamma, and delta, four intimin derivatives expressed by attaching and effacing microbial pathogens. *J. Clin. Microbiol.* **36**(3):662-8.
4. **Albert, M. J., K. Alam, M. Islam, J. Montanaro, A. S. Rahaman, K. Haider, M. A. Hossain, A. K. Kibriya, and S. Tzipori.** 1991. *Hafnia alvei*, a probable cause of diarrhea in humans. *Infect. Immun.* **59**(4):1507-13.
5. **Albert, M. J., S. M. Faruque, M. Ansaruzzaman, M. M. Islam, K. Haider, K. Alam, I. Kabir, and R. Robins-Browne.** 1992. Sharing of virulence-associated properties at the phenotypic and genetic levels between enteropathogenic *Escherichia coli* and *Hafnia alvei*. *J. Med. Microbiol.* **37**(5):310-4.
6. **Al-Hasani, K., B. Adler, K. Rajakumar, and H. Sakellaris.** 2001. Distribution and structural variation of the she pathogenicity island in enteric bacterial pathogens. *J. Med. Microbiol.* **50**(9):780-6.
7. **Al-Hasani, K., K. Rajakumar, D. Bulach, R. Robins-Browne, B. Adler, and H. Sakellaris.** 2001. Genetic organization of the she pathogenicity island in *Shigella flexneri* 2a. *Microb. Pathog.* **30**(1):1-8.

8. **Armstrong, K. A., R. Acosta, E. Ledner, Y. Machida, M. Pancotto, M. McCormick, H. Ohtsubo, and E. Ohtsubo.** 1984. A 37 X 10(3) molecular weight plasmid-encoded protein is required for replication and copy number control in the plasmid pSC101 and its temperature-sensitive derivative pHS1. *J. Mol. Biol.* **175(3):331-48.**
9. **Ausubel, F. M.** 1994. *Current protocols in molecular biology.* John Wiley & Sons, New York.
10. **Bach, S., C. Buchrieser, M. Prentice, A. Guiyoule, T. Msadek, and E. Carniel.** 1999. The high-pathogenicity island of *Yersinia enterocolitica* Ye8081 undergoes low-frequency deletion but not precise excision, suggesting recent stabilization in the genome. *Infect. Immun.* **67(10):5091-9.**
11. **Bach, S., A. de Almeida, and E. Carniel.** 2000. The *Yersinia* high-pathogenicity island is present in different members of the family Enterobacteriaceae. *FEMS Microbiol. Lett.* **183(2):289-94.**
12. **Badea, L., S. Doughty, L. Nicholls, J. Sloan, R. M. Robins-Browne, and E. L. Hartland.** 2003. Contribution of Efa1/LifA to the adherence of enteropathogenic *Escherichia coli* to epithelial cells. *Microb. Pathog.* **34(5):205-15.**
13. **Barthold, S. W., G. L. Coleman, P. N. Bhatt, G. W. Osbaldiston, and A. M. Jonas.** 1976. The etiology of transmissible murine colonic hyperplasia. *Lab. Anim. Sci.* **26(6 Pt 1):889-94.**
14. **Benkel, P., M. Hudel, and T. Chakraborty.** 1997. The localisation of the LEE-region in the O26:H- Shiga toxin-producing *Escherichia coli* strain 413/89-1. Abstract in the Kongreß der DGHM in Jena, Germany(P196):53.
15. **Blanc-Potard, A. B., F. Solomon, J. Kayser, and E. A. Groisman.** 1999. The SPI-3 pathogenicity island of *Salmonella enterica*. *J. Bacteriol.* **181(3):998-1004.**

16. **Blattner, F. R., G. Plunkett, 3rd, C. A. Bloch, N. T. Perna, V. Burland, M. Riley, J. Collado-Vides, J. D. Glasner, C. K. Rode, G. F. Mayhew, J. Gregor, N. W. Davis, H. A. Kirkpatrick, M. A. Goeden, D. J. Rose, B. Mau, and Y. Shao.** 1997. The complete genome sequence of *Escherichia coli* K-12. *Science*. **277**(5331):1453-74.
17. **Blum, G., V. Falbo, A. Caprioli, and J. Hacker.** 1995. Gene clusters encoding the cytotoxic necrotizing factor type 1, Prs- fimbriae and alpha-hemolysin form the pathogenicity island II of the uropathogenic *Escherichia coli* strain J96. *FEMS Microbiol. Lett.* **126**(2):189-95.
18. **Blum, G., M. Ott, A. Lischewski, A. Ritter, H. Imrich, H. Tschäpe, and J. Hacker.** 1994. Excision of large DNA regions termed pathogenicity islands from tRNA-specific loci in the chromosome of an *Escherichia coli* wild-type pathogen. *Infect. Immun.* **62**(2):606-14.
19. **Brenner, D. J., P. A. Grimont, A. G. Steigerwalt, G. R. Fanning, E. Ageron, and C. F. Riddle.** 1993. Classification of citrobacteria by DNA hybridization: designation of *Citrobacter farmeri* sp. nov., *Citrobacter youngae* sp. nov., *Citrobacter braakii* sp. nov., *Citrobacter werkmanii* sp. nov., *Citrobacter sedlakii* sp. nov., and three unnamed *Citrobacter* genomospecies. *Int. J. Syst. Bacteriol.* **43**(4):645-58.
20. **Bruner, W., and H. Karch.** 2000. Genome plasticity in *Enterobacteriaceae*. *Int. J. Med. Microbiol.* **290**(2):153-65.
21. **Buchrieser, C., R. Brosch, S. Bach, A. Guiyoule, and E. Carniel.** 1998. The high-pathogenicity island of *Yersinia pseudotuberculosis* can be inserted into any of the three chromosomal asn tRNA genes. *Mol. Microbiol.* **30**(5):965-78.
22. **Bustamante, V. H., F. J. Santana, E. Calva, and J. L. Puente.** 2001. Transcriptional regulation of type III secretion genes in enteropathogenic *Escherichia coli*: Ler antagonizes H-NS-dependent repression. *Mol. Microbiol.* **39**(3):664-78.

23. **Campos, L. C., T. S. Whittam, T. A. Gomes, J. R. Andrade, and L. R. Trabulsi.** 1994. *Escherichia coli* serogroup O111 includes several clones of diarrheagenic strains with different virulence properties. *Infect. Immun.* **62**(8):3282-8.
24. **Carniel, E., I. Guilvout, and M. Prentice.** 1996. Characterization of a large chromosomal "high-pathogenicity island" in biotype 1B *Yersinia enterocolitica*. *J. Bacteriol.* **178**(23):6743-51.
25. **Cavignac, Y.** 2002. Detection and characterisation of a genetic island inserted in the *pheU*-locus in the enterohemorrhagic *E. coli* (EHEC) strain RW1374 (O102:H2) and in other EHEC strains. In Berlin: Humboldt-Universität, Berlin, Germany.
26. **Censini, S., C. Lange, Z. Xiang, J. E. Crabtree, P. Ghiara, M. Borodovsky, R. Rappuoli, and A. Covacci.** 1996. *cag*, a pathogenicity island of *Helicobacter pylori*, encodes type I- specific and disease-associated virulence factors. *Proc. Natl. Acad. Sci. U. S. A.* **93**(25):14648-53.
27. **Cheetham, B. F., D. B. Tattersall, G. A. Bloomfield, J. I. Rood, and M. E. Katz.** 1995. Identification of a gene encoding a bacteriophage-related integrase in a *vap* region of the *Dichelobacter nodosus* genome. *Gene.* **162**(1):53-8.
28. **Cieslewicz, M., and E. Vimr.** 1997. Reduced polysialic acid capsule expression in *Escherichia coli* K1 mutants with chromosomal defects in *kpsF*. *Mol. Microbiol.* **26**(2):237-49.
29. **Cornelis, G. R., A. Boland, A. P. Boyd, C. Geuijen, M. Iriarte, C. Neyt, M. P. Sory, and I. Stainier.** 1998. The virulence plasmid of *Yersinia*, an antihost genome. *Microbiol. Mol. Biol. Rev.* **62**(4):1315-52.
30. **Cornelis, G. R., and H. Wolf-Watz.** 1997. The *Yersinia* Yop virulon: a bacterial system for subverting eukaryotic cells. *Mol. Microbiol.* **23**(5):861-7.

31. **Cravioto, A., R. J. Gross, S. M. Scotland, and B. Rowe.** 1979. An adhesive factor found in strains of *Escherichia coli* belonging to the traditional infantile enteropathogenic serotypes. *Curr. Microbiol.* **3**(1):95-99.
32. **Deng, W., Y. Li, B. A. Vallance, and B. B. Finlay.** 2001. Locus of enterocyte effacement from *Citrobacter rodentium*: sequence analysis and evidence for horizontal transfer among attaching and effacing pathogens. *Infect. Immun.* **69**(10):6323-35.
33. **Donnenberg, M. S., and J. B. Kaper.** 1992. Enteropathogenic *Escherichia coli*. *Infect. Immun.* **60**(10):3953-61.
34. **Donnenberg, M. S., L. C. Lai, and K. A. Taylor.** 1997. The locus of enterocyte effacement pathogenicity island of enteropathogenic *Escherichia coli* encodes secretion functions and remnants of transposons at its extreme right end. *Gene.* **184**(1):107-14.
35. **Donnenberg, M. S., C. O. Tacket, S. P. James, G. Losonsky, J. P. Nataro, S. S. Wasserman, J. B. Kaper, and M. M. Levine.** 1993. Role of the *eaeA* gene in experimental enteropathogenic *Escherichia coli* infection. *J. Clin. Invest.* **92**(3):1412-7.
36. **Donnenberg, M. S., and T. S. Whittam.** 2001. Pathogenesis and evolution of virulence in enteropathogenic and enterohemorrhagic *Escherichia coli*. *J. Clin. Invest.* **107**(5):539-48.
37. **Dower, W. J., J. F. Miller, and C. W. Ragsdale.** 1988. High efficiency transformation of *E. coli* by high voltage electroporation. *Nucleic. Acids. Res.* **16**(13):6127-45.
38. **Elliott, S. J., S. W. Hutcheson, M. S. Dubois, J. L. Mellies, L. A. Wainwright, M. Batchelor, G. Frankel, S. Knutton, and J. B. Kaper.** 1999. Identification of CesT, a chaperone for the type III secretion of Tir in enteropathogenic *Escherichia coli*. *Mol. Microbiol.* **33**(6):1176-89.

39. **Elliott, S. J., V. Sperandio, J. A. Giron, S. Shin, J. L. Mellies, L. Wainwright, S. W. Hutcheson, T. K. McDaniel, and J. B. Kaper.** 2000. The locus of enterocyte effacement (LEE)-encoded regulator controls expression of both LEE- and non-LEE-encoded virulence factors in enteropathogenic and enterohemorrhagic *Escherichia coli*. *Infect. Immun.* **68**(11):6115-26.
40. **Elliott, S. J., L. A. Wainwright, T. K. McDaniel, K. G. Jarvis, Y. K. Deng, L. C. Lai, B. P. McNamara, M. S. Donnenberg, and J. B. Kaper.** 1998. The complete sequence of the locus of enterocyte effacement (LEE) from enteropathogenic *Escherichia coli* E2348/69. *Mol. Microbiol.* **28**(1):1-4.
41. **Feng, P., M. Dey, A. Abe, and T. Takeda.** 2001. Isogenic strain of *Escherichia coli* O157:H7 that has lost both Shiga toxin 1 and 2 genes. *Clin. Diagn. Lab. Immunol.* **8**(4):711-7.
42. **Fetherston, J. D., P. Schuetze, and R. D. Perry.** 1992. Loss of the pigmentation phenotype in *Yersinia pestis* is due to the spontaneous deletion of 102 kb of chromosomal DNA which is flanked by a repetitive element. *Mol. Microbiol.* **6**(18):2693-704.
43. **Finlay, B. B., I. Rosenshine, M. S. Donnenberg, and J. B. Kaper.** 1992. Cytoskeletal composition of attaching and effacing lesions associated with enteropathogenic *Escherichia coli* adherence to HeLa cells. *Infect. Immun.* **60**(6):2541-3.
44. **Frankel, G., A. D. Phillips, I. Rosenshine, G. Dougan, J. B. Kaper, and S. Knutton.** 1998. Enteropathogenic and enterohaemorrhagic *Escherichia coli*: more subversive elements. *Mol. Microbiol.* **30**(5):911-21.
45. **Friedberg, D., T. Umanski, Y. Fang, and I. Rosenshine.** 1999. Hierarchy in the expression of the locus of enterocyte effacement genes of enteropathogenic *Escherichia coli*. *Mol. Microbiol.* **34**(5):941-52.

46. **Galan, J. E.** 1999. Interaction of *Salmonella* with host cells through the centisome 63 type III secretion system. *Curr. Opin. Microbiol.* **2**(1):46-50.
47. **Gómez-Duarte, O. G., and J. B. Kaper.** 1995. A plasmid-encoded regulatory region activates chromosomal *eaeA* expression in enteropathogenic *Escherichia coli*. *Infect. Immun.* **63**(5):1767-76.
48. **Griffin, P. M., and R. V. Tauxe.** 1991. The epidemiology of infections caused by *Escherichia coli* O157:H7, other enterohemorrhagic *E. coli*, and the associated hemolytic uremic syndrome. *Epidemiol. Rev.* **13**:60-98.
49. **Hacker, J., L. Bender, M. Ott, J. Wingender, B. Lund, R. Marre, and W. Goebel.** 1990. Deletions of chromosomal regions coding for fimbriae and hemolysins occur in vitro and in vivo in various extraintestinal *Escherichia coli* isolates. *Microb. Pathog.* **8**(3):213-25.
50. **Hacker, J., G. Blum-Oehler, I. Mühldorfer, and H. Tschäpe.** 1997. Pathogenicity islands of virulent bacteria: structure, function and impact on microbial evolution. *Mol. Microbiol.* **23**(6):1089-97.
51. **Hacker, J., and J. B. Kaper.** 1999. The concept of pathogenicity islands, p. 1-11. In J. B. Kaper and J. Hacker (ed.), *Pathogenicity islands and other mobile virulence elements*. ASM Press, Washington, D.C.
52. **Hacker, J., and J. B. Kaper.** 2000. Pathogenicity islands and the evolution of microbes. *Annu. Rev. Microbiol.* **54**:641-79.
53. **Hacker, J., S. Knapp, and W. Goebel.** 1983. Spontaneous deletions and flanking regions of the chromosomally inherited hemolysin determinant of an *Escherichia coli* O6 strain. *J. Bacteriol.* **154**(3):1145-52.
54. **Hanahan, D.** 1983. Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* **166**(4):557-80.

- 
55. **Hare, J. M., A. K. Wagner, and K. A. McDonough.** 1999. Independent acquisition and insertion into different chromosomal locations of the same pathogenicity island in *Yersinia pestis* and *Yersinia pseudotuberculosis*. *Mol. Microbiol.* **31**(1):291-303.
56. **Hartland, E. L., M. Batchelor, R. M. Delahay, C. Hale, S. Matthews, G. Dougan, S. Knutton, I. Connerton, and G. Frankel.** 1999. Binding of intimin from enteropathogenic *Escherichia coli* to Tir and to host cells. *Mol. Microbiol.* **32**(1):151-8.
57. **Hayashi, T., K. Makino, M. Ohnishi, K. Kurokawa, K. Ishii, K. Yokoyama, C. G. Han, E. Ohtsubo, K. Nakayama, T. Murata, M. Tanaka, T. Tobe, T. Iida, H. Takami, T. Honda, C. Sasakawa, N. Ogasawara, T. Yasunaga, S. Kuhara, T. Shiba, M. Hattori, and H. Shinagawa.** 2001. Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. *DNA Res.* **8**(1):11-22.
58. **Hensel, M., T. Nikolaus, and C. Egelseer.** 1999. Molecular and functional analysis indicates a mosaic structure of *Salmonella* pathogenicity island 2. *Mol. Microbiol.* **31**(2):489-98.
59. **Hensel, M., J. E. Shea, S. R. Waterman, R. Mundy, T. Nikolaus, G. Banks, A. Vazquez-Torres, C. Gleeson, F. C. Fang, and D. W. Holden.** 1998. Genes encoding putative effector proteins of the type III secretion system of *Salmonella* pathogenicity island 2 are required for bacterial virulence and proliferation in macrophages. *Mol. Microbiol.* **30**(1):163-74.
60. **Heuvelink, A. E., N. C. van de Kar, J. F. Meis, L. A. Monnens, and W. J. Melchers.** 1995. Characterization of verocytotoxin-producing *Escherichia coli* O157 isolates from patients with haemolytic uraemic syndrome in Western Europe. *Epidemiol. Infect.* **115**(1):1-14.
61. **Hochhut, B., K. Jahreis, J. W. Lengeler, and K. Schmid.** 1997. CTnscr94, a conjugative transposon found in enterobacteria. *J. Bacteriol.* **179**(7):2097-102.



- 
62. **Hou, Y. M.** 1999. Transfer RNAs and pathogenicity islands. *Trends Biochem. Sci.* **24**(8):295-8.
63. **Itoh, K., T. Matsui, K. Tsuji, T. Mitsuoka, and K. Ueda.** 1988. Genetic control in the susceptibility of germfree inbred mice to infection by *Escherichia coli* O115a,c:K(B). *Infect. Immun.* **56**(4):930-5.
64. **Janben, T., C. Schwarz, P. Preikschat, M. Voss, H. C. Philipp, and L. H. Wieler.** 2001. Virulence-associated genes in avian pathogenic *Escherichia coli* (APEC) isolated from internal organs of poultry having died from colibacillosis. *Int. J. Med. Microbiol.* **291**(5):371-8.
65. **Janda, J. M., S. L. Abbott, and M. J. Albert.** 1999. Prototypal diarrheagenic strains of *Hafnia alvei* are actually members of the genus *Escherichia*. *J. Clin. Microbiol.* **37**(8):2399-401.
66. **Jarvis, K. G., J. A. Giron, A. E. Jerse, T. K. McDaniel, M. S. Donnenberg, and J. B. Kaper.** 1995. Enteropathogenic *Escherichia coli* contains a putative type III secretion system necessary for the export of proteins involved in attaching and effacing lesion formation. *Proc. Natl. Acad. Sci. U. S. A.* **92**(17):7996-8000.
67. **Jermyn, W. S., and E. F. Boyd.** 2002. Characterization of a novel *Vibrio* pathogenicity island (VPI-2) encoding neuraminidase (nanH) among toxigenic *Vibrio cholerae* isolates. *Microbiology* **148**(Pt 11):3681-93.
68. **Jerse, A. E., J. Yu, B. D. Tall, and J. B. Kaper.** 1990. A genetic locus of enteropathogenic *Escherichia coli* necessary for the production of attaching and effacing lesions on tissue culture cells. *Proc. Natl. Acad. Sci. U. S. A.* **87**(20):7839-43.
69. **Jores, J., L. Rumer, S. Kiessling, J. B. Kaper, and L. H. Wieler.** 2001. A novel locus of enterocyte effacement (LEE) pathogenicity island inserted at *pheV* in bovine Shiga toxin-producing *Escherichia coli* strain O103:H2. *FEMS Microbiol. Lett.* **204**(1):75-9.

70. **Jores, J., Rumer, L., Eichberg, J., and Wieler, L. H.** 2003. Presented at the VTEC-Congress, 5th International Symposium on Shiga Toxin (Verocytotoxin)-Producing *Escherichia coli* Infections, Edinburgh, Scotland.
71. **Jores, J., K. Zehmke, J. Eichberg, L. Rumer, and L. H. Wieler.** 2003. Description of a Novel Intimin Variant (Type zeta) in the Bovine O84:NM Verotoxin-Producing *Escherichia coli* Strain 537/89 and the Diagnostic Value of Intimin Typing. *Exp. Biol. Med. (Maywood)* **228**(4):370-6.
72. **Kalman, D., O. D. Weiner, D. L. Goosney, J. W. Sedat, B. B. Finlay, A. Abo, and J. M. Bishop.** 1999. Enteropathogenic *E. coli* acts through WASP and Arp2/3 complex to form actin pedestals. *Nat. Cell. Biol.* **1**(6):389-91.
73. **Kaper, J. B., J. L. Mellies, and J. P. Nataro.** 1999. Pathogenicity Islands and Other Mobile Genetic Elements of Diarrheagenic *Escherichia coli*, p. 33-58. In J. B. Kaper and J. Hacker (ed.), *Pathogenicity Islands and Other Mobile Virulence Elements*. ASM Press, Washington, D.C.
74. **Karaolis, D. K., J. A. Johnson, C. C. Bailey, E. C. Boedeker, J. B. Kaper, and P. R. Reeves.** 1998. A *Vibrio cholerae* pathogenicity island associated with epidemic and pandemic strains. *Proc. Natl. Acad. Sci. U. S. A.* **95**(6):3134-9.
75. **Karaolis, D. K., S. Somara, D. R. Maneval, Jr., J. A. Johnson, and J. B. Kaper.** 1999. A bacteriophage encoding a pathogenicity island, a type-IV pilus and a phage receptor in cholera bacteria. *Nature.* **399**(6734):375-9.
76. **Karch, H., T. Meyer, H. Russmann, and J. Heesemann.** 1992. Frequent loss of Shiga-like toxin genes in clinical isolates of *Escherichia coli* upon subcultivation. *Infect. Immun.* **60**(8):3464-7.
77. **Karch, H., S. Schubert, D. Zhang, W. Zhang, H. Schmidt, T. Ölschlager, and J. Hacker.** 1999. A genomic island, termed high-pathogenicity island, is present in certain non-O157 Shiga toxin-producing *Escherichia coli* clonal lineages. *Infect. Immun.* **67**(11):5994-6001.

- 
78. **Kenny, B., R. DeVinney, M. Stein, D. J. Reinscheid, E. A. Frey, and B. B. Finlay.** 1997. Enteropathogenic *E. coli* (EPEC) transfers its receptor for intimate adherence into mammalian cells. *Cell*. **91**(4):511-20.
79. **Klapproth, J. M., I. C. Scaletsky, B. P. McNamara, L. C. Lai, C. Malström, S. P. James, and M. S. Donnenberg.** 2000. A large toxin from pathogenic *Escherichia coli* strains that inhibits lymphocyte activation. *Infect. Immun.* **68**(4):2148-55.
80. **Knutton, S., T. Baldwin, P. H. Williams, and A. S. McNeish.** 1989. Actin accumulation at sites of bacterial adhesion to tissue culture cells: basis of a new diagnostic test for enteropathogenic and enterohemorrhagic *Escherichia coli*. *Infect. Immun.* **57**(4):1290-8.
81. **Knutton, S., I. Rosenshine, M. J. Pallen, I. Nisan, B. C. Neves, C. Bain, C. Wolff, G. Dougan, and G. Frankel.** 1998. A novel EspA-associated surface organelle of enteropathogenic *Escherichia coli* involved in protein translocation into epithelial cells. *Embo. J.* **17**(8):2166-76.
82. **Kovach, M. E., M. D. Shaffer, and K. M. Peterson.** 1996. A putative integrase gene defines the distal end of a large cluster of ToxR-regulated colonization genes in *Vibrio cholerae*. *Microbiology.* **142**(Pt 8):2165-74.
83. **Lalioui, L., and C. Le Bouguenec.** 2001. afa-8 Gene cluster is carried by a pathogenicity island inserted into the tRNA(Phe) of human and bovine pathogenic *Escherichia coli* isolates. *Infect. Immun.* **69**(2):937-48.
84. **Lawrence, J. G., and H. Ochman.** 1998. Molecular archaeology of the *Escherichia coli* genome. *Proc. Natl. Acad. Sci. U. S. A.* **95**(16):9413-7.
85. **LeClerc, J. E., B. Li, W. L. Payne, and T. A. Cebula.** 1996. High mutation frequencies among *Escherichia coli* and *Salmonella* pathogens. *Science.* **274**(5290):1208-11.

- 
86. **Lee, C. A.** 1996. Pathogenicity islands and the evolution of bacterial pathogens. *Infect. Agents. Dis.* **5**(1):1-7.
87. **Levine, M. M., P. Ristaino, G. Marley, C. Smyth, S. Knutton, E. Boedeker, R. Black, C. Young, M. L. Clements, C. Cheney, and et al.** 1984. Coli surface antigens 1 and 3 of colonization factor antigen II- positive enterotoxigenic *Escherichia coli*: morphology, purification, and immune responses in humans. *Infect. Immun.* **44**(2):409-20.
88. **Lindsay, J. A., A. Ruzin, H. F. Ross, N. Kurepina, and R. P. Novick.** 1998. The gene for toxic shock toxin is carried by a family of mobile pathogenicity islands in *Staphylococcus aureus*. *Mol. Microbiol.* **29**(2):527-43.
89. **Low, D., V. David, D. Lark, G. Schoolnik, and S. Falkow.** 1984. Gene clusters governing the production of hemolysin and mannose- resistant hemagglutination are closely linked in *Escherichia coli* serotype O4 and O6 isolates from urinary tract infections. *Infect. Immun.* **43**(1):353-8.
90. **Luperchio, S. A., J. V. Newman, C. A. Dangler, M. D. Schrenzel, D. J. Brenner, A. G. Steigerwalt, and D. B. Schauer.** 2000. *Citrobacter rodentium*, the causative agent of transmissible murine colonic hyperplasia, exhibits clonality: synonymy of *C. rodentium* and mouse-pathogenic *Escherichia coli*. *J. Clin. Microbiol.* **38**:4343-4350
91. **Mainil, J.** 1999. Shiga/verocytotoxins and Shiga/verotoxigenic *Escherichia coli* in animals. *Vet. Res.* **30**(2-3):235-57.
92. **Mariani-Kurkdjian, P., E. Denamur, A. Milon, B. Picard, H. Cave, N. Lambert-Zechovsky, C. Loirat, P. Goulet, P. J. Sansonetti, and J. Elion.** 1993. Identification of a clone of *Escherichia coli* O103:H2 as a potential agent of hemolytic-uremic syndrome in France. *J. Clin. Microbiol.* **31**(2):296-301.
93. **McDaniel, T. K., K. G. Jarvis, M. S. Sonnenberg, and J. B. Kaper.** 1995. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc. Natl. Acad. Sci. U. S. A.* **92**(5):1664-8.

94. **McDaniel, T. K., and J. B. Kaper.** 1997. A cloned pathogenicity island from enteropathogenic *Escherichia coli* confers the attaching and effacing phenotype on *E. coli* K-12. *Mol. Microbiol.* **23**(2):399-407.
95. **McGraw, E. A., J. Li, R. K. Selander, and T. S. Whittam.** 1999. Molecular Evolution and Mosaic Structure of  $\alpha$ ,  $\beta$  and  $\gamma$  Intimins of Pathogenic *Escherichia coli*. *Mol. Biol. Evol.* **16**(1):12-22.
96. **McGraw E. A. and T. S. Whittam.** 1996. Clonal analysis of EPEC serogroups. *Rev. Microbiol.* **27**:7-16.
97. **Mellies, J. L., S. J. Elliott, V. Sperandio, M. S. Donnenberg, and J. B. Kaper.** 1999. The Per regulon of enteropathogenic *Escherichia coli*: identification of a regulatory cascade and a novel transcriptional activator, the locus of enterocyte effacement (LEE)-encoded regulator (Ler). *Mol. Microbiol.* **33**(2):296-306.
98. **Miller, J. H.** 1992. A short course in bacterial genetics: a laboratory manual and handbook for *Escherichia coli* and related bacteria. In J. H. Miller (ed.), A short course in bacterial genetics: a laboratory manual and handbook for *Escherichia coli* and related bacteria. Cold Spring Harbor Laboratory Press, Plainview, N.Y.
99. **Mills, D. M., V. Bajaj, and C. A. Lee.** 1995. A 40 kb chromosomal fragment encoding *Salmonella typhimurium* invasion genes is absent from the corresponding region of the *Escherichia coli* K-12 chromosome. *Mol. Microbiol.* **15**(4):749-59.
100. **Miold, S., W. Rabsch, M. Rohde, S. Stender, H. Tschäpe, H. Russmann, E. Igwe, and W. D. Hardt.** 1999. Isolation of a temperate bacteriophage encoding the type III effector protein SopE from an epidemic *Salmonella typhimurium* strain. *Proc. Natl. Acad. Sci. U. S. A.* **96**(17):9845-50.
101. **Moon, H. W., S. C. Whipp, R. A. Argenzio, M. M. Levine, and R. A. Giannella.** 1983. Attaching and effacing activities of rabbit and human enteropathogenic *Escherichia coli* in pig and rabbit intestines. *Infect. Immun.* **41**(3):1340-51.

102. **Morschhäuser, J., V. Vetter, L. Emody, and J. Hacker.** 1994. Adhesin regulatory genes within large, unstable DNA regions of pathogenic *Escherichia coli*: cross-talk between different adhesin gene clusters. *Mol. Microbiol.* **11**(3):555-66.
103. **Moss, J. E., T. J. Cardozo, A. Zychlinsky, and E. A. Groisman.** 1999. The *selC*-associated SHI-2 pathogenicity island of *Shigella flexneri*. *Mol. Microbiol.* **33**(1):74-83.
104. **Mullis, K. B., and F. A. Faloona.** 1987. Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. *Methods Enzymol.* **155**:335-50.
105. **Nataro, J. P., and J. B. Kaper.** 1998. Diarrheagenic *Escherichia coli*. *Clin. Microbiol. Rev.* **11**(1):142-201.
106. **Nataro, J. P., J. Seriwatana, A. Fasano, D. R. Maneval, L. D. Guers, F. Noriega, F. Dubovsky, M. M. Levine, and J. G. Morris, Jr.** 1995. Identification and cloning of a novel plasmid-encoded enterotoxin of enteroinvasive *Escherichia coli* and *Shigella* strains. *Infect. Immun.* **63**(12):4721-8.
107. **Neves, B. C., R. Mundy, L. Petrovska, G. Dougan, S. Knutton, and G. Frankel.** 2003. CesD2 of Enteropathogenic *Escherichia coli* Is a Second Chaperone for the Type III Secretion Translocator Protein EspD. *Infect. Immun.* **71**(4):2130-41.
108. **Nicholls, L., T. H. Grant, and R. M. Robins-Browne.** 2000. Identification of a novel genetic locus that is required for in vitro adhesion of a clinical isolate of enterohaemorrhagic *Escherichia coli* to epithelial cells. *Mol. Microbiol.* **35**(2):275-88.
109. **O'Brien, A. D., V. L. Tesh, A. Donohue-Rolfe, M. P. Jackson, S. Olsnes, K. Sandvig, A. A. Lindberg, and G. T. Keusch.** 1992. Shiga toxin: biochemistry, genetics, mode of action, and role in pathogenesis. *Curr. Top. Microbiol. Immunol.* **180**:65-94.
110. **Okutani, A., T. Tobe, C. Sasakawa, R. Nozu, K. Gotoh, A. Takakura, T. Itoh, Y. Miyamoto, and K. Itoh.** 2001. Comparison of bacteriological, genetic and

- pathological characters between *Escherichia coli* O115a,c:K(B) and *Citrobacter rodentium*. *Exp. Anim.* **50**(2):183-6.
111. **Oswald, E., H. Schmidt, S. Morabito, H. Karch, O. Marches, and A. Caprioli.** 2000. Typing of intimin genes in human and animal enterohemorrhagic and enteropathogenic *Escherichia coli*: characterization of a new intimin variant. *Infect. Immun.* **68**(1):64-71.
112. **Perna, N. T., G. F. Mayhew, G. Pósfai, S. Elliott, M. S. Sonnenberg, J. B. Kaper, and F. R. Blattner.** 1998. Molecular evolution of a pathogenicity island from enterohemorrhagic *Escherichia coli* O157:H7. *Infect. Immun.* **66**(8):3810-7.
113. **Perna, N. T., G. Plunkett, 3rd, V. Burland, B. Mau, J. D. Glasner, D. J. Rose, G. F. Mayhew, P. S. Evans, J. Gregor, H. A. Kirkpatrick, G. Pósfai, J. Hackett, S. Klink, A. Boutin, Y. Shao, L. Miller, E. J. Grotbeck, N. W. Davis, A. Lim, E. T. Dimalanta, K. D. Potamousis, J. Apodaca, T. S. Anantharaman, J. Lin, G. Yen, D. C. Schwartz, R. A. Welch, and F. R. Blattner.** 2001. Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature.* **409**(6819):529-33.
114. **Pósfai, G., M. D. Koob, H. A. Kirkpatrick, and F. R. Blattner.** 1997. Versatile insertion plasmids for targeted genome manipulations in bacteria: isolation, deletion, and rescue of the pathogenicity island LEE of the *Escherichia coli* O157:H7 genome. *J. Bacteriol.* **179**(13):4426-8.
115. **Purdy, G. E., and S. M. Payne.** 2001. The SHI-3 iron transport island of *Shigella boydii* 0-1392 carries the genes for aerobactin synthesis and transport. *J. Bacteriol.* **183**(14):4176-82.
116. **Rajakumar, K., D. Bulach, J. Davies, L. Ambrose, C. Sasakawa, and B. Adler.** 1997. Identification of a chromosomal *Shigella flexneri* multi-antibiotic resistance locus which shares sequence and organizational similarity with the resistance region of the plasmid NR1. *Plasmid.* **37**(3):159-68.

- 
117. **Rajakumar, K., C. Sasakawa, and B. Adler.** 1997. Use of a novel approach, termed island probing, identifies the *Shigella flexneri* she pathogenicity island which encodes a homolog of the immunoglobulin A protease-like family of proteins. *Infect. Immun.* **65**(11):4606-14.
118. **Rakin, A., C. Noelting, P. Schropp, and J. Heesemann.** 2001. Integrative module of the high-pathogenicity island of *Yersinia*. *Mol. Microbiol.* **39**(2):407-15.
119. **Rakin, A., C. Noelting, S. Schubert, and J. Heesemann.** 1999. Common and specific characteristics of the high-pathogenicity island of *Yersinia enterocolitica*. *Infect. Immun.* **67**(10):5265-74.
120. **Rakin, A., P. Urbitsch, and J. Heesemann.** 1995. Evidence for two evolutionary lineages of highly pathogenic *Yersinia* species. *J. Bacteriol.* **177**(9):2292-8.
121. **Raleigh, E. A., N. E. Murray, H. Revel, R. M. Blumenthal, D. Westaway, A. D. Reith, P. W. Rigby, J. Elhai, and D. Hanahan.** 1988. McrA and McrB restriction phenotypes of some *E. coli* strains and implications for gene cloning. *Nucleic. Acids. Res.* **16**(4):1563-75.
122. **Ravatn, R., S. Studer, D. Springael, A. J. Zehnder, and J. R. van der Meer.** 1998. Chromosomal integration, tandem amplification, and deamplification in *Pseudomonas putida* F1 of a 105-kilobase genetic element containing the chlorocatechol degradative genes from *Pseudomonas* sp. Strain B13. *J. Bacteriol.* **180**(17):4360-9.
123. **Reid, S. D., D. J. Betting, and T. S. Whittam.** 1999. Molecular detection and identification of intimin alleles in pathogenic *Escherichia coli* by multiplex PCR. *J. Clin. Microbiol.* **37**(8):2719-22.
124. **Reid, S. D., C. J. Herbelin, A. C. Bumbaugh, R. K. Selander, and T. S. Whittam.** 2000. Parallel evolution of virulence in pathogenic *Escherichia coli*. *Nature.* **406**(6791):64-7.



- 
125. **Reiter, W. D., P. Palm, and S. Yeats.** 1989. Transfer RNA genes frequently serve as integration sites for prokaryotic genetic elements. *Nucleic. Acids. Res.* **17**(5):1907-14.
126. **Riley, L. W., R. S. Remis, S. D. Helgerson, H. B. McGee, J. G. Wells, B. R. Davis, R. J. Hebert, E. S. Olcott, L. M. Johnson, N. T. Hargrett, P. A. Blake, and M. L. Cohen.** 1983. Hemorrhagic colitis associated with a rare *Escherichia coli* serotype. *N. Engl. J. Med.* **308**(12):681-5.
127. **Rumer, L., J. Jores, P. Kirsch, Y. Cavnac, K. Zehmke, and L. H. Wieler.** 2003. Dissemination of pheU- and pheV-located genomic islands among enteropathogenic (EPEC) and enterohemorrhagic (EHEC) *E. coli* and their possible role in the horizontal transfer of the locus of enterocyte effacement (LEE). *Int. J. Med. Microbiol.* **292**(7-8):463-75.
128. **Ruzin, A., J. Lindsay, and R. P. Novick.** 2001. Molecular genetics of SaPI1-a mobile pathogenicity island in *Staphylococcus aureus*. *Mol. Microbiol.* **41**(2):365-77.
129. **Saiki, R. K., S. Scharf, F. Faloona, K. B. Mullis, G. T. Horn, H. A. Erlich, and N. Arnheim.** 1985. Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science.* **230**(4732):1350-4.
130. **Sambrook, J., and D. W. Russell.** 2001. *Molecular Cloning: A Laboratory Manual*, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
131. **Sanchez-SanMartin, C., V. H. Bustamante, E. Calva, and J. L. Puente.** 2001. Transcriptional regulation of the orf19 gene and the tir-cesT-eae operon of enteropathogenic *Escherichia coli*. *J. Bacteriol.* **183**(9):2823-33.
132. **Sandner, L., L. E. Eguiarte, A. Navarro, A. Cravioto, and V. Souza.** 2001. The elements of the locus of enterocyte effacement in human and wild mammal isolates of *Escherichia coli*: evolution by assemblage or disruption?. *Microbiology.* **147**:3149-3158.

- 
133. **Schauer, D. B., and S. Falkow.** 1993. The *eae* gene of *Citrobacter freundii* biotype 4280 is necessary for colonization in transmissible murine colonic hyperplasia. *Infect. Immun.* **61**(11):4654-61.
134. **Schauer, D. B., B. A. Zabel, I. F. Pedraza, C. M. O'Hara, A. G. Steigerwalt, and D. J. Brenner.** 1995. Genetic and biochemical characterization of *Citrobacter rodentium* sp. nov. *J. Clin. Microbiol.* **33**(8):2064-8.
135. **Schmidt, H., W. L. Zhang, U. Hemmrich, S. Jelacic, W. Brunder, P. I. Tarr, U. Dobrindt, J. Hacker, and H. Karch.** 2001. Identification and characterization of a novel genomic island integrated at *selC* in locus of enterocyte effacement-negative, Shiga toxin-producing *Escherichia coli*. *Infect. Immun.* **69**(11):6863-73.
136. **Schubert, S., S. Cuenca, D. Fischer, and J. Heesemann.** 2000. High-pathogenicity island of *Yersinia pestis* in enterobacteriaceae isolated from blood cultures and urine samples: prevalence and functional expression. *J. Infect. Dis.* **182**(4):1268-71.
137. **Schubert, S., A. Rakin, D. Fischer, J. Sorsa, and J. Heesemann.** 1999. Characterization of the integration site of *Yersinia* high-pathogenicity island in *Escherichia coli*. *FEMS Microbiol. Lett.* **179**(2):409-14.
138. **Schubert, S., A. Rakin, H. Karch, E. Carniel, and J. Heesemann.** 1998. Prevalence of the "high-pathogenicity island" of *Yersinia* species among *Escherichia coli* strains that are pathogenic to humans. *Infect. Immun.* **66**(2):480-5.
139. **Schwartz, D. C., and C. R. Cantor.** 1984. Separation of yeast chromosome-sized DNAs by pulsed field gradient gel electrophoresis. *Cell.* **37**(1):67-75.
140. **Shea, J. E., M. Hensel, C. Gleeson, and D. W. Holden.** 1996. Identification of a virulence locus encoding a second type III secretion system in *Salmonella typhimurium*. *Proc. Natl. Acad. Sci. U. S. A.* **93**(6):2593-7.
141. **Shizuya, H., B. Birren, U. J. Kim, V. Mancino, T. Slepak, Y. Tachiiri, and M. Simon.** 1992. Cloning and stable maintenance of 300-kilobase-pair fragments of

- human DNA in *Escherichia coli* using an F-factor-based vector. Proc. Natl. Acad. Sci. U. S. A. **89**(18):8794-7.
142. **Slilaty, S. N., and S. Lebel.** 1998. Accurate insertional inactivation of lacZalpha: construction of pTrueBlue and M13TrueBlue cloning vectors. Gene. **213**(1-2):83-91.
143. **Southern, E. M.** 1975. Detection of specific sequences among DNA fragments separated by gel electrophoresis. J. Mol. Biol. **98**(3):503-17.
144. **Sperandio, V., J. B. Kaper, M. R. Bortolini, B. C. Neves, R. Keller, and L. R. Trabulsi.** 1998. Characterization of the locus of enterocyte effacement (LEE) in different enteropathogenic *Escherichia coli* (EPEC) and Shiga-toxin producing *Escherichia coli* (STEC) serotypes. FEMS Microbiol. Lett. **164**(1):133-9.
145. **Sperandio, V., C. C. Li, and J. B. Kaper.** 2002. Quorum-sensing *Escherichia coli* regulator A: a regulator of the LysR family involved in the regulation of the locus of enterocyte effacement pathogenicity island in enterohemorrhagic *E. coli*. Infect. Immun. **70**(6):3085-93.
146. **Sperandio, V., J. L. Mellies, R. M. Delahay, G. Frankel, J. A. Crawford, W. Nguyen, and J. B. Kaper.** 2000. Activation of enteropathogenic *Escherichia coli* (EPEC) LEE2 and LEE3 operons by Ler. Mol. Microbiol. **38**(4):781-93.
147. **Sperandio, V., J. L. Mellies, W. Nguyen, S. Shin, and J. B. Kaper.** 1999. Quorum sensing controls expression of the type III secretion gene transcription and protein secretion in enterohemorrhagic and enteropathogenic *Escherichia coli*. Proc. Natl. Acad. Sci. U. S. A. **96**(26):15196-201.
148. **Sperandio, V., A. G. Torres, J. A. Giron, and J. B. Kaper.** 2001. Quorum sensing is a global regulatory mechanism in enterohemorrhagic *Escherichia coli* O157:H7. J. Bacteriol. **183**(17):5187-97.
149. **Sperandio, V., A. G. Torres, and J. B. Kaper.** 2002. Quorum sensing *Escherichia coli* regulators B and C (QseBC): a novel two- component regulatory system involved

- in the regulation of flagella and motility by quorum sensing in *E. coli*. *Mol. Microbiol.* **43**(3):809-21.
150. **Stalker, D. M., R. Kolter, and D. R. Helinski.** 1982. Plasmid R6K DNA replication. I. Complete nucleotide sequence of an autonomously replicating segment. *J. Mol. Biol.* **161**(1):33-43.
151. **Stevens, M. P., O. Marches, J. Campbell, V. Huter, G. Frankel, A. D. Phillips, E. Oswald, and T. S. Wallis.** 2002. Intimin, tir, and shiga toxin 1 do not influence enteropathogenic responses to shiga toxin-producing *Escherichia coli* in bovine ligated intestinal loops. *Infect. Immun.* **70**(2):945-52.
152. **Stevens, M. P., P. M. van Diemen, G. Frankel, A. D. Phillips, and T. S. Wallis.** 2002. Efa1 influences colonization of the bovine intestine by shiga toxin-producing *Escherichia coli* serotypes O5 and O111. *Infect. Immun.* **70**(9):5158-66.
153. **Sullivan, J. T., and C. W. Ronson.** 1998. Evolution of rhizobia by acquisition of a 500-kb symbiosis island that integrates into a phe-tRNA gene. *Proc. Natl. Acad. Sci. U. S. A.* **95**(9):5145-9.
154. **Swenson, D. L., N. O. Bukanov, D. E. Berg, and R. A. Welch.** 1996. Two pathogenicity islands in uropathogenic *Escherichia coli* J96: cosmid cloning and sample sequencing. *Infect. Immun.* **64**(9):3736-43.
155. **Tao, Q., and H. B. Zhang.** 1998. Cloning and stable maintenance of DNA fragments over 300 kb in *Escherichia coli* with conventional plasmid-based vectors. *Nucleic Acids. Res.* **26**(21):4901-9.
156. **Tarr, C. L., and T. S. Whittam.** 2002. Molecular evolution of the intimin gene in O111 clones of pathogenic *Escherichia coli*. *J. Bacteriol.* **184**(2):479-87.
157. **Tauschek, M., R. A. Strugnell, and R. M. Robins-Browne.** 2002. Characterization and evidence of mobilization of the LEE pathogenicity island of rabbit-specific strains of enteropathogenic *Escherichia coli*. *Mol. Microbiol.* **44**(6):1533-50.

- 
158. **Taylor, K. A., P. W. Luther, and M. S. Sonnenberg.** 1999. Expression of the EspB protein of enteropathogenic *Escherichia coli* within HeLa cells affects stress fibers and cellular morphology. *Infect. Immun.* **67**(1):120-5.
159. **Tobe, T., G. K. Schoolnik, I. Sohel, V. H. Bustamante, and J. L. Puente.** 1996. Cloning and characterization of bfpTVW, genes required for the transcriptional activation of bfpA in enteropathogenic *Escherichia coli*. *Mol. Microbiol.* **21**(5):963-75.
160. **Vallance, B. A., and B. B. Finlay.** 2000. Exploitation of host cells by enteropathogenic *Escherichia coli*. *Proc. Natl. Acad. Sci. U. S. A.* **97**(16):8799-806.
161. **Vial, P. A., R. Robins-Browne, H. Lior, V. Prado, J. B. Kaper, J. P. Nataro, D. Maneval, A. Elsayed, and M. M. Levine.** 1988. Characterization of enteroadherent-aggregative *Escherichia coli*, a putative agent of diarrheal disease. *J. Infect. Dis.* **158**(1):70-9.
162. **Vokes, S. A., S. A. Reeves, A. G. Torres, and S. M. Payne.** 1999. The aerobactin iron transport system genes in *Shigella flexneri* are present within a pathogenicity island. *Mol. Microbiol.* **33**(1):63-73.
163. **Wainwright, L. A., and J. B. Kaper.** 1998. EspB and EspD require a specific chaperone for proper secretion from enteropathogenic *Escherichia coli*. *Mol. Microbiol.* **27**(6):1247-60.
164. **Weiss, B., A. Jacquemin-Sablon, T. R. Live, G. C. Fareed, and C. C. Richardson.** 1968. Enzymatic breakage and joining of deoxyribonucleic acid. VI. Further purification and properties of polynucleotide ligase from *Escherichia coli* infected with bacteriophage T4. *J. Biol. Chem.* **243**(17):4543-55.
165. **Whittam, T. S., M. L. Wolfe, I. K. Wachsmuth, F. Orskov, I. Orskov, and R. A. Wilson.** 1993. Clonal relationships among *Escherichia coli* strains that cause hemorrhagic colitis and infantile diarrhea. *Infect. Immun.* **61**(5):1619-29.

- 
166. **Wieler, L. H.** 1997. Bestimmung von Virulenzfaktoren boviner Shiga-Toxin-bildender *Escherichia coli*-(STEC-) Stämme als Bewertungsgrundlage ihrer klinischen Bedeutung für Rind und Mensch. Shaker Verlag.
167. **Wieler, L. H., R. Bauerfeind, and G. Baljer.** 1992. Characterization of Shiga-like toxin producing *Escherichia coli* (SLTEC) isolated from calves with and without diarrhoea. Zentralbl. Bakteriol. **276**(2):243-53.
168. **Wieler, L. H., T. K. McDaniel, T. S. Whittam, and J. B. Kaper.** 1997. Insertion site of the locus of enterocyte effacement in enteropathogenic and enterohemorrhagic *Escherichia coli* differs in relation to the clonal phylogeny of the strains. FEMS Microbiol. Lett. **156**(1):49-53.
169. **Wieler, L. H., E. Vieler, C. Erpenstein, T. Schlapp, H. Steinrück, R. Bauerfeind, A. Byomi, and G. Baljer.** 1996. Shiga toxin-producing *Escherichia coli* strains from bovines: association of adhesion with carriage of *eae* and other genes. J. Clin. Microbiol. **34**(12):2980-4.
170. **Wong, K. K., M. McClelland, L. C. Stillwell, E. C. Sisk, S. J. Thurston, and J. D. Saffer.** 1998. Identification and sequence analysis of a 27-kilobase chromosomal fragment containing a *Salmonella* pathogenicity island located at 92 minutes on the chromosome map of *Salmonella enterica* serovar typhimurium LT2. Infect. Immun. **66**(7):3365-71.
171. **Wood, M. W., M. A. Jones, P. R. Watson, S. Hedges, T. S. Wallis, and E. E. Galyov.** 1998. Identification of a pathogenicity island required for *Salmonella* enteropathogenicity. Mol. Microbiol. **29**(3):883-91.
172. **Zehmke, K., J. Jores, L. Roumer, S. Kiessling, S. Lautenschlaeger, and L. H. Wieler.** 2000. Presented at the 4th international symposium and workshop on "Shiga Toxin (Verotoxin) - producing *Escherichia coli* infections, Kyoto/ Japan, 29.10.-02.11.2000.

173. **Zhang, W. L., B. Kohler, E. Oswald, L. Beutin, H. Karch, S. Morabito, A. Caprioli, S. Suerbaum, and H. Schmidt.** 2002. Genetic diversity of intimin genes of attaching and effacing *Escherichia coli* strains. *J. Clin. Microbiol.* **40**(12):4486-92.
174. **Zhu, C., T. S. Agin, S. J. Elliott, L. A. Johnson, T. E. Thate, J. B. Kaper, and E. C. Boedeker.** 2001. Complete nucleotide sequence and analysis of the locus of enterocyte Effacement from rabbit diarrheagenic *Escherichia coli* RDEC-1. *Infect. Immun.* **69**(4):2107-15.
175. **Ziebuhr, W., V. Krimmer, S. Rachid, I. Lössner, F. Götz, and J. Hacker.** 1999. A novel mechanism of phase variation of virulence in *Staphylococcus epidermidis*: evidence for control of the polysaccharide intercellular adhesin synthesis by alternating insertion and excision of the insertion sequence element IS256. *Mol. Microbiol.* **32**(2):345-56.