

7. Literatur

- Abdel-Monhem,M., Dürwald,H., und Hoffmann-Berling,H. (1976). Enzymic unwinding of DNA II. Chain separation by an ATP-dependent DNA unwinding enzyme. *Eur. J. Biochem.* 65, 441-449.
- Arai,K. und Kornberg,A. (1979). A general priming system employing only DnaB and primase for DNA replication. *Proc. Natl. Acad. Sci. USA* 76, 4308-4312.
- Arai,K., Yasuda,S., und Kornberg,A. (1981). Mechanism of dnaB protein action. I. Crystallisation and properties of dnaB protein, an essential replication protein in *Escherichia coli*. *J. Biol. Chem.* 256, 5247-5252.
- Armengod,M. und Lambies,E. (1986). Overlapping arrangement of the *recF* and *dnaN* operons of *E. coli*; positive and negative control sequences. *Gene* 43, 183-196.
- Asada,K., Sugimoto,K., Oka,A., Takanami,M., und Hirota,Y. (1982). Structure of replication origin of the *Escherichia coli* K-12 chromosome: The presence of spacer sequence in the *ori* region carrying information for autonomous replication. *Nucl. Acids Res.* 10, 3745-3754.
- Asai,T., Takanami,M., und Imai,M. (1990). The AT richness and *gid* transcription determine the left border of the replication origin of the *E.coli* chromosome. *EMBO J.* 9, 4065-4072.
- Atlung,T., Clausen,E., und Hansen,F.G. (1985). Autoregulation of the *dnaA* gene of *Escherichia coli*. *Mol. Gen. Genet.* 200, 442-450.
- Augustin,L.B., Jacobson,B.A., und Fuchs,J.A. (1994). *Escherichia coli* Fis and DnaA proteins bind specifically to the *nrd* promoter region and affect expression of an *nrd-lac* fusion. *J. Bacteriol.* 176, 378-387.
- Bachmann,B.J. (1990). Linkage map of *Escherichia coli* K-12, edition 8. *Microbiol. Rev.* 54, 130-197.
- Baker,T.A., Funnell,B.E., und Kornberg,A. (1987). Helicase action of dnaB protein during replication from the *Escherichia coli* chromosomal origin *in vitro*. *J. Biol. Chem.* 262, 6877-6885.
- Baker,T.A., Sekimizu,K., Funnell,B.E., und Kornberg,A. (1986). Extensive unwinding of the plasmid template during staged enzymatic initiation of DNA replication from the origin of the *E. coli* chromosome. *Cell* 45, 53-64.
- Bernardi,A. und Bernardi,F. (1984). Complete Sequence of pSC101. *Nucl. Acids Res.* 12, 9415-9424.
- Biek,D.P. und Cohen,S.N. (1989a). Involvement of Integration Host Factor (IHF) in Maintenance of Plasmid pSC101 in *Escherichia coli*:Characterization of pSC101 Mutants that Replicate in the absence of IHF. *J. Bact.* 171, 1656-2065.
- Biek,D.P. und Cohen,S.N. (1989b). Involvement of Integration Host Factor (IHF) in Maintenance of Plasmid pSC101 in *Escherichia coli*:Mutations in the topA gene allow pSC101Replication in the absence of IHF. *J. Bact.* 171, 2066-2074.
- Birnboim,B.J. und Doly,J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucl. Acids Res.* 7, 1513-1523.
- Biswas,E.E. und Biswas,S.B. (1999b). Mechanism of DNA binding by the DnaB helicase of *Escherichia coli*: Analysis of the roles of domain gamma in DNA binding. *Biochemistry* 38, 10929-10939.
- Biswas,E.E. und Biswas,S.B. (1999a). Mechanism of DnaB helicase of *Escherichia coli*: Structural domains involved in ATP hydrolysis, DNA binding, and oligomerization. *Biochemistry* 38, 10919-10928.

- Blaesing,F., Weigel,C., und Messer,W. (2000). Analysis of the DNA binding domain of *Escherichia coli* DnaA protein. in press.
- Bradford,M.M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Analyt. Biochem.* 72, 248-254.
- Bramhill,D. und Kornberg,A. (1988). Duplex opening by dnaA protein at novel sequences in initiation of replication at the origin of the *E. coli* chromosome. *Cell* 52, 743-755.
- Braun,R.E., O'Day,K., und Wright,A. (1985). Autoregulation of the DNA replication gene *dnaA* in *E. coli*. *Cell* 40, 159-169.
- Bravo,A. und Salas,M. (1998). Polymerization of Bacteriophage phi 29 Replication Protein p1 into protofilament sheets. *EMBO J.* 17, 6096-6105.
- Buhk,H.J. und Messer,W. (1983). Replication origin region of *Escherichia coli*: Nucleotide sequence and functional units. *Gene* 24, 265-279.
- Bujalowski,W. und Jezewska,M.J. (2000). Kinetic Mechanism of the Single-stranded DNA Recognition by *Escherichia coli* Replicative Helicase DnaB Protein. Application of the Matrix Projection Operator Technique to Analyze Stopped-Flow Kinetics. *J. Mol. Biol.* 295, 831-852.
- Bujalowski,W. (1999). *E.coli* DnaB Helicase-DNA Complexes: Mechanism of Recognition and Structure. Workshop on Helicases as Molecular Motors in Nucleic Acid Strand Separation.
- Bujalowski,W. und Jezewska,M.J. (1995). Interactions of *Escherichia coli* primary replicative helicase DnaB protein with single-stranded DNA. The nucleic acid does not wrap around the protein hexamer. *Biochem.* 34, 8519.
- Bujalowski,W., Klonowska,M.M., und Jezewska,M.J. (1994). Oligomeric structure of *Escherichia coli* primary replicative helicase DnaB protein. *J. Biol. Chem.* 269, 31350-31358.
- Cairns,J. (1963). The bacterial chromosome and its manner of replication as seen by autoradiography. *J. Mol. Biol.* 6, 208-213.
- Carr,K.M. und Kaguni,J.M. (1996). The A184V missense mutation of the *dnaA5* and *dnaA46* alleles confers a defect in ATP binding and thermolability in initiation of *Escherichia coli* DNA replication. *Mol. Microbiol.* 20, 1307-1318.
- Cassler,M.R., Grimwade,J.E., und Leonard,A.C. (1995). Cell cycle-specific changes in nucleoprotein complexes at a chromosomal replication origin. *EMBO J.* 14, 5833-5841.
- Chakraborty,T., Yoshinaga,K., Lother,H., und Messer,W. (1982). Purification of the *E. coli* *dnaA* gene product. *EMBO J.* 1, 1545-1549.
- Chang,A.C.Y. und Cohen,S.N. (1978). Construction and characterization of amplifiable multicopy DNA cloning vehicles derived from the P15A cryptic miniplasmid. *J. Bact.* 134, 1141-1156.
- Chung,C.T., Niemela,S.L., und Miller,R.H. (1989). One-step preparation of competent *Escherichia coli*: transformation and storage of bacterial cells in the same solution. *Proceedings of the National Academy of Sciences of the United States of America* 86, 2172-2175.
- Cohen,S.N. und Chang,A.C.Y. (1973). Recircularization and autonomous Replication of a sheared R-Factor DNA segment in *Escherichia coli* transformants. *Proc. Natl. Acad. Sci.* 70, 1293-1297.
- Cooper,S. und Helmstetter,C.E. (1968). Chromosome replication and the division cycle of *Escherichia coli* B/r. *J. Mol. Biol.* 31, 519-540.
- Cronan,J.E. (1990). Biotinylation of proteins *in vivo*. A post-translational modification to label, purify, and study proteins. *J. Biol. Chem.* 265, 10327-10333.

- Crooke,E., Thresher,R., Hwang,D.S., Griffith,J., und Kornberg,A. (1993). Replicatively active complexes of DnaA protein and the *Escherichia coli* chromosomal origin observed in the electron microscope. J. Mol. Biol. 233, 16-24.
- Datta,H.J., Khatri,G.S., und Bastia,D. (1999). Mechanism of recruitment of DnaB helicase to the replication origin of the plasmid pSC101. Proc. Natl. Acad. Sci. USA 96, 73-78.
- Dean,F.B., Borowiec,J.A., Eki,T., und Hurwitz,J. (1992). The simian virus 40 T antigen double hexamer assembles around the DNA at the replication origin. J. Biol. Chem. 267, 14129-14137.
- Dixon,N.E. (1999). The Domains of *E.coli* DnaB: Structures, Insights and Prospects. Workshop on Helicases as Molecular Motors in Nucleic Acid Strand Separation.
- Dixon,N.E. und Kornberg,A. (1984). Protein HU in the enzymatic replication of the chromosomal origin of *Escherichia coli*. Proc. Natl. Acad. Sci. USA 81, 424-428.
- Dong,F., Gogol,E.P., und von Hippel,P.H. (1995). The phage T4-coded DNA replication helicase (gp41) forms a hexamer upon activation by nucleoside triphosphate. J. Biol. Chem. 270, 7462-7473.
- Dong,F. und von Hippel,P.H. (1996). The ATP-activated hexameric helicase of bacteriophage T4 (gp41) forms a stable primosome with a single subunit of T4-coded primase. J. Biol. Chem. 271, 19625-19631.
- Doran,K.S., Konieczny,I., und Helinski,D.R. (1998). Replication origin of the broad host range plasmid RK2 - Positioning of various motifs is critical for initiation of replication. J. Biol. Chem. 273, 8447-8453.
- Egelman,H.H., Yu,X., Wild,R., Hingorani,M.M., und Patel,S.S. (1995). Bacteriophage T7 helicase/primase proteins form rings around single-stranded DNA that suggest a general structure for hexameric helicases. Proc. Natl. Acad. Sci. 92, 3869-3873.
- Ellis,N.A., Groden,J., Ye,T.Z., Straughen,J., Lennon,D.J., Ciocci,S., Proytcheva,M., und German,J. (1995). The Bloom's Syndrome Gene product is homologous to RecQ Helicases. Cell 83, 655-666.
- Fang,L.H., Davey,M.J., und O'Donnell,M. (1999). Replisome assembly at oriC, the replication origin of E-coli, reveals an explanation for initiation sites outside an origin. Molecular Cell 4, 541-553.
- Fass,D., Bogden,C.E., und Berger,J.M. (1999). Crystal structure of the N-terminal domain of the DnaB hexameric helicase. Structure 7, 691-698.
- Fried,M. und Crothers,D.M. (1981). Equilibria and kinetics of lac repressor-operator interactions by polyacrylamide gel electrophoresis. Nucl. Acids Res. 9, 6505-6525.
- Fujita,M.Q., Yoshikawa,H., und Ogasawara,N. (1990). Structure of the dnaA region of *Micrococcus luteus*: Conservation and variations among eubacteria. Gene 93, 73-78.
- Fuller,R.S., Funnell,B.E., und Kornberg,A. (1984). The dnaA protein complex with the *E. coli* chromosomal origin (oriC) and other sites. Cell 38, 889-900.
- Fuller,R.S., Kaguni,J.M., und Kornberg,A. (1981). Enzymatic replication of the origin of the *E. coli* chromosome. Proc. Natl. Acad. Sci. USA 78, 7370-7374.
- Funnell,B.E., Baker,T.A., und Kornberg,A. (1987). *In vitro* assembly of a prepriming complex at the origin of the *Escherichia coli* chromosome. J. Biol. Chem. 262, 10327-10334.

- Garner,J. und Crooke,E. (1996). Membrane regulation of the chromosomal replication activity of *E. coli* DnaA requires a discrete site on the protein. EMBO J. 15, 3477-3485.
- Garner,M.M. und Revzin,A. (1981). A gel electrophoresis method for qualifying the binding of proteins to specific DNA regions: application to components of the *Escherichia coli* lactose operon regulatory system. Nucl. Acids Res. 9, 3047-3060.
- Gille,H. und Messer,W. (1991). Localized unwinding and structural perturbations in the origin of replication, *oriC*, of *Escherichia coli* *in vitro* and *in vivo*. EMBO J. 10, 1579-1584.
- Giraldo,R., Andreu,J.M., und Diaz-Orejas,R. (1998). Protein Domains and Conformational changes in the Activation of RepA, a DNA Replication Initiator. EMBO J. 17, 4511-4526.
- Gormley,E.P. und Davis,J. (1991). Transfer of Plasmid RSF1010 by Conjugation from *Escherichia coli* to *Streptomyces lividans* and *Mycobacterium smegmatis*. J. Bacteriol. 173, 6705-6708.
- Griffith (1976). Visualisation of prokaryotic DNA in a regulary condensed chromatin-like fiber. Proc. Natl. Acad. Sci. USA 73, 563-567.
- Guo,L., Katayama,T., Seyama,Y., Sekimizu,K., und Miki,T. (1999). Isolation and characterization of novel cold-sensitive dnaA mutants of *Escherichia coli*. FEMS Microbiol. Lett. 176, 357-366.
- Hansen,E.B., Hansen,F.G., und von Meyenburg,K. (1982a). The nucleotide sequence of the *dnaA* gene and the first part of the *dnaN* gene of *Escherichia coli* K12. Nucl. Acids Res. 10, 7373-7385.
- Hansen,E.B. und Yarmolinsky,M.B. (1986). Host participation in plasmid maintenance: Dependence upon *dnaA* of replicons derived from P1 and F. Proc. Natl. Acad. Sci. USA 83, 4423-4427.
- Hansen,F.G., Hansen,E.B., und Atlung,T. (1982b). The nucleotide sequence of the *dnaA* gene promoter and of the adjacent *rpmH* gene, coding for the ribosomal protein L34, of *Escherichia coli*. EMBO J. 1, 1043-1048.
- Hansen,F.G. und von Meyenburg,K. (1979). Characterization of the *dnaA*, *gyrB* and other genes in the *dnaA* region of the *Escherichia coli* chromosome on specialized transducing phages lambda-*tma*. Mol. Gen. Genet. 175, 135-144.
- Haring,V., Scholz,P., Frey,J., Derbyshire,K., Hatfull,G., Willets,N.S., und Bagdasarian,M. (1985). Protein RepC is involved in copy number control of the broad host range Plasmid RSF1010. Proc. Natl. Acad. Sci. 82, 6090-6094.
- Hasunuma,K. und Sekiguchi,M. (1977). Replication of plasmid pSC101 in *Escherichia coli* K12: requirement for dnaA function. Mol. Gen. Genet. 154, 225-230.
- Hiasa,H. und Marians,K.J. (1994). Primase couples leading- and lagging-strand DNA synthesis from *oriC*. J. Biol. Chem. 269, 6058-6063.
- Hiasa,H. und Marians,K.J. (1999). Initiation of bidirectional replication at the chromosomal origin is directed by the interaction between helicase and primase. J. Biol. Chem. 274, 27244-27248.
- Hidaka,M., obayashi,T., und Horiuchi,T. (1991). A newly identified DNA replication terminus site, TerE, on the *Escherichia coli* chromosome. J. Bacteriol. 173, 391-393.
- Hill,T.M., Henson,J.M., und Kuempel,P.K. (1987). The terminus region of the *Escherichia coli* chromosome contains two separate loci that exhibit polar inhibition of replication. Proc. Natl. Acad. Sci. USA 84, 1754-1758.

- Hill,T.M., Kopp,B.J., und Kuempel,P.L. (1988). Termination of DNA replication in *Escherichia coli* requires a trans-acting factor. J. Bacteriol. 170, 662-668.
- Hill,T.M. und Marians,K.J. (1990). *Escherichia coli* Tus protein acts to arrest the progression of DNA replication forks *in vitro*. Proc. Natl. Acad. Sci. USA 87, 2481-2485.
- Honda,Y., Nakamura,T., Tanaka,K., Higashi,A., Sakai,H., Komano,T., und Bagdasarian,M. (1992). DnaG-dependent Priming Signals can substitute for the two essential DNA Initiation Signals in oriV of the broad-host-range Plasmid RSF1010. Nucl. Acids Res. 20, 1733-1737.
- Honda,Y., Sakai,H., Hiasa,H., Tanaka,K., Komano,T., und Bagdasarian,M. (1991). Functional Division and Reconstruction of a Plasmid Replication Origin: Molecular Dissection of the oriV of the broad-host-range Plasmid RSF1010. Proc. Natl. Acad. Sci. 88, 179-183.
- Honda,Y., Sakai,H., und Komano,T. (1988). Two single-strand DNA Initiation Signals located in the oriV Region of Plasmid RSF1010. Gene 68, 221-228.
- Honda,Y., Sakai,H., Komano,T., und Bagdasarian,M. (1989). RepB' is required in trans for the two single-strand DNA Initiation Signals in oriV of Plasmid RSF1010. Gene 80, 155-159.
- Hsu,J., Bramhill,D., und Thompson,C.M. (1994). Open complex formation by DnaA initiation protein at the *E. coli* chromosomal origin requires the 13-mers precisely spaced relative to the 9-mers. Mol. Microbiol. 11, 903-911.
- Johnson,R.T. und Squires,S. (1992). The XPD complementation group. Insights into Xeroderma Pigmentosum, Cockayne's Syndrome and Trichothiodystrophy. Mutation Res. 273, 97-118.
- Jueterbock, W.-R. Das *dnaA*-Gen von *E.coli*: Reinigung des Genprodukts und Quantifizierung der Expression unter SOS-Bedingungen. 1991. Ref Type: Thesis/Dissertation
- Kaguni,J.M. (1997). *Escherichia coli* DnaA protein: the replication initiator. Molecules and Cells 7, 145-157.
- Kaguni,J.M. und Hwang,D.S. (1988). Purification and characterization of the *dnaA46* gene product. J. Biol. Chem. 263, 10625-10632.
- Kaguni,J.M. und Kornberg,A. (1984). Replication initiated at the origin (*oriC*) of the *E. coli* chromosome reconstituted with purified enzymes. Cell 38, 183-190.
- Katayama,T., Kubota,T., Kurokawa,K., Crooke,E., und Sekimizu,K. (1998). The initiator function of DnaA protein is negatively regulated by the sliding clamp of the *E.coli* chromosomal replicase. Cell 94, 61-71.
- Kawasaki,Y., Matsunaga,F., Kano,Y., Yura,T., und Wada,C. (1996). The localized melting of mini-F by the combined action of the mini-F initiator protein (RepE) and HU and DnaA of *Escherichia coli*. Mol. Gen. Genet. 253, 42-49.
- Kellenberger-Gujer,G., Podhajska,A.J., und Caro,L. (1978). A cold sensitive *dnaA* mutant of *E. coli* which overinitiates chromosome replication at low temperature. Mol. Gen. Genet. 162, 9-16.
- Kim,S., Dallmann,H.G., McHenry,C.S., und Marians,K.J. (1996). Coupling of a Replicative Polymerase and Helicase: a tau-DnaB Interaction mediates rapid Replication Fork Movement. Cell 84, 643-650.
- Klein, Lanka,E., und Schuster,H. (1980). Isolation of a complex between the P Protein of Phage lambda and the DnaB Protein of *Escherichia coli*. Eur. J. Biochem. 105, 1-6.
- Koch,C., Vandekerckhove,J., und Kahmann,R. (1988). *Escherichia coli* host factor for site-specific DNA inversion: Cloning and characterization of the *fis* gene. Proc. Natl. Acad. Sci. USA 85, 4237-4241.

- Konieczny,I., Doran,K.S., Helinski,D.R., und Blasina,A. (1997). Role of TrfA and DnaA proteins in origin opening during initiation of DNA replication of the broad host range plasmid RK2. *J. Biol. Chem.* 272, 20173-20178.
- Koonin,E.V. (1997). DnaC protein contains a modified ATP-binding motif and belongs to a novel family of ATPases including also DnaA. *Nucl. Acids Res.* 20.
- Kornberg,A. und Baker,T.A. (1992). DNA Replication. (New York: W. H. Freeman and Company).
- Kornberg,A., Baker,T.A., Bertsch,L.L., Bramhill,D., Funnell,B.E., Lasken,R.S., Maki,H., Maki,S., Sekimizu,K., und Wahle,E. (1987). Enzymatic studies of replication of *oriC* plasmids. In DNA Replication and Recombination. UCLA Symp. Molec. Cell. Biol., New Series, Vol. 47, R.McMacken and T.J.Kelly, eds. (New York: A. R. Liss), pp. 137-149.
- Krause,M., Lurz,R., Rückert,B., und Messer,W. (1997). Complexes at the replication origin of *Bacillus subtilis* with homologous and heterologous DnaA protein. *J. Mol. Biol.* 274, 365-380.
- Krause,M. und Messer,W. (1999). DnaA proteins of *Escherichia coli* and *Bacillus subtilis*: coordinate actions with single-stranded DNA-binding protein and interspecies inhibition during open complex formation at the replication origins. *Gene* 228, 123-132.
- Kubota,T., Katayama,T., Ito,Y., Mizushima,T., und Sekimizu,K. (1997). Conformational transition of DnaA protein by ATP: structural analysis of DnaA protein, the initiator of *Escherichia coli* chromosome replication. *Biochem. Biophys. Res. Comm.* 232, 130-135.
- Kücherer,C., Loher,H., Kölling,R., Schauzu,M.A., und Messer,W. (1986). Regulation of transcription of the chromosomal *dnaA* gene of *Escherichia coli*. *Mol. Gen. Genet.* 205, 115-121.
- Laemmli,U.K. (1970). Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. *Nature* 227, 680-685.
- Lanka,E., Edelbluth,C., Schlicht,M., und Schuster,H. (1978). *Escherichia coli* dnaB protein. Affinity chromatography on immobilized nucleotides. *J. Biol. Chem.* 253, 5847-5851.
- Lanka,E. und Schuster,H. (1983). The DnaC protein of *Escherichia coli*. Purification, physical properties and interaction with DnaB protein. *Nucl. Acids Res.* 11, 987-997.
- Lanzer,M. und Bujard,H. (1988). Promoters largely determine the efficiency of repressor action. *Proc. Natl. Acad. Sci. USA* 85, 8973-8977.
- Learn,B.A., Um,S.-J., Huang,L., und McMacken,R. (1997). Cryptic single-stranded-DNA binding activities of the phage λ P and *Escherichia coli* DnaC replication initiation proteins facilitate the transfer of *E. coli* DnaB helicase onto DNA. *Proc. Natl. Acad. Sci. USA* 94, 1154-1159.
- LeBowitz,J.H. und McMacken,R. (1986). The *Escherichia coli* dnaB replication protein is a DNA helicase. *J. Biol. Chem.* 261, 4738-4748.
- Li,X., Yoder,B.L., und Burgers,P.J.M. (1992). Three new helicases from *Saccharomyces cerevisiae*. *Chromosoma* 102, 93-99.
- Lohman,T.M. und Bjornson,K.P. (1996). Mechanism of helicase-catalysed DNA unwinding. *Ann. Rev. Biochem.* 65, 169-214.
- Louarn,J.-M., Funderberg,M., und Bird,E. (1974). More precise mapping of the replication origin in *Escherichia coli* K-12. *J. Bact.* 120, 1-5.
- Louarn,J.-M. und Patte,J. (1977). Evidence for a fixed termination site of chromosome replication in *Escherichia coli*. *J. Mol. Biol.* 115, 295-314.

- Lu,Y.B., Datta,H.J., und Bastia,D. (1998). Mechanistic studies of initiator-initiator interaction and replication initiation. *EMBO J.* 17, 5192-5200.
- Lu,Y.B., Ratnakar,P.V.A.L., Mohanty,B.K., und Bastia,D. (1996). Direct physical interaction between DnaG primase and DnaB helicase of *Escherichia coli* is necessary for optimal synthesis of primer RNA. *Proceedings of the National Academy of Sciences of the United States of America* 93 , 12902-12907.
- Maaloe,O. und Kjeldgaard,N.O. (1966). Control of Macromolecular Synthesis. (New York, Amsterdam: Benjamin).
- Maki,H., Maki,S., und Kornberg,A. (1988). DNA polymerase III holoenzyme of *Escherichia coli*. IV. The holoenzyme is an asymmetric dimer with twin active sites. *J. Biol. Chem.* 263, 6570-6578.
- Mallory,J.B., Alfano,C., und McMacken,R. (1990). Host Virus Interaction in the Initiation of Bacteriophage lambda DNA Replication. Recruitment of *Escherichia coli* DnaB Helicase by lambda P Replication Protein. *J. Biol. Chem.* 265, 13297-13307.
- Marians,K.J. (1996). Replication fork propagation. In *Escherichia coli and Salmonella*. Cellular and Molecular Biology, F.C.Neidhardt , R.Curtiss III, J.Ingraham, E.C.C.Lin, K.B.Low, B.Magasanik, W.S.Reznikoff, M.Riley, M.Schaechter, and H.E.Umbarger, eds. (Washington, D.C.: ASM Press), pp. 749-763.
- Marszalek,J. und Kaguni,J.M. (1994). DnaA protein directs the binding of DnaB protein in initiation of DNA replication in *Escherichia coli*. *J. Biol. Chem.* 269, 4883-4890.
- Masters,M. und Broda,P. (1971). Evidence for the bidirectional replication of the *E. coli* chromosome. *Nature New Biol.* 232, 137-140.
- Matson,S.W. (1991). DNA Helicases of *Escherichia coli*. *Progr. Nucl. Acids Res. & Mol. Biol.* 40, 289-326.
- Matson,S.W., Bean,D.W. und George,J.W. (1994). DNA helicases: enzymes with essential roles in all aspects of DNA metabolism. *BioEssays* 16, 13-22.
- Matson,S.W. und Kaiser-Rogers,K.A. (1990). DNA Helicases. *Ann. Rev. Biochem.* 59, 289-330.
- Matsui,M., Oka,A., Takanami,M., Yasuda,S., und Hirota,Y. (1985). Sites of dnaA protein-binding in the replication origin of the *E. coli* K-12 chromosome. *J. Mol. Biol.* 184, 529-533.
- McHenry,C.S. (1986). DNA polymerase III holoenzyme of *Escherichia coli*. Components and function of a true replicative complex. *Molec. Cell. Biochem.* 66, 71-85.
- McHenry,C.S. (1988). DNA Polymerase III holoenzyme of *Escherichia coli*. *Annual Rev. Biochem.* 57, 519-550.
- McMacken,R., Ueda,K., und Kornberg,A. (1977). Migration of *Escherichia coli* DnaB protein on the template DNA strand as a mechanism in initiating DNA replication. *Proc. Natl. Acad. Sci. USA* 74, 4190-4194.
- Meijer,M., Beck,E., Hansen,F.G., Bergmans,H.E., Messer,W., von Meyenburg,K., und Schaller,H. (1979). Nucleotide sequence of the origin of replication of the *E. coli* K-12 chromosome. *Proc. Natl. Acad. Sci. USA* 76, 580-584.
- Meselson,M. und Stahl,F. (1958). The replication of DNA in *E. coli*. *Proc. Natl. Acad. Sci. USA* 44, 671-682.
- Messer,W., Bergmans,H.E., Meijer,M., Womack,J.E., Hansen,F.G., und von Meyenburg,K. (1978). Minichromosomes: Plasmids which carry the *E. coli* replication origin. *Mol. Gen. Genet.* 162, 269-275.

- Messer,W., Blaesing,F., Majka,J., Nardmann,J., Schaper,S., Schmidt,A., Seitz,H., Speck,C., Tüngler,D., Wegrzyn,G., Weigel,C., Welzeck,M., und Zakrzewska-Czerwinska,J. (1999). Functional domains of DnaA proteins. *Biochimie* 81, 819-825.
- Messer,W., Egan,J.B., Gille,H., Holz,A., Schaefer,C., und Woelker,B. (1991). The complex of *oriC* DNA with the DnaA initiator protein. *Res. Microbiol.* 142, 119-125.
- Messer,W., Hartmann-Kühlein,H., Langer,U., Mahlow,E., Roth,A., Schaper,S., Urmoneit,B., und Woelker,B. (1992). The complex for replication initiation of *Escherichia coli*. *Chromosoma* 102, S1-S6.
- Messer,W. und Weigel,C. (1996). Initiation of chromosome replication. In *Escherichia coli* and *Salmonella*, Cellular and Molecular Biology, F.C.Neidhardt, R.Curtiss III, J.Ingraham, E.C.C.Lin, K.B.Low, B.Magasanik, W.S.Reznikoff, M.Riley, M.Schaechter, und H.E.Umbarger, eds. (Washington, D.C.: ASM), pp. 1579-1601.
- Miki,T., Kimura,M., Hiraga,S., Nagata,T., und Yura,T. (1979). Cloning and physical mapping of the *dnaA* region of the *Escherichia coli* chromosome. *J. Bact.* 140, 817-824.
- Mima,S., Yamaguchi,Y., Kondo,T., Tsuchiya,T., und Mizushima,T. (1999). Role of the amino-terminal region of the DnaA protein in opening of the duplex DNA at the *oriC* region. *FEMS Microbiol. Lett.* 176, 163-167.
- Mizushima,T., Nishida,S., Kurokawa,K., Katayama,T., Miki,T., und Sekimizu,K. (1997). Negative control of DNA replication by hydrolysis of ATP bound to DnaA protein, the initiator of chromosomal DNA replication in *Escherichia coli*. *EMBO J.* 16, 3724-3730.
- Mizushima,T., Takaki,T., Kubota,T., Tsuchiya,T., Miki,T., Katayama,T., und Sekimizu,K. (1998). Site-directed mutational analysis for the ATP binding of DnaA protein - Functions of two conserved amino acids (Lys-178 and Asp-235) located in the ATP-binding domain of DnaA protein *in vitro* and *in vivo*. *J. Biol. Chem.* 273, 20847-20851.
- Moore,K.J. und Lohman,T.M. (1995). Helicase-catalysed DNA unwinding: energy coupling by DNA motor proteins. *Biophys.* 68, 180-185.
- Moriya,S., Ogasawara,N., und Yoshikawa,H. (1985). Structure and function of the region of the replication origin of the *Bacillus subtilis* chromosome. III. Nucleotide sequence of some 10,000 base pairs in the origin region. *Nucl. Acids Res.* 13, 2251-2265.
- Nakayama,N., Arai,N., Kaziro,Y., und Arai,K. (1984). Structural and functional studies of the dnaB protein using limited proteolysis. Characterization of domains for DNA-dependent ATP hydrolysis and for protein association in the primosome. *J. Biol. Chem.* 259, 88-96.
- Oka,A., Sasaki,H., Sugimoto,K., und Takanami,M. (1984). Sequence organization of replication origin of the *Escherichia coli* K-12 chromosome. *J. Mol. Biol.* 176, 443-458.
- Oka,A., Sugimoto,K., Takanami,M., und Hirota,Y. (1980). Replication origin of the *Escherichia coli* K-12 chromosome: The size and structure of the minimum DNA segment carrying the information for autonomous replication. *Mol. Gen. Genet.* 178, 9-20.
- Okazaki,R., Okazaki,T., Sakabe,K., Sugimoto,K., und Sugino,A. (1968a). Mechanism of DNA chain growth. I. Possible discontinuity and unusual

- secondary structure of newly synthesized chains. Proc. Natl. Acad. Sci. 59, 598-605.
- Okazaki,R., Okazaki,T., Sakabe,K., Sugimoto,K., und Sugino,A. (1968b). Mechanism of DNA chain growth. I. Possible discontinuity and unusual secondary structure of newly synthesised chains. Proc. Natl. Acad. Sci. USA 59, 598-605.
- Patel,S.S. (1999). Steps Toward Understanding the Mechanism of the Helicase. Workshop on Helicases as Molecular Motors in Nucleic Acid Strand Separation.
- Polaczek,P. (1990). Bending of the origin of replication of *E. coli* by binding of IHF at a specific site. New Biol. 2, 265-271.
- Ratnakar,P.V.A.L., Mohanty,B.K., Lobert,M., und Bastia,D. (1996). The replication initiator protein pi of the plasmid R6K specifically interacts with the host-encoded helicase DnaB. Proc. Natl. Acad. Sci. USA 93, 5522-5526.
- Reha-Krantz,L.J. und Hurwitz,J. (1978). The *dnaB* gene product of *Escherichia coli*. II. Single stranded DNA-dependent ribonuclease triphosphatase activity. J. Biol. Chem. 253, 4051-4057.
- Richter,S., Hess,W.R., Krause,M., und Messer,W. (1998). Unique organization of the *dnaA* region from *Prochlorococcus marinus* CCMP1375, a marine cyanobacterium. Mol. Gen. Genet. 257, 534-541.
- Rossmann,M.G., Moras,D., und Olsen,K.W. (1974). Chemical and biological evolution of nucleotide-binding protein. Nature 250, 194-199.
- Roth,A. und Messer,W. (1995). The DNA binding domain of the initiator protein DnaA. EMBO J. 14, 2106-2111.
- Roth,A., Urmoneit,B., und Messer,W. (1994). Functions of histone-like proteins in the initiation of DNA replication at *oriC* of *Escherichia coli*. Biochimie 76, 917-923.
- Saiki,R.K., Gelfand,D.H., Stoffel,S., Scharf,S.J., Higuchi,R., Horn,G.T., Mullis,K.B., und Ehrlich,H.A. (1988). Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. Science 239, 487.
- Sakakibara,Y. und Mizukami,T. (1980). A temperature sensitive *Escherichia coli* mutant defective in DNA replication: *dnaN*, a new gene adjacent to the *dnaA* gene. Mol. Gen. Genet. 178, 541-553.
- Sambrook,J., Fritsch,E.F., und Maniatis,T. (1989). Molecular Cloning. (Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory Press).
- Samitt,C.E., Hansen,F.G., Miller,J.F., und Schaechter,M. (1989). *In vivo* studies of DnaA binding to the origin of replication of *Escherichia coli*. EMBO J. 8, 989-993.
- Samols,D., Thornton,C.G., Murtif,V.L., Kumar,G.K., Haase,F.C., und Wood,H.G. (1988). Evolutionary conservation among biotin enzymes. J. Biol. Chem. 263, 6461-6464.
- San Martin,C., Radermacher,M., Wolpensinger,B., Engel,A., Miles,C.S., Dixon,N.E., und Carazo,J.M. (1998). Three-dimensional reconstructions from cryoelectron microscopy images reveal an intimate complex between helicase DnaB and its loading partner DnaC. Structure 6, 501-509.
- San Martin,M.C., Stamford,N.P.J., Dammerova,N., Dixon,N.E., und Carazo,J.M. (1995). A structural model for the *Escherichia coli* DnaB helicase based on electron microscopy data. J. Struct. Biol. 114, 167-176.
- Sanger,F., Nicklen,S., und Coulson,R. (1977). DNA sequencing with chain-terminating inhibitors. Proc. Natl. Acad. Sci. USA 74, 5463-5467.

- Saraste,M., Sibbald,P.R., und Wittinghofer,A. (1990). The P-loop - a common motif in ATP- and GTP-binding proteins. *Trends Biochem. Sci.* 15, 430-434.
- Schaper,S. und Messer,W. (1995). Interaction of the initiator protein DnaA of *Escherichia coli* with its DNA target. *J. Biol. Chem.* 270, 17622-17626.
- Schaper,S. und Messer,W. (1997). Prediction of the structure of the replication initiator protein DnaA. *Proteins:Structure, Function, and Genetics* 28, 1-9.
- Scherzinger,E., Haring,V., Lurz,R., und Otto,S. (1991). Plasmid RSF1010 DNA Replication *in vitro* promoted by purified RSF1010 RepA, RepB and RepC proteins. *Nucl. Acids Res.* 19, 1203-1211.
- Scholz,P., Haring,V., Wittmann-Liebold,B., Ashman,K., Bagdasarian,M., und Scherzinger,E. (1989). Complete nucleotide sequence and gene organization of the broad-host-range plasmid RSF1010. *Gene* 75, 271-288.
- Scott,J.F., Eisenberg,S., Bertsch, L.L., und Kornberg,A. (1977). A mechanism of duplex DNA Replication revealed by enzymatic studies of phage Φ X174: catalytic strand separation in advance of replication. *Proc. Natl. Acad. Sci.* 74, 193-197.
- Sekimizu,K., Bramhill,D., und Kornberg,A. (1987). ATP activates dnaA protein in initiating replication of plasmids bearing the origin of the *E. coli* chromosome. *Cell* 50, 259-265.
- Seufert,W. und Messer,W. (1987). Start sites for bidirectional *in vitro* replication inside the replication origin, *oriC*, of *Escherichia coli*. *EMBO J.* 6, 2469-2472.
- Skarstad,K., Baker,T.A., und Kornberg,A. (1990). Strand separation required for initiation of replication at the chromosomal origin of *E. coli* is facilitated by a RNA-DNA hybrid. *EMBO J.* 9, 2341-2348.
- Skarstad,K. und Boye,E. (1994). The initiator protein DnaA: Evolution, properties and function. *Biochim. Biophys. Acta* 1217, 111-130.
- Skarstad,K., Boye,E., und Steen,H.B. (1986). Timing of initiation of chromosome replication in individual *E. coli* cells. *EMBO J.* 5, 1711-1717.
- Skarstad,K. und Wold,S. (1995). The speed of the *Escherichia coli* fork *in vivo* depends on the DnaB:DnaC ratio. *Mol. Microbiol.* 17, 825-831.
- Skovgaard,O., Olesen,K., und Wright,A. (1998). The central lysine in the P-loop motif of the *Escherichia coli* DnaA protein is essential for initiating DNA replication from the chromosomal origin, *oriC*, and the F factor origin, *oriS*, but is dispensable for initiation from the P1 plasmid origin, *oriR*. *Plasmid* 40, 91-99.
- Speck,C., Weigel,C., und Messer,W. (1997). From footprint to toeprint: a close-up of the DnaA box, the binding site for the initiator protein DnaA of *Escherichia coli*. *Nucl. Acids Res.* 25, 3242-3247.
- Speck,C., Weigel,C., und Messer,W. (1999). ATP- and ADP-DnaA protein, a molecular switch in gene regulation. *EMBO J.* 18, 6169-6176.
- Stasiak,A., Tsaneva,I.R., West,S.C., Benson,C.J., Yu,X., und Egelman,E.H. (1994). The *Escherichia coli* RuvB branch migration protein forms double hexameric rings around DNA. *Proceedings of the National Academy of Sciences of the United States of America* 91, 7618-7622.
- Stenzel,T.T., MacAllister,T., und Bastia,D. (1991). Cooperativity at a distance promoted by the combined action of two replication initiator proteins and a DNA bending protein at the replication origin of pSC101. *Genes Dev.* 5, 1453-1463.
- Stenzel,T.T., Patel,P., und Bastia,D. (1987). The integration host factor of *Escherichia coli* binds to bent DNA at the origin of replication of the plasmid pSC101. *Cell* 49, 709-717.

- Stordal,L. und Maurer,R. (1996). Defect in General Priming Conferred by Linker Region Mutants of *Escherichia coli dnaB*. J. Bact. 178, 4620-4627.
- Sugimoto,K., Oka,A., Sugisaki,H., Takanami,M., Nishimura,A., Yasuda,Y., und Hirota,Y. (1979). Nucleotide sequence of *Escherichia coli* replication origin. Proc. Natl. Acad. Sci. USA 76, 575-579.
- Sutton,M.D., Carr,K.M., Vicente,M., und Kaguni,J.M. (1998). *Escherichia coli* DnaA protein. The N-terminal domain and loading of DnaB helicase at the *E. coli* chromosomal origin. J. Biol. Chem. 273, 34255-34262.
- Sutton,M.D. und Kaguni,J.M. (1997b). Novel alleles of the *Escherichia coli dnaA* gene. J. Mol. Biol. 271, 693-703.
- Sutton,M.D. und Kaguni,J.M. (1997a). The *Escherichia coli dnaA* gene: Four functional domains. J. Mol. Biol. 274, 546-561.
- Sutton,M.D. und Kaguni,J.M. (1997c). Threonine 435 of *Escherichia coli* DnaA protein confers sequence-specific DNA binding activity. J. Biol. Chem. 272, 23017-23024.
- Taguchi,Y., Tanaka,K., Honda,Y., Miao,D.M., Sakai,H., Komano,T., und Bagdasarian,M. (1996). A dnaA box can functionally substitute for the priming signals in the *oriV* of the broad host-range plasmid RSF1010. FEBS Lett. 388, 169-172.
- Tucker,W.T., Miller,C.A., und Cohen,S.N. (1984). Structural and functional analysis of the *par* region of the pSC101 plasmid. Cell 38, 191-201.
- Ueda,K., McMacken,R., und Kornberg,A. (1978). dnaB protein of *Escherichia coli*. Purification and role in the replication of phiX174 DNA. J. Biol. Chem. 253, 261-269.
- van der Ende,A., Baker,T.A., Ogawa,T., und Kornberg,A. (1985). Initiation of enzymatic replication at the origin of the *E. coli* chromosome: Primase as the sole priming enzyme. Proc. Natl. Acad. Sci. USA 82, 3954-3958.
- Vocke,C. und Bastia,D. (1983). Primary structure of the essential replicon of the plasmid pSC101. Proc. Natl. Acad. Sci. USA 80, 6557-6581.
- von Meyenburg,K., Hansen,F.G., Riise,E., Bergmans,H.E., Meijer,M., und Messer,W. (1979). Origin of replication, *oriC*, of the *E. coli* K-12 chromosome: Genetic mapping and minichromosome replication. Cold Spring Harbor Symp. Quant. Biol. 43, 121-128.
- Wahle,E., Lasken,R.S., und Kornberg,A. (1989). The dnaB-dnaC replication protein complex of *Escherichia coli*. II. role of the complex in mobilizing dnaB functions. J. Biol. Chem. 264, 2469-2475.
- Wang,Q. und Kaguni,J.M. (1987). Transcriptional repression of the *dnaA* gene of *Escherichia coli* by dnaA protein. Mol. Gen. Genet. 209, 518-525.
- Wang,Q. und Kaguni,J.M. (1989). dnaA Protein regulates transcription of the *rpoH* gene of *Escherichia coli*. J. Biol. Chem. 264, 7338-7344.
- Weigel,C., Schmidt,A., Seitz,H., Tuengler,D., Welzeck,M., und Messer,W. (1999). The N-terminus promotes oligomerisation of the *Escherichia coli* initiator protein DnaA. Mol. Microbiol. 34, 53-66.
- Weigelt,J., Brown,S.E., Miles,C.S., Dixon,N.E., und Otting,G. (1999). NMR structure of the N-terminal domain of E-coli DnaB helicase: implications for structure rearrangements in the helicase hexamer. Structure 7, 681-690.
- Wickner,S., Wright,M., und Hurwitz,J. (1974). Association of DNA-dependent and -independent ribonucleoside triphosphatase activities with dnaB gene product of *Escherichia coli*. Proc. Natl. Acad. Sci. 71, 783-787.

- Wu,J., Sektas,M., Chen,D., und Filutowicz,M. (1997). Two forms of Replication Initiator Protein: positive and negative controls. Proc. Natl. Acad. Sci. 94, 13967-13972.
- Yamaguchi,Y., Hase,M., Makise,M., Mima,S., Yoshimi,T., Ishikawa,Y., Tsuchiya,T., und Mizushima,T. (1999). Involvement of Arg-328, Arg-334 and Arg-342 of DnaA protein in the functional interaction with acidic phospholipids. Biochem. J. 340, 433-438.
- Yasuda,S. und Hirota,Y. (1977). Cloning and mapping of the replication origin of *Escherichia coli*. Proc. Natl. Acad. Sci. USA 74, 5458-5462.
- Yoshikawa,H. und Ogasawara,N. (1991). Structure and function of DnaA and the DnaA-box in eubacteria: Evolutionary relationships of bacterial replication origins. Mol. Microbiol. 5, 2589-2597.
- Yoshikawa,H. und Wake,R.G. (1993). 36. Initiation and termination of chromosome replication. In *Bacillus subtilis* and Other Gram-Positive Bacteria: Biochemistry, Physiology, and Molecular Genetics, A.L.Sonenshein, J.A.Hoch, and R.Losick, eds. (Washington, D.C.: ASM), pp. 507-528.
- Yu,C.E., Fu,Y.H., Wijsman,E.M., Hisama,F., Alisch,R., Nakura,J., Miki,T., Martin,G.M., Mulligan,J., und Schellenberg,G.D. (1996). Positional cloning of the Werner's syndrome gene. Science 272, 258-262.
- Zyskind,J.W., Cleary,J.M., Brusilow,W.S.A., Harding,N.E., und Smith,D.W. (1983). Chromosomal replication origin from the marine bacterium *Vibrio harveyi* functions in *Escherichia coli*: *oriC* consensus sequence. Proc. Natl. Acad. Sci. USA 80, 1164-1168.
- Zyskind,J.W., Harding,N.E., Takeda,Y., Cleary,J.M., und Smith,D.W. (1981). The DNA replication origin region of the enterobacteriaceae. ICN-UCLA Symp. Mol. Cell. Biol. 22, 13-25.

