

Charité-Universitätsmedizin Berlin

Aus der Klinik für Angeborene Herzfehler und Kinderkardiologie im
Deutschen Herzzentrum Berlin

Dissertationsthema

**Noninvasive evaluation of right atrial function in patients with secundum
type atrial septal defect and in patients with corrected tetralogy of Fallot
using 2-D echocardiography and tissue Doppler imaging**

Inaugural-Dissertation

Zur

Erlangung der medizinischen Doktorwürde
an der Charité-Universitätsmedizin Berlin

Vorgelegt von: **Hui Wei**

aus: Shanghai, V.R. China

Referent: PD. Dr. Hashim Abdul-Khalil

Korreferent: Prof. Dr. K.-J. Bauknecht

Gedruckt mit Genehmigung der Charité – Universitätsmedizin Berlin

Campus Benjamin Franklin

Promoviert am: 17.12.2004

Contents

	Page
1 Abbreviations and Acronyms	5
2 Introduction	6
2.1 Studies of atrial function	6
2.2 Methods of evaluation of the three main atrial functions	7
2.3 Tissue Doppler imaging	8
3 Aims of the study	11
4 Patients	12
4.1 Right atrial and right ventricular function in patients before and after ASD closure (section 1)	12
4.2 Right atrial and right ventricular function in patients following corrective surgery of tetralogy of Fallot (section 2)	17
5 Methods	22
5.1 Two-dimensional echocardiography	22
5.2 Tissue Doppler imaging (TDI)	24
5.3 Magnetic resonance tomography (MRT)	26
5.4 Routine catheter examination and interventional closure of ASD with Amplatzer Septal Occluder	26
5.5 Surgical closure of ASD	27
5.6 Statistics	27
6 Results	29
6.1 ASD patients versus normal controls and comparison of the effects of catheter and surgical ASD closure on right atrial function (section 1)	29

6.1.1	Two-dimensional echocardiography	29
6.1.2	Tissue Doppler imaging	40
6.2	Right atrial pump function and its relation to right ventricular function in patients following corrective surgery of TOF (section 2)	47
6.2.1	Two-dimensional echocardiography	47
6.2.2	Tissue Doppler imaging	53
7	Discussion	58
7.1	Atrial strain rate curve derived from tissue Doppler imaging	58
7.2	Right atrial pump function in patients before ASD closure	59
7.3	Effects of surgical and catheter closure on global right atrial pump function	60
7.4	The value of tissue Doppler imaging on the evaluation of right atrial pump function in patients before and after surgical or catheter ASD closure	61
7.5	Global right atrial pump function in patients with corrected TOF	62
7.6	Regional right atrial performance in patients with corrected TOF	62
7.7	Relation between right atrial pump function and right ventricular systolic function in patients following corrective surgery of TOF	63
8	Conclusions	65
9	References	66
10	Summary	77
11	Zusammenfassung	79
12	Statement	82
13	Acknowledgement	83
14	Curriculum Vitae	84