

Potential use of superparamagnetic iron oxide nanoparticles for *in vitro* and *in vivo* bioimaging of human myoblasts.

Kamil R. Wierzbinski^{1#}, Tomasz Szymanski^{1,2,3#}, Natalia Rozwadowska¹, Jakub D. Rybka^{2,3}, Agnieszka Zimna¹, Tomasz Zalewski⁴, Karolina Nowicka-Bauer¹, Agnieszka Malcher¹, Magdalena Nowaczyk¹, Michal Krupinski⁵, Michal Fiedorowicz⁶, Piotr Bogorodzki⁶, Paweł Grieb⁶, Michał Giersig^{2,3,7}, Maciej K. Kurpisz^{1*}

¹ Institute of Human Genetics Polish Academy of Sciences, Poznan, Poland

² Faculty of Chemistry, Adam Mickiewicz University, Poznan, Poland

³ Wielkopolska Centre of Advanced Technologies, Adam Mickiewicz University, Poznan, Poland

⁴ NanoBioMedical Centre, Adam Mickiewicz University, Poznan, Poland

⁵ The Henryk Niewodniczanski Institute, Institute of Nuclear Physics Polish Academy of Sciences, Cracow, Poland

⁶ Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

⁷ Institute of Experimental Physics, Freie Universität Berlin, Berlin, Germany

Both authors equally contributed to the study

*Corresponding Author

Maciej K. Kurpisz MD PhD,

Institute of Human Genetics,

Polish Academy of Sciences, Poznan, Poland

e-mail: kurpimac@man.poznan.pl

Supplementary Figure S1: Diagrams representing results of ICP-MS analysis. **A-B** – iron concentrations in standard solutions of SPIONs in cell lysates. **C-D** – iron concentrations in SPION-treated cell lysates. **E-F** – calculated amounts of iron per single cell. Abbreviations: ICP-MS – inductively coupled plasma mass spectrometry. Asterisks indicate statistical significance (** p < 0.01; *** p < 0.001).

