

## **6 Literaturverzeichnis**

Adams, P.W., Lee, H.S., Waldman, W.J., Sedmak, D.D., Morgan, C.J., Ward, J.S., and Orosz, C.G. (1992). Alloantigenicity of human endothelial cells. 1. Frequency and phenotype of human T helper lymphocytes that can react to allogeneic endothelial cells. *J.Immunol.* 148, 3753-3760.

Alderson, M.R., Armitage, R.J., Tough, T.W., Strockbine, L., Fanslow, W.C., and Spriggs, M.K. (1993). CD40 expression by human monocytes: regulation by cytokines and activation of monocytes by the ligand for CD40. *J.Exp.Med.* 178, 669-674.

Allen, R.C., Armitage, R.J., Conley, M.E., Rosenblatt, H., Jenkins, N.A., Copeland, N.G., Bedell, M.A., Edelhoff, S., Disteche, C.M., Simoneaux, D.K., Fanslow, W.C., Belmont, J., and Spriggs, M.K. (1993). CD40 ligand gene defects responsible for X-linked hyper-IgM syndrome. *Science* 259, 990-993.

Armitage, R.J., Fanslow, W.C., Strockbine, L., Sato, T.A., Clifford, K.N., Macduff, B.M., Anderson, D.M., Gimpel, S.D., Davis-Smith, T., Maliszewski, C.R., Clark, E.A., Smith, C.A., Grabstein, K.H., Cosman, D., and Spriggs, M.K. (1992). Molecular and biological characterization of a murine ligand for CD40. *Nature* 357, 80-82.

Aruffo, A., Farrington, M., Hollenbaugh, D., Li, X., Milatovich, A., Nonoyama, S., Bajorath, J., Grosmaire, L.S., Stenkamp, R., Neubauer, M., Roberts, R.L., Noelle, R.J., Ledbetter, J.A., Francke, U., and Ochs, H.D. (1993). The CD40 ligand, gp39, is defective in activated T cells from patients with X-linked hyper-IgM syndrome. *Cell* 72, 291-300.

Aster, R.H. (1966). Pooling of platelets in the spleen: role in the pathogenesis of "hypersplenic" thrombocytopenia. *J.Clin.Invest.* 45, 645-657.

Banchereau, J. and Rousset, F. (1991). Growing human B lymphocytes in the CD40 system. *Nature* 353, 678-679.

Breslauer, K.J., Frank, R., Bloecker, H., and Marky, L.A. (1986). Predicting DNA duplex stability from the base sequence. *Proc.Natl.Acad.Sci.USA* 83, 3746-3750.

Brindley, L.L., Sweet, J.M., and Goetzl, E.J. (1983). Stimulation of histamine release from human basophils by human platelet factor 4. *J.Clin.Invest.* 72, 1218-1223.

Brown, M.P., Topham, D.J., Sangster, M.Y., Zhao, J., Flynn, K.J., Surman, S.L., Woodland, D.L., Doherty, P.C., Farr, A.G., Pattengale, P.K., and Brenner, M.K. (1998). Thymic lymphoproliferative disease after successful correction of CD40 ligand deficiency by gene transfer in mice. *Nat.Med.* 4, 1253-1260.

- Camussi, G., Aglietta, M., Malavasi, F., Tetta, C., Piacibello, W., Sanavio, F., and Bussolino, F. (1983). The release of platelet-activating factor from human endothelial cells in culture. *J.Immunol.* **131**, 2397-2403.
- Carbone, E., Ruggiero, G., Terrazzano, G., Palomba, C., Manzo, C., Fontana, S., Spits, H., Kaerre, K., and Zappacosta, S. (1997). A new mechanism of NK cell cytotoxicity activation: the CD40- CD40 ligand interaction. *J.Exp.Med.* **185**, 2053-2060.
- Casamayor-Palleja, M., Khan, M., and MacLennan, I.C. (1995). A subset of CD4<sup>+</sup> memory T cells contains preformed CD40 ligand that is rapidly but transiently expressed on their surface after activation through the T cell receptor complex. *J.Exp.Med.* **181**, 1293-1301.
- Caux, C., Massacrier, C., Vanbervliet, B., Dubois, B., van Kooten, C., Durand, I., and Banchereau, J. (1994). Activation of human dendritic cells through CD40 cross-linking. *J.Exp.Med.* **180**, 1263-1272.
- Chomczynski, P. and Sacchi, N. (1987). Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal.Biochem.* **162**, 156-159.
- Clark, E.A. and Ledbetter, J.A. (1986). Activation of human B cells mediated through two distinct cell surface differentiation antigens, Bp35 and Bp50. *Proc.Natl.Acad.Sci.USA* **83**, 4494-4498.
- Croft, M., Duncan, D.D., and Swain, S.L. (1992). Response of naive antigen-specific CD4<sup>+</sup> T cells in vitro: characteristics and antigen-presenting cell requirements. *J.Exp.Med.* **176**, 1431-1437.
- Curwen, K.D., Gimbrone, M.A., Jr., and Handin, R.I. (1980). In vitro studies of thromboresistance: the role of prostacyclin (PGI<sub>2</sub>) in platelet adhesion to cultured normal and virally transformed human vascular endothelial cells. *Lab.Invest.* **42**, 366-374.
- Dürkop, H., Anagnostopoulos, I. , Bulfone-Paus, S., and Stein, H. (1997). Expression of several members of the TNF-ligand and receptor family on tonsillar lymphoid B cells. *Br.J.Haematol.* **98**, 863-868.
- Dechanet, J., Grosset, C., Taupin, J.L., Merville, P., Banchereau, J., Ripoche, J., and Moreau, J.F. (1997). CD40 ligand stimulates proinflammatory cytokine production by human endothelial cells. *J.Immunol.* **159**, 5640-5647.
- Defrance, T., Vanbervliet, B., Briere, F., Durand, I., Rousset, F., and Banchereau, J. (1992). Interleukin 10 and transforming growth factor beta cooperate to induce anti-CD40-activated naive human B cells to secrete immunoglobulin A. *J.Exp.Med.* **175**, 671-682.

Deuel, T.F., Senior, R.M., Chang, D., Griffin, G.L., Heinrikson, R.L., and Kaiser, E.T. (1981). Platelet factor 4 is chemotactic for neutrophils and monocytes. *Proc.Natl.Acad.Sci.USA* 78, 4584-4587.

DiSanto, J.P., Bonnefoy, J.Y., Gauchat, J.F., Fischer, A., and de Saint Basile, G. (1993). CD40 ligand mutations in x-linked immunodeficiency with hyper- IgM. *Nature* 361, 541-543.

Eisen, S.A., Wedner, H.J., and Parker, C.W. (1972). Isolation of pure human peripheral blood T-lymphocytes using nylon wool columns. *Immunol.Commun.* 1, 571-577.

Fanslow, W.C., Srinivasan, S., Paxton, R., Gibson, M.G., Spriggs, M.K., and Armitage, R.J. (1994). Structural characteristics of CD40 ligand that determine biological function. *Semin.Immunol.* 6, 267-278.

Fries, K.M., Sempowski, G.D., Gaspari, A.A., Blieden, T., Looney, R.J., and Phipps, R.P. (1995). CD40 expression by human fibroblasts. *Clin.Immunol.Immunopathol.* 77, 42-51.

Fuleihan, R., Ramesh, N., Loh, R., Jabara, H., Rosen, R.S., Chatila, T., Fu, S.M., Stamenkovic, I., and Geha, R.S. (1993). Defective expression of the CD40 ligand in X chromosome-linked immunoglobulin deficiency with normal or elevated IgM. *Proc.Natl.Acad.Sci.USA* 90, 2170-2173.

Gauchat, J.F., Henchoz, S., Mazzei, G., Aubry, J.P., Brunner, T., Blasey, H., Life, P., Talabot, D., Flores-Romo, L., Thompson, J., Kishi, K., Butterfield, J., Dahinden, C., and Bonnefoy, J.Y. (1993). Induction of human IgE synthesis in B cells by mast cells and basophils. *Nature* 365, 340-343.

Gauchat, J.F., Henchoz, S., Fattah, D., Mazzei, G., Aubry, J.P., Jomotte, T., Dash, L., Page, K., Solari, R., Aldebert, D., Capron, M., Dahinden, C., and Bonnefoy, J.Y. (1995). CD40 ligand is functionally expressed on human eosinophils. *Eur.J.Immunol.* 25, 863-865.

George, J.N., Saucerman, S., Levine, S.P., Knieriem, L.K., and Bainton, D.F. (1985). Immunoglobulin G is a platelet alpha granule-secreted protein. *J.Clin.Invest.* 76, 2020-2025.

Gordon, J. (1995). CD40 and its ligand: central players in B lymphocyte survival, growth, and differentiation. *Blood Rev.* 9, 53-56.

Graf, D., Korthäuer, U., Mages, H.W., Senger, G., and Kroczeck, R.A. (1992). Cloning of TRAP, a ligand for CD40 on human T cells. *Eur.J.Immunol.* 22, 3191-3194.

Graf, D., Müller, S., Korthäuer, U., van Kooten, C., Weise, C., and Kroczeck, R.A. (1995). A soluble form of TRAP (CD40 ligand) is rapidly released after T cell activation. *Eur.J.Immunol.* 25, 1749-1754.

- Grewal, I.S. and Flavell, R.A. (1996). A central role of CD40 ligand in the regulation of CD4<sup>+</sup> T-cell responses. *Immunol.Today* 17, 410-414.
- Gruss, H.J. and Dower, S.K. (1995). Tumor necrosis factor ligand superfamily: involvement in the pathology of malignant lymphomas. *Blood* 85, 3378-3404.
- Hanahan, D., Jessee, J., and Bloom, F.R. (1991). Plasmid transformation of Escherichia coli and other bacteria. *Methods Enzymol.* 204, 63-113.
- Hawrylowicz, C.M., Santoro, S.A., Platt, F.M., and Unanue, E.R. (1989). Activated platelets express IL-1 activity. *J.Immunol.* 143, 4015-4018.
- Hawrylowicz, C.M., Howells, G.L., and Feldmann, M. (1991). Platelet-derived interleukin 1 induces human endothelial adhesion molecule expression and cytokine production. *J.Exp.Med.* 174, 785-790.
- He, M., Wilde, A., and Kaderbhai, M.A. (1990). A simple single-step procedure for small-scale preparation of Escherichia coli plasmids. *Nucleic Acids Res.* 18, 1660-1660.
- Herd, C.M. and Page, C.P. (1995). Do platelets have a role as inflammatory cells? In *Immunopharmacology of platelets*. M. Joseph, ed. (San Diego: Academic Press), pp. 1-20.
- Hollenbaugh, D., Mischel-Petty, N. , Edwards, C.P., Simon, J.C., Denfeld, R.W., Kiener, P.A., and Aruffo, A. (1995). Expression of functional CD40 by vascular endothelial cells. *J.Exp.Med.* 182, 33-40.
- Holmsen, H. (1994). Platelet secretion and energy metabolism. In *Hemostasis and Thrombosis* . R.W. Colman, J. Hirsh, V.J. Marder, and E.W. Salzman, eds. (Philadelphia: J.B. Lippincott Company), pp. 524-545.
- Innis, M.A., Gelfand, D.H., Sninsky, J.J., and White, T.J. (1990). PCR protocols: a guide to methods and applications (San Diego: Academic Press).
- Jaffe, E.A., Nachman, R.L., Becker, C.G., and Minick, C.R. (1973). Culture of human endothelial cells derived from umbilical veins. Identification by morphologic and immunologic criteria. *J.Clin.Invest.* 52, 2745-2756.
- Jaffe, E.A. (1994). Biochemistry, immunology and cell biology of endothelium. In *Hemostasis and Thrombosis*. R.W. Colman, J. Hirsh, V.J. Marder, and E.W. Salzman, eds. (Philadelphia: J.B. Lippincott Company), pp. 718-744.
- Kameyoshi, Y., Doerschner, A., Mallet, A.I., Christophers, E., and Schroeder, J.M. (1992). Cytokine RANTES released by thrombin-stimulated platelets is a potent attractant for human eosinophils. *J.Exp.Med.* 176, 587-592.

Kaplanski, G., Porat, R., Aiura, K., Erban, J.K., Gelfand, J.A., and Dinarello, C.A. (1993). Activated platelets induce endothelial secretion of interleukin- 8 in vitro via an interleukin-1-mediated event. *Blood* 81, 2492-2495.

Karmann, K., Hughes, C.C., Schechner, J., Fanslow, W.C., and Pober, J.S. (1995). CD40 on human endothelial cells: inducibility by cytokines and functional regulation of adhesion molecule expression. *Proc.Natl.Acad.Sci.USA* 92, 4342-4346.

Karmann, K., Min, W., Fanslow, W.C., and Pober, J.S. (1996a). Activation and homologous desensitization of human endothelial cells by CD40 ligand, tumor necrosis factor, and interleukin 1. *J.Exp.Med.* 184, 173-182.

Karmann, K., Hughes, C.C., Fanslow, W.C., and Pober, J.S. (1996b). Endothelial cells augment the expression of CD40 ligand on newly activated human CD4<sup>+</sup> T cells through a CD2/LFA-3 signaling pathway. *Eur.J.Immunol.* 26, 610-617.

Kehry, M.R. (1996). CD40-mediated signaling in B cells. Balancing cell survival, growth, and death. *J.Immunol.* 156, 2345-2348.

Kishimoto, T. (1989). The biology of interleukin-6. *Blood* 74, 1-10.

Klouche, M., Klinger, M.H., Kuehnel, W., and Wilhelm, D. (1997). Endocytosis, storage, and release of IgE by human platelets: differences in patients with type I allergy and nonatopic subjects. *J.Allergy Clin.Immunol.* 100, 235-241.

Komatsu, N., Nakauchi, H., Miwa, A., Ishihara, T., Eguchi, M., Moroi, M., Okada, M., Sato, Y., Wada, H., and et a (1991). Establishment and characterization of a human leukemic cell line with megakaryocytic features: dependency on granulocyte- macrophage colony-stimulating factor, interleukin 3, or erythropoietin for growth and survival. *Cancer Res* 51, 341-348.

Koppenhöfer, U., Brenner, B., Lang, F., and Gulbins, E. (1997). The CD40-ligand stimulates T-lymphocytes via the neutral sphingomyelinase: a novel function of the CD40-ligand as signalling molecule. *FEBS Lett.* 414, 444-448.

Korthäuer, U., Graf, D., Mages, H.W., Briere, F., Padayachee, M., Malcolm, S., Ugazio, A.G., Notarangelo, L.D., Levinsky, R.J., and Kroczeck, R.A. (1993). Defective expression of T-cell CD40 ligand causes X-linked immunodeficiency with hyper-IgM. *Nature* 361, 539-541.

Kuter, D.J., Beeler, D.L., and Rosenberg, R.D. (1994). The purification of megapoietin: a physiological regulator of megakaryocyte growth and platelet production. *Proc.Natl.Acad.Sci.USA* 91, 11104-11108.

Kyhse-Andersen, J. (1984). Electroblotting of multiple gels: a simple apparatus without buffer tank for rapid transfer of proteins from polyacrylamide to nitrocellulose. *J.Biochem.Biophys.Methods* 10, 203-209.

Laemmli, U.K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680-685.

Lauffer, L., Kanzy, E.J., Koehler, R., Kurrale, R., Enssle, K., and Seiler, F.R. (1995). Monomeric and dimeric forms of soluble receptors can differ in their neutralization potential. *Behring Inst.Mitt.* (96), 21-31.

Ledbetter, J.A., Shu, G., Gallagher, M., and Clark, E.A. (1987). Augmentation of normal and malignant B cell proliferation by monoclonal antibody to the B cell-specific antigen BP50 (CDW40). *J.Immunol.* 138, 788-794.

Ludewig, B., Henn, V., Schröder, J.M., Graf, D., and Kroczeck, R.A. (1996). Induction, regulation, and function of soluble TRAP (CD40 ligand) during interaction of primary CD4<sup>+</sup> CD45RA<sup>+</sup> T cells with dendritic cells. *Eur.J.Immunol.* 26, 3137-3143.

Mach, F., Schoenbeck, U., Sukhova, G.K., Bourcier, T., Bonnefoy, J.Y., Pober, J.S., and Libby, P. (1997). Functional CD40 ligand is expressed on human vascular endothelial cells, smooth muscle cells, and macrophages: implications for CD40-CD40 ligand signaling in atherosclerosis. *Proc.Natl.Acad.Sci.USA* 94, 1931-1936.

Mantovani, A., Bussolino, F., and Introna, M. (1997). Cytokine regulation of endothelial cell function: from molecular level to the bedside. *Immunol.Today* 18, 231-240.

Marcus, A.J., Safier, L.B., Hajjar, K.A., Ullman, H.L., Islam, N., Broekman, M.J., and Eiroa, A.M. (1991). Inhibition of platelet function by an aspirin-insensitive endothelial cell ADPase. Thromboregulation by endothelial cells. *J.Clin.Invest.* 88, 1690-1696.

Mazzei, G.J., Edgerton, M.D., Losberger, C., Lecoanet-Henchoz, S. , Gruber, P., Durandy, A., Gauchat, J.F., Bernard, A., Allet, B., and Bonnefoy, J.Y. (1995). Recombinant soluble trimeric CD40 ligand is biologically active. *J.Biol.Chem.* 270, 7025-7028.

Moncada, S., Herman, A.G., Higgs, E.A., and Vane, J.R. (1977). Differential formation of prostacyclin (PGX or PGI<sub>2</sub>) by layers of the arterial wall. An explanation for the anti-thrombotic properties of vascular endothelium. *Thromb.Res.* 11, 323-344.

Morley, J., Sanjar, S., and Page, C.P. (1984). The platelet in asthma. *Lancet* 2, 1142-1144.

Murray, A.G., Libby, P., and Pober, J.S. (1995). Human vascular smooth muscle cells poorly co-stimulate and actively inhibit allogeneic CD4+ T cell proliferation in vitro. *J.Immunol.* 154, 151-161.

Mustard, J.F. (1989). Isolation of human platelets from plasma by centrifugation and washing. *Methods Enzymol.* 169, 3-11.

Nagata, K., Tsuji, T., Todoroki, N., Katagiri, Y., Tanoue, K., Yamazaki, H., Hanai, N., and Irimura, T. (1993). Activated platelets induce superoxide anion release by monocytes and neutrophils through P-selectin (CD62). *J.Immunol.* 151, 3267-3273.

Nieuwenhuis, H.K., van Oosterhout, J.J., Rozemuller, E., van Iwaarden, F., and Sixma, J.J. (1987). Studies with a monoclonal antibody against activated platelets:evidence that a secreted 53,000-molecular weight lysosome-like granule protein is exposed on the surface of activated platelets in the circulation. *Blood* 70, 838-845.

Ogura, M., Morishima, Y., Ohno, R., Kato, Y., Hirabayashi, N., Nagura, H., and Saito, H. (1985). Establishment of a novel human megakaryoblastic leukemia cell line, MEG-01, with positive Philadelphia chromosome. *Blood* 66, 1384-1392.

Peguet-Navarro, J., Dalbiez-Gauthier, C., Moulon, C., Berthier, O., Reano, A., Gaucherand, M. , Banchereau, J., Rousset, F., and Schmitt, D. (1997). CD40 ligation of human keratinocytes inhibits their proliferation and induces their differentiation. *J.Immunol.* 158, 144-152.

Pietravalle, F., Lecoanet-Henchoz, S., Blasey, H., Aubry, J.P., Elson, G., Edgerton, M.D., Bonnefoy, J.Y., and Gauchat, J.F. (1996). Human native soluble CD40L is a biologically active trimer, processed inside microsomes. *J.Biol.Chem.* 271, 5965-5967.

Pinchuk, L.M., Klaus, S.J., Magaletti, D.M., Pinchuk, G.V., Norsen, J.P., and Clark, E.A. (1996). Functional CD40 ligand expressed by human blood dendritic cells is up-regulated by CD40 ligation. *J.Immunol.* 157, 4363-4370.

Qiagen (1992). The QIAexpressionist, high level purification system (Düsseldorf: Qiagen Inc.).

Radomski, M.W., Palmer, R.M., and Moncada, S. (1987). Endogenous nitric oxide inhibits human platelet adhesion to vascular endothelium. *Lancet* 2, 1057-1058.

Rohrer, M.J., Kestin, A.S., Ellis, P.A., Barnard, M.R., Rodino, L., Breckwoldt, W.L., Li, J.M., and Michelson, A.D. (1992). High-dose heparin suppresses platelet alpha granule secretion. *J.Vasc.Surg.* 15, 1000-8.

Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). Molecular cloning: a laboratory manual (New York: Cold Spring Harbor Laboratory Press).

Savage, B., Almus-Jacobs, F., and Ruggeri, Z.M. (1998). Specific synergy of multiple substrate-receptor interactions in platelet thrombus formation under flow. *Cell* 94, 657-666.

Schaegger, H. and von Jagow, G. (1987). Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Anal.Biochem.* 166, 368-379.

Schildkraut, C. (1965). Dependence of the melting temperature of DNA on salt concentration. *Biopolymers* 3, 195-208.

Schriever, F., Freedman, A.S., Freeman, G., Messner, E., Lee, G., Daley, J., and Nadler, L.M. (1989). Isolated human follicular dendritic cells display a unique antigenic phenotype. *J.Exp.Med.* 169, 2043-2058.

Sedlmayr, P., Blaschitz, A., Wilders-Truschnig, M., Tiran, A., and Dohr, G. (1995). Platelets contain interleukin-1 alpha and beta which are detectable on the cell surface after activation. *Scand.J.Immunol.* 42, 209-214.

Silverman, D.J. (1986). Adherence of platelets to human endothelial cells infected by Rickettsia rickettsii. *J.Infect.Dis.* 153, 694-700.

Snyder, F. (1990). Platelet-activating factor and related acetylated lipids as potent biologically active cellular mediators. *Am.J.Physiol.* 259 , Pt-708

Springer, T.A., Armitage, R.J., Strockbine, L., Clifford, K.N., Macduff, B.M., Sato, T.A., Maliszewski, C.R., and Fanslow, W.C. (1992). Recombinant human CD40 ligand stimulates B cell proliferation and immunoglobulin E secretion. *J.Exp.Med.* 176, 1543-1550.

Stamenkovic, I., Clark, E.A., and Seed, B. (1989). A B-lymphocyte activation molecule related to the nerve growth factor receptor and induced by cytokines in carcinomas. *Embo J.* 8, 1403-1410.

Steen, V.M., Holmsen, H., and Aarbakke, G. (1993). The platelet-stimulating effect of adrenaline through alpha 2-adrenergic receptors requires simultaneous activation by a true stimulatory platelet agonist. Evidence that adrenaline per se does not induce human platelet activation in vitro. *Thromb .Haemost* 70, 506-513.

Stenberg, P.E., McEver, R.P., Shuman, M.A., Jacques, Y.V., and Bainton, D.F. (1985). A platelet alpha-granule membrane protein (GMP-140) is expressed on the plasma membrane after activation. *J.Cell.Biol.* 101, 880-886.

Stout, R.D. and Suttles, J. (1996). The many roles of CD40 in cell-mediated inflammatory responses. *Immunol.Today* 17, 487-492.

Thuring, R.W., Sanders, J.P., and Borst, P. (1975). A freeze-squeeze method for recovering long DNA from agarose gels. *Anal.Biochem.* **66**, 213-220.

Toombs, C.F., Young, C.H., Glaspy, J.A., and Varnum, B.C. (1995). Megakaryocyte growth and development factor (MGDF) moderately enhances in-vitro platelet aggregation. *Thromb.Res.* **80**, 23-33.

Uckun, F.M., Schieven, G.L., Dibirdik, I., Chandan-Langlie, M., Tuel-Ahlgren, L., and Ledbetter, J.A. (1991). Stimulation of protein tyrosine phosphorylation, phosphoinositide turnover, and multiple previously unidentified serine/threonine-specific protein kinases by the Pan-B-cell receptor CD40/Bp50 at discrete developmental stages of human B- cell ontogeny. *J.Biol.Chem.* **266**, 17478-17485.

van Kooten, C. and Banchereau, J. (1996). CD40-CD40 ligand: a multifunctional receptor-ligand pair. *Adv.Immunol.* **61**, 1-77.

Venturini, C.M. and Kaplan, J.E. (1992). Thrombin induces platelet adhesion to endothelial cells. *Semin.Thromb.Hemost.* **18**, 275-283.

Walz, A., Dewald, B., von Tscharner, V., and Baggioolini, M. (1989). Effects of the neutrophil-activating peptide NAP-2, platelet basic protein, connective tissue-activating peptide III and platelet fycor 4 on human neutrophils. *J.Exp.Med.* **170**, 1745-1750.

Yellin, M.J., Sippel, K., Inghirami, G., Covey, L.R., Lee, J.J., Sining, J., Clark, E.A., Chess, L., and Lederman, S. (1994). CD40 molecules induce down-modulation and endocytosis of T cell surface T cell-B cell activating molecule/CD40-L. Potential role in regulating helper effector function. *J.Immunol.* **152**, 598-608.

Yellin, M.J., Winikoff, S., Fortune, S.M., Baum, D., Crow, M.K., Lederman, S., and Chess, L. (1995a). Ligation of CD40 on fibroblasts induces CD54 (ICAM-1) and CD106 (VCAM-1) up-regulation and IL-6 production and proliferation. *J.Leukoc.Biol.* **58**, 209-216.

Yellin, M.J., Brett, J., Baum, D., Matsushima, A., Szabolcs, M., Stern, D., and Chess, L. (1995b). Functional interactions of T cells with endothelial cells: the role of CD40L-CD40-mediated signals. *J.Exp.Med.* **182**, 1857-1864.

Zucker, M.B. (1980). The functioning of blood platelets. *Sci.Am.* **242**, 86-103.

## Danksagung

Herrn Prof. Dr. R. A. Kroczek danke ich für die Überlassung des Themas, sowie für die ausgezeichneten Arbeitsbedingungen, den unermüdlichen Einsatz und die vielen anregenden Diskussionen.

Herrn Dr. J. R. Slupsky danke ich herzlichst für die Einweisung in grundlegende Techniken, viele hilfreiche Ratschläge und ständige Ansprechbereitschaft.

Herrn Prof. Dr. F. Hucho danke ich für die freundliche Übernahme des Koreferats.

Herrn Dr. M. Gräfe danke ich für die Kultivierung der Endothelzellen, vielfältige Unterstützung und ständige Diskussionsbereitschaft.

Frau Dr. S. Müller danke ich dafür, daß sie mit grundlegenden Versuchen die Basis für den Erfolg dieser Arbeit gelegt hat.

Herrn Dr. R. Förster danke ich für die Durchführung der konfokalen Mikroskopie, Herrn Dr. I. Anagnostopoulos für die Durchführung der Immunhistologie.

Frau K. Ranke und Frau P. Jahn danke ich für die zuverlässige technische Unterstützung.

Vor allem möchte ich mich auch bei allen Mitarbeitern der Arbeitsgruppe bedanken, die mir stets mit Rat und Tat zur Seite standen und für eine hervorragende Arbeitsatmosphäre sorgten.

## Publikationen

Auszüge der vorliegenden Arbeit wurden veröffentlicht in:

Henn, V., Slupsky, J.R., Gräfe, M., Anagnostopoulos, I., Förster, R., Müller-Berghaus, G., and Kroczeck, R.A.

*CD40 ligand on activated platelets triggers an inflammatory reaction of endothelial cells*  
Nature (1998) 391, 591-594

Konferenzbeiträge:

Henn, V. and Kroczeck, R.A.

*CD40 on human platelets induces down-regulation of co-expressed TRAP (CD40 ligand)*  
Herbsttagung der Deutschen Gesellschaft für Immunologie, Freiburg 1998

Henn, V., Slupsky, J.R., Gräfe, M., Müller-Berghaus, G., and Kroczeck, R.A.

*TRAP (CD40 ligand) on activated platelets triggers a pro-inflammatory reaction of endothelial cells*  
Herbsttagung der Deutschen Gesellschaft für Immunologie, Würzburg 1997

**Lebenslauf****Persönliche Daten**

Name: Volker Henn  
Geburtsdatum: 27. März 1967  
Geburtsort: Klein Sisbeck, Kr. Helmstedt

**Schulausbildung**

1973-1977 Besuch der Albrecht-Dürer-Grundschule, Düsseldorf  
1977-1986 Besuch des Leibniz-Gymnasiums, Düsseldorf  
5.6.86 Erlangung der Allgemeinen Hochschulreife

**Hochschulausbildung**

April 1990 Immatrikulation im Fach Biochemie an der Freien Universität Berlin  
Februar-September 1995 Diplomarbeit bei Prof. R. A. Kroczek am Robert Koch-Institut, Berlin  
seit Oktober 1995 Dissertation bei Prof. R. A. Kroczek am Robert Koch-Institut, Berlin