

7 References

1. Pawson, T., Raina, M. and Nash, P. (2002). Interaction domains: from simple binding events to complex cellular behavior. *FEBS Lett* **513**, 2-10.
2. Ball, L.J., Jarchau, T., Oschkinat, H., *et al.* (2002). EVH1 domains: structure, function and interactions. *FEBS Lett* **513**, 45-52.
3. Pornillos, O., Alam, S.L., Rich, R.L., *et al.* (2002). Structure and functional interactions of the Tsg101 UEV domain. *Embo J* **21**, 2397-2406.
4. Zarrinpar, A., Bhattacharyya, R.P. and Lim, W.A. (2003). The structure and function of proline recognition domains. *Sci STKE* **2003**, RE8.
5. Schlessinger, J. (1994). SH2/SH3 signaling proteins. *Curr Opin Genet Dev* **4**, 25-30.
6. Pawson, T. (1995). Protein modules and signalling networks. *Nature* **373**, 573-580.
7. van der Geer, P. and Pawson, T. (1995). The PTB domain: a new protein module implicated in signal transduction. *Trends Biochem Sci* **20**, 277-280.
8. Yaffe, M.B. and Elia, A.E. (2001). Phosphoserine/threonine-binding domains. *Curr Opin Cell Biol* **13**, 131-138.
9. Sheng, M. and Sala, C. (2001). PDZ domains and the organization of supramolecular complexes. *Annu Rev Neurosci* **24**, 1-29.
10. van Ham, M. and Hendriks, W. (2003). PDZ domains-glue and guide. *Mol Biol Rep* **30**, 69-82.
11. Paoluzi, S., Castagnoli, L., Lauro, I., *et al.* (1998). Recognition specificity of individual EH domains of mammals and yeast. *Embo J* **17**, 6541-6550.
12. Cesareni, G., Panni, S., Nardelli, G., *et al.* (2002). Can we infer peptide recognition specificity mediated by SH3 domains? *FEBS Lett* **513**, 38-44.
13. Nguyen, J.T., Turck, C.W., Cohen, F.E., *et al.* (1998). Exploiting the basis of proline recognition by SH3 and WW domains: design of N-substituted inhibitors. *Science* **282**, 2088-2092.
14. Fesik, S.W. (2000). Insights into programmed cell death through structural biology. *Cell* **103**, 273-282.

15. Kim, C.A. and Bowie, J.U. (2003). SAM domains: uniform structure, diversity of function. *Trends Biochem Sci* **28**, 625-628.
16. Hillier, B.J., Christopherson, K.S., Prehoda, K.E., *et al.* (1999). Unexpected modes of PDZ domain scaffolding revealed by structure of nNOS-syntrophin complex. *Science* **284**, 812-815.
17. Pawson, T. and Nash, P. (2003). Assembly of cell regulatory systems through protein interaction domains. *Science* **300**, 445-452.
18. Venter, J.C., Adams, M.D., Myers, E.W., *et al.* (2001). The sequence of the human genome. *Science* **291**, 1304-1351.
19. Lander, E.S., Linton, L.M., Birren, B., *et al.* (2001). Initial sequencing and analysis of the human genome. *Nature* **409**, 860-921.
20. Yaffe, M.B., Leparc, G.G., Lai, J., *et al.* (2001). A motif-based profile scanning approach for genome-wide prediction of signaling pathways. *Nat Biotechnol* **19**, 348-353.
21. Tong, A.H., Drees, B., Nardelli, G., *et al.* (2002). A combined experimental and computational strategy to define protein interaction networks for peptide recognition modules. *Science* **295**, 321-324.
22. Kim, E., Niethammer, M., Rothschild, A., *et al.* (1995). Clustering of Shaker-type K⁺ channels by interaction with a family of membrane-associated guanylate kinases. *Nature* **378**, 85-88.
23. Woods, D.F. and Bryant, P.J. (1993). ZO-1, DlgA and PSD-95/SAP90: homologous proteins in tight, septate and synaptic cell junctions. *Mech Dev* **44**, 85-89.
24. Cho, K.O., Hunt, C.A. and Kennedy, M.B. (1992). The rat brain postsynaptic density fraction contains a homolog of the Drosophila discs-large tumor suppressor protein. *Neuron* **9**, 929-942.
25. Kennedy, M.B. (1995). Origin of PDZ (DHR, GLGF) domains. *Trends Biochem Sci* **20**, 350.
26. Ponting, C.P. (1997). Evidence for PDZ domains in bacteria, yeast, and plants. *Protein Sci* **6**, 464-468.

27. Nourry, C., Grant, S.G. and Borg, J.P. (2003). PDZ domain proteins: plug and play! *Sci STKE* **2003**, RE7.
28. Morais Cabral, J.H., Petosa, C., Sutcliffe, M.J., *et al.* (1996). Crystal structure of a PDZ domain. *Nature* **382**, 649-652.
29. Doyle, D.A., Lee, A., Lewis, J., *et al.* (1996). Crystal structures of a complexed and peptide-free membrane protein- binding domain: molecular basis of peptide recognition by PDZ. *Cell* **85**, 1067-1076.
30. Songyang, Z., Fanning, A.S., Fu, C., *et al.* (1997). Recognition of unique carboxyl-terminal motifs by distinct PDZ domains. *Science* **275**, 73-77.
31. Kozlov, G., Banville, D., Gehring, K., *et al.* (2002). Solution structure of the PDZ2 domain from cytosolic human phosphatase hPTP1E complexed with a peptide reveals contribution of the beta2-beta3 loop to PDZ domain-ligand interactions. *J Mol Biol* **320**, 813-820.
32. Karthikeyan, S., Leung, T., Birrane, G., *et al.* (2001). Crystal structure of the PDZ1 domain of human Na(+)/H(+) exchanger regulatory factor provides insights into the mechanism of carboxyl-terminal leucine recognition by class I PDZ domains. *J Mol Biol* **308**, 963-973.
33. Karthikeyan, S., Leung, T. and Ldias, J.A. (2002). Structural determinants of the Na⁺/H⁺ exchanger regulatory factor interaction with the beta 2 adrenergic and platelet-derived growth factor receptors. *J Biol Chem* **277**, 18973-18978.
34. Schultz, J., Hoffmuller, U., Krause, G., *et al.* (1998). Specific interactions between the syntrophin PDZ domain and voltage-gated sodium channels. *Nat Struct Biol* **5**, 19-24.
35. Brenman, J.E., Chao, D.S., Gee, S.H., *et al.* (1996). Interaction of nitric oxide synthase with the postsynaptic density protein PSD-95 and alpha1-syntrophin mediated by PDZ domains. *Cell* **84**, 757-767.
36. Tochio, H., Mok, Y.K., Zhang, Q., *et al.* (2000). Formation of nNOS/PSD-95 PDZ dimer requires a preformed beta-finger structure from the nNOS PDZ domain. *J Mol Biol* **303**, 359-370.

37. Im, Y.J., Park, S.H., Rho, S.H., *et al.* (2003). Crystal structure of GRIP1 PDZ6-peptide complex reveals the structural basis for class II PDZ target recognition and PDZ domain-mediated multimerization. *J Biol Chem* **278**, 8501-8507.
38. Zimmermann, P., Meerschaert, K., Reekmans, G., *et al.* (2002). PIP(2)-PDZ domain binding controls the association of syntenin with the plasma membrane. *Mol Cell* **9**, 1215-1225.
39. Harrison, S.C. (1996). Peptide-surface association: the case of PDZ and PTB domains. *Cell* **86**, 341-343.
40. Schepens, J., Cuppen, E., Wieringa, B., *et al.* (1997). The neuronal nitric oxide synthase PDZ motif binds to -G(D,E)XV* carboxyterminal sequences. *FEBS Lett* **409**, 53-56.
41. Maximov, A., Sudhof, T.C. and Bezprozvanny, I. (1999). Association of neuronal calcium channels with modular adaptor proteins. *J Biol Chem* **274**, 24453-24456.
42. Jaulin-Bastard, F., Saito, H., Le Bivic, A., *et al.* (2001). The ERBB2/HER2 receptor differentially interacts with ERBIN and PICK1 PSD-95/DLG/ZO-1 domain proteins. *J Biol Chem* **276**, 15256-15263.
43. Staudinger, J., Lu, J. and Olson, E.N. (1997). Specific interaction of the PDZ domain protein PICK1 with the COOH terminus of protein kinase C-alpha. *J Biol Chem* **272**, 32019-32024.
44. Irie, M., Hata, Y., Takeuchi, M., *et al.* (1997). Binding of neuroligins to PSD-95. *Science* **277**, 1511-1515.
45. Sattler, R., Xiong, Z., Lu, W.Y., *et al.* (1999). Specific coupling of NMDA receptor activation to nitric oxide neurotoxicity by PSD-95 protein. *Science* **284**, 1845-1848.
46. Brenman, J.E., Chao, D.S., Xia, H., *et al.* (1995). Nitric oxide synthase complexed with dystrophin and absent from skeletal muscle sarcolemma in Duchenne muscular dystrophy. *Cell* **82**, 743-752.
47. Thomas, G.D., Sander, M., Lau, K.S., *et al.* (1998). Impaired metabolic modulation of alpha-adrenergic vasoconstriction in dystrophin-deficient skeletal muscle. *Proc Natl Acad Sci U S A* **95**, 15090-15095.

48. Hsueh, Y.P., Kim, E. and Sheng, M. (1997). Disulfide-linked head-to-head multimerization in the mechanism of ion channel clustering by PSD-95. *Neuron* **18**, 803-814.
49. Dong, H., Zhang, P., Song, I., *et al.* (1999). Characterization of the glutamate receptor-interacting proteins GRIP1 and GRIP2. *J Neurosci* **19**, 6930-6941.
50. Srivastava, S., Osten, P., Vilim, F.S., *et al.* (1998). Novel anchorage of GluR2/3 to the postsynaptic density by the AMPA receptor-binding protein ABP. *Neuron* **21**, 581-591.
51. Xu, X.Z., Choudhury, A., Li, X., *et al.* (1998). Coordination of an array of signaling proteins through homo- and heteromeric interactions between PDZ domains and target proteins. *J Cell Biol* **142**, 545-555.
52. Cohen, N.A., Brenman, J.E., Snyder, S.H., *et al.* (1996). Binding of the inward rectifier K⁺ channel Kir 2.3 to PSD-95 is regulated by protein kinase A phosphorylation. *Neuron* **17**, 759-767.
53. Matsuda, S., Mikawa, S. and Hirai, H. (1999). Phosphorylation of serine-880 in GluR2 by protein kinase C prevents its C terminus from binding with glutamate receptor-interacting protein. *J Neurochem* **73**, 1765-1768.
54. Hall, R.A., Premont, R.T., Chow, C.W., *et al.* (1998). The beta2-adrenergic receptor interacts with the Na⁺/H⁺-exchanger regulatory factor to control Na⁺/H⁺ exchange. *Nature* **392**, 626-630.
55. Montell, C. (2000). A PDZ protein ushers in new links. *Nat Genet* **26**, 6-7.
56. Verpy, E., Leibovici, M., Zwaenepoel, I., *et al.* (2000). A defect in harmonin, a PDZ domain-containing protein expressed in the inner ear sensory hair cells, underlies Usher syndrome type 1C. *Nat Genet* **26**, 51-55.
57. Sherman, D.L., Fabrizi, C., Gillespie, C.S., *et al.* (2001). Specific disruption of a schwann cell dystrophin-related protein complex in a demyelinating neuropathy. *Neuron* **30**, 677-687.
58. Boerkoel, C.F., Takashima, H., Stankiewicz, P., *et al.* (2001). Periaxin mutations cause recessive Dejerine-Sottas neuropathy. *Am J Hum Genet* **68**, 325-333.

59. Prasad, R., Gu, Y., Alder, H., *et al.* (1993). Cloning of the ALL-1 fusion partner, the AF-6 gene, involved in acute myeloid leukemias with the t(6;11) chromosome translocation. *Cancer Res* **53**, 5624-5628.
60. Yamamoto, T., Harada, N., Kawano, Y., *et al.* (1999). In vivo interaction of AF-6 with activated Ras and ZO-1. *Biochem Biophys Res Commun* **259**, 103-107.
61. Quilliam, L.A., Castro, A.F., Rogers-Graham, K.S., *et al.* (1999). M-Ras/R-Ras3, a transforming ras protein regulated by Sos1, GRF1, and p120 Ras GTPase-activating protein, interacts with the putative Ras effector AF6. *J Biol Chem* **274**, 23850-23857.
62. Su, L., Hattori, M., Moriyama, M., *et al.* (2003). AF-6 controls integrin-mediated cell adhesion by regulating Rap1 activation through the specific recruitment of Rap1GTP and SPA-1. *J Biol Chem* **278**, 15232-15238.
63. Hock, B., Bohme, B., Karn, T., *et al.* (1998). PDZ-domain-mediated interaction of the Eph-related receptor tyrosine kinase EphB3 and the ras-binding protein AF6 depends on the kinase activity of the receptor. *Proc Natl Acad Sci U S A* **95**, 9779-9784.
64. Buchert, M., Schneider, S., Meskenaite, V., *et al.* (1999). The junction-associated protein AF-6 interacts and clusters with specific Eph receptor tyrosine kinases at specialized sites of cell-cell contact in the brain. *J Cell Biol* **144**, 361-371.
65. Takahashi, K., Nakanishi, H., Miyahara, M., *et al.* (1999). Nectin/PRR: an immunoglobulin-like cell adhesion molecule recruited to cadherin-based adherens junctions through interaction with Afadin, a PDZ domain-containing protein. *J Cell Biol* **145**, 539-549.
66. Ebnet, K., Schulz, C.U., Meyer Zu Brickwedde, M.K., *et al.* (2000). Junctional adhesion molecule interacts with the PDZ domain-containing proteins AF-6 and ZO-1. *J Biol Chem* **275**, 27979-27988.
67. Radziwill, G., Erdmann, R.A., Margelisch, U., *et al.* (2003). The Bcr kinase downregulates Ras signaling by phosphorylating AF-6 and binding to its PDZ domain. *Mol Cell Biol* **23**, 4663-4672.

68. Ikeda, W., Nakanishi, H., Miyoshi, J., *et al.* (1999). Afadin: A key molecule essential for structural organization of cell- cell junctions of polarized epithelia during embryogenesis. *J Cell Biol* **146**, 1117-1132.
69. Zhadanov, A.B., Provance, D.W., Jr., Speer, C.A., *et al.* (1999). Absence of the tight junctional protein AF-6 disrupts epithelial cell- cell junctions and cell polarity during mouse development. *Curr Biol* **9**, 880-888.
70. Borg, J.P., Marchetto, S., Le Bivic, A., *et al.* (2000). ERBIN: a basolateral PDZ protein that interacts with the mammalian ERBB2/HER2 receptor. *Nat Cell Biol* **2**, 407-414.
71. Klapper, L.N., Waterman, H., Sela, M., *et al.* (2000). Tumor-inhibitory antibodies to HER-2/ErbB-2 may act by recruiting c-Cbl and enhancing ubiquitination of HER-2. *Cancer Res* **60**, 3384-3388.
72. Yarden, Y. and Sliwkowski, M.X. (2001). Untangling the ErbB signalling network. *Nat Rev Mol Cell Biol* **2**, 127-137.
73. Huang, Y.Z., Zang, M., Xiong, W.C., *et al.* (2003). Erbin suppresses the MAP kinase pathway. *J Biol Chem* **278**, 1108-1114.
74. Jaulin-Bastard, F., Arsanto, J.P., Le Bivic, A., *et al.* (2002). Interaction between Erbin and a Catenin-related protein in epithelial cells. *J Biol Chem* **277**, 2869-2875.
75. Laura, R.P., Witt, A.S., Held, H.A., *et al.* (2002). The erbin PDZ domain binds with high affinity and specificity to the carboxyl-termini of d-catenin and ARVCF. *J Biol Chem* **30**, 12906-12914.
76. Bryant, P.J. and Huwe, A. (2000). LAP proteins: what's up with epithelia? *Nat Cell Biol* **2**, E141-143.
77. Bilder, D. and Perrimon, N. (2000). Localization of apical epithelial determinants by the basolateral PDZ protein Scribble. *Nature* **403**, 676-680.
78. Legouis, R., Gansmuller, A., Sookhareea, S., *et al.* (2000). LET-413 is a basolateral protein required for the assembly of adherens junctions in *Caenorhabditis elegans*. *Nat Cell Biol* **2**, 415-422.
79. Burshtyn, D.N., Yang, W., Yi, T., *et al.* (1997). A novel phosphotyrosine motif with a critical amino acid at position -2 for the SH2 domain-mediated activation of the tyrosine phosphatase SHP-1. *J Biol Chem* **272**, 13066-13072.

-
80. Favre, B., Fontao, L., Koster, J., *et al.* (2001). The hemidesmosomal protein bullous pemphigoid antigen 1 and the integrin beta 4 subunit bind to ERBIN. Molecular cloning of multiple alternative splice variants of ERBIN and analysis of their tissue expression. *J Biol Chem* **276**, 32427-32436.
 81. Adams, M.E., Dwyer, T.M., Dowler, L.L., *et al.* (1995). Mouse alpha 1- and beta 2-syntrophin gene structure, chromosome localization, and homology with a discs large domain. *J Biol Chem* **270**, 25859-25865.
 82. Straub, V. and Campbell, K.P. (1997). Muscular dystrophies and the dystrophin-glycoprotein complex. *Curr Opin Neurol* **10**, 168-175.
 83. Hasegawa, M., Cuenda, A., Spillantini, M.G., *et al.* (1999). Stress-activated protein kinase-3 interacts with the PDZ domain of alpha1-syntrophin. A mechanism for specific substrate recognition. *J Biol Chem* **274**, 12626-12631.
 84. Olalla, L., Aledo, J.C., Bannenberg, G., *et al.* (2001). The C-terminus of human glutaminase L mediates association with PDZ domain-containing proteins. *FEBS Lett* **488**, 116-122.
 85. Adams, M.E., Mueller, H.A. and Froehner, S.C. (2001). In vivo requirement of the alpha-syntrophin PDZ domain for the sarcolemmal localization of nNOS and aquaporin-4. *J Cell Biol* **155**, 113-122.
 86. Gee, S.H., Madhavan, R., Levinson, S.R., *et al.* (1998). Interaction of muscle and brain sodium channels with multiple members of the syntrophin family of dystrophin-associated proteins. *J Neurosci* **18**, 128-137.
 87. Rehm, T., Huber, R. and Holak, T.A. (2002). Application of NMR in structural proteomics: screening for proteins amenable to structural analysis. *Structure (Camb)* **10**, 1613-1618.
 88. Lian, L.Y., Barsukov, I.L., Derrick, J.P., *et al.* (1994). Mapping the interactions between streptococcal protein G and the Fab fragment of IgG in solution. *Nat Struct Biol* **1**, 355-357.
 89. Barsukov, I.L., Lian, L.Y., Ellis, J., *et al.* (1996). The conformation of coenzyme A bound to chloramphenicol acetyltransferase determined by transferred NOE experiments. *J Mol Biol* **262**, 543-558.

90. DeRisi, J.L., Iyer, V.R. and Brown, P.O. (1997). Exploring the metabolic and genetic control of gene expression on a genomic scale. *Science* **278**, 680-686.
91. Ito, T., Tashiro, K., Muta, S., *et al.* (2000). Toward a protein-protein interaction map of the budding yeast: A comprehensive system to examine two-hybrid interactions in all possible combinations between the yeast proteins. *Proc Natl Acad Sci USA* **97**, 1143-1147.
92. Ross-Macdonald, P., Coelho, P.S., Roemer, T., *et al.* (1999). Large-scale analysis of the yeast genome by transposon tagging and gene disruption. *Nature* **402**, 413-418.
93. Uetz, P., Giot, L., Cagney, G., *et al.* (2000). A comprehensive analysis of protein-protein interactions in *Saccharomyces cerevisiae*. *Nature* **403**, 623-627.
94. Winzeler, E.A., Shoemaker, D.D., Astromoff, A., *et al.* (1999). Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science* **285**, 901-906.
95. Xenarios, I., Salwinski, L., Duan, X.J., *et al.* (2002). DIP, the Database of Interacting Proteins: a research tool for studying cellular networks of protein interactions. *Nucleic Acids Res* **30**, 303-305.
96. Bader, G.D., Donaldson, I., Wolting, C., *et al.* (2001). BIND--The Biomolecular Interaction Network Database. *Nucleic Acids Res* **29**, 242-245.
97. Zanzoni, A., Montecchi-Palazzi, L., Quondam, M., *et al.* (2002). MINT: a Molecular INTeraction database. *FEBS Lett* **513**, 135-140.
98. Xenarios, I. and Eisenberg, D. (2001). Protein interaction databases. *Curr Opin Biotechnol* **12**, 334-339.
99. Valencia, A. and Pazos, F. (2002). Computational methods for the prediction of protein interactions. *Curr Opin Struct Biol* **12**, 368-373.
100. Brannetti, B., Via, A., Cestra, G., *et al.* (2000). SH3-SPOT: an algorithm to predict preferred ligands to different members of the SH3 gene family. *J Mol Biol* **298**, 313-328.
101. Fisinger, S., Serrano, L. and Lacroix, E. (2001). Computational estimation of specific side chain interaction energies in alpha helices. *Protein Sci* **10**, 809-818.

102. Lopez de la Paz, M., Lacroix, E., Ramirez-Alvarado, M., *et al.* (2001). Computer-aided design of beta-sheet peptides. *J Mol Biol* **312**, 229-246.
103. Reina, J., Lacroix, E., Hobson, S.D., *et al.* (2002). Computer-aided design of a PDZ domain to recognize new target sequences. *Nat Struct Biol* **9**, 621-627.
104. Hoffmüller, U., Russwurm, M., Kleinjung, F., *et al.* (1999). Interaktion einer PDZ-Proteindomäne mit einer synthetischen Bibliothek aller C-Termini humaner Proteine. *Angew. Chem.* **111**, 2180-2184.
105. Aasland, R., Abrams, C., Ampe, C., *et al.* (2002). Normalization of nomenclature for peptide motifs as ligands of modular protein domains. *FEBS Lett* **513**, 141-144.
106. Peranen, J., Rikkinen, M., Hyvonen, M., *et al.* (1996). T7 vectors with modified T7lac promoter for expression of proteins in Escherichia coli. *Anal Biochem* **236**, 371-373.
107. Sambrook, J. and Russell, D. (2001). *Molecular Cloning*, third edition Edition (New York: Cold Spring Harbor Laboratory Press).
108. Walker, P.A., Leong, L.E., Ng, P.W., *et al.* (1994). Efficient and rapid affinity purification of proteins using recombinant fusion proteases. *Biotechnology (N Y)* **12**, 601-605.
109. Cordingley, M.G., Callahan, P.L., Sardana, V.V., *et al.* (1990). Substrate requirements of human rhinovirus 3C protease for peptide cleavage in vitro. *J Biol Chem* **265**, 9062-9065.
110. Gill, S.C. and von Hippel, P.H. (1989). Calculation of protein extinction coefficients from amino acid sequence data. *Anal Biochem* **182**, 319-326.
111. Licha, K., Bhargava, S., Rheinländer, C., *et al.* (2000). Highly parallel nano-synthesis of cleavable peptide-dye conjugates on cellulose membranes. *Tetrahedron Letters* **41**, 1711-1715.
112. Kramer, A. and Schneider-Mergener, J. (1998). Synthesis and screening of peptide libraries on continuous cellulose membrane supports. *Methods Mol Biol* **87**, 25-39.
113. Frank, R. (1992). Spot synthesis an easy technique for positionally addressable, parallel chemical synthesis on a membrane support. *Tetrahedron* **48**, 9217-9232.

-
114. Schubert, M., Smalla, M., Schmieder, P., *et al.* (1999). MUSIC in triple-resonance experiments: amino acid type-selective (1)H- (15)N correlations. *J Magn Reson* **141**, 34-43.
 115. Schubert, M., Oschkinat, H. and Schmieder, P. (2001). MUSIC and aromatic residues: amino acid type-selective (1)H-(15)N correlations, III. *J Magn Reson* **153**, 186-192.
 116. Schubert, M., Oschkinat, H. and Schmieder, P. (2001). MUSIC, selective pulses, and tuned delays: amino acid type-selective (1)H-(15)N correlations, II. *J Magn Reson* **148**, 61-72.
 117. Schubert, M., Oschkinat, H. and Schmieder, P. (2001). Amino acid type-selective backbone 1H-15N-correlations for Arg and Lys. *J Biomol NMR* **20**, 379-384.
 118. Grzesiek, S., Dobeli, H., Gentz, R., *et al.* (1992). 1H, 13C, and 15N NMR backbone assignments and secondary structure of human interferon-gamma. *Biochemistry* **31**, 8180-8190.
 119. Grzesiek, S. and Bax, A. (1993). Amino acid type determination in the sequential assignment procedure of uniformly 13C/15N-enriched proteins. *J Biomol NMR* **3**, 185-204.
 120. Pires, J.R., Hong, X., Brockmann, C., *et al.* (2003). The ScPex13p SH3 domain exposes two distinct binding sites for Pex5p and Pex14p. *J Mol Biol* **326**, 1427-1435.
 121. Laskowski, R.A., Rullmann, J.A., MacArthur, M.W., *et al.* (1996). AQUA and PROCHECK-NMR: programs for checking the quality of protein structures solved by NMR. *J Biomol NMR* **8**, 477-486.
 122. Birrane, G., Chung, J. and Ldias, J.A. (2003). Novel mode of ligand recognition by the Erbin PDZ domain. *J Biol Chem* **278**, 1399-1402.
 123. AMBER: Assisted model building with energy refinement. (2002). Version 7, Department of Pharmaceutical Chemistry, University of California, San Francisco
 124. Cornell, W.D., Cieplak, P., Payly, C.I., *et al.* (1995). A second generation force field for the simulation of proteins, nucleic acids and organic molecules. *J Am Chem Soc* **117**, 5179-5197.

125. Koradi, R., Billeter, M. and Wuthrich, K. (1996). MOLMOL: a program for display and analysis of macromolecular structures. *J Mol Graph* **14**, 51-55, 29-32.
126. Venables, W.N. and Ripley, B.D. (2002). Modern Applied Statistics with S, Fourth edition Edition (New York: Springer).
127. Lehmann, G. (2002). Statistik, First edition Edition (Heidelberg: Spektrum Akademischer Verlag).
128. Laird, N.M. and Ware, J.H. (1982). Random-effects models for longitudinal data. *Biometrics* **38**, 963-974.
129. Team, R.D.C. (2003). R: A language and environment for statistical computing (Vienna, Austria: R Foundation for Statistical Computing).
130. Henn, V., Edemir, B., Stefan, E., *et al.* (2004). Identification of a novel A-kinase anchoring protein 18 isoform and evidence for its role in the vasopressin-induced aquaporin-2 shuttle in renal principal cells. *J Biol Chem* **279**, 26654-26665.
131. Cabilly, S. (1998). Methods in Molecular Biology, Volume 87 (New Jersey: Humana Press Inc.).
132. Songyang, Z., Blechner, S., Hoagland, N., *et al.* (1994). Use of an oriented peptide library to determine the optimal substrates of protein kinases. *Curr Biol* **4**, 973-982.
133. Otte, L., Wiedemann, U., Schlegel, B., *et al.* (2003). WW domain sequence activity relationships identified using ligand recognition propensities of 42 WW domains. *Protein Sci* **12**, 491-500.
134. Landgraf, c., Panni, S., Montecchi-Palazzi, L., *et al.* (2003). Protein interaction networks by proteome peptide scanning. *PLOS Biology* **2**, 94-103.
135. Rodriguez, M., Li, S.S., Harper, J.W., *et al.* (2003). An oriented peptide array library (OPAL) strategy to study protein-protein interactions. *J Biol Chem*.
136. Volkmer-Engert, R., Hoffmann, B. and Schneider-Mergener, J. (1997). Stable attachment of the HMB-linker to continuous cellulose membranes for parallel solid phase spot synthesis. *Tetrahedron Lett* **38**, 1029-1032.
137. Bhargava, S., Licha, K., Knaute, T., *et al.* (2002). A complete substitutional analysis of VIP for better tumor imaging properties. *J Mol Recognit* **15**, 145-153.

-
138. Reineke, U., Volkmer-Engert, R. and Schneider-Mergener, J. (2001). Applications of peptide arrays prepared by the SPOT-technology. *Curr Opin Biotechnol* **12**, 59-64.
 139. Hung, A.Y. and Sheng, M. (2002). PDZ domains: structural modules for protein complex assembly. *J Biol Chem* **277**, 5699-5702.
 140. Davies, M., Bonnat, M., Guiller, F., *et al.* (1998). Screening an inverted peptide library in water with a guanidinium-based tweezer receptor. *J Org Chem* **63**, 8696-8703.
 141. Davies, M. and Bradley, M. (1997). C-terminally modified peptides and peptide libraries - another end to peptide synthesis. *Angew Chem, Int. Ed. Engl.* **36**, 1097-1110.
 142. Kania, R.S., Zuckermann, R.N. and Marlove, C.K. (1994). Free C-Terminal Resin-Bound Peptides: Reversal of Peptide Orientation via a Cyclization/Cleavage Protocol. *J Am Chem Soc* **116**, 8835-8836.
 143. Frank, R. (2002). The SPOT-synthesis technique. Synthetic peptide arrays on membrane supports--principles and applications. *J Immunol Methods* **267**, 13-26.
 144. Carpino, L.A., Sadat-Aalaese, D., Chao, H.G., *et al.* (1990). [(9-Fluorenylmethyl)oxy]carbonyl (Fmoc) amino acid fluorides. Convenient new peptide coupling reagents applicable to the Fmoc/tert-butyl strategy for solution and solid-phase syntheses. *J Am Chem Soc* **112**, 9651-9652.
 145. Granitza, D., Beyermann, M., Wenschuh, H., *et al.* (1995). Efficient acylation of hydroxy functions by means of Fmoc amino acid fluorides. *J. Chem. Soc., Chem. Commun.*, 2223-2224.
 146. March, J. (1985). In Advanced Organic Chemistry. In *Esterification of acids*. pp. pp. 348-353, John Wiley & Sons: New York.
 147. Roberts, K.D., Lambert, J.N., Ede, N.K., *et al.* (1998). Efficient synthesis of thioether-based cyclic peptide libraries. *Tetrahedron Letters* **39**, 8357-8360.
 148. Niethammer, M., Valtschanoff, J.G., Kapoor, T.M., *et al.* (1998). CRIPT, a novel postsynaptic protein that binds to the third PDZ domain of PSD-95/SAP90. *Neuron* **20**, 693-707.

149. Lu, Q., Paredes, M., Medina, M., *et al.* (1999). delta-catenin, an adhesive junction-associated protein which promotes cell scattering. *J Cell Biol* **144**, 519-532.
150. Izawa, I., Nishizawa, M., Tomono, Y., *et al.* (2002). ERBIN associates with p0071, an armadillo protein, at cell-cell junctions of epithelial cells. *Genes Cells* **7**, 475-485.
151. Kolch, W. (2003). Erbin: sorting out ErbB2 receptors or giving Ras a break? *Sci STKE* **2003**, pe37.
152. Boguski, M.S. and McCormick, F. (1993). Proteins regulating Ras and its relatives. *Nature* **366**, 643-654.
153. Scheffzek, K., Ahmadian, M.R. and Wittinghofer, A. (1998). GTPase-activating proteins: helping hands to complement an active site. *Trends Biochem Sci* **23**, 257-262.
154. Bezprozvanny, I. and Maximov, A. (2001). Classification of PDZ domains. *FEBS Lett* **509**, 457-462.
155. Vaccaro, P. and Dente, L. (2002). PDZ domains: troubles in classification. *FEBS Lett* **512**, 345-349.
156. Fan, J.S. and Zhang, M. (2002). Signaling complex organization by PDZ domain proteins. *Neurosignals* **11**, 315-321.
157. Borrell-Pages, M., Fernandez-Larrea, J., Borroto, A., *et al.* (2000). The carboxy-terminal cysteine of the tetraspanin L6 antigen is required for its interaction with SITAC, a novel PDZ protein. *Mol Biol Cell* **11**, 4217-4225.
158. Harris, B.Z. and Lim, W.A. (2001). Mechanism and role of PDZ domains in signaling complex assembly. *J Cell Sci* **114**, 3219-3231.
159. Kimple, M.E., Siderovski, D.P. and Sondek, J. (2001). Functional relevance of the disulfide-linked complex of the N-terminal PDZ domain of InaD with NorpA. *Embo J* **20**, 4414-4422.
160. Boisguerin, P., Leben, R., Ay, B., *et al.* (2004). An improved method for the synthesis of cellulose membrane-bound peptides with free C termini is useful for PDZ domain binding studies. *Chem Biol* **11**, 449-459.

161. Skelton, N.J., Koehler, M.F., Zobel, K., *et al.* (2003). Origins of PDZ domain ligand specificity. Structure determination and mutagenesis of the Erbin PDZ domain. *J Biol Chem* **278**, 7645-7654.
162. Koike, G., Horiuchi, M., Yamada, T., *et al.* (1994). Human type 2 angiotensin II receptor gene: cloned, mapped to the X chromosome, and its mRNA is expressed in the human lung. *Biochem Biophys Res Commun* **203**, 1842-1850.
163. Murphy, T.J., Alexander, R.W., Griendling, K.K., *et al.* (1991). Isolation of a cDNA encoding the vascular type-1 angiotensin II receptor. *Nature* **351**, 233-236.
164. Timmermans, P.B., Benfield, P., Chiu, A.T., *et al.* (1992). Angiotensin II receptors and functional correlates. *Am J Hypertens* **5**, 221S-235S.
165. Schmitz, U., Ishida, M. and Berk, B.C. (1997). Angiotensin II stimulates tyrosine phosphorylation of phospholipase C-gamma-associated proteins. Characterization of a c-Src-dependent 97-kD protein in vascular smooth muscle cells. *Circ Res* **81**, 550-557.
166. Bottari, S.P., de Gasparo, M., Steckelings, U.M., *et al.* (1993). Angiotensin II receptor subtypes: characterization, signalling mechanisms, and possible physiological implications. *Front Neuroendocrinol* **14**, 123-171.
167. de Gasparo, M., Catt, K.J., Inagami, T., *et al.* (2000). International union of pharmacology. XXIII. The angiotensin II receptors. *Pharmacol Rev* **52**, 415-472.
168. Nouet, S. and Nahmias, C. (2000). Signal transduction from the angiotensin II AT2 receptor. *Trends Endocrinol Metab* **11**, 1-6.
169. Ichiki, T., Labosky, P.A., Shiota, C., *et al.* (1995). Effects on blood pressure and exploratory behaviour of mice lacking angiotensin II type-2 receptor. *Nature* **377**, 748-750.
170. Hein, L., Barsh, G.S., Pratt, R.E., *et al.* (1995). Behavioural and cardiovascular effects of disrupting the angiotensin II type-2 receptor in mice. *Nature* **377**, 744-747.
171. Nahmias, C. and Strosberg, A.D. (1995). The angiotensin AT2 receptor: searching for signal-transduction pathways and physiological function. *Trends Pharmacol Sci* **16**, 223-225.

-
172. Stoll, M. and Unger, T. (2001). Angiotensin and its AT2 receptor: new insights into an old system. *Regul Pept* **99**, 175-182.
 173. Zhu, M., Gelband, C.H., Moore, J.M., *et al.* (1998). Angiotensin II type 2 receptor stimulation of neuronal delayed-rectifier potassium current involves phospholipase A2 and arachidonic acid. *J Neurosci* **18**, 679-686.
 174. Horiuchi, M., Hayashida, W., Kambe, T., *et al.* (1997). Angiotensin type 2 receptor dephosphorylates Bcl-2 by activating mitogen-activated protein kinase phosphatase-1 and induces apoptosis. *J Biol Chem* **272**, 19022-19026.
 175. Bedecs, K., Elbaz, N., Sutren, M., *et al.* (1997). Angiotensin II type 2 receptors mediate inhibition of mitogen-activated protein kinase cascade and functional activation of SHP-1 tyrosine phosphatase. *Biochem J* **325 (Pt 2)**, 449-454.
 176. Buisson, B., Laflamme, L., Bottari, S.P., *et al.* (1995). A G protein is involved in the angiotensin AT2 receptor inhibition of the T-type calcium current in non-differentiated NG108-15 cells. *J Biol Chem* **270**, 1670-1674.
 177. Gallinat, S., Busche, S., Schutze, S., *et al.* (1999). AT2 receptor stimulation induces generation of ceramides in PC12W cells. *FEBS Lett* **443**, 75-79.
 178. Lehtonen, J.Y., Daviet, L., Nahmias, C., *et al.* (1999). Analysis of functional domains of angiotensin II type 2 receptor involved in apoptosis. *Mol Endocrinol* **13**, 1051-1060.
 179. Kang, J., Posner, P. and Summers, C. (1994). Angiotensin II type 2 receptor stimulation of neuronal K⁺ currents involves an inhibitory GTP binding protein. *Am J Physiol* **267**, C1389-1397.
 180. Masaki, H., Kurihara, T., Yamaki, A., *et al.* (1998). Cardiac-specific overexpression of angiotensin II AT2 receptor causes attenuated response to AT1 receptor-mediated pressor and chronotropic effects. *J Clin Invest* **101**, 527-535.
 181. Horiuchi, M., Akishita, M. and Dzau, V.J. (1999). Recent progress in angiotensin II type 2 receptor research in the cardiovascular system. *Hypertension* **33**, 613-621.
 182. Dulin, N.O., Alexander, L.D., Harwalkar, S., *et al.* (1998). Phospholipase A2-mediated activation of mitogen-activated protein kinase by angiotensin II. *Proc Natl Acad Sci U S A* **95**, 8098-8102.
-

183. AbdAlla, S., Lothar, H., Abdel-tawab, A.M., *et al.* (2001). The angiotensin II AT2 receptor is an AT1 receptor antagonist. *J Biol Chem* **276**, 39721-39726.
184. Huang, X.C., Summers, C. and Richards, E.M. (1996). Angiotensin II stimulates protein phosphatase 2A activity in cultured neuronal cells via type 2 receptors in a pertussis toxin sensitive fashion. *Adv Exp Med Biol* **396**, 209-215.
185. Horiuchi, M., Hayashida, W., Akishita, M., *et al.* (1999). Stimulation of different subtypes of angiotensin II receptors, AT1 and AT2 receptors, regulates STAT activation by negative crosstalk. *Circ Res* **84**, 876-882.
186. Hein, L., Meinel, L., Pratt, R.E., *et al.* (1997). Intracellular trafficking of angiotensin II and its AT1 and AT2 receptors: evidence for selective sorting of receptor and ligand. *Mol Endocrinol* **11**, 1266-1277.
187. Freyd, G., Kim, S.K. and Horvitz, H.R. (1990). Novel cysteine-rich motif and homeodomain in the product of the *Caenorhabditis elegans* cell lineage gene *lin-11*. *Nature* **344**, 876-879.
188. Karlsson, O., Thor, S., Norberg, T., *et al.* (1990). Insulin gene enhancer binding protein Isl-1 is a member of a novel class of proteins containing both a homeo- and a Cys-His domain. *Nature* **344**, 879-882.
189. Kloiber, K., Weiskirchen, R., Krautler, B., *et al.* (1999). Mutational analysis and NMR spectroscopy of quail cysteine and glycine-rich protein CRP2 reveal an intrinsic segmental flexibility of LIM domains. *J Mol Biol* **292**, 893-908.
190. Konrat, R., Krautler, B., Weiskirchen, R., *et al.* (1998). Structure of cysteine- and glycine-rich protein CRP2. Backbone dynamics reveal motional freedom and independent spatial orientation of the lim domains. *J Biol Chem* **273**, 23233-23240.
191. Sanchez-Garcia, I. and Rabbitts, T.H. (1994). The LIM domain: a new structural motif found in zinc-finger-like proteins. *Trends Genet* **10**, 315-320.
192. Dawid, I.B., Breen, J.J. and Toyama, R. (1998). LIM domains: multiple roles as adapters and functional modifiers in protein interactions. *Trends Genet* **14**, 156-162.
193. Warren, A.J., Colledge, W.H., Carlton, M.B., *et al.* (1994). The oncogenic cysteine-rich LIM domain protein *rbtn2* is essential for erythroid development. *Cell* **78**, 45-57.

194. Valge-Archer, V., Forster, A. and Rabbitts, T.H. (1998). The LMO1 AND LDB1 proteins interact in human T cell acute leukaemia with the chromosomal translocation t(11;14)(p15;q11). *Oncogene* **17**, 3199-3202.
195. Rabbitts, T.H. (1994). Chromosomal translocations in human cancer. *Nature* **372**, 143-149.
196. Murthy, K.K., Clark, K., Fortin, Y., *et al.* (1999). ZRP-1, a zyxin-related protein, interacts with the second PDZ domain of the cytosolic protein tyrosine phosphatase hPTP1E. *J Biol Chem* **274**, 20679-20687.
197. Cuppen, E., Gerrits, H., Pepers, B., *et al.* (1998). PDZ motifs in PTP-BL and RIL bind to internal protein segments in the LIM domain protein RIL. *Mol Biol Cell* **9**, 671-683.