

7 Literaturverzeichnis

- (1) DONALD I, MACVICAR J, BROWN TG. Investigation of abdominal masses by pulsed ultrasound. *Lancet* 1958 June 7;1(7032):1188-95.
- (2) Boyd PA, Chamberlain P, Hicks NR. 6-year experience of prenatal diagnosis in an unselected population in Oxford, UK. *Lancet* 1998 November 14;352(9140):1577-81.
- (3) Gotzmann L, Schonholzer SM, Kolble N et al. [Ultrasound examinations in the context of suspected fetal malformations: satisfaction of concerned women and their appraisals]. *Ultraschall Med* 2002 February;23(1):27-32.
- (4) Hedrick HL, Flake AW, Crombleholme TM et al. History of fetal diagnosis and therapy: Children's Hospital of Philadelphia experience. *Fetal Diagn Ther* 2003 March;18(2):65-82.
- (5) Stiller R, Huch R, Huch A, Zimmermann R. [Quality of praenatal diagnostic ultrasound - comparison of sonographically detected foetal anomalies with diagnostic findings verified post-partum in Switzerland]. *Ultraschall Med* 2001 October;22(5):225-30.
- (6) Queisser-Luft A, Stopfkuchen H, Stolz G et al. Prenatal diagnosis of major malformations: quality control of routine ultrasound examinations based on a five-year study of 20,248 newborn fetuses and infants. *Prenat Diagn* 1998 June;18(6):567-76.
- (7) Carrera JM, Torrents M, Mortera C et al. Routine prenatal ultrasound screening for fetal abnormalities: 22 years' experience. *Ultrasound Obstet Gynecol* 1995 March;5(3):174-9.
- (8) Merz E, Eichhorn KH, Hansmann M, Meinel K. [Quality demands on continuing differential diagnostic sonography in prenatal diagnostics (DEGUM stage II) during the 18th to 22nd weeks of gestation]. *Ultraschall Med* 2002 February;23(1):11-2.
- (9) Park YW. Diagnosis of fetal anomalies by sonography. *Yonsei Med J* 2001 December;42(6):660-8.
- (10) Kumar S, O'Brien A. Recent developments in fetal medicine. *BMJ* 2004 April 24;328(7446):1002-6.
- (11) Chaoui R, Heling KS. Three-dimensional ultrasound in prenatal diagnosis. *Curr Opin Obstet Gynecol* 2006 April;18(2):192-202.
- (12) Frates MC, Kumar AJ, Benson CB et al. Fetal anomalies: comparison of MR imaging and US for diagnosis. *Radiology* 2004 August;232(2):398-404.
- (13) Staebler M, Donner C, Van RN et al. Should determination of the karyotype be systematic for all malformations detected by obstetrical ultrasound? *Prenat Diagn* 2005 July;25(7):567-73.
- (14) Firth HV, Boyd PA, Chamberlain PF et al. Analysis of limb reduction defects in babies exposed to chorionic villus sampling. *Lancet* 1994 April 30;343(8905):1069-71.
- (15) Paek B, Goldberg JD, Albanese CT. Prenatal diagnosis. *World J Surg* 2003 January;27(1):27-37.
- (16) Matsuoka S, Takeuchi K, Yamanaka Y et al. Comparison of magnetic resonance imaging and ultrasonography in the prenatal diagnosis of congenital thoracic abnormalities. *Fetal Diagn Ther* 2003 November;18(6):447-53.
- (17) Garne E, Loane M, de VC et al. Prenatal diagnostic procedures used in pregnancies with congenital malformations in 14 regions of Europe. *Prenat Diagn* 2004 November;24(11):908-12.
- (18) Strauss A, Muller-Egloff S, Heer IM. [Obstetrical ultrasonography]. *Gynakol Geburtshilfliche Rundsch* 2002;42(2):75-83.

- (19) Raupach K, Zimmermann R. [False diagnosis in prenatal sonography - analysis of causes and formulation of conclusions for the quality management of prenatal sonographic diagnostics.]. *Ultraschall Med* 2004 December;25(6):438-43.
- (20) Skari H, Bjornland K, Bjornstad-Ostensen A et al. Consequences of prenatal ultrasound diagnosis: a preliminary report on neonates with congenital malformations. *Acta Obstet Gynecol Scand* 1998 July;77(6):635-42.
- (21) Stocker JT, Madewell JE, Drake RM. Congenital cystic adenomatoid malformation of the lung. Classification and morphologic spectrum. *Hum Pathol* 1977 March;8(2):155-71.
- (22) Berge SJ, Plath H, Reich RH, Hansmann M. [Significance of prenatal diagnosis of lip-jaw-palatal clefts]. *Mund Kiefer Gesichtschir* 2002 March;6(2):85-90.
- (23) Mammen L, Benson CB. Outcome of fetuses with clubfeet diagnosed by prenatal sonography. *J Ultrasound Med* 2004 April;23(4):497-500.
- (24) Barisic I, Clementi M, Hausler M et al. Evaluation of prenatal ultrasound diagnosis of fetal abdominal wall defects by 19 European registries. *Ultrasound Obstet Gynecol* 2001 October;18(4):309-16.
- (25) Chitty LS, Hunt GH, Moore J, Lobb MO. Effectiveness of routine ultrasonography in detecting fetal structural abnormalities in a low risk population. *BMJ* 1991 November 9;303(6811):1165-9.
- (26) Levi S, Schaaps JP, De HP et al. End-result of routine ultrasound screening for congenital anomalies: the Belgian Multicentric Study 1984-92. *Ultrasound Obstet Gynecol* 1995 June;5(6):366-71.
- (27) Richmond S, Atkins J. A population-based study of the prenatal diagnosis of congenital malformation over 16 years. *BJOG* 2005 October;112(10):1349-57.
- (28) Cristina MP, Ana G, Ines T et al. Perinatal results following the prenatal ultrasound diagnosis of single umbilical artery. *Acta Obstet Gynecol Scand* 2005 November;84(11):1068-74.
- (29) Stoll C, Alembik Y, Roth MP, Dott B. Study of 224 cases of oligohydramnios and congenital malformations in a series of 225,669 consecutive births. *Community Genet* 1998;1(2):71-7.
- (30) Lisi A, Botto LD, Rittler M et al. Sex and congenital malformations: an international perspective. *Am J Med Genet A* 2005 April 1;134(1):49-57.
- (31) Kuller JA, Katz VL, Wells SR et al. Cesarean delivery for fetal malformations. *Obstet Gynecol Surv* 1996 June;51(6):371-5.
- (32) Arizawa M, Imai S, Suehara N, Nakayama M. [Prenatal diagnosis of laryngeal atresia]. *Nippon Sanka Fujinka Gakkai Zasshi* 1989 July;41(7):907-10.
- (33) Onderoglu L, Saygan KB, Bulun A et al. Prenatal diagnosis of laryngeal atresia. *Prenat Diagn* 2003 April;23(4):277-80.
- (34) Kalache KD, Chaoui R, Tennstedt C, Bollmann R. Prenatal diagnosis of laryngeal atresia in two cases of congenital high airway obstruction syndrome (CHAOS). *Prenat Diagn* 1997 June;17(6):577-81.
- (35) Kassanos D, Christodoulou CN, Agapitos E et al. Prenatal ultrasonographic detection of the tracheal atresia sequence. *Ultrasound Obstet Gynecol* 1997 August;10(2):133-6.
- (36) Scott JN, Trevenen CL, Wiseman DA, Elliott PD. Tracheal atresia: ultrasonographic and pathologic correlation. *J Ultrasound Med* 1999 May;18(5):375-7.
- (37) Meizner I, Sherizly I, Mashiah R et al. Prenatal sonographic diagnosis of laryngeal atresia in association with single umbilical artery. *J Clin Ultrasound* 2000 October;28(8):435-8.

- (38) Achiron R, Hegesh J, Yagel S. Fetal lung lesions: a spectrum of disease. New classification based on pathogenesis, two-dimensional and color Doppler ultrasound. *Ultrasound Obstet Gynecol* 2004 August;24(2):107-14.
- (39) Calvert JK, Boyd PA, Chamberlain PC et al. Outcome of antenatally suspected congenital cystic adenomatoid malformation of the lung: 10 years' experience 1991-2001. *Arch Dis Child Fetal Neonatal Ed* 2005 August 30.
- (40) Monni G, Paladini D, Iriba RM et al. Prenatal ultrasound diagnosis of congenital cystic adenomatoid malformation of the lung: a report of 26 cases and review of the literature. *Ultrasound Obstet Gynecol* 2000 August;16(2):159-62.
- (41) Wilson RD, Hedrick HL, Liechty KW et al. Cystic adenomatoid malformation of the lung: Review of genetics, prenatal diagnosis, and in utero treatment. *Am J Med Genet A* 2006 January 15;140(2):151-5.
- (42) Adzick NS, Harrison MR, Glick PL et al. Fetal cystic adenomatoid malformation: prenatal diagnosis and natural history. *J Pediatr Surg* 1985 October;20(5):483-8.
- (43) Kahler C, Schulze E, Eichhorn KH, Seewald HJ. [Fetal hyperechogenic and cystic pulmonary masses: sonographic findings, antenatal management and outcome of 12 cases]. *Z Geburtshilfe Neonatol* 2002 November;206(6):205-10.
- (44) Gornall AS, Budd JL, Draper ES et al. Congenital cystic adenomatoid malformation: accuracy of prenatal diagnosis, prevalence and outcome in a general population. *Prenat Diagn* 2003 December 15;23(12):997-1002.
- (45) Bunduki V, Ruano R, da Silva MM et al. Prognostic factors associated with congenital cystic adenomatoid malformation of the lung. *Prenat Diagn* 2000 June;20(6):459-64.
- (46) Vijayaraghavan SB, Rao PS, Selvarasu CD, Rao TM. Prenatal sonographic features of intralobar bronchopulmonary sequestration. *J Ultrasound Med* 2003 May;22(5):541-4.
- (47) Frober R, Klemm P, Kahler C et al. [The extralobar sequestration of the fetal lung]. *Ann Anat* 2001 November;183(6):519-25.
- (48) Tsao K, Albanese CT, Harrison MR. Prenatal therapy for thoracic and mediastinal lesions. *World J Surg* 2003 January;27(1):77-83.
- (49) Ruano R, Benachi A, Aubry MC et al. Prenatal diagnosis of pulmonary sequestration using three-dimensional power Doppler ultrasound. *Ultrasound Obstet Gynecol* 2005 February;25(2):128-33.
- (50) Davenport M, Warne SA, Cacciaguerra S et al. Current outcome of antenally diagnosed cystic lung disease. *J Pediatr Surg* 2004 April;39(4):549-56.
- (51) Chen JS, Walford N, Yan YL et al. Foetal intralobar lung sequestration: antenatal diagnosis and management. *Singapore Med J* 2003 December;44(12):630-4.
- (52) Pumberger W, Moroder W, Wiesbauer P. Intraabdominal extralobar pulmonary sequestration exhibiting cystic adenomatoid malformation: prenatal diagnosis and characterization of a left suprarenal mass in the newborn. *Abdom Imaging* 2001 January;26(1):28-31.
- (53) Chandran H, Upadhyay V, Pease PW. Congenital cystic adenomatoid malformation and extralobar sequestration occurring independently in the ipsilateral hemithorax. *Pediatr Surg Int* 2000;16(1-2):102-3.
- (54) Becmeur F, Horta-Geraud P, Donato L, Sauvage P. Pulmonary sequestrations: prenatal ultrasound diagnosis, treatment, and outcome. *J Pediatr Surg* 1998 March;33(3):492-6.
- (55) Pumberger W, Hormann M, Deutinger J et al. Longitudinal observation of antenatally detected congenital lung malformations (CLM): natural history, clinical outcome and long-term follow-up. *Eur J Cardiothorac Surg* 2003 November;24(5):703-11.

- (56) Roggin KK, Breuer CK, Carr SR et al. The unpredictable character of congenital cystic lung lesions. *J Pediatr Surg* 2000 May;35(5):801-5.
- (57) Usui N, Kamata S, Sawai T et al. Outcome predictors for infants with cystic lung disease. *J Pediatr Surg* 2004 April;39(4):603-6.
- (58) Duncombe GJ, Dickinson JE, Kikios CS. Prenatal diagnosis and management of congenital cystic adenomatoid malformation of the lung. *Am J Obstet Gynecol* 2002 October;187(4):950-4.
- (59) Bagolan P, Nahom A, Giorlandino C et al. Cystic adenomatoid malformation of the lung: clinical evolution and management. *Eur J Pediatr* 1999 November;158(11):879-82.
- (60) Hedrick HL, Flake AW, Crombleholme TM et al. History of fetal diagnosis and therapy: Children's Hospital of Philadelphia experience. *Fetal Diagn Ther* 2003 March;18(2):65-82.
- (61) Lopoo JB, Goldstein RB, Lipshutz GS et al. Fetal pulmonary sequestration: a favorable congenital lung lesion. *Obstet Gynecol* 1999 October;94(4):567-71.
- (62) Crombleholme TM, Coleman B, Hedrick H et al. Cystic adenomatoid malformation volume ratio predicts outcome in prenatally diagnosed cystic adenomatoid malformation of the lung. *J Pediatr Surg* 2002 March;37(3):331-8.
- (63) Thorpe-Beeston JG, Nicolaides KH. Cystic adenomatoid malformation of the lung: prenatal diagnosis and outcome. *Prenat Diagn* 1994 August;14(8):677-88.
- (64) Harrison MR, Adzick NS, Jennings RW et al. Antenatal intervention for congenital cystic adenomatoid malformation. *Lancet* 1990 October 20;336(8721):965-7.
- (65) De SM, Masini L, Noia G et al. Congenital cystic adenomatoid malformation of the lung: antenatal ultrasound findings and fetal-neonatal outcome. Fifteen years of experience. *Fetal Diagn Ther* 2000 July;15(4):246-50.
- (66) Ishikawa S, Kamata S, Usui N et al. Ultrasonographic prediction of clinical pulmonary hypoplasia: measurement of the chest/trunk-length ratio in fetuses. *Pediatr Surg Int* 2003 May;19(3):172-5.
- (67) Bush A. Congenital lung disease: a plea for clear thinking and clear nomenclature. *Pediatr Pulmonol* 2001 October;32(4):328-37.
- (68) Heling KS, Tennstedt C, Chaoui R et al. Reliability of prenatal sonographic lung biometry in the diagnosis of pulmonary hypoplasia. *Prenat Diagn* 2001 August;21(8):649-57.
- (69) Wayne C, Cook K, Sairam S et al. Sensitivity and accuracy of routine antenatal ultrasound screening for isolated facial clefts. *Br J Radiol* 2002 July;75(895):584-9.
- (70) Berge SJ, Plath H, von Lindern JJ et al. Natural history of 70 fetuses with a prenatally diagnosed orofacial cleft. *Fetal Diagn Ther* 2002 July;17(4):247-51.
- (71) Sohan K, Freer M, Mercer N et al. Prenatal detection of facial clefts. *Fetal Diagn Ther* 2001 July;16(4):196-9.
- (72) Robinson JN, McElrath TF, Benson CB et al. Prenatal ultrasonography and the diagnosis of fetal cleft lip. *J Ultrasound Med* 2001 November;20(11):1165-70.
- (73) Johnson N, Sandy R. Prenatal diagnosis of cleft lip and palate. *Cleft Palate Craniofac J* 2003 March;40(2):186-9.
- (74) Clementi M, Tenconi R, Bianchi F, Stoll C. Evaluation of prenatal diagnosis of cleft lip with or without cleft palate and cleft palate by ultrasound: experience from 20 European registries. EUROSCAN study group. *Prenat Diagn* 2000 November;20(11):870-5.
- (75) Blumenfeld Z, Blumenfeld I, Bronshtein M. The early prenatal diagnosis of cleft lip and the decision-making process. *Cleft Palate Craniofac J* 1999 March;36(2):105-7.

- (76) Bronshtein M, Blumenfeld I, Kohn J, Blumenfeld Z. Detection of cleft lip by early second-trimester transvaginal sonography. *Obstet Gynecol* 1994 July;84(1):73-6.
- (77) Jones MC. Prenatal diagnosis of cleft lip and palate: detection rates, accuracy of ultrasonography, associated anomalies, and strategies for counseling. *Cleft Palate Craniofac J* 2002 March;39(2):169-73.
- (78) Nyberg DA, Sickler GK, Hegge FN et al. Fetal cleft lip with and without cleft palate: US classification and correlation with outcome. *Radiology* 1995 June;195(3):677-84.
- (79) Depaepe A, Dolk H, Lechat MF. The epidemiology of tracheo-oesophageal fistula and oesophageal atresia in Europe. EUROCAT Working Group. *Arch Dis Child* 1993 June;68(6):743-8.
- (80) Kluth D, Fiegel H. The embryology of the foregut. *Semin Pediatr Surg* 2003 February;12(1):3-9.
- (81) Pameijer CR, Hubbard AM, Coleman B, Flake AW. Combined pure esophageal atresia, duodenal atresia, biliary atresia, and pancreatic ductal atresia: prenatal diagnostic features and review of the literature. *J Pediatr Surg* 2000 May;35(5):745-7.
- (82) Langer JC, Hussain H, Khan A et al. Prenatal diagnosis of esophageal atresia using sonography and magnetic resonance imaging. *J Pediatr Surg* 2001 May;36(5):804-7.
- (83) Has R, Gunay S. Upper neck pouch sign in prenatal diagnosis of esophageal atresia. *Arch Gynecol Obstet* 2004 July;270(1):56-8.
- (84) Kalache KD, Chaoui R, Mau H, Bollmann R. The upper neck pouch sign: a prenatal sonographic marker for esophageal atresia. *Ultrasound Obstet Gynecol* 1998 February;11(2):138-40.
- (85) Shulman A, Mazkereth R, Zalel Y et al. Prenatal identification of esophageal atresia: the role of ultrasonography for evaluation of functional anatomy. *Prenat Diagn* 2002 August;22(8):669-74.
- (86) Satoh S, Takashima T, Takeuchi H et al. Antenatal sonographic detection of the proximal esophageal segment: specific evidence for congenital esophageal atresia. *J Clin Ultrasound* 1995 September;23(7):419-23.
- (87) D'Elia A, Pighetti M, Nappi C. Prenatal ultrasonographic appearance of esophageal atresia. *Eur J Obstet Gynecol Reprod Biol* 2002 October 10;105(1):77-8.
- (88) Eyheremendy E, Pfister M. Antenatal real-time diagnosis of esophageal atresias. *J Clin Ultrasound* 1983 September;11(7):395-7.
- (89) Stringer MD, McKenna KM, Goldstein RB et al. Prenatal diagnosis of esophageal atresia. *J Pediatr Surg* 1995 September;30(9):1258-63.
- (90) Sparey C, Jawaheer G, Barrett AM, Robson SC. Esophageal atresia in the Northern Region Congenital Anomaly Survey, 1985-1997: prenatal diagnosis and outcome. *Am J Obstet Gynecol* 2000 February;182(2):427-31.
- (91) Barnewolt CE. Congenital abnormalities of the gastrointestinal tract. *Semin Roentgenol* 2004 April;39(2):263-81.
- (92) Corteville JE, Gray DL, Langer JC. Bowel abnormalities in the fetus--correlation of prenatal ultrasonographic findings with outcome. *Am J Obstet Gynecol* 1996 September;175(3 Pt 1):724-9.
- (93) Haeusler MC, Berghold A, Stoll C et al. Prenatal ultrasonographic detection of gastrointestinal obstruction: results from 18 European congenital anomaly registries. *Prenat Diagn* 2002 July;22(7):616-23.
- (94) Skupski DW. Prenatal diagnosis of gastrointestinal anomalies with ultrasound. What have we learned? *Ann N Y Acad Sci* 1998 June 18;847:53-8.
- (95) Veyrac C, Couture A, Saguintah M, Baud C. [MRI of fetal GI tract]. *J Gynecol Obstet Biol Reprod (Paris)* 2005 February;34(1 Suppl):S14-S17.

- (96) Romero R, Ghidini A, Costigan K et al. Prenatal diagnosis of duodenal atresia: does it make any difference? *Obstet Gynecol* 1988 May;71(5):739-41.
- (97) Escobar MA, Ladd AP, Grosfeld JL et al. Duodenal atresia and stenosis: long-term follow-up over 30 years. *J Pediatr Surg* 2004 June;39(6):867-71.
- (98) Lawrence MJ, Ford WD, Furness ME et al. Congenital duodenal obstruction: early antenatal ultrasound diagnosis. *Pediatr Surg Int* 2000;16(5-6):342-5.
- (99) Bittencourt DG, Barini R, Marba S, Sbragia L. Congenital duodenal obstruction: does prenatal diagnosis improve the outcome? *Pediatr Surg Int* 2004 August;20(8):582-5.
- (100) Weiss H, Sherer DM, Manning FA. Ultrasonography of fetal annular pancreas. *Obstet Gynecol* 1999 November;94(5 Pt 2):852.
- (101) Berrocal T, Lamas M, Gutierrez J et al. Congenital anomalies of the small intestine, colon, and rectum. *Radiographics* 1999 September;19(5):1219-36.
- (102) Lam YH, Shek T, Tang MH. Sonographic features of anal atresia at 12 weeks. *Ultrasound Obstet Gynecol* 2002 May;19(5):523-4.
- (103) Redkar R, Davenport M, Howard ER. Antenatal diagnosis of congenital anomalies of the biliary tract. *J Pediatr Surg* 1998 May;33(5):700-4.
- (104) Rousseau T, Couvreur S, Senet-Lacombe E et al. Prenatal diagnosis of enteric duplication cyst of the tongue. *Prenat Diagn* 2004 February;24(2):98-100.
- (105) Richards DS, Langham MR, Anderson CD. The prenatal sonographic appearance of enteric duplication cysts. *Ultrasound Obstet Gynecol* 1996 January;7(1):17-20.
- (106) Dewbury KC, Aluwihare AP, Birch SJ, Freeman NV. Prenatal ultrasound demonstration of a choledochal cyst. *Br J Radiol* 1980 September;53(633):906-7.
- (107) Lee HJ, Lee SM, Park WH, Choi SO. Objective criteria of triangular cord sign in biliary atresia on US scans. *Radiology* 2003 November;229(2):395-400.
- (108) Bronshtein M, Blazer S. Prenatal diagnosis of liver calcifications. *Obstet Gynecol* 1995 November;86(5):739-43.
- (109) Cohen SB, Lipitz S, Mashiach S et al. In utero ultrasonographic diagnosis of an aberrant umbilical vein associated with fetal hepatic hyperechogenicity. *Prenat Diagn* 1997 October;17(10):978-82.
- (110) Simchen MJ, Toi A, Bona M et al. Fetal hepatic calcifications: prenatal diagnosis and outcome. *Am J Obstet Gynecol* 2002 December;187(6):1617-22.
- (111) Avni EF, Rypens F, Donner C et al. Hepatic cysts and hyperechogenicities: perinatal assessment and unifying theory on their origin. *Pediatr Radiol* 1994;24(8):569-72.
- (112) Stein B, Bromley B, Michlewitz H et al. Fetal liver calcifications: sonographic appearance and postnatal outcome. *Radiology* 1995 November;197(2):489-92.
- (113) Achiron R, Seidman DS, Afek A et al. Prenatal ultrasonographic diagnosis of fetal hepatic hyperechogenicities: clinical significance and implications for management. *Ultrasound Obstet Gynecol* 1996 April;7(4):251-5.
- (114) Koopman E, Wladimiroff JW. Fetal intrahepatic hyperechogenic foci: prenatal ultrasound diagnosis and outcome. *Prenat Diagn* 1998 April;18(4):339-42.
- (115) Blumenthal DH, Rushovich AM, Williams RK, Rochester D. Prenatal sonographic findings of meconium peritonitis with pathologic correlation. *J Clin Ultrasound* 1982 September;10(7):350-2.

- (116) Chan KL, Tang MH, Tse HY et al. Meconium peritonitis: prenatal diagnosis, postnatal management and outcome. *Prenat Diagn* 2005 August;25(8):676-82.
- (117) Chalubinski K, Deutinger J, Bernaschek G. Meconium peritonitis: extrusion of meconium and different sonographical appearances in relation to the stage of the disease. *Prenat Diagn* 1992 August;12(8):631-6.
- (118) Shyu MK, Shih JC, Lee CN et al. Correlation of prenatal ultrasound and postnatal outcome in meconium peritonitis. *Fetal Diagn Ther* 2003 July;18(4):255-61.
- (119) Konje JC, de CR, MacFadyen U, Taylor DJ. Antenatal diagnosis and management of meconium peritonitis: a case report and review of the literature. *Ultrasound Obstet Gynecol* 1995 July;6(1):66-9.
- (120) Dirkes K, Crombleholme TM, Craig SD et al. The natural history of meconium peritonitis diagnosed in utero. *J Pediatr Surg* 1995 July;30(7):979-82.
- (121) Kamata S, Nose K, Ishikawa S et al. Meconium peritonitis in utero. *Pediatr Surg Int* 2000;16(5-6):377-9.
- (122) Eckoldt F, Heling KS, Woderich R et al. Meconium peritonitis and pseudo-cyst formation: prenatal diagnosis and post-natal course. *Prenat Diagn* 2003 November;23(11):904-8.
- (123) Tseng JJ, Chou MM, Ho ES. Meconium peritonitis in utero: prenatal sonographic findings and clinical implications. *J Chin Med Assoc* 2003 June;66(6):355-9.
- (124) Lorimer WS, Jr., Ellis DG. Meconium peritonitis. *Surgery* 1966 August;60(2):470-5.
- (125) Foster MA, Nyberg DA, Mahony BS et al. Meconium peritonitis: prenatal sonographic findings and their clinical significance. *Radiology* 1987 December;165(3):661-5.
- (126) PAYNE RM, NIELSEN AM. Meconium peritonitis. *Am Surg* 1962 April;28:224-31.
- (127) Tibboel D, Gaillard JL, Molenaar JC. The importance of mesenteric vascular insufficiency in meconium peritonitis. *Hum Pathol* 1986 April;17(4):411-6.
- (128) Fordham LA. Congenital abnormalities of the musculoskeletal system: perinatal evaluation and long-term outcome. *Semin Roentgenol* 2004 April;39(2):304-22.
- (129) Kevern L, Warwick D, Wellesley D et al. Prenatal ultrasound: detection and diagnosis of limb abnormalities. *J Pediatr Orthop* 2003 March;23(2):251-3.
- (130) Lapaire O, Schiesser M, Peukert R et al. Split hand and foot malformation: ultrasound detection in the first trimester. *Ultrasound Obstet Gynecol* 2002 November;20(5):511-2.
- (131) Blaicher W, Mittermayer C, Messerschmidt A et al. Fetal skeletal deformities - the diagnostic accuracy of prenatal ultrasonography and fetal magnetic resonance imaging. *Ultraschall Med* 2004 June;25(3):195-9.
- (132) Edmonds LD, James LM. Temporal trends in the prevalence of congenital malformations at birth based on the birth defects monitoring program, United States, 1979-1987. *MMWR CDC Surveill Summ* 1990 December;39(4):19-23.
- (133) Castilla EE, Lugarinho R, da Graca DM, Salgado LJ. Associated anomalies in individuals with polydactyly. *Am J Med Genet* 1998 December 28;80(5):459-65.
- (134) Ryu JK, Cho JY, Choi JS. Prenatal sonographic diagnosis of focal musculoskeletal anomalies. *Korean J Radiol* 2003 October;4(4):243-51.
- (135) Froster-Iskenius UG, Baird PA. Limb reduction defects in over one million consecutive livebirths. *Teratology* 1989 February;39(2):127-35.
- (136) Stoll C, Wiesel A, Queisser-Luft A et al. Evaluation of the prenatal diagnosis of limb reduction deficiencies. *EUROSCAN Study Group. Prenat Diagn* 2000 October;20(10):811-8.

- (137) Stoll C, Alembik Y, Dott B, Roth MP. Evaluation of prenatal diagnosis of limb reduction defects by a registry of congenital anomalies. *Prenat Diagn* 1994 September;14(9):781-6.
- (138) Malone FD, Marino T, Bianchi DW et al. Isolated clubfoot diagnosed prenatally: is karyotyping indicated? *Obstet Gynecol* 2000 March;95(3):437-40.
- (139) Keret D, Ezra E, Lokiec F et al. Efficacy of prenatal ultrasonography in confirmed club foot. *J Bone Joint Surg Br* 2002 September;84(7):1015-9.
- (140) Shipp TD, Benacerraf BR. The significance of prenatally identified isolated clubfoot: is amniocentesis indicated? *Am J Obstet Gynecol* 1998 March;178(3):600-2.
- (141) Barry M. Prenatal assessment of foot deformity. *Early Hum Dev* 2005 October;81(10):793-6.
- (142) Treadwell MC, Stanitski CL, King M. Prenatal sonographic diagnosis of clubfoot: implications for patient counseling. *J Pediatr Orthop* 1999 January;19(1):8-10.
- (143) Bar-On E, Mashiach R, Inbar O et al. Prenatal ultrasound diagnosis of club foot: outcome and recommendations for counselling and follow-up. *J Bone Joint Surg Br* 2005 July;87(7):990-3.
- (144) Sobel E, Giorgini RJ, Michel R, Cohen SI. The natural history and longitudinal study of the surgically corrected clubfoot. *J Foot Ankle Surg* 2000 September;39(5):305-20.
- (145) Tillett RL, Fisk NM, Murphy K, Hunt DM. Clinical outcome of congenital talipes equinovarus diagnosed antenatally by ultrasound. *J Bone Joint Surg Br* 2000 August;82(6):876-80.
- (146) Jeanty P, Romero R, d'Alton M et al. In utero sonographic detection of hand and foot deformities. *J Ultrasound Med* 1985 November;4(11):595-601.
- (147) Bakalis S, Sairam S, Homfray T et al. Outcome of antenatally diagnosed talipes equinovarus in an unselected obstetric population. *Ultrasound Obstet Gynecol* 2002 September;20(3):226-9.
- (148) Pagnotta G, Maffulli N, Aureli S et al. Antenatal sonographic diagnosis of clubfoot: a six-year experience. *J Foot Ankle Surg* 1996 January;35(1):67-71.
- (149) Katz K, Meizner I, Mashiach R, Soudry M. The contribution of prenatal sonographic diagnosis of clubfoot to preventive medicine. *J Pediatr Orthop* 1999 January;19(1):5-7.
- (150) Woodrow N, Tran T, Umstad M et al. Mid-trimester ultrasound diagnosis of isolated talipes equinovarus: accuracy and outcome for infants. *Aust N Z J Obstet Gynaecol* 1998 August;38(3):301-5.
- (151) Parilla BV, Leeth EA, Kambich MP et al. Antenatal detection of skeletal dysplasias. *J Ultrasound Med* 2003 March;22(3):255-8.
- (152) Doray B, Favre R, Viville B et al. Prenatal sonographic diagnosis of skeletal dysplasias. A report of 47 cases. *Ann Genet* 2000 July;43(3-4):163-9.
- (153) Dudkiewicz I, Achiron R, Ganel A. Prenatal diagnosis of distal arthrogryposis type 1. *Skeletal Radiol* 1999 April;28(4):233-5.
- (154) Au-Yeung JY, Chan KL. Prenatal surgery for congenital diaphragmatic hernia. *Asian J Surg* 2003 October;26(4):240-3.
- (155) Bedoyan JK, Blackwell SC, Treadwell MC et al. Congenital diaphragmatic hernia: associated anomalies and antenatal diagnosis. Outcome-related variables at two Detroit hospitals. *Pediatr Surg Int* 2004 March;20(3):170-6.
- (156) Cannon C, Dildy GA, Ward R et al. A population-based study of congenital diaphragmatic hernia in Utah: 1988-1994. *Obstet Gynecol* 1996 June;87(6):959-63.

- (157) Deprest J, Jani J, Gratacos E et al. Fetal intervention for congenital diaphragmatic hernia: the European experience. *Semin Perinatol* 2005 April;29(2):94-103.
- (158) Dillon E, Renwick M, Wright C. Congenital diaphragmatic herniation: antenatal detection and outcome. *Br J Radiol* 2000 April;73(868):360-5.
- (159) Enns GM, Cox VA, Goldstein RB et al. Congenital diaphragmatic defects and associated syndromes, malformations, and chromosome anomalies: a retrospective study of 60 patients and literature review. *Am J Med Genet* 1998 September 23;79(3):215-25.
- (160) Graham G, Devine PC. Antenatal diagnosis of congenital diaphragmatic hernia. *Semin Perinatol* 2005 April;29(2):69-76.
- (161) Harrison MR, Adzick NS, Estes JM, Howell LJ. A prospective study of the outcome for fetuses with diaphragmatic hernia. *JAMA* 1994 February 2;271(5):382-4.
- (162) Hubbard AM, Crombleholme TM, Adzick NS et al. Prenatal MRI evaluation of congenital diaphragmatic hernia. *Am J Perinatol* 1999;16(8):407-13.
- (163) Rottier R, Tibboel D. Fetal lung and diaphragm development in congenital diaphragmatic hernia. *Semin Perinatol* 2005 April;29(2):86-93.
- (164) Ruano R, Benachi A, Joubin L et al. Three-dimensional ultrasonographic assessment of fetal lung volume as prognostic factor in isolated congenital diaphragmatic hernia. *BJOG* 2004 May;111(5):423-9.
- (165) Ruano R, Joubin L, Sonigo P et al. Fetal lung volume estimated by 3-dimensional ultrasonography and magnetic resonance imaging in cases with isolated congenital diaphragmatic hernia. *J Ultrasound Med* 2004 March;23(3):353-8.
- (166) Skari H, Bjornland K, Haugen G et al. Congenital diaphragmatic hernia: a meta-analysis of mortality factors. *J Pediatr Surg* 2000 August;35(8):1187-97.
- (167) Varlet F, Bousquet F, Clemenson A et al. Congenital diaphragmatic hernia. Two cases with early prenatal diagnosis and increased nuchal translucency. *Fetal Diagn Ther* 2003 January;18(1):33-5.
- (168) Harrison MR, Bjordal RI, Langmark F, Knutrud O. Congenital diaphragmatic hernia: the hidden mortality. *J Pediatr Surg* 1978 June;13(3):227-30.
- (169) Gallot D, Coste K, Francannet C et al. Antenatal detection and impact on outcome of congenital diaphragmatic hernia: A 12-year experience in Auvergne (France). *Eur J Obstet Gynecol Reprod Biol* 2005 August 11.
- (170) Harrison MR, Adzick NS, Bullard KM et al. Correction of congenital diaphragmatic hernia in utero VII: a prospective trial. *J Pediatr Surg* 1997 November;32(11):1637-42.
- (171) Stevens TP, Chess PR, McConnochie KM et al. Survival in early- and late-term infants with congenital diaphragmatic hernia treated with extracorporeal membrane oxygenation. *Pediatrics* 2002 September;110(3):590-6.
- (172) Bohn D. Congenital diaphragmatic hernia. *Am J Respir Crit Care Med* 2002 October 1;166(7):911-5.
- (173) Tonks A, Wyldes M, Somerset DA et al. Congenital malformations of the diaphragm: findings of the West Midlands Congenital Anomaly Register 1995 to 2000. *Prenat Diagn* 2004 August;24(8):596-604.
- (174) Garne E, Haeusler M, Barisic I et al. Congenital diaphragmatic hernia: evaluation of prenatal diagnosis in 20 European regions. *Ultrasound Obstet Gynecol* 2002 April;19(4):329-33.
- (175) Bollmann R, Kalache K, Mau H et al. Associated malformations and chromosomal defects in congenital diaphragmatic hernia. *Fetal Diagn Ther* 1995 January;10(1):52-9.

- (176) Metkus AP, Filly RA, Stringer MD et al. Sonographic predictors of survival in fetal diaphragmatic hernia. *J Pediatr Surg* 1996 January;31(1):148-51.
- (177) Hedrick HL, Crombleholme TM, Flake AW et al. Right congenital diaphragmatic hernia: Prenatal assessment and outcome. *J Pediatr Surg* 2004 March;39(3):319-23.
- (178) Walsh DS, Hubbard AM, Olutoye OO et al. Assessment of fetal lung volumes and liver herniation with magnetic resonance imaging in congenital diaphragmatic hernia. *Am J Obstet Gynecol* 2000 November;183(5):1067-9.
- (179) Puri P, Wester T. Historical aspects of congenital diaphragmatic hernia. *Pediatr Surg Int* 1997 March 21;12(2/3):95-100.
- (180) Geary MP, Chitty LS, Morrison JJ et al. Perinatal outcome and prognostic factors in prenatally diagnosed congenital diaphragmatic hernia. *Ultrasound Obstet Gynecol* 1998 August;12(2):107-11.
- (181) Hatch EI, Jr., Kendall J, Blumhagen J. Stomach position as an in utero predictor of neonatal outcome in left-sided diaphragmatic hernia. *J Pediatr Surg* 1992 June;27(6):778-9.
- (182) Heling KS, Wauer RR, Hammer H et al. Reliability of the lung-to-head ratio in predicting outcome and neonatal ventilation parameters in fetuses with congenital diaphragmatic hernia. *Ultrasound Obstet Gynecol* 2005 February;25(2):112-8.
- (183) Lipshutz GS, Albanese CT, Feldstein VA et al. Prospective analysis of lung-to-head ratio predicts survival for patients with prenatally diagnosed congenital diaphragmatic hernia. *J Pediatr Surg* 1997 November;32(11):1634-6.
- (184) Okazaki T, Kohno S, Hasegawa S et al. Congenital diaphragmatic hernia: efficacy of ultrasound examination in its management. *Pediatr Surg Int* 2003 May;19(3):176-9.
- (185) CARTER RE, WATERSTON DJ, ABERDEEN E. Hernia and eventration of the diaphragm in childhood. *Lancet* 1962 March 31;1:656-9.
- (186) Cass DL. Fetal surgery for congenital diaphragmatic hernia: the North American experience. *Semin Perinatol* 2005 April;29(2):104-11.
- (187) Vettraino IM, Lee W, Comstock CH. The evolving appearance of a congenital diaphragmatic hernia. *J Ultrasound Med* 2002 January;21(1):85-9.
- (188) Albanese CT, Lopoo J, Goldstein RB et al. Fetal liver position and perinatal outcome for congenital diaphragmatic hernia. *Prenat Diagn* 1998 November;18(11):1138-42.
- (189) Deprest J, Jani J, Van SD et al. Current consequences of prenatal diagnosis of congenital diaphragmatic hernia. *J Pediatr Surg* 2006 February;41(2):423-30.
- (190) Stoll C, Alembik Y, Dott B, Roth MP. Risk factors in congenital abdominal wall defects (omphalocele and gastroschisis): a study in a series of 265,858 consecutive births. *Ann Genet* 2001 October;44(4):201-8.
- (191) Weber TR, Au-Fliegner M, Downard CD, Fishman SJ. Abdominal wall defects. *Curr Opin Pediatr* 2002 August;14(4):491-7.
- (192) Davenport M, Haugen S, Greenough A, Nicolaides K. Closed gastroschisis: Antenatal and postnatal features. *J Pediatr Surg* 2001 December;36(12):1834-7.
- (193) Japaraj RP, Hockey R, Chan FY. Gastroschisis: can prenatal sonography predict neonatal outcome? *Ultrasound Obstet Gynecol* 2003 April;21(4):329-33.
- (194) Kilby MD, Lander A, Usher-Somers M. Exomphalos (omphalocele). *Prenat Diagn* 1998 December;18(12):1283-8.

- (195) Rankin J, Dillon E, Wright C. Congenital anterior abdominal wall defects in the north of England, 1986-1996: occurrence and outcome. *Prenat Diagn* 1999 July;19(7):662-8.
- (196) Reid KP, Dickinson JE, Doherty DA. The epidemiologic incidence of congenital gastroschisis in Western Australia. *Am J Obstet Gynecol* 2003 September;189(3):764-8.
- (197) Salvesen KA. Fetal abdominal wall defects--easy to diagnose--and then what? *Ultrasound Obstet Gynecol* 2001 October;18(4):301-4.
- (198) Saxena AK, Hulskamp G, Schleef J et al. Gastroschisis: a 15-year, single-center experience. *Pediatr Surg Int* 2002 September;18(5-6):420-4.
- (199) Biard JM, Wilson RD, Johnson MP et al. Prenatally diagnosed giant omphaloceles: short- and long-term outcomes. *Prenat Diagn* 2004 June;24(6):434-9.
- (200) Huang J, Kurkchubasche AG, Carr SR et al. Benefits of term delivery in infants with antenatally diagnosed gastroschisis. *Obstet Gynecol* 2002 October;100(4):695-9.
- (201) Axt R, Quijano F, Boos R et al. Omphalocele and gastroschisis: prenatal diagnosis and peripartal management. A case analysis of the years 1989-1997 at the Department of Obstetrics and Gynecology, University of Homburg/Saar. *Eur J Obstet Gynecol Reprod Biol* 1999 November;87(1):47-54.
- (202) Holland AJ, Ford WD, Linke RJ et al. Influence of antenatal ultrasound on the management of fetal exomphalos. *Fetal Diagn Ther* 1999 July;14(4):223-8.
- (203) Weir E. Congenital abdominal wall defects. *CMAJ* 2003 October 14;169(8):809-10.
- (204) Kalache KD, Bierlich A, Hammer H, Bollmann R. Is unexplained third trimester intrauterine death of fetuses with gastroschisis caused by umbilical cord compression due to acute extra-abdominal bowel dilatation? *Prenat Diagn* 2002 August;22(8):715-7.
- (205) Durfee SM, Downard CD, Benson CB, Wilson JM. Postnatal outcome of fetuses with the prenatal diagnosis of gastroschisis. *J Ultrasound Med* 2002 March;21(3):269-74.
- (206) Tawil A, Comstock CH, Chang Ch CH. Prenatal closure of abdominal defect in gastroschisis: case report and review of the literature. *Pediatr Dev Pathol* 2001 November;4(6):580-4.
- (207) Kaiser MM, Kahl F, von SC, Halsband H. [Omphalocele and gastroschisis. Outcome--complications--follow-up--quality of life]. *Chirurg* 2000 October;71(10):1256-62.
- (208) ina-Mumuney AJ, Fischer AC, Blakemore KJ et al. A dilated fetal stomach predicts a complicated postnatal course in cases of prenatally diagnosed gastroschisis. *Am J Obstet Gynecol* 2004 May;190(5):1326-30.
- (209) Pelizzo G, Maso G, Dell'oste C et al. Giant omphaloceles with a small abdominal defect: prenatal diagnosis and neonatal management. *Ultrasound Obstet Gynecol* 2005 December;26(7):786-8.
- (210) Moir CR, Ramsey PS, Ogburn PL et al. A prospective trial of elective preterm delivery for fetal gastroschisis. *Am J Perinatol* 2004 July;21(5):289-94.
- (211) Hausler MC, Fasching G, Zotter H et al. [Effect of the mode of delivery on the newborn infant with gastroschisis. With data from the Styrian Malformation Register]. *Gynakol Geburtshilfliche Rundsch* 1997;37(1):25-9.
- (212) Molik KA, Gingalewski CA, West KW et al. Gastroschisis: a plea for risk categorization. *J Pediatr Surg* 2001 January;36(1):51-5.
- (213) Stoll C, Clementi M. Prenatal diagnosis of dysmorphic syndromes by routine fetal ultrasound examination across Europe. *Ultrasound Obstet Gynecol* 2003 June;21(6):543-51.
- (214) Batukan C, Schwabe M, Heling KS et al. [Prenatal diagnosis of right atrial isomerism (asplenia-syndrome): case report]. *Ultraschall Med* 2005 June;26(3):234-8.

- (215) IVEMARK BI. Implications of agenesis of the spleen on the pathogenesis of conotruncus anomalies in childhood; an analysis of the heart malformations in the splenic agenesis syndrome, with fourteen new cases. *Acta Paediatr* 1955 November;44(Suppl 104):7-110.
- (216) Britz-Cunningham SH, Shah MM, Zuppan CW, Fletcher WH. Mutations of the Connexin43 gap-junction gene in patients with heart malformations and defects of laterality. *N Engl J Med* 1995 May 18;332(20):1323-9.
- (217) Noack F, Sayk F, Ressel A et al. Ivemark syndrome with agenesis of the corpus callosum: a case report with a review of the literature. *Prenat Diagn* 2002 November;22(11):1011-5.
- (218) Stroustrup SA, Grable I, Levine D. Case 66: caudal regression syndrome in the fetus of a diabetic mother. *Radiology* 2004 January;230(1):229-33.
- (219) Pena SD, Shokeir MH. Autosomal recessive cerebro-oculo-facio-skeletal (COFS) syndrome. *Clin Genet* 1974;5(4):285-93.
- (220) Tongsong T, Chanprapaph P, Khunamornpong S. Prenatal ultrasound of regional akinesia with Pena-Shokier phenotype. *Prenat Diagn* 2000 May;20(5):422-5.
- (221) Ajayi RA, Keen CE, Knott PD. Ultrasound diagnosis of the Pena Shokeir phenotype at 14 weeks of pregnancy. *Prenat Diagn* 1995 August;15(8):762-4.
- (222) Horrow MM, Rosenberg HK, Schneider AS et al. US case of the day. Pena-Shokeir phenotype. *Radiographics* 1995 May;15(3):726-9.
- (223) Hall JG. Analysis of Pena Shokeir phenotype. *Am J Med Genet* 1986 September;25(1):99-117.
- (224) Fujishiro E, Suzuki Y, Sato T et al. Characteristic findings for diagnosis of baby complicated with both the VACTERL association and duodenal atresia. *Fetal Diagn Ther* 2004 March;19(2):134-7.
- (225) Fordham LA. Congenital abnormalities of the musculoskeletal system: perinatal evaluation and long-term outcome. *Semin Roentgenol* 2004 April;39(2):304-22.
- (226) Botto LD, Khouri MJ, Mastroiacovo P et al. The spectrum of congenital anomalies of the VATER association: an international study. *Am J Med Genet* 1997 July 11;71(1):8-15.
- (227) Miller OF, Kolon TF. Prenatal diagnosis of VACTERL association. *J Urol* 2001 December;166(6):2389-91.
- (228) Teoh M, Meagher S. First-trimester diagnosis of micrognathia as a presentation of Pierre Robin syndrome. *Ultrasound Obstet Gynecol* 2003 June;21(6):616-8.
- (229) Rotten D, Levaillant JM, Martinez H et al. The fetal mandible: a 2D and 3D sonographic approach to the diagnosis of retrognathia and micrognathia. *Ultrasound Obstet Gynecol* 2002 February;19(2):122-30.
- (230) Hsieh YY, Chang CC, Tsai HD et al. The prenatal diagnosis of Pierre-Robin sequence. *Prenat Diagn* 1999 June;19(6):567-9.
- (231) Kuo JS, Casey SO, Thompson L, Truwit CL. Pallister-Hall syndrome: clinical and MR features. *AJNR Am J Neuroradiol* 1999 November;20(10):1839-41.
- (232) Roscioli T, Kennedy D, Cui J et al. Pallister-Hall syndrome: unreported skeletal features of a GLI3 mutation. *Am J Med Genet A* 2005 August 1;136(4):390-4.
- (233) Ars E, Kruyer H, Gaona A et al. Prenatal diagnosis of sporadic neurofibromatosis type 1 (NF1) by RNA and DNA analysis of a splicing mutation. *Prenat Diagn* 1999 August;19(8):739-42.
- (234) De VC, Baena N, Cariati E et al. Contribution of ultrasonographic examination to the prenatal detection of chromosomal abnormalities in 19 centres across Europe. *Ann Genet* 2001 October;44(4):209-17.
- (235) Estroff JA. Prenatal diagnosis and imaging of genetic syndromes. *Semin Roentgenol* 2004 April;39(2):323-35.

- (236) Whittle M. Ultrasonographic "soft markers" of fetal chromosomal defects. *BMJ* 1997 March 29;314(7085):918.
- (237) Picklesimer AH, Moise KJ, Jr., Wolfe HM. The impact of gestational age on the sonographic detection of aneuploidy. *Am J Obstet Gynecol* 2005 September;193(3 Pt 2):1243-7.
- (238) Nyberg DA, Luthy DA, Resta RG et al. Age-adjusted ultrasound risk assessment for fetal Down's syndrome during the second trimester: description of the method and analysis of 142 cases. *Ultrasound Obstet Gynecol* 1998 July;12(1):8-14.
- (239) Egan JF, Benn P, Borgida AF et al. Efficacy of screening for fetal Down syndrome in the United States from 1974 to 1997. *Obstet Gynecol* 2000 December;96(6):979-85.
- (240) Chitty LS. Antenatal screening for aneuploidy. *Curr Opin Obstet Gynecol* 1998 April;10(2):91-6.
- (241) Krampl E. [Screening of chromosome anomalies during the first trimester]. *Gynakol Geburtshilfliche Rundsch* 2005 April;45(2):86-92.
- (242) Nicolaides KH, Spencer K, Avgidou K et al. Multicenter study of first-trimester screening for trisomy 21 in 75 821 pregnancies: results and estimation of the potential impact of individual risk-orientated two-stage first-trimester screening. *Ultrasound Obstet Gynecol* 2005 March;25(3):221-6.
- (243) Avgidou K, Papageorghiou A, Bindra R et al. Prospective first-trimester screening for trisomy 21 in 30,564 pregnancies. *Am J Obstet Gynecol* 2005 June;192(6):1761-7.
- (244) Lee W, DeVore GR, Comstock CH et al. Nasal bone evaluation in fetuses with Down syndrome during the second and third trimesters of pregnancy. *J Ultrasound Med* 2003 January;22(1):55-60.
- (245) Wald NJ, Cuckle HS, Densem JW et al. Maternal serum screening for Down's syndrome in early pregnancy. *BMJ* 1988 October 8;297(6653):883-7.
- (246) Wald NJ, Hackshaw AK. Combining ultrasound and biochemistry in first-trimester screening for Down's syndrome. *Prenat Diagn* 1997 September;17(9):821-9.
- (247) Gasiorek-Wiens A, Tercanli S, Kozlowski P et al. Screening for trisomy 21 by fetal nuchal translucency and maternal age: a multicenter project in Germany, Austria and Switzerland. *Ultrasound Obstet Gynecol* 2001 December;18(6):645-8.
- (248) Zoppi MA, Ibba RM, Putzolu M et al. Assessment of risk for chromosomal abnormalities at 10-14 weeks of gestation by nuchal translucency and maternal age in 5,210 fetuses at a single centre. *Fetal Diagn Ther* 2000 May;15(3):170-3.
- (249) Thilaganathan B, Sairam S, Michailidis G, Wathen NC. First trimester nuchal translucency: effective routine screening for Down's syndrome. *Br J Radiol* 1999 October;72(862):946-8.
- (250) Schwarzler P, Carvalho JS, Senat MV et al. Screening for fetal aneuploidies and fetal cardiac abnormalities by nuchal translucency thickness measurement at 10-14 weeks of gestation as part of routine antenatal care in an unselected population. *Br J Obstet Gynaecol* 1999 October;106(10):1029-34.
- (251) Theodoropoulos P, Lolis D, Papageorgiou C et al. Evaluation of first-trimester screening by fetal nuchal translucency and maternal age. *Prenat Diagn* 1998 February;18(2):133-7.
- (252) Wellesley D, De VC, Baena N et al. Contribution of ultrasonographic examination to the prenatal detection of trisomy 21: experience from 19 European registers. *Ann Genet* 2004 October;47(4):373-80.
- (253) Grandjean H, Larroque D, Levi S. Detection of chromosomal abnormalities, an outcome of ultrasound screening. The Eurofetus Team. *Ann N Y Acad Sci* 1998 June 18;847:136-40.
- (254) Cicero S, Curcio P, Papageorghiou A et al. Absence of nasal bone in fetuses with trisomy 21 at 11-14 weeks of gestation: an observational study. *Lancet* 2001 November 17;358(9294):1665-7.

- (255) Snijders RJ, Platt LD, Greene N et al. Femur length and trisomy 21: impact of gestational age on screening efficiency. *Ultrasound Obstet Gynecol* 2000 August;16(2):142-5.
- (256) Grandjean H, Larroque D, Levi S. Detection of chromosomal abnormalities, an outcome of ultrasound screening. The Eurofetus Team. *Ann N Y Acad Sci* 1998 June 18;847:136-40.
- (257) Greenberg F. DiGeorge syndrome: an historical review of clinical and cytogenetic features. *J Med Genet* 1993 October;30(10):803-6.
- (258) Chaoui R, Kalache KD, Heling KS et al. Absent or hypoplastic thymus on ultrasound: a marker for deletion 22q11.2 in fetal cardiac defects. *Ultrasound Obstet Gynecol* 2002 December;20(6):546-52.
- (259) Glover TW. CATCHing a break on 22. *Nat Genet* 1995 July;10(3):257-8.
- (260) Baena N, De VC, Cariati E et al. Turner syndrome: evaluation of prenatal diagnosis in 19 European registries. *Am J Med Genet A* 2004 August 15;129(1):16-20.
- (261) Bronshtein M, Zimmer EZ, Blazer S. A characteristic cluster of fetal sonographic markers that are predictive of fetal Turner syndrome in early pregnancy. *Am J Obstet Gynecol* 2003 April;188(4):1016-20.
- (262) Surerus E, Huggon IC, Allan LD. Turner's syndrome in fetal life. *Ultrasound Obstet Gynecol* 2003 September;22(3):264-7.
- (263) Bojesen A, Juul S, Gravholt CH. Prenatal and postnatal prevalence of Klinefelter syndrome: a national registry study. *J Clin Endocrinol Metab* 2003 February;88(2):622-6.
- (264) Khoury-Collado F, Wehbeh AN, Fisher AJ et al. Prenatal diagnosis of 47,XXX. *Am J Obstet Gynecol* 2005 May;192(5):1469-71.
- (265) Moore SW, Satge D, Sasco AJ et al. The epidemiology of neonatal tumours. Report of an international working group. *Pediatr Surg Int* 2003 September;19(7):509-19.
- (266) Chen WY, Lin CN, Chao CS et al. Prenatal diagnosis of congenital mesoblastic nephroma in mid-second trimester by sonography and magnetic resonance imaging. *Prenat Diagn* 2003 November;23(11):927-31.
- (267) Rasidaki M, Sifakis S, Vardaki E, Koumantakis E. Prenatal diagnosis of a fetal chest wall cystic lymphangioma using ultrasonography and MRI: a case report with literature review. *Fetal Diagn Ther* 2005 November;20(6):504-7.
- (268) Gorincour G, Dugougeat-Pilleul F, Bouvier R et al. Prenatal presentation of cervical congenital neuroblastoma. *Prenat Diagn* 2003 August;23(8):690-3.
- (269) Heling KS, Chaoui R, Hartung J et al. Prenatal diagnosis of congenital neuroblastoma. Analysis of 4 cases and review of the literature. *Fetal Diagn Ther* 1999 January;14(1):47-52.
- (270) Acharya S, Jayabose S, Kogan SJ et al. Prenatally diagnosed neuroblastoma. *Cancer* 1997 July 15;80(2):304-10.
- (271) Fenart D, Deville A, Donzeau M, Bruneton JN. [Retroperitoneal neuroblastoma diagnosed in utero. Apropos of 1 case]. *J Radiol* 1983 May;64(5):359-61.
- (272) Saylors RL, III, Cohn SL, Morgan ER, Brodeur GM. Prenatal detection of neuroblastoma by fetal ultrasonography. *Am J Pediatr Hematol Oncol* 1994 November;16(4):356-60.
- (273) Kerbl R, Urban CE, Lackner H et al. Connatal localized neuroblastoma. The case to delay treatment. *Cancer* 1996 April 1;77(7):1395-401.
- (274) Goldstein I, Shoshani G, Ben-Harush E, Sujoy P. Prenatal diagnosis of congenital mesoblastic nephroma. *Ultrasound Obstet Gynecol* 2002 February;19(2):209-11.

- (275) Fuchs IB, Henrich W, Brauer M et al. Prenatal diagnosis of congenital mesoblastic nephroma in 2 siblings. *J Ultrasound Med* 2003 August;22(8):823-7.
- (276) Viora E, Grassi PP, Comoglio F et al. Ultrasonographic detection of fetal cranio-facial hemangioma: case report and review of the literature. *Ultrasound Obstet Gynecol* 2000 May;15(5):431-4.
- (277) Boon LM, Enjolras O, Mulliken JB. Congenital hemangioma: evidence of accelerated involution. *J Pediatr* 1996 March;128(3):329-35.
- (278) Shaub M, Wilson R, Collea J. Fetal cystic lymphangioma (cystic hygroma): prepartum ultrasonic findings. *Radiology* 1976 November;121(2):449-50.
- (279) Isaacs H, Jr. Perinatal (fetal and neonatal) germ cell tumors. *J Pediatr Surg* 2004 July;39(7):1003-13.
- (280) Olutoye OO, Johnson MP, Coleman BG et al. Abnormal umbilical cord Doppler sonograms may predict impending demise in fetuses with sacrococcygeal teratoma. A report of two cases. *Fetal Diagn Ther* 2004 January;19(1):35-9.
- (281) Bond SJ, Harrison MR, Schmidt KG et al. Death due to high-output cardiac failure in fetal sacrococcygeal teratoma. *J Pediatr Surg* 1990 December;25(12):1287-91.
- (282) Hedrick HL, Flake AW, Crombleholme TM et al. Sacrococcygeal teratoma: prenatal assessment, fetal intervention, and outcome. *J Pediatr Surg* 2004 March;39(3):430-8.