

6 Literaturverzeichnis

- Abremski, K., Hoess, R. und Sternberg, N.** (1983). Studies on the properties of P1 site-specific recombination: evidence for topologically unlinked products following recombination. *Cell* **32**, 1301-11.
- Achim, C. L., Katyal, S., Wiley, C. A., Shiratori, M., Wang, G., Oshika, E., Petersen, B. E., Li, J. M. und Michalopoulos, G. K.** (1997). Expression of HGF and cMet in the developing and adult brain. *Brain Res Dev Brain Res* **102**, 299-303.
- Arnheiter, H., Skuntz, S., Noteborn, M., Chang, S. und Meier, E.** (1990). Transgenic mice with intracellular immunity to influenza virus. *Cell* **62**, 51-61.
- Arnold, H. H. und Braun, T.** (2000). Genetics of muscle determination and development. *Curr Top Dev Biol* **48**, 129-64.
- Berger, S. L. und Kimmel, A. R.** (1987). Guide To Molecular Cloning Techniques. San Diego, CA 92101-4495, USA: Academic Press.
- Beviglia, L., Matsumoto, K., Lin, C. S., Ziober, B. L. und Kramer, R. H.** (1997). Expression of the c-Met/HGF receptor in human breast carcinoma: correlation with tumor progression. *Int J Cancer* **74**, 301-9.
- Birchmeier, C., Birchmeier, W., Gherardi, E. und Vande Woude, G. F.** (2003). Met, metastasis, motility and more. *Nat Rev Mol Cell Biol* **4**, 915-25.
- Birchmeier, C. und Brohmann, H.** (2000). Genes that control the development of migrating muscle precursor cells. *Curr Opin Cell Biol* **12**, 725-30.
- Birnboim, H. C. und Doly, J.** (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* **7**, 1513-23.
- Bladt, F., Riethmacher, D., Isenmann, S., Aguzzi, A. und Birchmeier, C.** (1995). Essential role for the c-met receptor in the migration of myogenic precursor cells into the limb bud. *Nature* **376**, 768-71.
- Bober, E., Franz, T., Arnold, H. H., Gruss, P. und Tremblay, P.** (1994). Pax-3 is required for the development of limb muscles: a possible role for the migration of dermomyotomal muscle progenitor cells. *Development* **120**, 603-12.
- Borycki, A. G. und Emerson, C. P., Jr.** (2000). Study of skeletal myogenesis in cultures of unsegmented paraxial mesoderm. *Methods Mol Biol* **137**, 351-7.
- Bottaro, D. P., Rubin, J. S., Faletto, D. L., Chan, A. M., Kmiecik, T. E., Vande Woude, G. F. und Aaronson, S. A.** (1991). Identification of the hepatocyte growth factor receptor as the c-met proto-oncogene product. *Science* **251**, 802-4.
- Bradford, M. M.** (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**, 248-54.

- Bradley, A. und Robertson, E.** (1986). Embryo-derived stem cells: a tool for elucidating the developmental genetics of the mouse. *Curr Top Dev Biol* **20**, 357-71.
- Branda, C. S. und Dymecki, S. M.** (2004). Talking about a revolution: The impact of site-specific recombinases on genetic analyses in mice. *Dev Cell* **6**, 7-28.
- Breathnach, R., Benoist, C., O'Hare, K., Gannon, F. und Chambon, P.** (1978). Ovalbumin gene: evidence for a leader sequence in mRNA and DNA sequences at the exon-intron boundaries. *Proc Natl Acad Sci U S A* **75**, 4853-7.
- Brohmann, H., Jagla, K. und Birchmeier, C.** (2000). The role of Lbx1 in migration of muscle precursor cells. *Development* **127**, 437-45.
- Buchholz, F., Angrand, P. O. und Stewart, A. F.** (1998). Improved properties of FLP recombinase evolved by cycling mutagenesis. *Nat Biotechnol* **16**, 657-62.
- Caton, A., Hacker, A., Naeem, A., Livet, J., Maina, F., Bladt, F., Klein, R., Birchmeier, C. und Guthrie, S.** (2000). The branchial arches and HGF are growth-promoting and chemoattractant for cranial motor axons. *Development* **127**, 1751-66.
- Choi, J., Costa, M. L., Mermelstein, C. S., Chagas, C., Holtzer, S. und Holtzer, H.** (1990). MyoD converts primary dermal fibroblasts, chondroblasts, smooth muscle, and retinal pigmented epithelial cells into striated mononucleated myoblasts and multinucleated myotubes. *Proc Natl Acad Sci U S A* **87**, 7988-92.
- Christ, B., Jacob, H. J. und Jacob, M.** (1977). Experimental analysis of the origin of the wing musculature in avian embryos. *Anat Embryol (Berl)* **150**, 171-86.
- Christ, B. und Ordahl, C. P.** (1995). Early stages of chick somite development. *Anat Embryol (Berl)* **191**, 381-96.
- Cinnamon, Y., Kahane, N. und Kalcheim, C.** (1999). Characterization of the early development of specific hypaxial muscles from the ventrolateral myotome. *Development* **126**, 4305-15.
- Cooper, C. S., Park, M., Blair, D. G., Tainsky, M. A., Huebner, K., Croce, C. M. und Vande Woude, G. F.** (1984). Molecular cloning of a new transforming gene from a chemically transformed human cell line. *Nature* **311**, 29-33.
- Danilkovitch-Miagkova, A. und Zbar, B.** (2002). Dysregulation of Met receptor tyrosine kinase activity in invasive tumors. *J Clin Invest* **109**, 863-7.
- Daston, G., Lamar, E., Olivier, M. und Goulding, M.** (1996). Pax-3 is necessary for migration but not differentiation of limb muscle precursors in the mouse. *Development* **122**, 1017-27.
- Denhardt, D. T.** (1966). A membrane-filter technique for the detection of complementary DNA. *Biochem Biophys Res Commun* **23**, 641-6.
- Dietrich, S., Abou-Rebyeh, F., Brohmann, H., Bladt, F., Sonnenberg-Riethmacher, E., Yamaai, T., Lumsden, A., Brand-Saberi, B. und Birchmeier, C.** (1999). The role of SF/HGF and c-Met in the development of skeletal muscle. *Development* **126**, 1621-9.

- Dietrich, S., Schubert, F. R., Healy, C., Sharpe, P. T. und Lumsden, A.** (1998). Specification of the hypaxial musculature. *Development* **125**, 2235-49.
- Donate, L. E., Gherardi, E., Srinivasan, N., Sowdhamini, R., Aparicio, S. und Blundell, T. L.** (1994). Molecular evolution and domain structure of plasminogen-related growth factors (HGF/SF and HGF1/MSP). *Protein Sci* **3**, 2378-94.
- Duprez, D.** (2002). Signals regulating muscle formation in the limb during embryonic development. *Int J Dev Biol* **46**, 915-25.
- Dymecki, S. M.** (1996a). Flp recombinase promotes site-specific DNA recombination in embryonic stem cells and transgenic mice. *Proc Natl Acad Sci U S A* **93**, 6191-6.
- Dymecki, S. M.** (1996b). A modular set of Flp, FRT and lacZ fusion vectors for manipulating genes by site-specific recombination. *Gene* **171**, 197-201.
- Dymecki, S. M. und Tomasiewicz, H.** (1998). Using Flp-recombinase to characterize expansion of Wnt1-expressing neural progenitors in the mouse. *Dev Biol* **201**, 57-65.
- Ebens, A., Brose, K., Leonardo, E. D., Hanson, M. G., Jr., Bladt, F., Birchmeier, C., Barres, B. A. und Tessier-Lavigne, M.** (1996). Hepatocyte growth factor/scatter factor is an axonal chemoattractant and a neurotrophic factor for spinal motor neurons. *Neuron* **17**, 1157-72.
- Epstein, D. J., Vekemans, M. und Gros, P.** (1991). Splotch (Sp2H), a mutation affecting development of the mouse neural tube, shows a deletion within the paired homeodomain of Pax-3. *Cell* **67**, 767-74.
- Epstein, J. A., Shapiro, D. N., Cheng, J., Lam, P. Y. und Maas, R. L.** (1996). Pax3 modulates expression of the c-Met receptor during limb muscle development. *Proc Natl Acad Sci U S A* **93**, 4213-8.
- Fan, C. M. und Tessier-Lavigne, M.** (1994). Patterning of mammalian somites by surface ectoderm and notochord: evidence for sclerotome induction by a hedgehog homolog. *Cell* **79**, 1175-86.
- Fan, S., Ma, Y. X., Gao, M., Yuan, R. Q., Meng, Q., Goldberg, I. D. und Rosen, E. M.** (2001). The multisubstrate adapter Gab1 regulates hepatocyte growth factor (scatter factor)-c-Met signaling for cell survival and DNA repair. *Mol Cell Biol* **21**, 4968-84.
- Farley, F. W., Soriano, P., Steffen, L. S. und Dymecki, S. M.** (2000). Widespread recombinase expression using FLPeR (flipper) mice. *Genesis* **28**, 106-10.
- Feinberg, A. P. und Vogelstein, B.** (1983). A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal Biochem* **132**, 6-13.
- Fiering, S., Epner, E., Robinson, K., Zhuang, Y., Telling, A., Hu, M., Martin, D. I., Enver, T., Ley, T. J. und Groudine, M.** (1995). Targeted deletion of 5'HS2 of the murine beta-globin LCR reveals that it is not essential for proper regulation of the beta-globin locus. *Genes Dev* **9**, 2203-13.

- Fixman, E. D., Fournier, T. M., Kamikura, D. M., Naujokas, M. A. und Park, M.** (1996). Pathways downstream of Shc and Grb2 are required for cell transformation by the tpr-Met oncoprotein. *J Biol Chem* **271**, 13116-22.
- Franz, T., Kothary, R., Surani, M. A., Halata, Z. und Grim, M.** (1993). The Splotch mutation interferes with muscle development in the limbs. *Anat Embryol (Berl)* **187**, 153-60.
- Fujiwara, K., Nagoshi, S., Ohno, A., Hirata, K., Ohta, Y., Mochida, S., Tomiya, T., Higashio, K. und Kurokawa, K.** (1993). Stimulation of liver growth by exogenous human hepatocyte growth factor in normal and partially hepatectomized rats. *Hepatology* **18**, 1443-9.
- Furge, K. A., Zhang, Y. W. und Vande Woude, G. F.** (2000). Met receptor tyrosine kinase: enhanced signaling through adapter proteins. *Oncogene* **19**, 5582-9.
- Garcia-Guzman, M., Dolfi, F., Zeh, K. und Vuori, K.** (1999). Met-induced JNK activation is mediated by the adapter protein Crk and correlates with the Gab1 - Crk signaling complex formation. *Oncogene* **18**, 7775-86.
- Gaudino, G., Follenzi, A., Naldini, L., Collesi, C., Santoro, M., Gallo, K. A., Godowski, P. J. und Comoglio, P. M.** (1994). RON is a heterodimeric tyrosine kinase receptor activated by the HGF homologue MSP. *Embo J* **13**, 3524-32.
- Gavrieli, Y., Sherman, Y. und Ben-Sasson, S. A.** (1992). Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. *J Cell Biol* **119**, 493-501.
- Gherardi, E., Gray, J., Stoker, M., Perryman, M. und Furlong, R.** (1989). Purification of scatter factor, a fibroblast-derived basic protein that modulates epithelial interactions and movement. *Proc Natl Acad Sci U S A* **86**, 5844-8.
- Gherardi, E., Youles, M. E., Miguel, R. N., Blundell, T. L., Iamele, L., Gough, J., Bandyopadhyay, A., Hartmann, G. und Butler, P. J.** (2003). Functional map and domain structure of MET, the product of the c-met protooncogene and receptor for hepatocyte growth factor/scatter factor. *Proc Natl Acad Sci U S A* **100**, 12039-44.
- Gorski, J. A. und Jones, K. R.** (1999). Efficient bicistronic expression of cre in mammalian cells. *Nucleic Acids Res* **27**, 2059-61.
- Goulding, M., Lumsden, A. und Paquette, A. J.** (1994). Regulation of Pax-3 expression in the dermomyotome and its role in muscle development. *Development* **120**, 957-71.
- Goulding, M. und Paquette, A.** (1994). Pax genes and neural tube defects in the mouse. *Ciba Found Symp* **181**, 103-13; discussion 113-7.
- Goulding, M. D., Chalepakis, G., Deutsch, U., Erselius, J. R. und Gruss, P.** (1991). Pax-3, a novel murine DNA binding protein expressed during early neurogenesis. *Embo J* **10**, 1135-47.

- Graziani, A., Gramaglia, D., Cantley, L. C. und Comoglio, P. M.** (1991). The tyrosine-phosphorylated hepatocyte growth factor/scatter factor receptor associates with phosphatidylinositol 3-kinase. *J Biol Chem* **266**, 22087-90.
- Graziani, A., Gramaglia, D., dalla Zonca, P. und Comoglio, P. M.** (1993). Hepatocyte growth factor/scatter factor stimulates the Ras-guanine nucleotide exchanger. *J Biol Chem* **268**, 9165-8.
- Gross, M. K., Moran-Rivard, L., Velasquez, T., Nakatsu, M. N., Jagla, K. und Goulding, M.** (2000). Lbx1 is required for muscle precursor migration along a lateral pathway into the limb. *Development* **127**, 413-24.
- Gu, H., Marth, J. D., Orban, P. C., Mossmann, H. und Rajewsky, K.** (1994). Deletion of a DNA polymerase beta gene segment in T cells using cell type-specific gene targeting. *Science* **265**, 103-6.
- Gu, H., Pratt, J. C., Burakoff, S. J. und Neel, B. G.** (1998). Cloning of p97/Gab2, the major SHP2-binding protein in hematopoietic cells, reveals a novel pathway for cytokine-induced gene activation. *Mol Cell* **2**, 729-40.
- Han, S., Stuart, L. A. und Degen, S. J.** (1991). Characterization of the DNF15S2 locus on human chromosome 3: identification of a gene coding for four kringle domains with homology to hepatocyte growth factor. *Biochemistry* **30**, 9768-80.
- Hartmann, G., Prospero, T., Brinkmann, V., Ozcelik, C., Winter, G., Hepple, J., Batley, S., Bladt, F., Sachs, M., Birchmeier, C. et al.** (1998). Engineered mutants of HGF/SF with reduced binding to heparan sulphate proteoglycans, decreased clearance and enhanced activity in vivo. *Curr Biol* **8**, 125-34.
- Hartmann, G., Weidner, K. M., Schwarz, H. und Birchmeier, W.** (1994). The motility signal of scatter factor/hepatocyte growth factor mediated through the receptor tyrosine kinase met requires intracellular action of Ras. *J Biol Chem* **269**, 21936-9.
- Hebert, J. M. und McConnell, S. K.** (2000). Targeting of cre to the Foxg1 (BF-1) locus mediates loxP recombination in the telencephalon and other developing head structures. *Dev Biol* **222**, 296-306.
- Herbst, R., Carroll, P. M., Allard, J. D., Schilling, J., Raabe, T. und Simon, M. A.** (1996). Daughter of sevenless is a substrate of the phosphotyrosine phosphatase Corkscrew and functions during sevenless signaling. *Cell* **85**, 899-909.
- Hoess, R. H. und Abremski, K.** (1985). Mechanism of strand cleavage and exchange in the Cre-lox site-specific recombination system. *J Mol Biol* **181**, 351-62.
- Hogan, B., Beddington, R., Constantini, F. und Lacy, E.** (1994). Manipulating The Mouse Embryo - A Laboratory Manual. Cold Spring Harbor, New York 11803-2500, USA: Cold Spring Harbor Laboratory Press.
- Holgado-Madruga, M., Emlet, D. R., Moscatello, D. K., Godwin, A. K. und Wong, A. J.** (1996). A Grb2-associated docking protein in EGF- and insulin-receptor signalling. *Nature* **379**, 560-4.

- Hug, B. A., Wesselschmidt, R. L., Fiering, S., Bender, M. A., Epner, E., Groudine, M. und Ley, T. J.** (1996). Analysis of mice containing a targeted deletion of beta-globin locus control region 5' hypersensitive site 3. *Mol Cell Biol* **16**, 2906-12.
- Hug, H., Costas, M., Staeheli, P., Aebi, M. und Weissmann, C.** (1988). Organization of the murine Mx gene and characterization of its interferon- and virus-inducible promoter. *Mol Cell Biol* **8**, 3065-79.
- Hunter, T.** (1998). The role of tyrosine phosphorylation in cell growth and disease. *Harvey Lect* **94**, 81-119.
- Hunter, T., Lindberg, R. A., Middlemas, D. S., Tracy, S. und van der Geer, P.** (1992). Receptor protein tyrosine kinases and phosphatases. *Cold Spring Harb Symp Quant Biol* **57**, 25-41.
- Ieraci, A., Forni, P. E. und Ponzetto, C.** (2002). Viable hypomorphic signaling mutant of the Met receptor reveals a role for hepatocyte growth factor in postnatal cerebellar development. *Proc Natl Acad Sci U S A* **99**, 15200-5.
- Imai, T., Chambon, P. und Metzger, D.** (2000). Inducible site-specific somatic mutagenesis in mouse hepatocytes. *Genesis* **26**, 147-8.
- Innis, M. A., Gelfand, D. H. und Sninsky, J. J.** (1989). PCR Protocols: A Guide To Methods And Applications, (ed., pp. 482. San Diego, CA 92101-4495, USA: Academic Press.
- Inoue, H., Nojima, H. und Okayama, H.** (1990). High efficiency transformation of Escherichia coli with plasmids. *Gene* **96**, 23-8.
- Itoh, M., Yoshida, Y., Nishida, K., Narimatsu, M., Hibi, M. und Hirano, T.** (2000). Role of Gab1 in heart, placenta, and skin development and growth factor- and cytokine-induced extracellular signal-regulated kinase mitogen-activated protein kinase activation. *Mol Cell Biol* **20**, 3695-704.
- Jagla, K., Dolle, P., Mattei, M. G., Jagla, T., Schuhbaur, B., Dretzen, G., Bellard, F. und Bellard, M.** (1995). Mouse Lbx1 and human LBX1 define a novel mammalian homeobox gene family related to the Drosophila lady bird genes. *Mech Dev* **53**, 345-56.
- Jeffers, M., Rong, S. und Vande Woude, G. F.** (1996). Enhanced tumorigenicity and invasion-metastasis by hepatocyte growth factor/scatter factor-met signalling in human cells concomitant with induction of the urokinase proteolysis network. *Mol Cell Biol* **16**, 1115-25.
- Jerpseth, B., Greener, A., Short, J. M., Viola, J. und Kretz, P.** (1992). XL1-Blue MRF' E. coli cells: McrA-, McrCB-, McrF-, Mrr-, HsdR- derivative of XL1-Blue cells. *Strategies* **5**, 81-83.
- Johnson, R. L., Laufer, E., Riddle, R. D. und Tabin, C.** (1994). Ectopic expression of Sonic hedgehog alters dorsal-ventral patterning of somites. *Cell* **79**, 1165-73.
- Joyner, A. L.** (1999). Gene Targeting: A Practical Approach. Oxford, United Kingdom: Oxford University Press.

- Jung, W., Castren, E., Odenthal, M., Vande Woude, G. F., Ishii, T., Dienes, H. P., Lindholm, D. und Schirmacher, P.** (1994). Expression and functional interaction of hepatocyte growth factor-scatter factor and its receptor c-met in mammalian brain. *J Cell Biol* **126**, 485-94.
- Kang, J. Y., Dolled-Filhart, M., Ocal, I. T., Singh, B., Lin, C. Y., Dickson, R. B., Rimm, D. L. und Camp, R. L.** (2003). Tissue microarray analysis of hepatocyte growth factor/Met pathway components reveals a role for Met, matriptase, and hepatocyte growth factor activator inhibitor 1 in the progression of node-negative breast cancer. *Cancer Res* **63**, 1101-5.
- Kawaguchi, T., Qin, L., Shimomura, T., Kondo, J., Matsumoto, K., Denda, K. und Kitamura, N.** (1997). Purification and cloning of hepatocyte growth factor activator inhibitor type 2, a Kunitz-type serine protease inhibitor. *J Biol Chem* **272**, 27558-64.
- Kawaida, K., Matsumoto, K., Shimazu, H. und Nakamura, T.** (1994). Hepatocyte growth factor prevents acute renal failure and accelerates renal regeneration in mice. *Proc Natl Acad Sci U S A* **91**, 4357-61.
- Kellendonk, C., Tronche, F., Monaghan, A. P., Angrand, P. O., Stewart, F. und Schutz, G.** (1996). Regulation of Cre recombinase activity by the synthetic steroid RU 486. *Nucleic Acids Res* **24**, 1404-11.
- Kiefer, J. C. und Hauschka, S. D.** (2001). Myf-5 is transiently expressed in nonmuscle mesoderm and exhibits dynamic regional changes within the presegmented mesoderm and somites I-IV. *Dev Biol* **232**, 77-90.
- Kitta, K., Day, R. M., Ikeda, T. und Suzuki, Y. J.** (2001). Hepatocyte growth factor protects cardiac myocytes against oxidative stress-induced apoptosis. *Free Radic Biol Med* **31**, 902-10.
- Komada, M. und Kitamura, N.** (1993). The cell dissociation and motility triggered by scatter factor/hepatocyte growth factor are mediated through the cytoplasmic domain of the c-Met receptor. *Oncogene* **8**, 2381-90.
- Kuhn, R., Rajewsky, K. und Muller, W.** (1991). Generation and analysis of interleukin-4 deficient mice. *Science* **254**, 707-10.
- Kuhn, R., Schwenk, F., Aguet, M. und Rajewsky, K.** (1995). Inducible gene targeting in mice. *Science* **269**, 1427-9.
- Laemmli, U. K.** (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-5.
- Lamorte, L., Kamikura, D. M. und Park, M.** (2000). A switch from p130Cas/Crk to Gab1/Crk signaling correlates with anchorage independent growth and JNK activation in cells transformed by the Met receptor oncoprotein. *Oncogene* **19**, 5973-81.
- Ley, T. J., Hug, B., Fiering, S., Epner, E., Bender, M. A. und Groudine, M.** (1998). Reduced beta-globin gene expression in adult mice containing deletions of locus control region 5' HS-2 or 5' HS-3. *Ann N Y Acad Sci* **850**, 45-53.

- Liu, Y. und Rohrschneider, L. R.** (2002). The gift of Gab. *FEBS Lett* **515**, 1-7.
- Lobe, C. G., Koop, K. E., Kreppner, W., Lomeli, H., Gertsenstein, M. und Nagy, A.** (1999). Z/AP, a double reporter for cre-mediated recombination. *Dev Biol* **208**, 281-92.
- Lock, L. S., Maroun, C. R., Naujokas, M. A. und Park, M.** (2002). Distinct recruitment and function of Gab1 and Gab2 in Met receptor-mediated epithelial morphogenesis. *Mol Biol Cell* **13**, 2132-46.
- Lock, L. S., Royal, I., Naujokas, M. A. und Park, M.** (2000). Identification of an atypical Grb2 carboxyl-terminal SH3 domain binding site in Gab docking proteins reveals Grb2-dependent and -independent recruitment of Gab1 to receptor tyrosine kinases. *J Biol Chem* **275**, 31536-45.
- Lokker, N. A., Mark, M. R., Luis, E. A., Bennett, G. L., Robbins, K. A., Baker, J. B. und Godowski, P. J.** (1992). Structure-function analysis of hepatocyte growth factor: identification of variants that lack mitogenic activity yet retain high affinity receptor binding. *Embo J* **11**, 2503-10.
- MacGregor, G. R. und Caskey, C. T.** (1989). Construction of plasmids that express *E. coli* beta-galactosidase in mammalian cells. *Nucleic Acids Res* **17**, 2365.
- Maina, F., Casagranda, F., Audero, E., Simeone, A., Comoglio, P. M., Klein, R. und Ponzetto, C.** (1996). Uncoupling of Grb2 from the Met receptor in vivo reveals complex roles in muscle development. *Cell* **87**, 531-42.
- Maina, F., Hilton, M. C., Andres, R., Wyatt, S., Klein, R. und Davies, A. M.** (1998). Multiple roles for hepatocyte growth factor in sympathetic neuron development. *Neuron* **20**, 835-46.
- Maina, F., Hilton, M. C., Ponzetto, C., Davies, A. M. und Klein, R.** (1997). Met receptor signaling is required for sensory nerve development and HGF promotes axonal growth and survival of sensory neurons. *Genes Dev* **11**, 3341-50.
- Maina, F. und Klein, R.** (1999). Hepatocyte growth factor, a versatile signal for developing neurons. *Nat Neurosci* **2**, 213-7.
- Maroun, C. R., Holgado-Madruga, M., Royal, I., Naujokas, M. A., Fournier, T. M., Wong, A. J. und Park, M.** (1999a). The Gab1 PH domain is required for localization of Gab1 at sites of cell-cell contact and epithelial morphogenesis downstream from the met receptor tyrosine kinase. *Mol Cell Biol* **19**, 1784-99.
- Maroun, C. R., Moscatello, D. K., Naujokas, M. A., Holgado-Madruga, M., Wong, A. J. und Park, M.** (1999b). A conserved inositol phospholipid binding site within the pleckstrin homology domain of the Gab1 docking protein is required for epithelial morphogenesis. *J Biol Chem* **274**, 31719-26.
- Maroun, C. R., Naujokas, M. A., Holgado-Madruga, M., Wong, A. J. und Park, M.** (2000). The tyrosine phosphatase SHP-2 is required for sustained activation of extracellular signal-regulated kinase and epithelial morphogenesis downstream from the met receptor tyrosine kinase. *Mol Cell Biol* **20**, 8513-25.

- Mars, W. M., Zarnegar, R. und Michalopoulos, G. K.** (1993). Activation of hepatocyte growth factor by the plasminogen activators uPA and tPA. *Am J Pathol* **143**, 949-58.
- Maulik, G., Shrikhande, A., Kijima, T., Ma, P. C., Morrison, P. T. und Salgia, R.** (2002). Role of the hepatocyte growth factor receptor, c-Met, in oncogenesis and potential for therapeutic inhibition. *Cytokine Growth Factor Rev* **13**, 41-59.
- McDonell, M. W., Simon, M. N. und Studier, F. W.** (1977). Analysis of restriction fragments of T7 DNA and determination of molecular weights by electrophoresis in neutral and alkaline gels. *J Mol Biol* **110**, 119-46.
- McMahon, J. A., Takada, S., Zimmerman, L. B., Fan, C. M., Harland, R. M. und McMahon, A. P.** (1998). Noggin-mediated antagonism of BMP signaling is required for growth and patterning of the neural tube and somite. *Genes Dev* **12**, 1438-52.
- Meyers, E. N., Lewandoski, M. und Martin, G. R.** (1998). An Fgf8 mutant allelic series generated by Cre- and Flp-mediated recombination. *Nat Genet* **18**, 136-41.
- Miyazawa, K., Shimomura, T., Kitamura, A., Kondo, J., Morimoto, Y. und Kitamura, N.** (1993). Molecular cloning and sequence analysis of the cDNA for a human serine protease responsible for activation of hepatocyte growth factor. Structural similarity of the protease precursor to blood coagulation factor XII. *J Biol Chem* **268**, 10024-8.
- Nagy, A., Moens, C., Ivanyi, E., Pawling, J., Gertsenstein, M., Hadjantonakis, A. K., Pirity, M. und Rossant, J.** (1998). Dissecting the role of N-myc in development using a single targeting vector to generate a series of alleles. *Curr Biol* **8**, 661-4.
- Nagy, A., Rossant, J., Nagy, R., Abramow-Newerly, W. und Roder, J. C.** (1993). Derivation of completely cell culture-derived mice from early-passage embryonic stem cells. *Proc Natl Acad Sci U S A* **90**, 8424-8.
- Nakamura, T., Mizuno, S., Matsumoto, K., Sawa, Y. und Matsuda, H.** (2000). Myocardial protection from ischemia/reperfusion injury by endogenous and exogenous HGF. *J Clin Invest* **106**, 1511-9.
- Nakamura, T., Nishizawa, T., Hagiya, M., Seki, T., Shimonishi, M., Sugimura, A., Tashiro, K. und Shimizu, S.** (1989). Molecular cloning and expression of human hepatocyte growth factor. *Nature* **342**, 440-3.
- Naldini, L., Vigna, E., Narsimhan, R. P., Gaudino, G., Zarnegar, R., Michalopoulos, G. K. und Comoglio, P. M.** (1991). Hepatocyte growth factor (HGF) stimulates the tyrosine kinase activity of the receptor encoded by the proto-oncogene c-MET. *Oncogene* **6**, 501-4.
- Nguyen, L., Holgado-Madruga, M., Maroun, C., Fixman, E. D., Kamikura, D., Fournier, T., Charest, A., Tremblay, M. L., Wong, A. J. und Park, M.** (1997). Association of the multisubstrate docking protein Gab1 with the hepatocyte growth factor receptor requires a functional Grb2 binding site involving tyrosine 1356. *J Biol Chem* **272**, 20811-9.

- Nishida, K., Yoshida, Y., Itoh, M., Fukada, T., Ohtani, T., Shirogane, T., Atsumi, T., Takahashi-Tezuka, M., Ishihara, K., Hibi, M. et al.** (1999). Gab-family adapter proteins act downstream of cytokine and growth factor receptors and T- and B-cell antigen receptors. *Blood* **93**, 1809-16.
- Noden, D. M.** (1983). The embryonic origins of avian cephalic and cervical muscles and associated connective tissues. *Am J Anat* **168**, 257-76.
- Orban, P. C., Chui, D. und Marth, J. D.** (1992). Tissue- and site-specific DNA recombination in transgenic mice. *Proc Natl Acad Sci U S A* **89**, 6861-5.
- Ordahl, C. P. und Le Douarin, N. M.** (1992). Two myogenic lineages within the developing somite. *Development* **114**, 339-53.
- Ott, M. O., Bober, E., Lyons, G., Arnold, H. und Buckingham, M.** (1991). Early expression of the myogenic regulatory gene, myf-5, in precursor cells of skeletal muscle in the mouse embryo. *Development* **111**, 1097-107.
- Palka, H. L., Park, M. und Tonks, N. K.** (2003). Hepatocyte growth factor receptor tyrosine kinase met is a substrate of the receptor protein-tyrosine phosphatase DEP-1. *J Biol Chem* **278**, 5728-35.
- Park, M., Dean, M., Cooper, C. S., Schmidt, M., O'Brien, S. J., Blair, D. G. und Vande Woude, G. F.** (1986). Mechanism of met oncogene activation. *Cell* **45**, 895-904.
- Patel, K., Christ, B. und Stockdale, F. E.** (2002). Control of muscle size during embryonic, fetal, and adult life. *Results Probl Cell Differ* **38**, 163-86.
- Pelicci, G., Giordano, S., Zhen, Z., Salcini, A. E., Lanfrancone, L., Bardelli, A., Panayotou, G., Waterfield, M. D., Ponzetto, C., Pelicci, P. G. et al.** (1995). The motogenic and mitogenic responses to HGF are amplified by the Shc adaptor protein. *Oncogene* **10**, 1631-8.
- Ponzetto, C., Bardelli, A., Zhen, Z., Maina, F., dalla Zonca, P., Giordano, S., Graziani, A., Panayotou, G. und Comoglio, P. M.** (1994). A multifunctional docking site mediates signaling and transformation by the hepatocyte growth factor/scatter factor receptor family. *Cell* **77**, 261-71.
- Potempa, S. und Ridley, A. J.** (1998). Activation of both MAP kinase and phosphatidylinositide 3-kinase by Ras is required for hepatocyte growth factor/scatter factor-induced adherens junction disassembly. *Mol Biol Cell* **9**, 2185-200.
- Pownall, M. E. und Emerson, C. P., Jr.** (1992). Sequential activation of three myogenic regulatory genes during somite morphogenesis in quail embryos. *Dev Biol* **151**, 67-79.
- Puelles, E., Acampora, D., Lacroix, E., Signore, M., Annino, A., Tuorto, F., Filosa, S., Corte, G., Wurst, W., Ang, S. L. et al.** (2003). Otx dose-dependent integrated control of antero-posterior and dorso-ventral patterning of midbrain. *Nat Neurosci* **6**, 453-60.

- Raabe, T., Riesgo-Escovar, J., Liu, X., Bausenwein, B. S., Deak, P., Maroy, P. und Hafen, E.** (1996). DOS, a novel pleckstrin homology domain-containing protein required for signal transduction between sevenless and Ras1 in Drosophila. *Cell* **85**, 911-20.
- Ramirez-Solis, R., Rivera-Perez, J., Wallace, J. D., Wims, M., Zheng, H. und Bradley, A.** (1992). Genomic DNA microextraction: a method to screen numerous samples. *Anal Biochem* **201**, 331-5.
- Rappolee, D. A., Iyer, A. und Patel, Y.** (1996). Hepatocyte growth factor and its receptor are expressed in cardiac myocytes during early cardiogenesis. *Circ Res* **78**, 1028-36.
- Reese, M. G., Eeckman, F. H., Kulp, D. und Haussler, D.** (1997). Improved splice site detection in Genie. *J Comput Biol* **4**, 311-23.
- Relaix, F., Rocancourt, D., Mansouri, A. und Buckingham, M.** (2004). Divergent functions of murine Pax3 and Pax7 in limb muscle development. *Genes Dev* **18**, 1088-105.
- Ridley, A. J., Comoglio, P. M. und Hall, A.** (1995). Regulation of scatter factor/hepatocyte growth factor responses by Ras, Rac, and Rho in MDCK cells. *Mol Cell Biol* **15**, 1110-22.
- Rodrigues, G. A. und Park, M.** (1993). Dimerization mediated through a leucine zipper activates the oncogenic potential of the met receptor tyrosine kinase. *Mol Cell Biol* **13**, 6711-22.
- Rodrigues, G. A. und Park, M.** (1994). Autophosphorylation modulates the kinase activity and oncogenic potential of the Met receptor tyrosine kinase. *Oncogene* **9**, 2019-27.
- Rodrigues, G. A., Park, M. und Schlessinger, J.** (1997). Activation of the JNK pathway is essential for transformation by the Met oncogene. *Embo J* **16**, 2634-45.
- Rodriguez, C. I., Buchholz, F., Galloway, J., Sequerra, R., Kasper, J., Ayala, R., Stewart, A. F. und Dymecki, S. M.** (2000). High-efficiency deleter mice show that FLPe is an alternative to Cre-loxP. *Nat Genet* **25**, 139-40.
- Rosario, M. und Birchmeier, W.** (2003). How to make tubes: signaling by the Met receptor tyrosine kinase. *Trends Cell Biol* **13**, 328-35.
- Rosen, E. M., Goldberg, I. D., Kacinski, B. M., Buckholz, T. und Vinter, D. W.** (1989). Smooth muscle releases an epithelial cell scatter factor which binds to heparin. *In Vitro Cell Dev Biol* **25**, 163-73.
- Royal, I., Lamarche-Vane, N., Lamorte, L., Kaibuchi, K. und Park, M.** (2000). Activation of cdc42, rac, PAK, and rho-kinase in response to hepatocyte growth factor differentially regulates epithelial cell colony spreading and dissociation. *Mol Biol Cell* **11**, 1709-25.

- Sachs, M., Brohmann, H., Zechner, D., Muller, T., Hulsken, J., Walther, I., Schaeper, U., Birchmeier, C. und Birchmeier, W.** (2000). Essential role of Gab1 for signaling by the c-Met receptor in vivo. *J Cell Biol* **150**, 1375-84.
- Sachs, M., Weidner, K. M., Brinkmann, V., Walther, I., Obermeier, A., Ullrich, A. und Birchmeier, W.** (1996). Motogenic and morphogenic activity of epithelial receptor tyrosine kinases. *J Cell Biol* **133**, 1095-1107.
- Saiki, R. K., Scharf, S., Faloona, F., Mullis, K. B., Horn, G. T., Erlich, H. A. und Arnheim, N.** (1985). Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* **230**, 1350-4.
- Sakkab, D., Lewitzky, M., Posern, G., Schaeper, U., Sachs, M., Birchmeier, W. und Feller, S. M.** (2000). Signaling of hepatocyte growth factor/scatter factor (HGF) to the small GTPase Rap1 via the large docking protein Gab1 and the adapter protein CRKL. *J Biol Chem* **275**, 10772-8.
- Sambrook, J. und Russell, D. W.** (2001). Molecular Cloning: A Laboratory Manual. Cold Spring Harbor, New York 11803-2500, USA: Cold Spring Harbor Laboratory Press.
- Sanger, F., Nicklen, S. und Coulson, A. R.** (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* **74**, 5463-7.
- Sassoon, D. A.** (1993). Myogenic regulatory factors: dissecting their role and regulation during vertebrate embryogenesis. *Dev Biol* **156**, 11-23.
- Sato, T., Tani, Y., Murao, S., Fujieda, H., Sato, H., Matsumoto, M., Takeuchi, T. und Ohtsuki, Y.** (2001). Focal enhancement of expression of c-Met/hepatocyte growth factor receptor in the myocardium in human myocardial infarction. *Cardiovasc Pathol* **10**, 235-40.
- Sauer, B. und Henderson, N.** (1988). Site-specific DNA recombination in mammalian cells by the Cre recombinase of bacteriophage P1. *Proc Natl Acad Sci U S A* **85**, 5166-70.
- Scaal, M., Bonafede, A., Dathe, V., Sachs, M., Cann, G., Christ, B. und Brand-Saberi, B.** (1999). SF/HGF is a mediator between limb patterning and muscle development. *Development* **126**, 4885-93.
- Schaeper, U., Gehring, N. H., Fuchs, K. P., Sachs, M., Kempkes, B. und Birchmeier, W.** (2000). Coupling of Gab1 to c-Met, Grb2, and Shp2 mediates biological responses. *J Cell Biol* **149**, 1419-32.
- Schafer, K. und Braun, T.** (1999). Early specification of limb muscle precursor cells by the homeobox gene Lbx1h. *Nat Genet* **23**, 213-6.
- Schlessinger, J.** (2000). Cell signaling by receptor tyrosine kinases. *Cell* **103**, 211-25.
- Schmidt, C., Bladt, F., Goedelke, S., Brinkmann, V., Zschiesche, W., Sharpe, M., Gherardi, E. und Birchmeier, C.** (1995). Scatter factor/hepatocyte growth factor is essential for liver development. *Nature* **373**, 699-702.

- Schmidt, L., Duh, F. M., Chen, F., Kishida, T., Glenn, G., Choyke, P., Scherer, S. W., Zhuang, Z., Lubensky, I., Dean, M. et al.** (1997). Germline and somatic mutations in the tyrosine kinase domain of the MET proto-oncogene in papillary renal carcinomas. *Nat Genet* **16**, 68-73.
- Schultzman, J. L., Borland, C. Z., Newman, J. C., Robinson, M. K., Kokel, M. und Stern, M. J.** (2001). The *Caenorhabditis elegans* EGL-15 signaling pathway implicates a DOS-like multisubstrate adaptor protein in fibroblast growth factor signal transduction. *Mol Cell Biol* **21**, 8104-16.
- Sham, M. H., Vesque, C., Nonchev, S., Marshall, H., Frain, M., Gupta, R. D., Whiting, J., Wilkinson, D., Charnay, P. und Krumlauf, R.** (1993). The zinc finger gene Krox20 regulates HoxB2 (Hox2.8) during hindbrain segmentation. *Cell* **72**, 183-96.
- Shi, Z. Q., Lu, W. und Feng, G. S.** (1998). The Shp-2 tyrosine phosphatase has opposite effects in mediating the activation of extracellular signal-regulated and c-Jun NH₂-terminal mitogen-activated protein kinases. *J Biol Chem* **273**, 4904-8.
- Shimomura, T., Kondo, J., Ochiai, M., Naka, D., Miyazawa, K., Morimoto, Y. und Kitamura, N.** (1993). Activation of the zymogen of hepatocyte growth factor activator by thrombin. *J Biol Chem* **268**, 22927-32.
- Simeone, A., Acampora, D., Gulisano, M., Stornaiuolo, A. und Boncinelli, E.** (1992). Nested expression domains of four homeobox genes in developing rostral brain. *Nature* **358**, 687-90.
- Simeone, A., Acampora, D., Mallamaci, A., Stornaiuolo, A., D'Apice, M. R., Nigro, V. und Boncinelli, E.** (1993). A vertebrate gene related to orthodenticle contains a homeodomain of the bicoid class and demarcates anterior neuroectoderm in the gastrulating mouse embryo. *Embo J* **12**, 2735-47.
- Skeel, A., Yoshimura, T., Showalter, S. D., Tanaka, S., Appella, E. und Leonard, E. J.** (1991). Macrophage stimulating protein: purification, partial amino acid sequence, and cellular activity. *J Exp Med* **173**, 1227-34.
- Sonnenberg, E., Meyer, D., Weidner, K. M. und Birchmeier, C.** (1993a). Scatter factor/hepatocyte growth factor and its receptor, the c-met tyrosine kinase, can mediate a signal exchange between mesenchyme and epithelia during mouse development. *J Cell Biol* **123**, 223-35.
- Sonnenberg, E., Weidner, K. M. und Birchmeier, C.** (1993b). Expression of the met-receptor and its ligand, HGF-SF during mouse embryogenesis. *Exs* **65**, 381-94.
- Sorge, J. A.** (1988). Bacteriophage lambda cloning vectors. *Biotechnology* **10**, 43-60.
- Southern, E. M.** (1975). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* **98**, 503-17.
- Sternberg, N. und Cohen, G.** (1989). Genetic analysis of the lytic replicon of bacteriophage P1. II. Organization of replicon elements. *J Mol Biol* **207**, 111-33.

- Sternberg, N., Hamilton, D., Austin, S., Yarmolinsky, M. und Hoess, R.** (1981). Site-specific recombination and its role in the life cycle of bacteriophage P1. *Cold Spring Harb Symp Quant Biol* **45 Pt 1**, 297-309.
- Stockdale, F. E. und Holtzer, H.** (1961). DNA synthesis and myogenesis. *Exp Cell Res* **24**, 508-20.
- Stoker, M., Gherardi, E., Perryman, M. und Gray, J.** (1987). Scatter factor is a fibroblast-derived modulator of epithelial cell mobility. *Nature* **327**, 239-42.
- Tabor, S. und Richardson, C. C.** (1987). DNA sequence analysis with a modified bacteriophage T7 DNA polymerase. *Proc Natl Acad Sci U S A* **84**, 4767-71.
- Takaishi, K., Sasaki, T., Kato, M., Yamochi, W., Kuroda, S., Nakamura, T., Takeichi, M. und Takai, Y.** (1994). Involvement of Rho p21 small GTP-binding protein and its regulator in the HGF-induced cell motility. *Oncogene* **9**, 273-9.
- Takayama, H., La Rochelle, W. J., Anver, M., Bockman, D. E. und Merlino, G.** (1996). Scatter factor/hepatocyte growth factor as a regulator of skeletal muscle and neural crest development. *Proc Natl Acad Sci U S A* **93**, 5866-71.
- Testa, G. und Stewart, A. F.** (2000). Creating a transloxation. Engineering interchromosomal translocations in the mouse. *EMBO Rep* **1**, 120-1.
- Thewke, D. P. und Seeds, N. W.** (1999). The expression of mRNAs for hepatocyte growth factor/scatter factor, its receptor c-met, and one of its activators tissue-type plasminogen activator show a systematic relationship in the developing and adult cerebral cortex and hippocampus. *Brain Res* **821**, 356-67.
- Thiery, J. P.** (2002). Epithelial-mesenchymal transitions in tumour progression. *Nat Rev Cancer* **2**, 442-54.
- Thorey, I. S., Muth, K., Russ, A. P., Otte, J., Reffelmann, A. und von Melchner, H.** (1998). Selective disruption of genes transiently induced in differentiating mouse embryonic stem cells by using gene trap mutagenesis and site-specific recombination. *Mol Cell Biol* **18**, 3081-8.
- Tremblay, P., Dietrich, S., Mericskay, M., Schubert, F. R., Li, Z. und Paulin, D.** (1998). A crucial role for Pax3 in the development of the hypaxial musculature and the long-range migration of muscle precursors. *Dev Biol* **203**, 49-61.
- Ueda, H., Nakamura, T., Matsumoto, K., Sawa, Y. und Matsuda, H.** (2001). A potential cardioprotective role of hepatocyte growth factor in myocardial infarction in rats. *Cardiovasc Res* **51**, 41-50.
- Uehara, Y., Minowa, O., Mori, C., Shiota, K., Kuno, J., Noda, T. und Kitamura, N.** (1995). Placental defect and embryonic lethality in mice lacking hepatocyte growth factor/scatter factor. *Nature* **373**, 702-5.
- Urlinger, S., Baron, U., Thellmann, M., Hasan, M. T., Bujard, H. und Hillen, W.** (2000). Exploring the sequence space for tetracycline-dependent transcriptional

activators: novel mutations yield expanded range and sensitivity. *Proc Natl Acad Sci U S A* **97**, 7963-8.

Vogelstein, B. und Gillespie, D. (1979). Preparative and analytical purification of DNA from agarose. *Proc Natl Acad Sci U S A* **76**, 615-9.

Wang, M. H., Ronsin, C., Gesnel, M. C., Coupey, L., Skeel, A., Leonard, E. J. und Breathnach, R. (1994). Identification of the ron gene product as the receptor for the human macrophage stimulating protein. *Science* **266**, 117-9.

Wang, X., DeFrances, M. C., Dai, Y., Pediaditakis, P., Johnson, C., Bell, A., Michalopoulos, G. K. und Zarnegar, R. (2002). A mechanism of cell survival: sequestration of Fas by the HGF receptor Met. *Mol Cell* **9**, 411-21.

Weidner, K. M., Arakaki, N., Hartmann, G., Vandekerckhove, J., Weingart, S., Rieder, H., Fonatsch, C., Tsubouchi, H., Hishida, T., Daikuhara, Y. et al. (1991). Evidence for the identity of human scatter factor and human hepatocyte growth factor. *Proc Natl Acad Sci U S A* **88**, 7001-5.

Weidner, K. M., Behrens, J., Vandekerckhove, J. und Birchmeier, W. (1990). Scatter factor: molecular characteristics and effect on the invasiveness of epithelial cells. *J Cell Biol*.

Weidner, K. M., Di Cesare, S., Sachs, M., Brinkmann, V., Behrens, J. und Birchmeier, W. (1996). Interaction between Gab1 and the c-Met receptor tyrosine kinase is responsible for epithelial morphogenesis. *Nature* **384**, 173-6.

Weidner, K. M., Sachs, M. und Birchmeier, W. (1993). The Met receptor tyrosine kinase transduces motility, proliferation, and morphogenic signals of scatter factor/hepatocyte growth factor in epithelial cells. *J Cell Biol* **121**, 145-54.

Weintraub, H., Tapscott, S. J., Davis, R. L., Thayer, M. J., Adam, M. A., Lassar, A. B. und Miller, A. D. (1989). Activation of muscle-specific genes in pigment, nerve, fat, liver, and fibroblast cell lines by forced expression of MyoD. *Proc Natl Acad Sci U S A* **86**, 5434-8.

Wilkinson, D. G. (1992). In situ Hybridization: A Practical Approach. Oxford, United Kingdom: Oxford University Press.

Williams, B. A. und Ordahl, C. P. (1994). Pax-3 expression in segmental mesoderm marks early stages in myogenic cell specification. *Development* **120**, 785-96.

Wolf, I., Jenkins, B. J., Liu, Y., Seiffert, M., Custodio, J. M., Young, P. und Rohrschneider, L. R. (2002). Gab3, a new DOS/Gab family member, facilitates macrophage differentiation. *Mol Cell Biol* **22**, 231-44.

Wu, C. J., O'Rourke, D. M., Feng, G. S., Johnson, G. R., Wang, Q. und Greene, M. I. (2001). The tyrosine phosphatase SHP-2 is required for mediating phosphatidylinositol 3-kinase/Akt activation by growth factors. *Oncogene* **20**, 6018-25.

- Wunderlich, F. T., Wildner, H., Rajewsky, K. und Edenhofer, F.** (2001). New variants of inducible Cre recombinase: a novel mutant of Cre-PR fusion protein exhibits enhanced sensitivity and an expanded range of inducibility. *Nucleic Acids Res* **29**, E47.
- Xiao, G. H., Jeffers, M., Bellacosa, A., Mitsuuchi, Y., Vande Woude, G. F. und Testa, J. R.** (2001). Anti-apoptotic signaling by hepatocyte growth factor/Met via the phosphatidylinositol 3-kinase/Akt and mitogen-activated protein kinase pathways. *Proc Natl Acad Sci U S A* **98**, 247-52.
- Yamamoto, Y., Livet, J., Pollock, R. A., Garces, A., Arce, V., deLapeyriere, O. und Henderson, C. E.** (1997). Hepatocyte growth factor (HGF/SF) is a muscle-derived survival factor for a subpopulation of embryonic motoneurons. *Development* **124**, 2903-13.
- Yang, X. M. und Park, M.** (1995). Expression of the hepatocyte growth factor/scatter factor receptor tyrosine kinase is localized to epithelia in the adult mouse. *Lab Invest* **73**, 483-91.
- Yang, X. M., Vogan, K., Gros, P. und Park, M.** (1996). Expression of the met receptor tyrosine kinase in muscle progenitor cells in somites and limbs is absent in Splotch mice. *Development* **122**, 2163-71.
- Zarnegar, R. und Michalopoulos, G.** (1989). Purification and biological characterization of human hepatopoietin A, a polypeptide growth factor for hepatocytes. *Cancer Res* **49**, 3314-20.
- Zhao, C., Yu, D. H., Shen, R. und Feng, G. S.** (1999). Gab2, a new pleckstrin homology domain-containing adapter protein, acts to uncouple signaling from ERK kinase to Elk-1. *J Biol Chem* **274**, 19649-54.
- Zhu, H., Naujokas, M. A., Fixman, E. D., Torossian, K. und Park, M.** (1994). Tyrosine 1356 in the carboxyl-terminal tail of the HGF/SF receptor is essential for the transduction of signals for cell motility and morphogenesis. *J Biol Chem* **269**, 29943-8.