

## 8. Literatur

Aberle, H., Bauer, A., Stappert, J., Kispert, A. and Kemler, R. (1997) beta-catenin is a target for the ubiquitin-proteasome pathway. *Embo J*, 16, 3797-3804.

Agou, F., Ye, F., Goffinont, S., Courtois, G., Yamaoka, S., Israel, A. and Veron, M. (2002) NEMO trimerizes through its coiled-coil C-terminal domain. *J Biol Chem*, 277, 17464-17475.

Andres, D., Diez-Fernandez, C., Castrillo, A. and Cascales, M. (2002) Relationship between the activation of heat shock factor and the suppression of nuclear factor-kappaB activity in rat hepatocyte cultures treated with cyclosporine A. *Biochem Pharmacol*, 64, 247-256.

Arenzana-Seisdedos, F., Turpin, P., Rodriguez, M., Thomas, D., Hay, R.T., Virelizier, J.L. and Dargemont, C. (1997) Nuclear localization of I kappa B alpha promotes active transport of NF-kappa B from the nucleus to the cytoplasm. *J Cell Sci*, 110, 369-378.

Baeuerle, P.A. (1991) The inducible transcription activator NF-kappa B: regulation by distinct protein subunits. *Biochim Biophys Acta*, 1072, 63-80.

Baeuerle, P.A. and Baltimore, D. (1988a) Activation of DNA-binding activity in an apparently cytoplasmic precursor of the NF-kappa B transcription factor. *Cell*, 53, 211-217.

Baeuerle, P.A. and Baltimore, D. (1988b) I kappa B: a specific inhibitor of the NF-kappa B transcription factor. *Science*, 242, 540-546.

Baldwin, A.S., Jr. (1996) The NF-kappa B and I kappa B proteins: new discoveries and insights. *Annu Rev Immunol*, 14, 649-683.

Bardwell, J.C. and Craig, E.A. (1987) Eukaryotic Mr 83,000 heat shock protein has a homologue in Escherichia coli. *Proc Natl Acad Sci U S A*, 84, 5177-5181.

Bargou, R.C., Leng, C., Krappmann, D., Emmerich, F., Mapara, M.Y., Bommert, K., Royer, H.D., Scheidereit, C. and Dorken, B. (1996) High-level nuclear NF-kappa B and Oct-2 is a common feature of cultured Hodgkin/Reed-Sternberg cells. *Blood*, 87, 4340-4347.

Bargou, R.C., Emmerich, F., Krappmann, D., Bommert, K., Mapara, M.Y., Arnold, W., Royer, H.D., Grinstein, E., Greiner, A., Scheidereit, C. and Dorken, B. (1997) Constitutive nuclear factor-kappaB-RelA activation is required for proliferation and survival of Hodgkin's disease tumor cells. *J Clin Invest*, 100, 2961-2969.

Barroga, C.F., Stevenson, J.K., Schwarz, E.M. and Verma, I.M. (1995) Constitutive phosphorylation of I kappa B alpha by casein kinase II. *Proc Natl Acad Sci U S A*, 92, 7637-7641.

Basso, A.D., Solit, D.B., Chiosis, G., Giri, B., Tsihchlis, P. and Rosen, N. (2002) Akt forms an intracellular complex with heat shock protein 90 (Hsp90) and Cdc37 and is destabilized by inhibitors of Hsp90 function. *J Biol Chem*, 277, 39858-39866.

Baud, V., Liu, Z.G., Bennett, B., Suzuki, N., Xia, Y. and Karin, M. (1999) Signaling by proinflammatory cytokines: oligomerization of TRAF2 and TRAF6 is sufficient for JNK and IKK activation and target gene induction via an amino-terminal effector domain. *Genes Dev*, 13, 1297-1308.

- Beg, A.A., Ruben, S.M., Scheinman, R.I., Haskill, S., Rosen, C.A. and Baldwin, A.S., Jr. (1992) I kappa B interacts with the nuclear localization sequences of the subunits of NF-kappa B: a mechanism for cytoplasmic retention. *Genes Dev*, 6, 1899-1913.
- Bender, K., Gottlicher, M., Whiteside, S., Rahmsdorf, H.J. and Herrlich, P. (1998) Sequential DNA damage-independent and -dependent activation of NF-kappaB by UV. *Embo J*, 17, 5170-5181.
- Bergerat, A., de Massy, B., Gadelle, D., Varoutas, P.C., Nicolas, A. and Forterre, P. (1997) An atypical topoisomerase II from Archaea with implications for meiotic recombination. *Nature*, 386, 414-417.
- Betts, J.C. and Nabel, G.J. (1996) Differential regulation of NF-kappaB2(p100) processing and control by amino-terminal sequences. *Mol Cell Biol*, 16, 6363-6371.
- Bijlmakers, M.J. and Marsh, M. (2000) Hsp90 is essential for the synthesis and subsequent membrane association, but not the maintenance, of the Src-kinase p56(lck). *Mol Biol Cell*, 11, 1585-1595.
- Birbach, A., Gold, P., Binder, B.R., Hofer, E., de Martin, R. and Schmid, J.A. (2002) Signaling molecules of the NF-kappa B pathway shuttle constitutively between cytoplasm and nucleus. *J Biol Chem*, 277, 10842-10851.
- Birnby, D.A., Link, E.M., Vowels, J.J., Tian, H., Colacurcio, P.L. and Thomas, J.H. (2000) A transmembrane guanylyl cyclase (DAF-11) and Hsp90 (DAF-21) regulate a common set of chemosensory behaviors in caenorhabditis elegans. *Genetics*, 155, 85-104.
- Bonizzi, G. and Karin, M. (2004) The two NF-kappaB activation pathways and their role in innate and adaptive immunity. *Trends Immunol*, 25, 280-288.
- Borkovich, K.A., Farrelly, F.W., Finkelstein, D.B., Taulien, J. and Lindquist, S. (1989) hsp82 is an essential protein that is required in higher concentrations for growth of cells at higher temperatures. *Mol Cell Biol*, 9, 3919-3930.
- Bours, V., Azarenko, V., Dejardin, E. and Siebenlist, U. (1994) Human RelB (I-Rel) functions as a kappa B site-dependent transactivating member of the family of Rel-related proteins. *Oncogene*, 9, 1699-1702.
- Bouwmeester, T., Bauch, A., Ruffner, H., Angrand, P.O., Bergamini, G., Croughton, K., Cruciat, C., Eberhard, D., Gagneur, J., Ghidelli, S., Hopf, C., Huhse, B., Mangano, R., Michon, A.M., Schirle, M., Schlegl, J., Schwab, M., Stein, M.A., Bauer, A., Casari, G., Drewes, G., Gavin, A.C., Jackson, D.B., Joberty, G., Neubauer, G., Rick, J., Kuster, B. and Superti-Furga, G. (2004) A physical and functional map of the human TNF-alpha/NF-kappa B signal transduction pathway. *Nat Cell Biol*, 6, 97-105.
- Brockman, J.A., Scherer, D.C., McKinsey, T.A., Hall, S.M., Qi, X., Lee, W.Y. and Ballard, D.W. (1995) Coupling of a signal response domain in I kappa B alpha to multiple pathways for NF-kappa B activation. *Mol Cell Biol*, 15, 2809-2818.
- Brown, K., Park, S., Kanno, T., Franzoso, G. and Siebenlist, U. (1993) Mutual regulation of the transcriptional activator NF-kappa B and its inhibitor, I kappa B-alpha. *Proc Natl Acad Sci U S A*, 90, 2532-2536.
- Brown, K., Gerstberger, S., Carlson, L., Franzoso, G. and Siebenlist, U. (1995) Control of I kappa B-alpha proteolysis by site-specific, signal-induced phosphorylation. *Science*, 267, 1485-1488.

- Brown, K., Franzoso, G., Baldi, L., Carlson, L., Mills, L., Lin, Y.C., Gerstberger, S. and Siebenlist, U. (1997) The signal response of I $\kappa$ B $\alpha$  is regulated by transferable N- and C-terminal domains. *Mol Cell Biol*, 17, 3021-3027.
- Buchner, J. (1996) Supervising the fold: functional principles of molecular chaperones. *Faseb J*, 10, 10-19.
- Bull, P., Morley, K.L., Hoekstra, M.F., Hunter, T. and Verma, I.M. (1990) The mouse c-rel protein has an N-terminal regulatory domain and a C-terminal transcriptional transactivation domain. *Mol Cell Biol*, 10, 5473-5485.
- Byrd, C.A., Bornmann, W., Erdjument-Bromage, H., Tempst, P., Pavletich, N., Rosen, N., Nathan, C.F. and Ding, A. (1999) Heat shock protein 90 mediates macrophage activation by Taxol and bacterial lipopolysaccharide. *Proc Natl Acad Sci U S A*, 96, 5645-5650.
- Cabannes, E., Khan, G., Aillet, F., Jarrett, R.F. and Hay, R.T. (1999) Mutations in the I $\kappa$ B $\alpha$  gene in Hodgkin's disease suggest a tumour suppressor role for I $\kappa$ B $\alpha$ . *Oncogene*, 18, 3063-3070.
- Cahir McFarland, E.D., Izumi, K.M. and Mosialos, G. (1999) Epstein-barr virus transformation: involvement of latent membrane protein 1-mediated activation of NF- $\kappa$ B. *Oncogene*, 18, 6959-6964.
- Cao, Z., Henzel, W.J. and Gao, X. (1996a) IRAK: a kinase associated with the interleukin-1 receptor. *Science*, 271, 1128-1131.
- Cao, Z., Xiong, J., Takeuchi, M., Kurama, T. and Goeddel, D.V. (1996b) TRAF6 is a signal transducer for interleukin-1. *Nature*, 383, 443-446.
- Cechetto, J.D. and Gupta, R.S. (2000) Immunoelectron microscopy provides evidence that tumor necrosis factor receptor-associated protein 1 (TRAP-1) is a mitochondrial protein which also localizes at specific extramitochondrial sites. *Exp Cell Res*, 260, 30-39.
- Chen, F.E., Huang, D.B., Chen, Y.Q. and Ghosh, G. (1998) Crystal structure of p50/p65 heterodimer of transcription factor NF- $\kappa$ B bound to DNA. *Nature*, 391, 410-413.
- Chen, G., Cao, P. and Goeddel, D.V. (2002) TNF-induced recruitment and activation of the IKK complex require Cdc37 and Hsp90. *Mol Cell*, 9, 401-410.
- Chen, S. and Smith, D.F. (1998) Hop as an adaptor in the heat shock protein 70 (Hsp70) and hsp90 chaperone machinery. *J Biol Chem*, 273, 35194-35200.
- Chen, Z.J., Parent, L. and Maniatis, T. (1996) Site-specific phosphorylation of I $\kappa$ B $\alpha$  by a novel ubiquitination-dependent protein kinase activity. *Cell*, 84, 853-862.
- Chinnaiyan, A.M., O'Rourke, K., Tewari, M. and Dixit, V.M. (1995) FADD, a novel death domain-containing protein, interacts with the death domain of Fas and initiates apoptosis. *Cell*, 81, 505-512.
- Chiosis, G., Lucas, B., Huezo, H., Solit, D., Basso, A. and Rosen, N. (2003) Development of purine-scaffold small molecule inhibitors of Hsp90. *Curr Cancer Drug Targets*, 3, 371-376.
- Chow, J.C., Young, D.W., Golenbock, D.T., Christ, W.J. and Gusovsky, F. (1999) Toll-like receptor-4 mediates lipopolysaccharide-induced signal transduction. *J Biol Chem*, 274, 10689-10692.

- Citri, A., Gan, J., Mosesson, Y., Vereb, G., Szollosi, J. and Yarden, Y. (2004) Hsp90 restrains ErbB-2/HER2 signalling by limiting heterodimer formation. *EMBO Rep*, 5, 1165-1170.
- Claudio, E., Brown, K., Park, S., Wang, H. and Siebenlist, U. (2002) BAFF-induced NEMO-independent processing of NF-kappa B2 in maturing B cells. *Nat Immunol*, 3, 958-965.
- Cogswell, P.C., Scheinman, R.I. and Baldwin, A.S., Jr. (1993) Promoter of the human NF-kappa B p50/p105 gene. Regulation by NF-kappa B subunits and by c-REL. *J Immunol*, 150, 2794-2804.
- Cohen, L., Henzel, W.J. and Baeuerle, P.A. (1998) IKAP is a scaffold protein of the IkappaB kinase complex. *Nature*, 395, 292-296.
- Connell, P., Ballinger, C.A., Jiang, J., Wu, Y., Thompson, L.J., Hohfeld, J. and Patterson, C. (2001) The co-chaperone CHIP regulates protein triage decisions mediated by heat-shock proteins. *Nat Cell Biol*, 3, 93-96.
- Connelly, M.A. and Marcu, K.B. (1995) CHUK, a new member of the helix-loop-helix and leucine zipper families of interacting proteins, contains a serine-threonine kinase catalytic domain. *Cell Mol Biol Res*, 41, 537-549.
- Cross, S.L., Halden, N.F., Lenardo, M.J. and Leonard, W.J. (1989) Functionally distinct NF-kappa B binding sites in the immunoglobulin kappa and IL-2 receptor alpha chain genes. *Science*, 244, 466-469.
- Curry, H.A., Clemens, R.A., Shah, S., Bradbury, C.M., Botero, A., Goswami, P. and Gius, D. (1999) Heat shock inhibits radiation-induced activation of NF-kappaB via inhibition of I-kappaB kinase. *J Biol Chem*, 274, 23061-23067.
- Cutforth, T. and Rubin, G.M. (1994) Mutations in Hsp83 and cdc37 impair signaling by the sevenless receptor tyrosine kinase in *Drosophila*. *Cell*, 77, 1027-1036.
- de Carcer, G. (2004) Heat shock protein 90 regulates the metaphase-anaphase transition in a polo-like kinase-dependent manner. *Cancer Res*, 64, 5106-5112.
- Dejardin, E., Droin, N.M., Delhase, M., Haas, E., Cao, Y., Makris, C., Li, Z.W., Karin, M., Ware, C.F. and Green, D.R. (2002) The lymphotoxin-beta receptor induces different patterns of gene expression via two NF-kappaB pathways. *Immunity*, 17, 525-535.
- Delhase, M., Hayakawa, M., Chen, Y. and Karin, M. (1999) Positive and negative regulation of IkappaB kinase activity through IKKbeta subunit phosphorylation. *Science*, 284, 309-313.
- Deng, L., Wang, C., Spencer, E., Yang, L., Braun, A., You, J., Slaughter, C., Pickart, C. and Chen, Z.J. (2000) Activation of the IkappaB kinase complex by TRAF6 requires a dimeric ubiquitin-conjugating enzyme complex and a unique polyubiquitin chain. *Cell*, 103, 351-361.
- Denis, M. and Gustafsson, J.A. (1989) The Mr approximately 90,000 heat shock protein: an important modulator of ligand and DNA-binding properties of the glucocorticoid receptor. *Cancer Res*, 49, 2275s-2281s.
- Devin, A., Cook, A., Lin, Y., Rodriguez, Y., Kelliher, M. and Liu, Z. (2000) The distinct roles of TRAF2 and RIP in IKK activation by TNF-R1: TRAF2 recruits IKK to TNF-R1 while RIP mediates IKK activation. *Immunity*, 12, 419-429.

- DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E. and Karin, M. (1997) A cytokine-responsive I $\kappa$ B kinase that activates the transcription factor NF- $\kappa$ B. *Nature*, 388, 548-554.
- Dittmar, K.D. and Pratt, W.B. (1997) Folding of the glucocorticoid receptor by the reconstituted Hsp90-based chaperone machinery. The initial hsp90.p60.hsp70-dependent step is sufficient for creating the steroid binding conformation. *J Biol Chem*, 272, 13047-13054.
- Dittmar, K.D., Banach, M., Galigniana, M.D. and Pratt, W.B. (1998) The role of DnaJ-like proteins in glucocorticoid receptor.hsp90 heterocomplex assembly by the reconstituted hsp90.p60.hsp70 foldosome complex. *J Biol Chem*, 273, 7358-7366.
- Dobrzanski, P., Ryseck, R.P. and Bravo, R. (1993) Both N- and C-terminal domains of RelB are required for full transactivation: role of the N-terminal leucine zipper-like motif. *Mol Cell Biol*, 13, 1572-1582.
- Donze, O., Abbas-Terki, T. and Picard, D. (2001) The Hsp90 chaperone complex is both a facilitator and a repressor of the dsRNA-dependent kinase PKR. *Embo J*, 20, 3771-3780.
- Drexler, H.G. (1993) Recent results on the biology of Hodgkin and Reed-Sternberg cells. II. Continuous cell lines. *Leuk Lymphoma*, 9, 1-25.
- Ducut Sigala, J.L., Bottero, V., Young, D.B., Shevchenko, A., Mercurio, F. and Verma, I.M. (2004) Activation of transcription factor NF- $\kappa$ B requires ELKS, an I $\kappa$ B kinase regulatory subunit. *Science*, 304, 1963-1967.
- Ellis, J. (1987) Proteins as molecular chaperones. *Nature*, 328, 378-379.
- Emmerich, F., Meiser, M., Hummel, M., Demel, G., Foss, H.D., Jundt, F., Mathas, S., Krappmann, D., Scheidereit, C., Stein, H. and Dorken, B. (1999) Overexpression of I $\kappa$ B $\alpha$  without inhibition of NF- $\kappa$ B activity and mutations in the I $\kappa$ B $\alpha$  gene in Reed-Sternberg cells. *Blood*, 94, 3129-3134.
- Erbse, A., Mayer, M.P. and Bukau, B. (2004) Mechanism of substrate recognition by Hsp70 chaperones. *Biochem Soc Trans*, 32, 617-621.
- Fan, C.M. and Maniatis, T. (1991) Generation of p50 subunit of NF- $\kappa$ B by processing of p105 through an ATP-dependent pathway. *Nature*, 354, 395-398.
- Felts, S.J., Owen, B.A., Nguyen, P., Trepel, J., Donner, D.B. and Toft, D.O. (2000) The hsp90-related protein TRAP1 is a mitochondrial protein with distinct functional properties. *J Biol Chem*, 275, 3305-3312.
- Fink, A.L. (1999) Chaperone-mediated protein folding. *Physiol Rev*, 79, 425-449.
- Fisher, D.L., Mandart, E. and Doree, M. (2000) Hsp90 is required for c-Mos activation and biphasic MAP kinase activation in *Xenopus* oocytes. *Embo J*, 19, 1516-1524.
- Fliss, A.E., Fang, Y., Boschelli, F. and Caplan, A.J. (1997) Differential in vivo regulation of steroid hormone receptor activation by Cdc37p. *Mol Biol Cell*, 8, 2501-2509.
- Fontana, J., Fulton, D., Chen, Y., Fairchild, T.A., McCabe, T.J., Fujita, N., Tsuruo, T. and Sessa, W.C. (2002) Domain mapping studies reveal that the M domain of hsp90 serves as a molecular scaffold to regulate Akt-dependent phosphorylation of endothelial nitric oxide synthase and NO release. *Circ Res*, 90, 866-873.

- Frydman, J. and Hohfeld, J. (1997) Chaperones get in touch: the Hip-Hop connection. *Trends Biochem Sci*, 22, 87-92.
- Garnier, C., Barbier, P., Gilli, R., Lopez, C., Peyrot, V. and Briand, C. (1998) Heat-shock protein 90 (hsp90) binds in vitro to tubulin dimer and inhibits microtubule formation. *Biochem Biophys Res Commun*, 250, 414-419.
- Garnier, C., Lafitte, D., Tsvetkov, P.O., Barbier, P., Leclerc-Devin, J., Millot, J.M., Briand, C., Makarov, A.A., Catelli, M.G. and Peyrot, V. (2002) Binding of ATP to heat shock protein 90: evidence for an ATP-binding site in the C-terminal domain. *J Biol Chem*, 277, 12208-12214.
- Ghosh, G., van Duyne, G., Ghosh, S. and Sigler, P.B. (1995) Structure of NF-kappa B p50 homodimer bound to a kappa B site. *Nature*, 373, 303-310.
- Ghosh, S. and Karin, M. (2002) Missing pieces in the NF-kappaB puzzle. *Cell*, 109, S81-96.
- Ghosh, S., May, M.J. and Kopp, E.B. (1998) NF-kappa B and Rel proteins: evolutionarily conserved mediators of immune responses. *Annu Rev Immunol*, 16, 225-260.
- Gilmore, T.D. (1999) Multiple mutations contribute to the oncogenicity of the retroviral oncoprotein v-Rel. *Oncogene*, 18, 6925-6937.
- Gires, O., Zimmer-Strobl, U., Gonnella, R., Ueffing, M., Marschall, G., Zeidler, R., Pich, D. and Hammerschmidt, W. (1997) Latent membrane protein 1 of Epstein-Barr virus mimics a constitutively active receptor molecule. *Embo J*, 16, 6131-6140.
- Glover, J.R. and Lindquist, S. (1998) Hsp104, Hsp70, and Hsp40: a novel chaperone system that rescues previously aggregated proteins. *Cell*, 94, 73-82.
- Goes, F.S. and Martin, J. (2001) Hsp90 chaperone complexes are required for the activity and stability of yeast protein kinases Mik1, Wee1 and Swe1. *Eur J Biochem*, 268, 2281-2289.
- Goloubinoff, P., Mogk, A., Zvi, A.P., Tomoyasu, T. and Bukau, B. (1999) Sequential mechanism of solubilization and refolding of stable protein aggregates by a bichaperone network. *Proc Natl Acad Sci U S A*, 96, 13732-13737.
- Graham, F.L. and van der Eb, A.J. (1973) A new technique for the assay of infectivity of human adenovirus 5 DNA. *Virology*, 52, 456-467.
- Grammatikakis, N., Lin, J.H., Grammatikakis, A., Tsiachlis, P.N. and Cochran, B.H. (1999) p50(cdc37) acting in concert with Hsp90 is required for Raf-1 function. *Mol Cell Biol*, 19, 1661-1672.
- Greenfeder, S.A., Nunes, P., Kwee, L., Labow, M., Chizzonite, R.A. and Ju, G. (1995) Molecular cloning and characterization of a second subunit of the interleukin 1 receptor complex. *J Biol Chem*, 270, 13757-13765.
- Griffin, G.E., Leung, K., Folks, T.M., Kunkel, S. and Nabel, G.J. (1989) Activation of HIV gene expression during monocyte differentiation by induction of NF-kappa B. *Nature*, 339, 70-73.
- Gupta, R.S. (1995) Phylogenetic analysis of the 90 kD heat shock family of protein sequences and an examination of the relationship among animals, plants, and fungi species. *Mol Biol Evol*, 12, 1063-1073.

- Guzhova, I.V., Darieva, Z.A., Melo, A.R. and Margulis, B.A. (1997) Major stress protein Hsp70 interacts with NF-kB regulatory complex in human T-lymphoma cells. *Cell Stress Chaperones*, 2, 132-139.
- Hannink, M. and Temin, H.M. (1989) Transactivation of gene expression by nuclear and cytoplasmic rel proteins. *Mol Cell Biol*, 9, 4323-4336.
- Hansen, L.K., Houchins, J.P. and O'Leary, J.J. (1991) Differential regulation of HSC70, HSP70, HSP90 alpha, and HSP90 beta mRNA expression by mitogen activation and heat shock in human lymphocytes. *Exp Cell Res*, 192, 587-596.
- Hartl, F.U. (1995) Principles of chaperone-mediated protein folding. *Philos Trans R Soc Lond B Biol Sci*, 348, 107-112.
- Haskill, S., Beg, A.A., Tompkins, S.M., Morris, J.S., Yurochko, A.D., Sampson-Johannes, A., Mondal, K., Ralph, P. and Baldwin, A.S., Jr. (1991) Characterization of an immediate-early gene induced in adherent monocytes that encodes I kappa B-like activity. *Cell*, 65, 1281-1289.
- Haslbeck, M. (2002) sHsps and their role in the chaperone network. *Cell Mol Life Sci*, 59, 1649-1657.
- Hawkes, N.A., Otero, G., Winkler, G.S., Marshall, N., Dahmus, M.E., Krappmann, D., Scheidereit, C., Thomas, C.L., Schiavo, G., Erdjument-Bromage, H., Tempst, P. and Svejstrup, J.Q. (2002) Purification and characterization of the human elongator complex. *J Biol Chem*, 277, 3047-3052.
- Hayden, M.S. and Ghosh, S. (2004) Signaling to NF-kappaB. *Genes Dev*, 18, 2195-2224.
- Heissmeyer, V., Krappmann, D., Hatada, E.N. and Scheidereit, C. (2001) Shared pathways of I kappa B kinase-induced SCF(betaTrCP)-mediated ubiquitination and degradation for the NF-kappaB precursor p105 and I kappa Balpha. *Mol Cell Biol*, 21, 1024-1035.
- Hernandez, M.P., Chadli, A. and Toft, D.O. (2002) HSP40 binding is the first step in the HSP90 chaperoning pathway for the progesterone receptor. *J Biol Chem*, 277, 11873-11881.
- Herrero, J.A., Mathew, P. and Paya, C.V. (1995) LMP-1 activates NF-kappa B by targeting the inhibitory molecule I kappa B alpha. *J Virol*, 69, 2168-2174.
- Hershko, A. and Ciechanover, A. (1998) The ubiquitin system. *Annu Rev Biochem*, 67, 425-479.
- Hideshima, T., Chauhan, D., Richardson, P., Mitsiades, C., Mitsiades, N., Hayashi, T., Munshi, N., Dang, L., Castro, A., Palombella, V., Adams, J. and Anderson, K.C. (2002) NF-kappa B as a therapeutic target in multiple myeloma. *J Biol Chem*, 277, 16639-16647.
- Hinz, M., Loser, P., Mathas, S., Krappmann, D., Dorken, B. and Scheidereit, C. (2001) Constitutive NF-kappaB maintains high expression of a characteristic gene network, including CD40, CD86, and a set of antiapoptotic genes in Hodgkin/Reed-Sternberg cells. *Blood*, 97, 2798-2807.
- Hinz, M., Lemke, P., Anagnostopoulos, I., Hacker, C., Krappmann, D., Mathas, S., Dorken, B., Zenke, M., Stein, H. and Scheidereit, C. (2002) Nuclear factor kappaB-dependent gene expression profiling of Hodgkin's disease tumor cells, pathogenetic

significance, and link to constitutive signal transducer and activator of transcription 5a activity. *J Exp Med*, 196, 605-617.

Horie, R., Watanabe, T., Ito, K., Morisita, Y., Watanabe, M., Ishida, T., Higashihara, M. and Kadin, M. (2002b) Cytoplasmic aggregation of TRAF2 and TRAF5 proteins in the Hodgkin-Reed-Sternberg cells. *Am J Pathol*, 160, 1647-1654.

Horie, R., Watanabe, T., Morishita, Y., Ito, K., Ishida, T., Kanegae, Y., Saito, I., Higashihara, M., Mori, S. and Kadin, M.E. (2002a) Ligand-independent signaling by overexpressed CD30 drives NF-kappaB activation in Hodgkin-Reed-Sternberg cells. *Oncogene*, 21, 2493-2503.

Hostein, I., Robertson, D., DiStefano, F., Workman, P. and Clarke, P.A. (2001) Inhibition of signal transduction by the Hsp90 inhibitor 17-allylamino-17-demethoxygeldanamycin results in cytostasis and apoptosis. *Cancer Res*, 61, 4003-4009.

Hsu, H., Xiong, J. and Goeddel, D.V. (1995) The TNF receptor 1-associated protein TRADD signals cell death and NF-kappa B activation. *Cell*, 81, 495-504

Hsu, H., Huang, J., Shu, H.B., Baichwal, V. and Goeddel, D.V. (1996a) TNF-dependent recruitment of the protein kinase RIP to the TNF receptor-1 signaling complex. *Immunity*, 4, 387-396.

Hsu, H., Shu, H.B., Pan, M.G. and Goeddel, D.V. (1996b) TRADD-TRAF2 and TRADD-FADD interactions define two distinct TNF receptor 1 signal transduction pathways. *Cell*, 84, 299-308.

Hu, M.C., Lee, D.F., Xia, W., Golfman, L.S., Ou-Yang, F., Yang, J.Y., Zou, Y., Bao, S., Hanada, N., Saso, H., Kobayashi, R. and Hung, M.C. (2004) IkappaB kinase promotes tumorigenesis through inhibition of forkhead FOXO3a. *Cell*, 117, 225-237.

Hu, Y., Baud, V., Delhase, M., Zhang, P., Deerinck, T., Ellisman, M., Johnson, R. and Karin, M. (1999) Abnormal morphogenesis but intact IKK activation in mice lacking the IKKalpha subunit of IkappaB kinase. *Science*, 284, 316-320.

Huang, T.T., Kudo, N., Yoshida, M. and Miyamoto, S. (2000) A nuclear export signal in the N-terminal regulatory domain of IkappaBalpha controls cytoplasmic localization of inactive NF-kappaB/IkappaBalpha complexes. *Proc Natl Acad Sci U S A*, 97, 1014-1019.

Huang, T.T. and Miyamoto, S. (2001) Postrepression activation of NF-kappaB requires the amino-terminal nuclear export signal specific to IkappaBalpha. *Mol Cell Biol*, 21, 4737-4747.

Huxford, T., Huang, D.B., Malek, S. and Ghosh, G. (1998) The crystal structure of the IkappaBalpha/NF-kappaB complex reveals mechanisms of NF-kappaB inactivation. *Cell*, 95, 759-770.

Inoue, H., Nojima, H. and Okayama, H. (1990) High efficiency transformation of *Escherichia coli* with plasmids. *Gene*, 96, 23-28.

Isaacs, J.S., Xu, W. and Neckers, L. (2003) Heat shock protein 90 as a molecular target for cancer therapeutics. *Cancer Cell*, 3, 213-217.

Ito, C.Y., Kazantsev, A.G. and Baldwin, A.S., Jr. (1994) Three NF-kappa B sites in the Ikappa B-alpha promoter are required for induction of gene expression by TNF alpha. *Nucleic Acids Res*, 22, 3787-3792.



- Jacob, J., Kelsoe, G., Rajewsky, K. and Weiss, U. (1991) Intraclonal generation of antibody mutants in germinal centres. *Nature*, 354, 389-392.
- Jacobs, M.D. and Harrison, S.C. (1998) Structure of an I kappa B alpha/NF-kappa B complex. *Cell*, 95, 749-758.
- Jakob, U. and Buchner, J. (1994) Assisting spontaneity: the role of Hsp90 and small Hsps as molecular chaperones. *Trends Biochem Sci*, 19, 205-211.
- Johnson, B.D., Schumacher, R.J., Ross, E.D. and Toft, D.O. (1998) Hop modulates Hsp70/Hsp90 interactions in protein folding. *J Biol Chem*, 273, 3679-3686.
- Johnson, C., Van Antwerp, D. and Hope, T.J. (1999) An N-terminal nuclear export signal is required for the nucleocytoplasmic shuttling of I kappa B alpha. *Embo J*, 18, 6682-6693.
- Kamal, A., Thao, L., Sensintaffar, J., Zhang, L., Boehm, M.F., Fritz, L.C. and Burrows, F.J. (2003) A high-affinity conformation of Hsp90 confers tumour selectivity on Hsp90 inhibitors. *Nature*, 425, 407-410.
- Kanayama, A., Seth, R.B., Sun, L., Ea, C.K., Hong, M., Shaito, A., Chiu, Y.H., Deng, L. and Chen, Z.J. (2004) TAB2 and TAB3 activate the NF-kappa B pathway through binding to polyubiquitin chains. *Mol Cell*, 15, 535-548.
- Kapahi, P., Takahashi, T., Natoli, G., Adams, S.R., Chen, Y., Tsien, R.Y. and Karin, M. (2000) Inhibition of NF-kappa B activation by arsenite through reaction with a critical cysteine in the activation loop of I kappa B kinase. *J Biol Chem*, 275, 36062-36066.
- Karin, M. and Ben-Neriah, Y. (2000) Phosphorylation meets ubiquitination: the control of NF-[kappa]B activity. *Annu Rev Immunol*, 18, 621-663.
- Karin, M., Yamamoto, Y. and Wang, Q.M. (2004) The IKK NF-kappa B system: a treasure trove for drug development. *Nat Rev Drug Discov*, 3, 17-26.
- Kato, T., Jr., Delhase, M., Hoffmann, A. and Karin, M. (2003) CK2 Is a C-Terminal I kappa B Kinase Responsible for NF-kappa B Activation during the UV Response. *Mol Cell*, 12, 829-839.
- Kelliher, M.A., Grimm, S., Ishida, Y., Kuo, F., Stanger, B.Z. and Leder, P. (1998) The death domain kinase RIP mediates the TNF-induced NF-kappa B signal. *Immunity*, 8, 297-303.
- Kitamura, H., Kanehira, K., Okita, K., Morimatsu, M. and Saito, M. (2000) MAIL, a novel nuclear I kappa B protein that potentiates LPS-induced IL-6 production. *FEBS Lett*, 485, 53-56.
- Kopp, E. and Ghosh, S. (1994) Inhibition of NF-kappa B by sodium salicylate and aspirin. *Science*, 265, 956-959.
- Kopp, E., Medzhitov, R., Carothers, J., Xiao, C., Douglas, I., Janeway, C.A. and Ghosh, S. (1999) ECSIT is an evolutionarily conserved intermediate in the Toll/IL-1 signal transduction pathway. *Genes Dev*, 13, 2059-2071.
- Kosano, H., Stensgard, B., Charlesworth, M.C., McMahon, N. and Toft, D. (1998) The assembly of progesterone receptor-hsp90 complexes using purified proteins. *J Biol Chem*, 273, 32973-32979.

- Krappmann, D., Emmerich, F., Kordes, U., Scharschmidt, E., Dorken, B. and Scheidereit, C. (1999) Molecular mechanisms of constitutive NF-kappaB/Rel activation in Hodgkin/Reed-Sternberg cells. *Oncogene*, 18, 943-953.
- Krappmann, D., Hatada, E.N., Tegethoff, S., Li, J., Klippel, A., Giese, K., Baeuerle, P.A. and Scheidereit, C. (2000) The I kappa B kinase (IKK) complex is tripartite and contains IKK gamma but not IKAP as a regular component. *J Biol Chem*, 275, 29779-29787.
- Krappmann, D., Wegener, E., Sunami, Y., Esen, M., Thiel, A., Mordmuller, B. and Scheidereit, C. (2004) The IkappaB kinase complex and NF-kappaB act as master regulators of lipopolysaccharide-induced gene expression and control subordinate activation of AP-1. *Mol Cell Biol*, 24, 6488-6500.
- Krishna, P. and Gloor, G. (2001) The Hsp90 family of proteins in Arabidopsis thaliana. *Cell Stress Chaperones*, 6, 238-246.
- Lai, B.T., Chin, N.W., Stanek, A.E., Keh, W. and Lanks, K.W. (1984) Quantitation and intracellular localization of the 85K heat shock protein by using monoclonal and polyclonal antibodies. *Mol Cell Biol*, 4, 2802-2810.
- Le Bail, O., Schmidt-Ullrich, R. and Israel, A. (1993) Promoter analysis of the gene encoding the I kappa B-alpha/MAD3 inhibitor of NF-kappa B: positive regulation by members of the rel/NF-kappa B family. *Embo J*, 12, 5043-5049.
- Lee, P., Rao, J., Fliss, A., Yang, E., Garrett, S. and Caplan, A.J. (2002) The Cdc37 protein kinase-binding domain is sufficient for protein kinase activity and cell viability. *J Cell Biol*, 159, 1051-1059.
- Lenardo, M. and Siebenlist, U. (1994) Bcl-3-mediated nuclear regulation of the NF-kappa B trans-activating factor. *Immunol Today*, 15, 145-147.
- Lewis, J., Devin, A., Miller, A., Lin, Y., Rodriguez, Y., Neckers, L. and Liu, Z.G. (2000) Disruption of hsp90 function results in degradation of the death domain kinase, receptor-interacting protein (RIP), and blockage of tumor necrosis factor-induced nuclear factor-kappaB activation. *J Biol Chem*, 275, 10519-10526.
- Li, N. and Karin, M. (1998) Ionizing radiation and short wavelength UV activate NF-kappaB through two distinct mechanisms. *Proc Natl Acad Sci U S A*, 95, 13012-13017.
- Li, Q., Lu, Q., Hwang, J.Y., Buscher, D., Lee, K.F., Izpisua-Belmonte, J.C. and Verma, I.M. (1999b) IKK1-deficient mice exhibit abnormal development of skin and skeleton. *Genes Dev*, 13, 1322-1328.
- Li, Q., Van Antwerp, D., Mercurio, F., Lee, K.F. and Verma, I.M. (1999a) Severe liver degeneration in mice lacking the IkappaB kinase 2 gene. *Science*, 284, 321-325.
- Liang, P. and MacRae, T.H. (1997) Molecular chaperones and the cytoskeleton. *J Cell Sci*, 110, 1431-1440.
- Lin, R., Beauparlant, P., Makris, C., Meloche, S. and Hiscott, J. (1996) Phosphorylation of IkappaBalpha in the C-terminal PEST domain by casein kinase II affects intrinsic protein stability. *Mol Cell Biol*, 16, 1401-1409.
- Ling, L., Cao, Z. and Goeddel, D.V. (1998) NF-kappaB-inducing kinase activates IKK-alpha by phosphorylation of Ser-176. *Proc Natl Acad Sci U S A*, 95, 3792-3797.

- Liptay, S., Schmid, R.M., Nabel, E.G. and Nabel, G.J. (1994) Transcriptional regulation of NF-kappa B2: evidence for kappa B-mediated positive and negative autoregulation. *Mol Cell Biol*, 14, 7695-7703.
- Lucas, P.C., McAllister-Lucas, L.M. and Nunez, G. (2004) NF-kappaB signaling in lymphocytes: a new cast of characters. *J Cell Sci*, 117, 31-39.
- MacLean, M. and Picard, D. (2003) Cdc37 goes beyond Hsp90 and kinases. *Cell Stress Chaperones*, 8, 114-119.
- Malhotra, V. and Wong, H.R. (2002) Interactions between the heat shock response and the nuclear factor-kappa B signaling pathway. *Crit Care Med*, 30, S89-95.
- Malinin, N.L., Boldin, M.P., Kovalenko, A.V. and Wallach, D. (1997) MAP3K-related kinase involved in NF-kappaB induction by TNF, CD95 and IL-1. *Nature*, 385, 540-544.
- Maloney, A. and Workman, P. (2002) HSP90 as a new therapeutic target for cancer therapy: the story unfolds. *Expert Opin Biol Ther*, 2, 3-24.
- Marcu, M.G., Chadli, A., Bouhouche, I., Catelli, M. and Neckers, L.M. (2000) The heat shock protein 90 antagonist novobiocin interacts with a previously unrecognized ATP-binding domain in the carboxyl terminus of the chaperone. *J Biol Chem*, 275, 37181-37186.
- Margottin, F., Bour, S.P., Durand, H., Selig, L., Benichou, S., Richard, V., Thomas, D., Strebel, K. and Benarous, R. (1998) A novel human WD protein, h-beta TrCp, that interacts with HIV-1 Vpu connects CD4 to the ER degradation pathway through an F-box motif. *Mol Cell*, 1, 565-574.
- Maruya, M., Sameshima, M., Nemoto, T. and Yahara, I. (1999) Monomer arrangement in HSP90 dimer as determined by decoration with N and C-terminal region specific antibodies. *J Mol Biol*, 285, 903-907.
- Mathas, S., Hinz, M., Anagnostopoulos, I., Krappmann, D., Lietz, A., Jundt, F., Bommert, K., Mechta-Grigoriou, F., Stein, H., Dorken, B. and Scheidereit, C. (2002) Aberrantly expressed c-Jun and JunB are a hallmark of Hodgkin lymphoma cells, stimulate proliferation and synergize with NF-kappa B. *Embo J*, 21, 4104-4113.
- Mathas, S., Lietz, A., Janz, M., Hinz, M., Jundt, F., Scheidereit, C., Bommert, K. and Dorken, B. (2003) Inhibition of NF-kappaB essentially contributes to arsenic-induced apoptosis. *Blood*, 102, 1028-1034.
- Mathas, S., Lietz, A., Anagnostopoulos, I., Hummel, F., Wiesner, B., Janz, M., Jundt, F., Hirsch, B., Johrens-Leder, K., Vornlocher, H.P., Bommert, K., Stein, H. and Dorken, B. (2004) c-FLIP mediates resistance of Hodgkin/Reed-Sternberg cells to death receptor-induced apoptosis. *J Exp Med*, 199, 1041-1052.
- McElhinny, J.A., Trushin, S.A., Bren, G.D., Chester, N. and Paya, C.V. (1996) Casein kinase II phosphorylates I kappa B alpha at S-283, S-289, S-293, and T-291 and is required for its degradation. *Mol Cell Biol*, 16, 899-906.
- McLaughlin, S.H., Smith, H.W. and Jackson, S.E. (2002) Stimulation of the weak ATPase activity of human hsp90 by a client protein. *J Mol Biol*, 315, 787-798.
- McLaughlin, S.H., Ventouras, L.A., Lobbezoo, B. and Jackson, S.E. (2004) Independent ATPase activity of Hsp90 subunits creates a flexible assembly platform. *J Mol Biol*, 344, 813-826.

- Meng, X., Devin, J., Sullivan, W.P., Toft, D., Baulieu, E.E. and Catelli, M.G. (1996) Mutational analysis of Hsp90 alpha dimerization and subcellular localization: dimer disruption does not impede 'in vivo' interaction with estrogen receptor. *J Cell Sci*, 109, 1677-1687.
- Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J., Young, D.B., Barbosa, M., Mann, M., Manning, A. and Rao, A. (1997) IKK-1 and IKK-2: cytokine-activated I $\kappa$ B kinases essential for NF- $\kappa$ B activation. *Science*, 278, 860-866.
- Meyer, P., Prodromou, C., Hu, B., Vaughan, C., Roe, S.M., Panaretou, B., Piper, P.W. and Pearl, L.H. (2003) Structural and functional analysis of the middle segment of hsp90: implications for ATP hydrolysis and client protein and cochaperone interactions. *Mol Cell*, 11, 647-658.
- Meyer, R., Hatada, E.N., Hohmann, H.P., Haiker, M., Bartsch, C., Rothlisberger, U., Lahm, H.W., Schlaeger, E.J., van Loon, A.P. and Scheidereit, C. (1991) Cloning of the DNA-binding subunit of human nuclear factor kappa B: the level of its mRNA is strongly regulated by phorbol ester or tumor necrosis factor alpha. *Proc Natl Acad Sci U S A*, 88, 966-970.
- Miller, B.S. and Zandi, E. (2001) Complete reconstitution of human I $\kappa$ B kinase (IKK) complex in yeast. Assessment of its stoichiometry and the role of IKK $\gamma$  on the complex activity in the absence of stimulation. *J Biol Chem*, 276, 36320-36326.
- Miyata, Y. and Nishida, E. (2004) CK2 controls multiple protein kinases by phosphorylating a kinase-targeting molecular chaperone, Cdc37. *Mol Cell Biol*, 24, 4065-4074.
- Muller, C.W., Rey, F.A., Sodeoka, M., Verdine, G.L. and Harrison, S.C. (1995) Structure of the NF- $\kappa$ B p50 homodimer bound to DNA. *Nature*, 373, 311-317.
- Munster, P.N., Srethapakdi, M., Moasser, M.M. and Rosen, N. (2001) Inhibition of heat shock protein 90 function by ansamycins causes the morphological and functional differentiation of breast cancer cells. *Cancer Res*, 61, 2945-2952.
- Muschen, M., Rajewsky, K., Brauninger, A., Baur, A.S., Oudejans, J.J., Roers, A., Hansmann, M.L. and Kuppers, R. (2000) Rare occurrence of classical Hodgkin's disease as a T cell lymphoma. *J Exp Med*, 191, 387-394.
- Naumann, M., Nieters, A., Hatada, E.N. and Scheidereit, C. (1993) NF- $\kappa$ B precursor p100 inhibits nuclear translocation and DNA binding of NF- $\kappa$ B/rel-factors. *Oncogene*, 8, 2275-2281.
- Nimmanapalli, R., O'Bryan, E. and Bhalla, K. (2001) Geldanamycin and its analogue 17-allylamino-17-demethoxygeldanamycin lowers Bcr-Abl levels and induces apoptosis and differentiation of Bcr-Abl-positive human leukemic blasts. *Cancer Res*, 61, 1799-1804.
- Nollen, E.A. and Morimoto, R.I. (2002) Chaperoning signaling pathways: molecular chaperones as stress-sensing 'heat shock' proteins. *J Cell Sci*, 115, 2809-2816.
- Owen, B.A., Sullivan, W.P., Felts, S.J. and Toft, D.O. (2002) Regulation of heat shock protein 90 ATPase activity by sequences in the carboxyl terminus. *J Biol Chem*, 277, 7086-7091.
- Pahl, H.L. (1999) Activators and target genes of Rel/NF- $\kappa$ B transcription factors. *Oncogene*, 18, 6853-6866.

- Palombella, V.J., Rando, O.J., Goldberg, A.L. and Maniatis, T. (1994) The ubiquitin-proteasome pathway is required for processing the NF-kappa B1 precursor protein and the activation of NF-kappa B. *Cell*, 78, 773-785.
- Panaretou, B., Siligardi, G., Meyer, P., Maloney, A., Sullivan, J.K., Singh, S., Millson, S.H., Clarke, P.A., Naaby-Hansen, S., Stein, R., Cramer, R., Mollapour, M., Workman, P., Piper, P.W., Pearl, L.H. and Prodromou, C. (2002) Activation of the ATPase activity of hsp90 by the stress-regulated cochaperone aha1. *Mol Cell*, 10, 1307-1318.
- Park, K.J., Gaynor, R.B. and Kwak, Y.T. (2003) Heat shock protein 27 association with the I kappa B kinase complex regulates tumor necrosis factor alpha-induced NF-kappa B activation. *J Biol Chem*, 278, 35272-35278.
- Pearl, L.H. (2005) Hsp90 and Cdc37 - a chaperone cancer conspiracy. *Curr Opin Genet Dev*, 15, 55-61.
- Picard, D., Khursheed, B., Garabedian, M.J., Fortin, M.G., Lindquist, S. and Yamamoto, K.R. (1990) Reduced levels of hsp90 compromise steroid receptor action in vivo. *Nature*, 348, 166-168.
- Piper, P.W. (2001) The Hsp90 chaperone as a promising drug target. *Curr Opin Investig Drugs*, 2, 1606-1610.
- Pommier, Y., Sordet, O., Antony, S., Hayward, R.L. and Kohn, K.W. (2004) Apoptosis defects and chemotherapy resistance: molecular interaction maps and networks. *Oncogene*, 23, 2934-2949.
- Pratt, W.B. (1997) The role of the hsp90-based chaperone system in signal transduction by nuclear receptors and receptors signaling via MAP kinase. *Annu Rev Pharmacol Toxicol*, 37, 297-326.
- Pratt, W.B. and Toft, D.O. (1997) Steroid receptor interactions with heat shock protein and immunophilin chaperones. *Endocr Rev*, 18, 306-360.
- Prince, T. and Matts, R.L. (2004) Definition of protein kinase sequence motifs that trigger high affinity binding of Hsp90 and Cdc37. *J Biol Chem*, 279, 39975-39981.
- Prodromou, C., Panaretou, B., Chohan, S., Siligardi, G., O'Brien, R., Ladbury, J.E., Roe, S.M., Piper, P.W. and Pearl, L.H. (2000) The ATPase cycle of Hsp90 drives a molecular 'clamp' via transient dimerization of the N-terminal domains. *Embo J*, 19, 4383-4392.
- Prodromou, C., Roe, S.M., O'Brien, R., Ladbury, J.E., Piper, P.W. and Pearl, L.H. (1997) Identification and structural characterization of the ATP/ADP-binding site in the Hsp90 molecular chaperone. *Cell*, 90, 65-75.
- Prodromou, C., Siligardi, G., O'Brien, R., Woolfson, D.N., Regan, L., Panaretou, B., Ladbury, J.E., Piper, P.W. and Pearl, L.H. (1999) Regulation of Hsp90 ATPase activity by tetratricopeptide repeat (TPR)-domain co-chaperones. *Embo J*, 18, 754-762.
- Ran, R., Lu, A., Zhang, L., Tang, Y., Zhu, H., Xu, H., Feng, Y., Han, C., Zhou, G., Rigby, A.C. and Sharp, F.R. (2004) Hsp70 promotes TNF-mediated apoptosis by binding IKK gamma and impairing NF-kappa B survival signaling. *Genes Dev*, 18, 1466-1481.

- Rao, J., Lee, P., Benzeno, S., Cardozo, C., Albertus, J., Robins, D.M. and Caplan, A.J. (2001) Functional interaction of human Cdc37 with the androgen receptor but not with the glucocorticoid receptor. *J Biol Chem*, 276, 5814-5820.
- Rayet, B. and Gelinas, C. (1999) Aberrant rel/nfkb genes and activity in human cancer. *Oncogene*, 18, 6938-6947.
- Raynal, P. and Pollard, H.B. (1994) Annexins: the problem of assessing the biological role for a gene family of multifunctional calcium- and phospholipid-binding proteins. *Biochim Biophys Acta*, 1197, 63-93.
- Reed, S.I. (1980) The selection of *S. cerevisiae* mutants defective in the start event of cell division. *Genetics*, 95, 561-577.
- Regnier, C.H., Song, H.Y., Gao, X., Goeddel, D.V., Cao, Z. and Rothe, M. (1997) Identification and characterization of an I kappa B kinase. *Cell*, 90, 373-383.
- Rice, N.R., Copeland, T.D., Simek, S., Oroszlan, S. and Gilden, R.V. (1986) Detection and characterization of the protein encoded by the v-rel oncogene. *Virology*, 149, 217-229.
- Rice, N.R., MacKichan, M.L. and Israel, A. (1992) The precursor of NF-kappa B p50 has I kappa B-like functions. *Cell*, 71, 243-253.
- Richter, K. and Buchner, J. (2001) Hsp90: chaperoning signal transduction. *J Cell Physiol*, 188, 281-290.
- Richter, K., Muschler, P., Hainzl, O., Reinstein, J. and Buchner, J. (2003) Sti1 is a non-competitive inhibitor of the Hsp90 ATPase. Binding prevents the N-terminal dimerization reaction during the atpase cycle. *J Biol Chem*, 278, 10328-10333.
- Ripley, B.J., Stephanou, A., Isenberg, D.A. and Latchman, D.S. (1999) Interleukin-10 activates heat-shock protein 90beta gene expression. *Immunology*, 97, 226-231.
- Rodriguez, M.S., Wright, J., Thompson, J., Thomas, D., Baleux, F., Virelizier, J.L., Hay, R.T. and Arenzana-Seisdedos, F. (1996) Identification of lysine residues required for signal-induced ubiquitination and degradation of I kappa B-alpha in vivo. *Oncogene*, 12, 2425-2435.
- Roe, S.M., Ali, M.M., Meyer, P., Vaughan, C.K., Panaretou, B., Piper, P.W., Prodromou, C. and Pearl, L.H. (2004) The Mechanism of Hsp90 regulation by the protein kinase-specific cochaperone p50(cdc37). *Cell*, 116, 87-98.
- Roe, S.M., Prodromou, C., O'Brien, R., Ladbury, J.E., Piper, P.W. and Pearl, L.H. (1999) Structural basis for inhibition of the Hsp90 molecular chaperone by the antitumor antibiotics radicicol and geldanamycin. *J Med Chem*, 42, 260-266.
- Roeder, R.G. (1996) The role of general initiation factors in transcription by RNA polymerase II. *Trends Biochem Sci*, 21, 327-335.
- Rothwarf, D.M. and Karin, M. (1999) The NF-kappa B activation pathway: a paradigm in information transfer from membrane to nucleus. *Sci STKE*, 5, 26.
- Rothwarf, D.M., Zandi, E., Natoli, G. and Karin, M. (1998) IKK-gamma is an essential regulatory subunit of the I kappa B kinase complex. *Nature*, 395, 297-300.
- Sato, S., Fujita, N. and Tsuruo, T. (2000) Modulation of Akt kinase activity by binding to Hsp90. *Proc Natl Acad Sci U S A*, 97, 10832-10837.

- Scheibel, T., Weikl, T. and Buchner, J. (1998) Two chaperone sites in Hsp90 differing in substrate specificity and ATP dependence. *Proc Natl Acad Sci U S A*, 95, 1495-1499.
- Schmitz, G., Schmidt, M. and Feierabend, J. (1996) Characterization of a plastid-specific HSP90 homologue: identification of a cDNA sequence, phylogenetic descendance and analysis of its mRNA and protein expression. *Plant Mol Biol*, 30, 479-492.
- Schmitz, M.L., Bacher, S. and Kracht, M. (2001) I kappa B-independent control of NF-kappa B activity by modulatory phosphorylations. *Trends Biochem Sci*, 26, 186-190.
- Schmitz, M.L., dos Santos Silva, M.A., Altmann, H., Czisch, M., Holak, T.A. and Baeuerle, P.A. (1994) Structural and functional analysis of the NF-kappa B p65 C terminus. An acidic and modular transactivation domain with the potential to adopt an alpha-helical conformation. *J Biol Chem*, 269, 25613-25620.
- Scholz, G., Hartson, S.D., Cartledge, K., Hall, N., Shao, J., Dunn, A.R. and Matts, R.L. (2000) p50(Cdc37) can buffer the temperature-sensitive properties of a mutant of Hck. *Mol Cell Biol*, 20, 6984-6995.
- Scholz, G.M., Cartledge, K. and Hall, N.E. (2001) Identification and characterization of Hrc, a novel Hsp90-associating relative of Cdc37. *J Biol Chem*, 276, 30971-30979.
- Schulte, T.W. and Neckers, L.M. (1998) The benzoquinone ansamycin 17-allylamino-17-demethoxygeldanamycin binds to HSP90 and shares important biologic activities with geldanamycin. *Cancer Chemother Pharmacol*, 42, 273-279.
- Sen, R. and Baltimore, D. (1986b) Inducibility of kappa immunoglobulin enhancer-binding protein Nf-kappa B by a posttranslational mechanism. *Cell*, 47, 921-928.
- Sen, R. and Baltimore, D. (1986a) Multiple nuclear factors interact with the immunoglobulin enhancer sequences. *Cell*, 46, 705-716.
- Senftleben, U., Cao, Y., Xiao, G., Greten, F.R., Krahn, G., Bonizzi, G., Chen, Y., Hu, Y., Fong, A., Sun, S.C. and Karin, M. (2001) Activation by IKKalpha of a second, evolutionary conserved, NF-kappa B signaling pathway. *Science*, 293, 1495-1499.
- Shao, J., Grammatikakis, N., Scroggins, B.T., Uma, S., Huang, W., Chen, J.J., Hartson, S.D. and Matts, R.L. (2001) Hsp90 regulates p50(cdc37) function during the biogenesis of the active conformation of the heme-regulated eIF2 alpha kinase. *J Biol Chem*, 276, 206-214.
- Shao, J., Irwin, A., Hartson, S.D. and Matts, R.L. (2003a) Functional dissection of cdc37: characterization of domain structure and amino acid residues critical for protein kinase binding. *Biochemistry*, 42, 12577-12588.
- Shao, J., Prince, T., Hartson, S.D. and Matts, R.L. (2003b) Phosphorylation of serine 13 is required for the proper function of the Hsp90 co-chaperone, Cdc37. *J Biol Chem*, 278, 38117-38120.
- Shinkura, R., Kitada, K., Matsuda, F., Tashiro, K., Ikuta, K., Suzuki, M., Kogishi, K., Serikawa, T. and Honjo, T. (1999) Alymphoplasia is caused by a point mutation in the mouse gene encoding Nf-kappa b-inducing kinase. *Nat Genet*, 22, 74-77.

- Siligardi, G., Panaretou, B., Meyer, P., Singh, S., Woolfson, D.N., Piper, P.W., Pearl, L.H. and Prodromou, C. (2002) Regulation of Hsp90 ATPase activity by the co-chaperone Cdc37p/p50cdc37. *J Biol Chem*, 277, 20151-20159.
- Silverstein, A.M., Grammatikakis, N., Cochran, B.H., Chinkers, M. and Pratt, W.B. (1998) p50(cdc37) binds directly to the catalytic domain of Raf as well as to a site on hsp90 that is topologically adjacent to the tetratricopeptide repeat binding site. *J Biol Chem*, 273, 20090-20095.
- Simeoni, L., Kliche, S., Lindquist, J. and Schraven, B. (2004) Adaptors and linkers in T and B cells. *Curr Opin Immunol*, 16, 304-313.
- Skinnider, B.F., Elia, A.J., Gascoyne, R.D., Trumper, L.H., von Bonin, F., Kapp, U., Patterson, B., Snow, B.E. and Mak, T.W. (2001) Interleukin 13 and interleukin 13 receptor are frequently expressed by Hodgkin and Reed-Sternberg cells of Hodgkin lymphoma. *Blood*, 97, 250-255.
- Smith, D.F., Schowalter, D.B., Kost, S.L. and Toft, D.O. (1990) Reconstitution of progesterone receptor with heat shock proteins. *Mol Endocrinol*, 4, 1704-1711.
- Solan, N.J., Miyoshi, H., Carmona, E.M., Bren, G.D. and Paya, C.V. (2002) RelB cellular regulation and transcriptional activity are regulated by p100. *J Biol Chem*, 277, 1405-1418.
- Song, H.Y., Dunbar, J.D., Zhang, Y.X., Guo, D. and Donner, D.B. (1995) Identification of a protein with homology to hsp90 that binds the type 1 tumor necrosis factor receptor. *J Biol Chem*, 270, 3574-3581.
- Soti, C., Vermes, A., Haystead, T.A. and Csermely, P. (2003) Comparative analysis of the ATP-binding sites of Hsp90 by nucleotide affinity cleavage: a distinct nucleotide specificity of the C-terminal ATP-binding site. *Eur J Biochem*, 270, 2421-2428.
- Stanger, B.Z., Leder, P., Lee, T.H., Kim, E. and Seed, B. (1995) RIP: a novel protein containing a death domain that interacts with Fas/APO-1 (CD95) in yeast and causes cell death. *Cell*, 81, 513-523.
- Staudt, L.M. (2000) The molecular and cellular origins of Hodgkin's disease. *J Exp Med*, 191, 207-212.
- Stebbins, C.E., Russo, A.A., Schneider, C., Rosen, N., Hartl, F.U. and Pavletich, N.P. (1997) Crystal structure of an Hsp90-geldanamycin complex: targeting of a protein chaperone by an antitumor agent. *Cell*, 89, 239-250.
- Stephens, R.M., Rice, N.R., Hiebsch, R.R., Bose, H.R., Jr. and Gilden, R.V. (1983) Nucleotide sequence of v-rel: the oncogene of reticuloendotheliosis virus. *Proc Natl Acad Sci U S A*, 80, 6229-6233.
- Studier, F.W. and Moffatt, B.A. (1986) Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes. *J Mol Biol*, 189, 113-130.
- Sugita, T., Tanaka, S., Murakami, T., Miyoshi, H. and Ohnuki, T. (1999) Immunosuppressive effects of the heat shock protein 90-binding antibiotic geldanamycin. *Biochem Mol Biol Int*, 47, 587-595.
- Sun, L., Deng, L., Ea, C.K., Xia, Z.P. and Chen, Z.J. (2004) The TRAF6 ubiquitin ligase and TAK1 kinase mediate IKK activation by BCL10 and MALT1 in T lymphocytes. *Mol Cell*, 14, 289-301.



- Sun, S.C., Ganchi, P.A., Ballard, D.W. and Greene, W.C. (1993) NF-kappa B controls expression of inhibitor I kappa B alpha: evidence for an inducible autoregulatory pathway. *Science*, 259, 1912-1915.
- Sun, S.C. and Ballard, D.W. (1999) Persistent activation of NF-kappaB by the tax transforming protein of HTLV-1: hijacking cellular IkappaB kinases. *Oncogene*, 18, 6948-6958.
- Supko, J.G., Hickman, R.L., Grever, M.R. and Malspeis, L. (1995) Preclinical pharmacologic evaluation of geldanamycin as an antitumor agent. *Cancer Chemother Pharmacol*, 36, 305-315.
- Takaesu, G., Kishida, S., Hiyama, A., Yamaguchi, K., Shibuya, H., Irie, K., Ninomiya-Tsuji, J. and Matsumoto, K. (2000) TAB2, a novel adaptor protein, mediates activation of TAK1 MAPKKK by linking TAK1 to TRAF6 in the IL-1 signal transduction pathway. *Mol Cell*, 5, 649-658.
- Takeda, K., Takeuchi, O., Tsujimura, T., Itami, S., Adachi, O., Kawai, T., Sanjo, H., Yoshikawa, K., Terada, N. and Akira, S. (1999) Limb and skin abnormalities in mice lacking IKKalpha. *Science*, 284, 313-316.
- Tamaru, J., Hummel, M., Zemlin, M., Kalvelage, B. and Stein, H. (1994) Hodgkin's disease with a B-cell phenotype often shows a VDJ rearrangement and somatic mutations in the VH genes. *Blood*, 84, 708-715.
- Tanaka, M., Fuentes, M.E., Yamaguchi, K., Durnin, M.H., Dalrymple, S.A., Hardy, K.L. and Goeddel, D.V. (1999) Embryonic lethality, liver degeneration, and impaired NF-kappa B activation in IKK-beta-deficient mice. *Immunity*, 10, 421-429.
- Tatebe, H. and Shiozaki, K. (2003) Identification of Cdc37 as a novel regulator of the stress-responsive mitogen-activated protein kinase. *Mol Cell Biol*, 23, 5132-5142.
- Tegethoff, S., Behlke, J. and Scheidereit, C. (2003) Tetrameric oligomerization of IkappaB kinase gamma (IKKgamma) is obligatory for IKK complex activity and NF-kappaB activation. *Mol Cell Biol*, 23, 2029-2041.
- Ten, R.M., Paya, C.V., Israel, N., Le Bail, O., Mattei, M.G., Virelizier, J.L., Kourilsky, P. and Israel, A. (1992) The characterization of the promoter of the gene encoding the p50 subunit of NF-kappa B indicates that it participates in its own regulation. *Embo J*, 11, 195-203.
- Thome, M. (2004) CARMA1, BCL-10 and MALT1 in lymphocyte development and activation. *Nat Rev Immunol*, 4, 348-359.
- Torgerson, T.R., Colosia, A.D., Donahue, J.P., Lin, Y.Z. and Hawiger, J. (1998) Regulation of NF-kappa B, AP-1, NFAT, and STAT1 nuclear import in T lymphocytes by noninvasive delivery of peptide carrying the nuclear localization sequence of NF-kappa B p50. *J Immunol*, 161, 6084-6092.
- Towbin, H., Staehelin, T. and Gordon, J. (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci U S A*, 76, 4350-4354.
- Trumper, L.H., Brady, G., Bagg, A., Gray, D., Loke, S.L., Griesser, H., Wagman, R., Brazier, R., Gascoyne, R.D., Vicini, S. and et al. (1993) Single-cell analysis of Hodgkin and Reed-Sternberg cells: molecular heterogeneity of gene expression and p53 mutations. *Blood*, 81, 3097-3115.

- Uma, S., Hartson, S.D., Chen, J.J. and Matts, R.L. (1997) Hsp90 is obligatory for the heme-regulated eIF-2 $\alpha$  kinase to acquire and maintain an activable conformation. *J Biol Chem*, 272, 11648-11656.
- Ungermann, C., Neupert, W. and Cyr, D.M. (1994) The role of Hsp70 in conferring unidirectionality on protein translocation into mitochondria. *Science*, 266, 1250-1253.
- van der Straten, A., Rommel, C., Dickson, B. and Hafen, E. (1997) The heat shock protein 83 (Hsp83) is required for Raf-mediated signalling in Drosophila. *Embo J*, 16, 1961-1969.
- Vanden Berghe, T., Kalai, M., van Loo, G., Declercq, W. and Vandenabeele, P. (2003) Disruption of HSP90 function reverts tumor necrosis factor-induced necrosis to apoptosis. *J Biol Chem*, 278, 5622-5629.
- Verma, I.M., Stevenson, J.K., Schwarz, E.M., Van Antwerp, D. and Miyamoto, S. (1995) Rel/NF-kappa B/I kappa B family: intimate tales of association and dissociation. *Genes Dev*, 9, 2723-2735.
- Voss, A.K., Thomas, T. and Gruss, P. (2000) Mice lacking HSP90 $\beta$  fail to develop a placental labyrinth. *Development*, 127, 1-11.
- Wahl, C., Liptay, S., Adler, G. and Schmid, R.M. (1998) Sulfasalazine: a potent and specific inhibitor of nuclear factor kappa B. *J Clin Invest*, 101, 1163-1174.
- Walter, S. and Buchner, J. (2002) Molecular chaperones--cellular machines for protein folding. *Angew Chem Int Ed Engl*, 41, 1098-1113.
- Waterfield, M., Jin, W., Reiley, W., Zhang, M. and Sun, S.C. (2004) I $\kappa$ B kinase is an essential component of the Tpl2 signaling pathway. *Mol Cell Biol*, 24, 6040-6048.
- Weiss, L.M., Strickler, J.G., Warnke, R.A., Purtle, D.T. and Sklar, J. (1987) Epstein-Barr viral DNA in tissues of Hodgkin's disease. *Am J Pathol*, 129, 86-91.
- Wesche, H., Henzel, W.J., Shillinglaw, W., Li, S. and Cao, Z. (1997) MyD88: an adapter that recruits IRAK to the IL-1 receptor complex. *Immunity*, 7, 837-847.
- Whitesell, L., Shifrin, S.D., Schwab, G. and Neckers, L.M. (1992) Benzoquinonoid ansamycins possess selective tumoricidal activity unrelated to src kinase inhibition. *Cancer Res*, 52, 1721-1728.
- Whitesell, L., Mimnaugh, E.G., De Costa, B., Myers, C.E., Neckers, L.M., Shifrin, S.D. and Schwab, G. (1994) Inhibition of heat shock protein HSP90-pp60v-src heteroprotein complex formation by benzoquinone ansamycins: essential role for stress proteins in oncogenic transformation. *Proc Natl Acad Sci U S A*, 91, 8324-8328.
- Wigley, D.B., Davies, G.J., Dodson, E.J., Maxwell, A. and Dodson, G. (1991) Crystal structure of an N-terminal fragment of the DNA gyrase B protein. *Nature*, 351, 624-629.
- Wood, K.M., Roff, M. and Hay, R.T. (1998) Defective I $\kappa$ B $\alpha$  in Hodgkin cell lines with constitutively active NF-kappaB. *Oncogene*, 16, 2131-2139.
- Workman, P. (2003) Auditing the pharmacological accounts for Hsp90 molecular chaperone inhibitors: unfolding the relationship between pharmacokinetics and pharmacodynamics. *Mol Cancer Ther*, 2, 131-138.

- Woronicz, J.D., Gao, X., Cao, Z., Rothe, M. and Goeddel, D.V. (1997) I $\kappa$ B kinase- $\beta$ : NF- $\kappa$ B activation and complex formation with I $\kappa$ B kinase- $\alpha$  and NIK. *Science*, 278, 866-869.
- Wulczyn, F.G., Naumann, M. and Scheidereit, C. (1992) Candidate proto-oncogene bcl-3 encodes a subunit-specific inhibitor of transcription factor NF- $\kappa$ B. *Nature*, 358, 597-599.
- Xia, Y., Makris, C., Su, B., Li, E., Yang, J., Nemerow, G.R. and Karin, M. (2000) MEK kinase 1 is critically required for c-Jun N-terminal kinase activation by proinflammatory stimuli and growth factor-induced cell migration. *Proc Natl Acad Sci U S A*, 97, 5243-5248.
- Xiao, G., Fong, A. and Sun, S.C. (2004) Induction of p100 processing by NF- $\kappa$ B-inducing kinase involves docking I $\kappa$ B kinase  $\alpha$  (IKK $\alpha$ ) to p100 and IKK $\alpha$ -mediated phosphorylation. *J Biol Chem*, 279, 30099-30105.
- Xiao, G., Harhaj, E.W. and Sun, S.C. (2001) NF- $\kappa$ B-inducing kinase regulates the processing of NF- $\kappa$ B2 p100. *Mol Cell*, 7, 401-409.
- Xu, Y. and Lindquist, S. (1993) Heat-shock protein hsp90 governs the activity of pp60v-src kinase. *Proc Natl Acad Sci U S A*, 90, 7074-7078.
- Yamaoka, S., Courtois, G., Bessia, C., Whiteside, S.T., Weil, R., Agou, F., Kirk, H.E., Kay, R.J. and Israel, A. (1998) Complementation cloning of NEMO, a component of the I $\kappa$ B kinase complex essential for NF- $\kappa$ B activation. *Cell*, 93, 1231-1240.
- Yang, J., Lin, Y., Guo, Z., Cheng, J., Huang, J., Deng, L., Liao, W., Chen, Z., Liu, Z. and Su, B. (2001) The essential role of MEK3 in TNF-induced NF- $\kappa$ B activation. *Nat Immunol*, 2, 620-624.
- Yaron, A., Hatzubai, A., Davis, M., Lavon, I., Amit, S., Manning, A.M., Andersen, J.S., Mann, M., Mercurio, F. and Ben-Neriah, Y. (1998) Identification of the receptor component of the I $\kappa$ B $\alpha$ -ubiquitin ligase. *Nature*, 396, 590-594.
- Yin, L., Wu, L., Wesche, H., Arthur, C.D., White, J.M., Goeddel, D.V. and Schreiber, R.D. (2001) Defective lymphotoxin- $\beta$  receptor-induced NF- $\kappa$ B transcriptional activity in NIK-deficient mice. *Science*, 291, 2162-2165.
- Yorgin, P.D., Hartson, S.D., Fella, A.M., Scroggins, B.T., Huang, W., Katsanis, E., Couchman, J.M., Matts, R.L. and Whitesell, L. (2000) Effects of geldanamycin, a heat-shock protein 90-binding agent, on T cell function and T cell nonreceptor protein tyrosine kinases. *J Immunol*, 164, 2915-2923.
- Young, J.C. and Hartl, F.U. (2000) Polypeptide release by Hsp90 involves ATP hydrolysis and is enhanced by the co-chaperone p23. *Embo J*, 19, 5930-5940.
- Yue, L., Karr, T.L., Nathan, D.F., Swift, H., Srinivasan, S. and Lindquist, S. (1999) Genetic analysis of viable Hsp90 alleles reveals a critical role in Drosophila spermatogenesis. *Genetics*, 151, 1065-1079.
- Zandi, E., Chen, Y. and Karin, M. (1998) Direct phosphorylation of I $\kappa$ B by IKK $\alpha$  and IKK $\beta$ : discrimination between free and NF- $\kappa$ B-bound substrate. *Science*, 281, 1360-1363.
- Zandi, E., Rothwarf, D.M., Delhase, M., Hayakawa, M. and Karin, M. (1997) The I $\kappa$ B kinase complex (IKK) contains two kinase subunits, IKK $\alpha$  and IKK $\beta$ , necessary for I $\kappa$ B phosphorylation and NF- $\kappa$ B activation. *Cell*, 91, 243-252.

Zhang, H., Wu, W., Du, Y., Santos, S.J., Conrad, S.E., Watson, J.T., Grammatikakis, N. and Gallo, K.A. (2004) Hsp90/p50cdc37 is required for mixed-lineage kinase (MLK) 3 signaling. *J Biol Chem*, 279, 19457-19463.

Zhang, S.L., Yu, J., Cheng, X.K., Ding, L., Heng, F.Y., Wu, N.H. and Shen, Y.F. (1999) Regulation of human hsp90alpha gene expression. *FEBS Lett*, 444, 130-135.

Zhao, Q., Boschelli, F., Caplan, A.J. and Arndt, K.T. (2004) Identification of a conserved sequence motif that promotes Cdc37 and cyclin D1 binding to Cdk4. *J Biol Chem*, 279, 12560-12564.

Zou, J., Guo, Y., Guettouche, T., Smith, D.F. and Voellmy, R. (1998) Repression of heat shock transcription factor HSF1 activation by HSP90 (HSP90 complex) that forms a stress-sensitive complex with HSF1. *Cell*, 94, 471-480.