

6 Literaturverzeichnis

1. Aalto M, Saksanen R, Laine P, Forsstrom R, Raikaa M, Kiviluoto M, Seppa K, Sillanaukee P: Brief intervention for female heavy drinkers in routine general practice: a 3-year randomized, controlled study. *Alc Clin Exp Res* 2000;24:1680-6.
2. Aalto M, Seppa K, Mattila P, Mustonen H, Ruuth K, Hyvarinen H, Pulkkinen H, Alho H, Sillanaukee P: Brief intervention for male heavy drinkers in routine general practice: a three-year randomized controlled study. *Alc Alcohol* 2001;36:224-30.
3. Aalto M, Pekuri P, Seppa K: Obstacles to carrying out brief intervention for heavy drinkers in primary health care: a focus group study. *Drug Alcohol Rev* 2003;22(2):169-73.
4. Allen JP, Litten RZ, Fertig JB, Babor T: Review of research on the Alcohol Use Disorder Identification Test (AUDIT). *Alc Clin Exp Res* 1997;21:613-9.
5. AWMF – Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften: Akutbehandlung alkoholbezogener Störungen. *Sucht* 2003;49(3):147-67.
6. AWMF – Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften: Riskanter schädlicher und abhängiger Alkoholkonsum: Screening, Diagnostik, Kurzintervention. *Sucht* 2004;50(2):102-12.
7. Babor TF, Higgins-Biddle JC, Saunders JB, Monteiro MG: Alcohol Use Disorder Identification Test: Guidelines for use in Primary Health Care, Second Edition, 2001. (Zugriff am 14.03.2006 auf <http://www.dass.stir.ac.uk/DRUGS/pdf/audit.pdf>)
8. Babor TF, Kadden RM: Screening and interventions for alcohol and drug problems in medical settings: what works? *J Trauma* 2005;59(3 Suppl):80-7.
9. Baker SP, O'Neill B, Haddon W Jr, Long WB: The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma* 1974;14(3):187-96.
10. Bazargan-Hejazi S, Bing E, Bazargan M, Der-Martirosian C, Hardin E, Bernstein J, Bernstein E: Evaluation of a brief intervention in an inner-city emergency department. *Ann Emerg Med* 2005;46(1):67-76.
11. Beich A, Thorsen T, Rollnick S: Screening in brief intervention trials targeting excessive drinkers in general practice: systematic review and meta-analysis. *BMJ* 2003;327:536-42.
12. Bendtsen P, Timka T: Acceptability of computerized self-report of alcohol habits: a patient perspective. *Alcohol Alcohol* 1999;34:575-80.
13. Bertholet N, Daepfen JB, Wietlisbach V, Fleming M, Burnand B: Reduction of Alcohol Consumption by Brief Alcohol Intervention in Primary Care. *Arch Intern Med* 2005;165: 986-95.
14. Bien TH, Miller WR, Tonigan JS: Brief intervention for alcohol problems: a review. *Addiction* 1993;88:315-35.

15. Bischof G, Rumpf HJ, Hapke U, Meyer C, John U: Types of natural recovery from alcohol dependence: a cluster analytic approach. *Addiction* 2003;98:1737-46.
16. Brismar B, Bergmann B: The significance of alcohol for violence and accidents. *Alc Clin Exp Res* 1998;22:299-306.
17. BMA - British Medical Association (1995): Guidelines on sensible drinking. London
18. Bundesministerium für Bildung und Forschung: Grund- und Strukturdaten 2001/2002. (Zugriff am 14.03.2006 auf http://www.bmbf.de/pub/GuS2002_ges_dt.pdf)
19. Bühringer G, Augustin R, Bergmann E, Bloomfield K, Funk W, Junge B, Kraus L, Merfert-Diete C, Rumpf HJ, Simon R, Töppich J et al.: Alkoholkonsum und alkoholbezogene Störungen. Schriftenreihe des Bundesministeriums für Gesundheit, Band 128, Baden-Baden: Nomos 2000.
20. Cloud RN, Peacock PL: Internet screening and interventions for problem drinking: Results from the www.carebetter.com pilot study. *Alcoholism Treatment Quarterly* 2001;19:23-44.
21. Chan-Pensley E: Alcohol Use Disorder Identification Test: A comparison between paper and pencil and computerized versions. *Alcohol Alcohol* 1999;34:882-5.
22. Cherpitel CJ: Drinking patterns and problems: a comparison of ER patients in an HMO and in the general population. *Alc Clin Exp Res* 1992;16(6):1104-9.
23. Cherpitel CJ: Screening for alcohol problems in the emergency department. *Ann Emerg Med* 1995;26:158-66.
24. Cherpitel CJ: Changes in Substance Use Associated with Emergency Room and Primary Care Services Utilization in the United States General Population: 1995-2000. *Am J Drug Alcohol Abuse* 2003;29(4):789-802.
25. Cherpitel CJ, Ye Y, Bond J, Rehm J, Poznyak V, Macdonald S, Stafström M, Hao W: Multi-Level analysis of alcohol-related injury among emergency department patients: a cross-national study. *Addiction* 2005;100(12):1840-50.
26. Chiauzzi E, Green TC, Lord S, Thum C, Goldstein M: My student body: a high-risk drinking prevention web site for college students. *J Am Coll Health* 2005;53(6):263-74.
27. Cryer HG: Barriers to Intervention for Alcohol Problems in Trauma Centers. *J Trauma* 2005;59:S104-S111.
28. CSAT – Center for Substance Abuse Treatment: Alcohol and Other Drug Screening of Hospitalized Trauma Patients, Treatment Improvement Protocol (TIP) Series 16, Services Administration, Center for Substance Abuse Treatment Rockwall II, 5600 Fishers Lane, Rockville, MD 20857 (DHHS Publication No. (SMA) 1995-3039.)

29. Corrao G, Bagnardi V, Zambon A, Arico S: Exploring the dose-response relationship between alcohol consumption and the risk of several alcohol-related conditions: a meta-analysis. *Addiction* 1999;94(10):1551-73.
30. Cunningham JA, Lin E, Ross HE, Walsh GW: Factors associated with untreated remissions from alcohol abuse or dependence. *Addict Behav* 2000;25:317-21.
31. D'Onofrio G, Bernstein F: Patients with alcohol problems in the Emergency Department, part 1: improving detection. SAEM Substance abuse Task Force. Society for Academic Emergency Medicine. *Acad Emerg Med* 1998;12:200-9.
32. D'Onofrio G, Degutis L: Preventive care in the emergency department: screening and brief intervention for alcohol problems in the emergency department: a systematic review. *Acad Emerg Med* 2002;9:627-38.
33. Danielsson PE, Rivara C, Heitmann E: Reasons why trauma surgeons fail to screen for alcohol problems. *Arch Surg* 1999;134:564-8.
34. Davidson P, Koziol-Mc Lain J, Harrison L, Timken D, Lowenstein SR: Intoxicated ED-patients: a 5-year follow up of morbidity and mortality. *Ann Emerg Med* 1997;30:593-7.
35. Demmel R: Motivational Interviewing: Ein Literaturüberblick. *Sucht* 2001;47:171-188.
36. DHS – Deutsche Hauptstelle für Suchtfragen: DHS – Grundsatzpapier Gender Mainstreaming in der Suchtarbeit: Chancen und Notwendigkeiten. (Zugriff am 14.03.2006 auf <http://www.dhs-intern.de/pdf/gender.pdf>.)
37. DiClemente CC: The Challenge of Change. *J Trauma* 2005;59:S3-S4.
38. DIMDI - Deutsches Institut für Medizinische Dokumentation und Information: Internationale Statistische Klassifikation der Krankheiten und verwandter Gesundheitsprobleme. 10. Revision, Version 1.3, Köln 2003 (Zugriff am 14.03.2006 auf <http://www.dimdi.de/dynamic/de/klassi/diagnosen/icd10/htmlamtl/fr-icd.htm>.)
39. Dinh-Zarr T, DiGiuseppi C, Heitman E, Roberts I: Interventions for preventing injuries in problem drinkers. *Cochrane Database Syst Rev.* 2004;(3):CD001857.
40. Dunn CW, Zatzick DF, Russo J: Hazardous drinking by trauma patients during the year after trauma. *J Trauma* 2003;54:707-12.
41. Ellickson PL, Tucker JS, Klein DJ: Ten-year prospective study of public health problems associated with early drinking. *Pediatrics* 2003;111:949-55.
42. Feunekes GI, van't Veer P, van Staren WA, Kok FJ: Alcohol intake assessment: the sober facts. *Am J Epidemiol* 1999;150: 105-12.
43. Gentilello LM: Alcohol interventions in a trauma center as a mean of reducing the risk of injury recurrence. *Ann Surg* 1999;230:473-80.

44. Graham, DM, Maio RF: Emergency physicians attitudes concerning intervention for alcohol abuse/dependence delivered in the emergency department: a brief report. *J Addict Dis* 2000;9:45-53.
45. Hannover W, Rumpf HJ, Meyer C, Hapke U, John U: Der Fragebogen zur Änderungsbereitschaft bei Alkoholkonsum (RCQ-G). In A. Glöckner-Rist, F. Rist, & H. Küfner (Hrsg.), *Elektronisches Handbuch zu Erhebungsinstrumenten im Suchtbereich (EHES)*. Version 3.00. Mannheim: Zentrum für Umfragen, Methoden und Analysen, 2003.
46. Heather N, Rollnick S, Bell A: Predictive validity of the readiness to change questionnaire. *Addiction* 1993;88:1667-77.
47. Heatherton TF, Kozlowski LT, Frecker RC, Fagerström KO (1991): The Fagerström Test for Nicotine Dependence: a revision of the Fagerström Tolerance Questionnaire. *Br J Addict* 1991;86:1119-27.
48. Heatherton TF, Kozlowski LT, Frecker RC, Rickert W, Robinson J: Measuring the heaviness of smoking: using self-reported time to the first cigarette of the day and number of cigarettes smoked per day. *Br J Addict* 1989;84:791-99.
49. Hester RK, Squires DD, Delaney HD: The Drinker's Check-up: 12-month outcomes of a controlled clinical trial of a stand-alone software program for problem drinkers. *J Subst Abuse Treat* 2005;28(2):159-69.
50. Horn K, Leontieva L, Williams JM, Furbee PM, Helmkamp JC, Manley WG 3rd: Alcohol problems among young adult emergency department patients: making predictions using routine sociodemographic information. *J Crit Care* 2002;17:212-20.
51. Hoyt DB, Coimbra R, Potenza BM: Trauma Systems, Triage, and Transport. In: Moore EE; Feliciano DV; Mattox KL: *Trauma*, 5th ed. Mc Graw Hill, New York, 2000:57-83.
52. Hungerford DW, Pollock DA: Emergency department services for patients with alcohol problems: research directions. *Acad Emerg Med* 2003;10:79-84.
53. Karlsson A, Bendtsen P (2005a): Acceptability of a computerized alcohol screening and advice routine in an emergency department setting--a patient perspective. *Addict Behav* 2005a;30(4):767-76.
54. Karlsson A, Johansson K, Nordqvist C, Bendtsen P: Feasibility of a computerized alcohol screening and personalized written advice in the ED: opportunities and obstacles. *Accid Emerg Nurs* 2005b;13(1):44-53.
55. Kox WJ, Spies CD: *Check-up Anästhesiologie*. Springer Verlag, Berlin, Heidelberg, New York, Hongkong, London, Mailand, Paris, Tokio, 2003.
56. Kypri K, Saunders JB, Williams SM, McGee RO, Langley JD, Cashell-Smith ML, Gallagher SJ: Web-based screening and brief intervention for hazardous drinking: a double-blind randomized controlled trial. *Addiction* 2004;99(11):1410-7.

57. Leontieva L, Horn K, Haque A, Helmkamp J, Ehrlich P, Williams J: Readiness to change problematic drinking assessed in the emergency department as a predictor of change. *J Crit Care* 2005;20(3):251-6.
58. Longabaugh R, Woolard RE, Nirenberg TD, Minugh AP, Becker B, Clifford PR, Carty K, Licsw, Sparadeo F, Gogineni A: Evaluating the effects of a brief motivational intervention for injured drinkers in the emergency department. *J Stud Alcohol* 2001;62:806-16.
59. Maio RF, Shope JT, Blow FC, Gregor MA, Zakrajsek JS, Weber JE, Nypaver MM: A randomized controlled trial of an emergency department-based interactive computer program to prevent alcohol misuse among injured adolescent. *Ann Emerg Med* 2005;45(4):420-9.
60. Mann KF: Neue ärztliche Aufgaben bei Alkoholproblemen. *Deutsches Ärzteblatt* 2002;99(10): A-632/B-515/C-486.
61. McDonald AJ 3rd, Wang N, Camargo CA Jr: US emergency department visits for alcohol-related diseases and injuries between 1992 and 2000. *Arch Intern Med* 2004;164(5):531-7.
62. Mello MJ, Nirenberg TD, Longabaugh R, Woolard R, Minugh A, Becker B, Baird J, Stein L: Emergency department brief motivational interventions for alcohol with motor vehicle crash patients. *Ann Emerg Med* 2005;45(6):620-5.
63. Miller WR, Rollnick S: Motivierende Gesprächsführung - ein Konzept zur Beratung von Menschen mit Suchtproblemen. Hrsg. und Deutsche Ausgabe bearbeitet von Georg Kremer Freiburg i. Brg., 1999.
64. Monti PM, Colby SM, Barnett NP, Spirito A, Rohsenow DJ, Myers M, Woolard R, Lewander W: Brief intervention for harm reduction with alcohol-positive older adolescents in a hospital emergency department. *J Consult Clin Psychol* 1999;67:989-94.
65. Moore EE: Alcohol and trauma: the perfect storm. *J Trauma* 2005;59(3 Suppl):S53-6.
66. Moyer A, Finney JW, Swearingen CE, Vergun P: Brief interventions for alcohol problems: a meta-analytic review of controlled investigations in treatment-seeking and non-treatment-seeking populations. *Addiction* 2002;97:279-92.
67. Neighbors C, Larimer ME, Lewis MA: Targeting misperceptions of descriptive drinking norms: efficacy of a computer-delivered personalized normative feedback intervention. *J Consult Clin Psychol* 2004;72(3):434-47.
68. Neumann T, Neuner B, Gentilello LM, Weiss-Gerlach E, Mentz H, Rettig JS, Schröder T, Wauer H, Müller C, Schütz M, Mann K, Siebert G, Dettling M, Müller JM, Kox WJ, Spies CD: Gender differences in the performance of a computerized version of the alcohol use disorders identification test in subcritically injured patients who are admitted to the emergency department. *Alcohol Clin Exp Res* 2004;28(11):1693-701.
69. Neumann T, Kox W, Spies CD: The anesthesiologist's approach to patients with addictive disorders. *Anästhesiologisches Vorgehen bei Sucht- und Begleiterkrankungen in Suchtmedizin in Forschung und Praxis* 2003;5:13-20.

70. Neuner B, Miller PM, Felsmann B, Weiss-Gerlach E, Neumann T, Wernecke KD, Spies CD: Health-related quality of life in urban surgical emergency department patients: comparison with a representative German population sample. *Health Qual Life Outcomes* 2005;3:77.
71. Piccinelli M, Tessari E, Bortolomasi M, Piasere O, Semenzin M, Garzotto N: Efficacy of the alcohol use disorders identification test as a screening tool for hazardous alcohol intake and related disorders in primary care: a validity study. *BMJ* 1997;314:420-4.
72. Rehm J: Measuring quantity, frequency and volume of drinking. *Alcohol Clin Exp Res* 1998;22(2 Suppl):4S-14S
73. Rehm J, Patra J, Popova S: Alcohol-attributable mortality and potential years of life lost in Canada 2001: implications for prevention and policy. *Addiction* 2006;101(3):373-84.
74. Reinert DF, Allen JP: The Alcohol Use Disorder Identification Test (AUDIT): A review of recent research. *Alcohol Clin Exp Res* 2002;26:272-9.
75. Rhodes KV, Lauderdale DS, Stocking CB, Howes DS, Roizen MF, Levinson W: Better health while you wait: a controlled trial of a computer-based intervention for screening and health promotion in the emergency department. *Ann Emerg Med* 2001;37(3):284-91.
76. Roche AM, Freeman T, Skinner N: From data to evidence, to action: Findings from a systematic review of hospital screening studies for high risk alcohol consumption. *Drug Alcohol Depend.* in Druck, angenommen 11/2005 (Stand 03/2006).
77. Rollnick S, Heather N, Gold R, Hall W: Development of a short 'readiness to change' questionnaire for use in brief, opportunistic interventions among excessive drinkers. *Br J Addict* 1992;87:743-54.
78. Rumpf HJ, Meyer C, Hapke U, John U: Deutsche Version des Alcohol Use Disorders Identification Test (AUDIT-G-L). In A. Glöckner-Rist, F. Rist, & H. Küfner (Hrsg.), *Elektronisches Handbuch zu Erhebungsinstrumenten im Suchtbereich (EHES)*. Version 3.00. Mannheim: Zentrum für Umfragen, Methoden und Analysen, 2003.
79. Runge JW, Hargazun S (2001): Recommended Best Practice of Emergency Medical Care for the Alcohol-Impaired Patient: Screening and Brief Intervention for Alcohol Problems in the Emergency Department, 2001. (Zugriff am 14.03.2006 auf <http://www.nhtsa.dot.gov/people/injury/alcohol/EmergCare/recommended.htm>.)
80. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M: Development of the Alcohol Use Disorder Identification Test (AUDIT): WHO Collaborated Project on Early Detection of Persons with Harmful Alcohol Consumption--II. *Addiction* 1993;88:791-804.
81. Saunders JB, Kypri K: Approaches to Brief Intervention for Hazardous Drinking in Young People. *Alcohol Clin Exp Res* 2004;28(2):322-9.
82. Schermer CR, Moyers TB, Miller WR, Bloomfield LA: Trauma center brief interventions for alcohol disorders decrease subsequent driving under the influence arrests. *J Trauma* 2006;60(1):29-34.

83. Schinke SP, Schwinn TM, Di Noia J, Cole KC: Reducing the risks of alcohol use among urban youth: three-year effects of a computer-based intervention with and without parent involvement. *J Stud Alcohol* 2004;65(4):443-9.
84. Schmidt LG: Frühdiagnostik und Kurzintervention beim beginnenden Alkoholismus. *Deutsches Ärzteblatt* 1997;94(44):A-2905/B-2481/C-2305.
85. Schumann A, Rumpf HJ, Meyer C, Hapke U, John U: Deutsche Version des Fagerström-Test for Nicotine Dependence (FTND-G) und des Heaviness of Smoking Index (HSI-G). In A. Glöckner-Rist, F. Rist, & H. Kufner (Hrsg.), *Elektronisches Handbuch zu Erhebungsinstrumenten im Suchtbereich (EHES)*. Version 3.00. Mannheim: Zentrum für Umfragen, Methoden und Analysen, 2003.
86. Seppä K, Makela R, Sillanaukee P: Effectiveness of the Alcohol Use Disorders Identification Test in occupational health screenings. *Alcohol Clin Exp Res* 1995;19:999-1003.
87. SIGN - Scottish Intercollegiate Guidelines Network: Management of harmful drinking and alcohol dependence in primary care. A national clinical guideline, 2004. (Zugriff am 14.03.2006 auf <http://www.sign.ac.uk/pdf/sign74.pdf>.)
88. Sobell LC, Maisto SA, Sobell MB, Cooper AM: Reliability of alcohol abusers' self-reports of drinking behavior. *Behav Res Ther* 1979;17:157-60.
89. Soderstrom CA, Dischinger PC, Kerns TJ, Kufera JA, McDuff DR, Gorelick DA, Smith GS: Screening trauma patients for alcoholism according to NIAAA guidelines with alcohol use disorders identification test questions. *Alcohol Clin Exp Res* 1998;22:1470-5.
90. Soderstrom CA, Cole FJ, Porter JM, Injury Control and Violence Prevention Committee: Injury in America: the role of alcohol and other drugs- an EAST position paper prepared by the injury control and violence preventing committee. *J Trauma* 2001;50(1):1-12.
91. Spirito A, Monti PM, Barnett NP, Colby SM, Sindelar H, Rohsenow DJ, Lewander W, Myers M: A randomized clinical trial of a brief motivational intervention for alcohol-positive adolescents treated in an emergency department. *J Pediatr* 2004;145(3):396-402.
92. Spies CD, Neuner B, Neumann T, Blum S, Müller C, Rommelspacher H, Rieger A, Sanft C, Specht M, Hannemann L, Striebel HW, Schaffartzik W.: Intercurrent complications in chronic alcoholic man admitted to the intensive care unit following trauma. *Intensive Care Med* 1996;22(4):286-93.
93. Statistisches Bundesamt Wiesbaden: Alterspyramide 2001. (Zugriff am 14.03.2006 auf <http://www.destatis.de/basis/d/bevoe/bevoegra2.htm>.)
94. Veltrup C: Motivationale Interventionen bei Menschen mit Alkoholproblemen. In *Deutsche Hauptstelle gegen Suchtgefahren* (Hrsg.) *Alkoholabhängigkeit-Motivation und Diagnose*. Lambertus, Freiburg i. Brg., 1996.
95. Williams S, Brown A, Patton R, Crawford MJ, Touquet R: The half of the "teachable moment" for alcohol misusing patients in the emergency department. *Drug Alcohol Depend* 2005;77:205-8.

96. WHO - World Health Organization: International Guide for Monitoring Alcohol Consumption and Related Harm, 2000. (Zugriff am 14.03.2006 auf http://whqlibdoc.who.int/hq/2000/WHO_MSD_MSB_00.4.pdf.)
97. WHO - World Health Organization: WHO Collaborative Study on Alcohol and Injuries. Protocol, 2001. (Zugriff am 14.03.2006 auf http://www.who.int/substance_abuse/activities/en/InjuriesProtocolEnglish.pdf.)
98. WHO - World Health Organization: Global Status Report on Alcohol 2004. (Zugriff am 14.03.2006 auf http://www.who.int/substance_abuse/publications/global_status_report_2004_overview.pdf.)