

6. REFERENCES

- Alexander R.W., Muralikrishna P. and Cooperman B.S.** (1994) Ribosomal components neighboring the conserved 518-533 loop of 16S rRNA in 30S subunits. *Biochemistry*, 33: 12109-12118
- Altamura S., Sanz J.L., Amils R., Cammarano P. and Londei, P.** (1988) The antibiotic sensitivity spectra of ribosomes from the thermoproteales phylogenetic depth and distribution of antibiotic binding sites. *Syst. Appl. Microbiol.*, 10: 218-225
- Auerbach T., Pioletti M., Avila H., Anagnostopoulos K., Weinstein S., Franceschi F. and Yonath A.** (2000) Genetic and biochemical manipulations of the small ribosomal subunit from *Thermus thermophilus* HB8. *Journal of Biomolecular Structure & Dynamics*, 17(4): 617-628
- Ban N., Nissen P., Hansen J., Moore P.B. and Steitz T.A.** (2000) The complete atomic structure of the large ribosomal subunit at 2.4 angstrom resolution. *Science*, 289: 905-920
- Berkhout B., van der Laken C.J. and van Knippenberg P.H.** (1986) Formylmethionyl-tRNA binding to 30S ribosomes programmed with homopolynucleotides and the effect of translational initiation factor 3. *Biochimica et Biophysica Acta*. 866(2-3): 144-153
- Bhangu R. and Wollenzien P.** (1992) The mRNA binding track in the *Escherichia coli* ribosome for mRNAs of different sequences. *Biochemistry*, 31: 5937-5944
- Biou V., Shu F. and Ramakrishnan V.** (1995) X-ray crystallography shows that translational initiation factor IF3 consists of two compact alpha/beta domains linked by an alpha-helix. *EMBO Journal*, 14(16): 4056-4064
- Bradford M.M.** (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72: 248-254.
- Brock T.D. and Freeze H.** (1969) *Thermus aquaticus* gen.n. and sp.n., a non-sprulating extreme thermophile. *J. Bacteriology*, 98: 289-297
- Brodersen D.E, Clemons W.M, Carter A.P, Morgan-Warren R.J., Wimberly B.T. and Ramakrishnan V.** (2000) The structural basis for the action of the antibiotics tetracycline, pactamycin, and hygromycin B on the 30S ribosomal subunit. *Cell*, 103: 1143-1154

- Bruhns J. and Gualerzi C.** (1980) Structure-function relationship in *Escherichia coli* initiation factors: role of tyrosine residues in ribosomal binding and functional activity of IF3. *Biochemistry*, 19(8): 1670-1676
- Burkhardt N., Junemann R., Spahn CM. And Nierhaus K.H.** (1998) Ribosomal tRNA binding sites: three-site models of translation. *Critical Reviews in Biochemistry Molecular Biology*, 33(2): 95-149
- Butler J.S., Springer M., Dondon J., Graffe M. and Grunberg-Manago M.** (1986) *Escherichia coli* protein synthesis initiation factor IF3 controls its own gene expression at the translational level in vivo. *Journal of Molecular Biology*, 192(4): 767-780
- Butler J.S., Springer M., Grunberg-Manago M.** (1987) AUU-to-AUG mutation in the initiator codon of the translation initiation factor IF3 abolishes translational autocontrol of its own gene (*infC*) in vivo. *Proc. Nat. Ac. Sci. USA*, 84(12): 4022-4025
- Carter A., Clemons W., Brodersen D., Morgan-Warren R., Wimberly B. and Ramakrishnan, V.** (2000) Functional insights from the structure of the 30S ribosomal subunit and its interactions with antibiotics. *Nature*, 407: 340-348
- Carter A., Clemons W., Brodersen D., Morgan-Warren R., Hartsch T. Wimberly B. and Ramakrishnan V.** (2001) Crystal structure of an initiation factor bound to the 30S ribosomal subunit. *Science*, 291: 498-501
- Cate J.H., Yusupov M.M., Yusupova G.Z., Earnest T.N. and Noller H.F.** (1999) X-ray crystal structure of 70S ribosome functional complexes. *Science*, 285: 2095-2104
- Choli T., Franceschi F., Yonath A. and Wittmann-Liebold B.** (1993) Isolation and characterization of a new ribosomal protein from the thermophilic eubacteria, *Thermus thermophilus*, *T. aquaticus* and *T. flavus*. *Biological Chemistry Hoppe-Seyler*, 374(6): 377-383
- Cline J., Braman J.C. and Hogrefe, H.H.** (1996) PCR fidelity of *Pfu* DNA polymerase and other thermostable DNA polymerases. *Nucleic Acids Reserch*, 24: 3546-3551
- Cummings H.S. and Hershey J.W.** (1994) Translation initiation factor IF1 is essential for cell viability in *Escherichia coli*. *Journal of Bacteriology*, 176(1): 198-205
- Dantley K.A., Dannelly H.K. and Burdett V.** (1998) Binding interaction between Tet(M) and the ribosome: requirements for binding. *J. Bacteriol.*, 180: 4089-4092
- De Bellis D., Liveris D., Goss D., Ringquist S. and Schwartz I.** (1992) Structure-function analysis of *Escherichia coli* translation initiation factor IF3: tyrosine 107 and lysine 110 are required for ribosome binding. *Biochemistry*, 31(48): 11984-11990

de Cock E., Springer M., Dardel F. (1999) The interdomain linker of *Escherichia coli* initiation factor IF3: a possible trigger of translation initiation specificity. *Molecular Microbiology*, 32(1): 193-202

Deisenhofer J. (1981) Crystallographic refinement and atomic models of a human Fc fragment and its complex with fragment B of protein A from *Staphylococcus aureus* at 2.9- and 2.8-Å resolution. *Biochemistry*, 20(9): 2361-2370

Demeshkina N., Repkova M., Ven'yaminova A., Graifer D. and Karpova G. (2000) Nucleotides of 18S rRNA surrounding mRNA codons at the human ribosomal A, P, and E sites: A crosslinking study with mRNA analogs carrying an aryl azide group at either the uracil or the guanine residue. *RNA*, 6: 1727-1736

Dower W.J., Miller J.F. and Ragsdale C.W. (1988) High efficacy transformation of *E. coli* by high voltage electroporation. *Nucleic Acids Research*, 16(13): 6127-6145

Dunford R.P. and Dyer T.A. (1997) A method for preparing PCR products for restriction-enzymes-directed cloning without enzymatic digestion. *Technical Tips Online*.

Ehresmann C., Moine H., Mougél M., Dondon J., Grunberg-Manago M., Ebel J.P. and Ehresmann B. (1986) Cross-linking of initiation factor IF3 to *Escherichia coli* 30S ribosomal subunit by trans-diamminedichloroplatinum(II): characterization of two cross-linking sites in 16S rRNA; a possible way of functioning for IF3. *Nucleic Acids Research*, 14(12): 4803-4821

Eikenberry E.F., Bickle T.A., Traut R.R. and Price C.A. (1970) Separation of large quantities of ribosomal subunits by zonal ultracentrifugation. *European Journal of Biochemistry*, 12(1): 113-116

Eisenstein M., Shariv I., Koren G., Friesem A.A. and Katchalski-Katzir E. (1997) Modeling supra-molecular helices: extension of the molecular surface recognition algorithm and application to the protein coat of the tobacco mosaic virus. *J. Mol. Biol.*, 266: 135-143

Firpo M.A., Connelly M.B., Goss D.J. and Dahlberg AE. (1996) Mutations at two invariant nucleotides in the 3'-minor domain of *Escherichia coli* 16 S rRNA affecting translational initiation and initiation factor 3 function. *Journal of Biological Chemistry*, 271(9): 4693-4698

Fourmy D., Recht M.I., Blanchard S.C. and Puglisi J.D. (1996) Structure of the A site of *Escherichia coli* 16S ribosomal RNA complexed with an aminoglycoside antibiotic. *Science*, 274(5291): 1367-1371

Gabashvili I.S., Agrawal R.K., Grassucci R. and Frank J. (1999a) Structure and structural variations of the *Escherichia coli* 30S ribosomal subunit as revealed by three-dimensional cryo- electron microscopy. *J. Mol. Biol.*, 286: 1285-1291

Gabashvili I.S., Agrawal R.K., Grassucci R., Squires C.L., Dahlberg A.E. and Frank J. (1999b) Major rearrangements in the 70S ribosomal 3D structure caused by a conformational switch in 16S ribosomal RNA. *Embo J.*, 18: 6501-6507

Gale E.F., Cundliffe E., Reynolds P.E., Richmond M.H. and Warning M.J. (1981) The molecular basis of antibiotic action. London: John Wiley & Sons.

Garcia C., Fortier P.L., Blanquet S., Lallemand J.Y. and Dardel F. (1995a) 1H and 15N resonance assignments and structure of the N-terminal domain of *Escherichia coli* initiation factor 3. *European Journal of Biochemistry*, 228(2): 395-402

Garcia C., Fortier P.L., Blanquet S., Lallemand J.Y., Dardel F. (1995b) Solution structure of the ribosome-binding domain of *E. coli* translation initiation factor IF3. Homology with the U1A protein of the eukaryotic spliceosome. *Journal of Molecular Biology*, 254(2): 247-259

Gill S.C. and von Hippel P.H. (1989) Calculation of protein extinction coefficients from amino acid sequence data. *Analytical Biochemistry*, 182(2): 319-326

Gilliland G.L., Tung M., Blakeslee D.M. and Ladner J. (1994) The biological macromolecule crystallization database, version 3.0: new feature, data, and the NASA archive for protein crystal growth data. *Acta Crystallographica-D*, 50: 408-413

Godefroy-Colburn T., Wolfe A.D., Dondon J., Grunberg-Manago M., Desse P. and Pantaloni D. (1975) Light-scattering studies showing effect of initiation-factors on reversible dissociation of *Escherichia coli* ribosomes. *J. Mol. Biol.*, 94(3): 461-478

Gold L., Stormo G. and Saunders R. (1984) *Escherichia coli* translational initiation factor IF3: a unique case of translational regulation. *Proc. Nat. Ac. Sci. USA*, 81(22): 7061-7065

Goldman R.A., Hasan T., Hall C.C., Strycharz W.A. and Cooperman B.S. (1983) Photoincorporation of tetracycline into *Escherichia coli* ribosomes. Identification of the major proteins photolabeled by native tetracycline and tetracycline photoproducts and implications for the inhibitory action of tetracycline on protein synthesis. *Biochemistry*, 22(2): 359-368

Gordon J. (1969) Hydrolysis of guanosine 5'-triphosphate associated with binding of aminoacyl transfer ribonucleic acid to ribosomes. *J. Biol. Chem.*, 244: 5680-5686

Gualerzi C. and Pon C.L. (1990) Initiation of mRNA translation in prokaryotes. *Biochemistry*, 29: 5881-5889

Gutell R.R., Subashchandran S., Schnare M., Du Y., Lin N., Madabusi L., Muller K., Pande N., Yu N., Shang Z., Date S., Konings D., Schweiker V., Weiser B. and Cannone J.J. (1999). Comparative sequence analysis and the prediction of RNA structure, and the Web. <http://www.rna.icmb.utexas.edu/>

Harms J., Tocilj A., Levin I., Agmon I., Kölln I., Stark H., van Heel M., Cuff M., Schlünzen F., Bashan A., Franceschi F. and Yonath A. (1999) Elucidating the medium resolution structure of ribosomal particles: an interplay between electron-cryo-microscopy and X-ray crystallography. *Structure*, 7: 931-941

Hartz D., Binkley J., Hollingsworth T. and Gold L. (1990) Domains of initiator tRNA and initiation codon crucial for initiator tRNA selection by *Escherichia coli* IF3. *Genes & Development*, 4(10): 1790-1800

Hartz D., McPheeters D.S. and Gold L. (1989) Selection of the initiator tRNA by *Escherichia coli* initiation factors. *Genes & Development*, 3(12A): 1899-1912

Held W.A., Ballou B., Mizushima S and Nomura M. (1974) Assembly mapping of 30S ribosomal-proteins from *Escherichia coli* - Further studies. *J. Biol. Chem.*, 249: 3103-3111

Herrera F., Moreno N. and Martinez J.A. (1984) Increased ribosomal affinity for messenger-RNA causes resistance to edeine in a mutant of *Saccharomyces cerevisiae*. *Eur. J. Biochem.*, 145: 339-344

Hill W.E., Camp D.G., Tappich W.E. and Tassanakajohn A. (1988) Probing ribosome structure and function using short oligodeoxyribonucleotides. *Methods in Enzymol.*, 164: 401-419

Jerinic O. and Joseph S. (2000) Conformational changes in the ribosome induced by translational miscoding agents. *J. Mol. Biol.*, 304: 707-713

Kaji A., Teyssier E. and Hirokawa G. (1998) Disassembly of the post-termination complex and reduction of translational error by ribosome recycling factor (RRF)-A possible new target for antibacterial agents. *Biochemical & Biophysical Res. Com.*, 250(1): 1-4

Kim K.K. Min K. and Suh S.W. (2000) Crystal structure of the ribosome recycling factor from *Escherichia coli*. *EMBO Journal*, 19(10):2362-70

Kozak M. (1999) Initiation of translation in prokaryotes and eukaryotes. *Gene*, 234(2): 187-208

Kozak M. and Shatkin A.J. (1978) Migration of 40S ribosomal subunits on messenger RNA in the presence of edeine. *J. Biol. Chem.*, 253: 6568-6577

Kurylo-Borowska Z. (1975) Biosynthesis of Edeine: II. Localization of edeine synthetase within *Bacillus brevis* Vm4. *Biochim. Biophys. Acta*, 399: 31-41

La Teana A., Pon C.L. and Gualerzi C.O. (1993) Translation of mRNAs with degenerate initiation triplet AUU displays high initiation factor 2 dependence and is subject to initiation factor 3 repression. *Proc. Nat. Ac. Sci. USA*, 90(9): 4161-4165

Lodmell J.S. and Dahlberg A.E. (1997) A conformational switch in *Escherichia coli* 16S ribosomal RNA during decoding of messenger RNA. *Science*, **277**, 1262-1267

MacKeen L.A., Kahan L., Wahba A.J. and Schwartz I. (1980) Photochemical cross-linking of initiation factor 3 to *Escherichia coli* 30S ribosomal subunits. *Journal of Biological Chemistry*, 255(21): 10526-10531

Marmur J. (1961) A procedure for the isolation of deoxyribonucleic acid from microorganisms. *J. Mol. Biol.*, 3: 208-218

McCutcheon J.P., Agrawal R.K., Philips S.M., Grassucci R.A., Gerchman S.E., Clemons W.M., Ramakrishnan V. and Frank J. (1999) Location of translational initiation factor IF3 on the small ribosomal subunit. *Proc. Nat. Acad. Sci. USA*, 96(8): 4301-4306

Meinzel T., Sacerdot C., Graffe M., Blanquet S. and Springer M. (1999) Discrimination by *Escherichia coli* initiation factor IF3 against initiation on non-canonical codons relies on complementarity rules. *J. Mol. Biol.*, 290: 825-837

Moazed D. and Noller H.F. (1987) Interaction of antibiotics with functional sites in 16S ribosomal RNA. *Nature*, 327(6121): 389-394

Moazed D., Samaha R.R., Gualerzi C. and Noller H.F. (1995) Specific protection of 16S rRNA by translational initiation factors. *Journal of Molecular Biology*, 248(2): 207-210

Moreau M., de Cock E., Fortier P.L., Garcia C., Albaret C., Blanquet S., Lallemand J.Y. and Dardel F. (1997) Heteronuclear NMR studies of *E. coli* translation initiation factor IF3. Evidence that the inter-domain region is disordered in solution. *Journal of Molecular Biology*, 266(1): 15-22

Moreno J.M., Drskjotersen L., Kristensen J.E., Mortensen K.K. and Sperling-Petersen H.U. (1999) Characterization of the domains of *E. coli* initiation factor IF2 responsible for recognition of the ribosome. *FEBS Letters*, 455(1-2): 130-134

Mueller F. and Brimacombe R. (1997) A new model for the three-dimensional folding of *Escherichia coli* 16S ribosomal RNA. I. Fitting the RNA to a 3D electron microscopic map at 20 Å. *Journal of Molecular Biology*, 271(4): 524-544

Mullis K., Faloona F., Scharf S., Saiki R., Horn G. and Erlich H. (1986) Specific enzymatic amplification of DNA in vitro: the polymerase chain reaction. *Cold Spring Harbor Symposia on Quantitative Biology*, 51: 263-273

Nakamura Y., Ito K. and Ehrenberg M. (2000) Mimicry grasps reality in translation termination. *Cell*, 101(4): 349-352

Neu H.C. (1992) The crisis in antibiotic resistance. *Science*, 257(5073): 1064-1073

Nissen P., Hansen J., Ban N., Moore P.B. and Steitz T.A. (2000) The structural basis of ribosome activity in peptide bond synthesis. *Science*, 289: 920-930

Noah J.W., Dolan M.A., Babin P. and Wollenzien P. (1999) Effects of tetracycline and spectinomycin on the tertiary structure of ribosomal RNA in the *Escherichia coli* 30 S ribosomal subunit. *Journal of Biological Chemistry*, 274(23): 16576-16581

Nowotny V. and Nierhaus K.H. (1988) Assembly of the 30S subunit from *Escherichia coli* ribosomes occurs via 2 assembly domains which are initiated by S4 and S7. *Biochemistry*, 27: 7051-7055

Oakes, M.I., Clark, M.W., Henderson, E. and Lake, J.A. (1986) DNA hybridization electron microscopy: ribosomal RNA nucleotides 1392-1407 are exposed in the cleft of the small subunit. *Proc. Nat. Acad. Sci. USA*, 83(2): 275-279

O'Connor M., Brunelli C.A., Firpo M.A., Gregory S.T., Lieberman K.R., Lodmell J.S., Moine H., VanRyk D.I. and Dahlberg A.E. (1995) Genetic probes of ribosomal RNA function. *Biochem. Cell Biol.*, 73: 859-868

O'Connor M., Thomas C.L., Zimmermann R.A. and Dahlberg A.E. (1997) Decoding fidelity at the ribosomal A and P sites: Influence of mutations in three different regions of the decoding domain in 16S rRNA. *Nucleic Acids Res.*, 25: 1185-1193

Odon O.W., Kramer G., Henderson A.B., Pinphanichakarn P. and Hardesty, B. (1978) GTP hydrolysis during methionyl-tRNA^f binding to 40 S ribosomal subunits and the site of edeine inhibition. *J. Biol. Chem.*, 253: 1807-1816

Oehler R., Polacek N., Steiner G. and Barta A. (1997) Interaction of tetracycline with RNA: photoincorporation into ribosomal RNA of *Escherichia coli*. *Nucleic Acids Research*, 25(6): 1219-1224

Ohsawa H. and Gualerzi C. (1981) Structure-function relationship in *Escherichia coli* initiation factors. Identification of a lysine residue in the ribosomal binding site of initiation factor by site-specific chemical modification with pyridoxal phosphate. *Journal of Biological Chemistry*, 256(10): 4905-4912

Orth P., Saenger W. and Hinrichs W. (1999) Tetracycline-chelated Mg²⁺ ion initiates helix unwinding in Tet repressor induction. *Biochemistry*, 38: 191-198

Oshima T. and Imahori K. (1971) Isolation of an extreme thermophile and thermostability of its transfer ribonucleic acid and ribosome. *J. Gen. Appl. Microbiol.*, 17: 513-517

Otwinowski Z. and Minor W. (1997) Processing of X-ray diffraction data collected in oscillation mode. *Macromolecular Crystallography Pt A*, 276: 307-326

Pavlov M.Y., Freistroffer D.V., Heurgue-Hamard V., Buckingham R.H. and Ehrenberg M. (1997) Release factor RF3 abolishes competition between release factor RF1 and ribosome recycling factor (RRF) for a ribosome binding site. *Journal of Molecular Biology*, 273(2): 389-401

Ricker R.D. and Kaji A. (1991) Use of single-stranded DNA oligonucleotides in programming ribosomes for translation. *Nucl Acid Res.*, 19: 6573-6578

Ringquist S., Cunningham P., Weitzmann C., Formenoy L., Pleij C., Ofengand J. and Gold L. (1993) Translation initiation complex-formation with 30S ribosomal particles mutated at conserved positions in the 3'-minor domain of 16S RNA. *J. Mol. Biol.*, 234: 14-27

Risuleo G., Gualerzi C. and Pon C. (1976) Specificity and properties of the destabilisation, induced by initiation factor IF3, of ternary complexes of the 30S ribosomal subunit, aminoacyl-tRNA and polynucleotides. *European Journal of Biochemistry*, 67(2): 603-613

Roberts M.C. (1996) Tetracycline resistance determinants: mechanisms of action, regulation of expression, genetic mobility, and distribution. *FEMS Microbiol. Rev.*, 19: 1-24

Ross J.I., Eady E.A., Cove J.H. and Cunliffe W.J. (1998) 16S rRNA mutation associated with tetracycline resistance in a gram-positive bacterium. *Antimicrobial Agents & Chemotherapy*, 42(7): 1702-1705

Sacerdot C., Fayat G., Dessen P., Springer M., Plumbridge J.A., Grunberg-Manago M. and Blanquet S. (1982) Sequence of a 1.26 Kb DNA fragment containing the structural gene for *E. coli* initiation factor IF3: presence of an AUU initiator codon. *EMBO Journal*, 1(3): 311-315

Santer M., Bennettguerrero E., Byahatti S., Czarnecki S., Oconnell D., Meyer M., Khoury J., Cheng X., Schwartz I. and McLaughlin J. (1990) Base changes at position-792 of *Escherichia coli* 16S ribosomal- RNA affect assembly of 70S ribosomes. *Proc. Nat. Acad. Sci. USA*, 87: 3700-3704

Schlünzen F., Gluehmann M., Janell D., Levin I., Bashan A., Harms J., Bartels H., Auerbach T., Pioletti M., Avila H., Anagnostopoulos K., Hansen H.A.S., Bennett W.S., Agmon I., Kessler M., Tocilj A., Peretz M., Weinstein S., Franceschi F. and Yonath A. (1999) The identification of selected components in electron density maps of prokaryotic ribosomes at 7 Å resolution. *J. Syn. Radiation*, 6: 928-941

Schlünzen F., Tocilj A., Zarivach R., Harms J., Gluehmann M., Janell D., Bashan A., Bartels H., Agmon I., Franceschi F., Yonath A. (2000) Structure of functionally activated small ribosomal subunit at 3.3 Å resolution. *Cell*, 102(5): 615-623

Sergiev P.V., Lavrik I.N., Wlasoff V.A., Dokudovskaya S.S., Dontsova O.A., Bogdanov A.A. and Brimacombe R. (1997) The path of mRNA through the bacterial ribosome: A site-directed crosslinking study using new photoreactive derivatives of guanosine and uridine. *RNA*, 3: 464-475

Sette M., Spurio R., van Tilborg P., Gualerzi C.O. and Boelens R. (1999) Identification of the ribosome binding sites of translation initiation factor IF3 by multidimensional heteronuclear NMR spectroscopy. *Rna*, 5(1): 82-92

Shapkina T.G., Dolan M.A., Babin P. and Wollenzien P. (2000) Initiation factor 3 induced structural changes in the 30S ribosomal subunit and in complexes containing tRNA(f)(Met) and mRNA. *Journal of Molecular Biology*, 299(3): 615-628

Spahn C.M.T. and Prescott C.D. (1996) Throwing a spanner in the works: Antibiotics and the translation apparatus. *J. Mol. Med.*, 74: 423-439

Srivastava S., Verschoor A. and Frank J. (1992) Eukaryotic initiation factor 3 does not prevent association through physical blockage of the ribosomal subunit-subunit interface. *Journal of Molecular Biology*, 226(2): 301-304

Stark H., Orlova E.V., Rinke-Appel J., Junke N., Mueller F., Rodnina M., Wintermeyer W., Brimacombe R. and van Heel M. (1997) Arrangement of tRNAs in pre- and post-translocational ribosomes revealed by electron cryomicroscopy. *Cell*, 88(1): 19-28

Studier F.W., Rosenberg A.H., Dunn J.J. and Dubendorff J.W. (1990) Use of T7 RNA polymerase to direct expression of cloned genes. *Meth. Enzymol.*, 185: 60-89

Tapprich W.E., Goss D.J. and Dahlberg A.E. (1989) Mutation at position 791 in *Escherichia coli* 16S ribosomal-RNA affects processes involved in the initiation of protein synthesis. *Proc. Nat. Acad. Sci. USA*, 86: 4927-4931

Tapprich W.E. and Hill W.E. (1986) Involvement of bases 787-795 of *E. coli* 16S rRNA in ribosomal subunit association. *Proc. Nat. Acad. Sci. USA*, 83: 556-560

Tedin K., Moll I., Grill S., Resch A., Graschopf A., Gualerzi C.O. and Blasi U. (1999) Translation initiation factor 3 antagonizes authentic start codon selection on leaderless mRNAs. *Mol. Microbiol.*, 31: 67-77

Tocilj A., Schlünzen F., Janell D., Glühmann M., Hansen H.A.S., Harms J., Bashan A., Bartels H., Agmon I., Franceschi F., Yonath A. (1999) The small ribosomal subunit from *Thermus thermophilus* at 4.5 Å resolution: Pattern fittings and the identification of a functional site. *Proc. Natl. Acad. Sci. USA*, 96: 14252-14257

Towbin H., Staehlin T. and Dordon J. (1979) Electrophoretic transfer of proteins from polyacrilamide gels to nitrocellulose sheets: procedure and some applications. *Proc. Natl. Acad. Sci. USA*, 59: 777-784

Tsiboli P., Herfurth E. and Choli T. (1994) Purification and characterization of the 30S ribosomal proteins from the bacterium *Thermus thermophilus*. *European Journal of Biochemistry*, 226(1): 169-177

VanLoock M.S., Agrawal R.K., Gabashvili I.S., Qi L., Frank J. and Harvey S.C. (2000) Movement of the decoding region of the 16S ribosomal RNA accompanies tRNA translocation. *J. Mol. Biol.*, 304: 507-515

Vogelstein, B. and Gillespie, D. (1979) Preparative and analytical purification of DNA from agarose. *Proc. Natl. Acad. Sci.*, 76: 615-619

Walsh C. (2000) Molecular mechanisms that confer antibacterial drug resistance. *Nature*, 406: 775-781

Wang R., Alexander R.W., VanLoock M., Vladimirov S., Bukhtiyarov Y., Harvey S.C. and Cooperman B.S. (1999) Three-dimensional placement of the conserved 530 loop of 16S rRNA and of its neighbouring components in the 30S subunit. *Journal of Molecular Biology*, 286(2): 521-540

Weiel J. and Hershey J.W. (1981) Fluorescence polarization studies of the interaction of *Escherichia coli* protein synthesis initiation factor 3 with 30S ribosomal subunits. *Biochemistry*, 20(20): 5859-5865

Wilson K.S. and Noller H.F. (1998) Molecular movement inside the translational engine. *Cell*, 92: 337-349

Wimberly B.T., Brodersen D.E., Clemons W.M., Morgan-Warren R.J., Carter A.P., Vornrhein C., Hartsch T. and Ramakrishnan V. (2000) Structure of the 30S ribosomal subunit. *Nature*, 407: 327-339

Wittman H.G., Mussig J., Piefke J., Gewitz H.S., Rheinberger H.J. and Yonath A. (1982) Crystallisation of *E. coli* ribosomes. *FEBS letters*, 146(1): 217-220

Woodcock J., Moazed D., Cannon M., Davies J. and Noller H.F. (1991) Interaction of antibiotics with A-site-specific and P-site-specific bases in 16S ribosomal RNA. *Embo J.*, 10: 3099-3104

Wu X.Q., Iyengar P. and Rajbhandary U.L. (1996) Ribosome-initiator tRNA complex as an intermediate in translation initiation in *Escherichia coli* revealed by use of mutant initiator tRNAs and specialised ribosomes. *EMBO J.*, 15(17): 4734-4739

Yonath A., Glotz C., Gewitz H.S., Bartels K.S., von Bohlen K., Makowski I. And Wittmann H.G. (1988) Characterization of crystals of small ribosomal subunits. *Journal of Molecular Biology*, 203(3): 831-834

Yonath A., Harms J., Hansen H.A.S., Bashan A., Peretz M., Bartels H., Schlünzen F., Koelln I., Bennett W.S., Levin I., Krumbholz S., Avila H. Morlang S., Tocilj A., Weinstein S., Agmon I., Pioletti M., Auerbach T. and Franceschi F. (1998) The quest for the molecular structure of a large macromolecular assembly exhibiting severe non-isomorphism, extreme beam sensitivity and no internal symmetry. *Acta Cryst.*, 54A: 945-955

Zamir A., Miskin R., Elson D. (1971) Inactivation and reactivation of ribosomal subunits: amino acyl-transfer RNA binding activity of the 30S subunit of *Escherichia coli*. *Journal of Molecular Biology*, 60(2): 347-364