

## Literatur

- Abel R., Rybak J. und Menzel R.. Structure and Response Patterns of Olfactory Interneurons in the Honeybee, *Apis mellifera*. *J.comp.Neurol* 437:363-383, 2001.
- Abel R.. Das olfaktorische System der Honigbiene: elektrophysiologische und morphologische Charakterisierung von Antennallobus Neuronen und deren Beteiligung beim olfaktorischen Lernen. *PhD-Thesis* : 1-155, 1997.
- Arnold G., Masson C. und Budharugsa S.. Comparative study of the antennal lobes and their afferent pathway in the worker bee and the drone (*Apis mellifera*). *Cell Tissue Res* 242:593-605, 1985.
- Bicker G., Kreissl S. und Hofbauer A.. Monoclonal antibody labels olfactory and visual pathways in *Drosophila* and *Apis* brains. *Journal of Comparative Neurology* 335:413-424, 1993.
- Bicker G.. Chemical architecture of antennal pathways mediating proboscis extension learning in the honeybee. *Apidologie* 24:235-248, 1993.
- Bicker G.. Histochemistry of classical neurotransmitters in antennal lobes and mushroom bodies of the honeybee. *Microsc Res Tech* 45 (3):174-183, 1999.
- Bitterman M. E., Menzel R., Fietz A. und Schäfer S.. Classical conditioning of proboscis extension in honeybees (*Apis mellifera*). *Journal of Comparative Psychology* 97:107-119, 1983.
- Bornhauser B. C. und Meyer E. P.. Histamine-like immunoreactivity in the visual system and brain of an orthopteran and a hymenopteran insect. *Cell and Tissue Research* 287:211-221, 1997.

- Breer H. und Sattelle D. B.. Molecular forms and subunits of the acetylcholine receptors in the central nervous system of insects. *J.Neurosci.* 5: 3386-3392, 1985.
- Brennan P.A., Schellinck H.M., de la Riva C., Kendrick K.M. und Keverne E.B.. Changes in neurotransmitter release in the main olfactory bulb following an olfactory conditioning procedure in mice. *Neuroscience* 87 (3):583-590, 1998.
- Christensen T. A., Waldrop B. R., Harrow I. D. und Hildebrand J. G.. Local interneurons and information processing in the olfactory glomeruli of the moth *Manduca sexta*. *J.Comp.Physiol.[A]* 173:385-399, 1993.
- de Bruyne M., Foster K. und Carlson J.R.. Odor coding in the *Drosophila* antenna. *Neuron* 30:537-552, 2001.
- Erber J., Masuhr T. und Menzel R.. Localization of short-term memory in the brain of the bee, *Apis mellifera*. *Physiol.Entomol.* 5:343-358, 1980.
- Esslen J. und Kaissling K. E.. Zahl und Verteilung antennaler Sensillen bei der Honigbiene (*Apis mellifera* L.). *Zoomorphol.* 83:227-251, 1976.
- Esteban J.A., Shi S-H, Wilson C., Nuriya M., Haganir R.L. und Malinow R. PKA phosphorylation of AMPA receptor subunits controls synaptic trafficking underlying plasticity. *nature neuroscience* 6 (2):136-143, 2003.
- Faber T., Joerges J. und Menzel R.. Associative learning modifies neural representations of odors in the insect brain. *nature neuroscience* 2 (1):74-78, 1999.
- Faber T.. Veränderung der neuronalen Repräsentation von Düften im Gehirn der Biene durch Lernen. *PhD-Thesis.* 1-125, 1999.

- Farooqui T., Robinson K., Vaessin H. und Smith B.H.. Modulation of early olfactory processing by an octopaminergic reinforcement pathway in the honeybee. *The Journal of Neuroscience* 23 (12):5370-5380, 2003.
- Flanagan D. und Mercer A. R.. An atlas and 3-D reconstruction of the antennal lobes in the worker honey bee, *Apis mellifera* L. (Hymenoptera: Apidae). *Int.J.Insect Morphol.& Embryol.* 18:145-159, 1989a.
- Flanagan D. und Mercer A. R.. Morphology and response characteristics of neurones in the deutocerebrum of the brain in the honeybee *Apis mellifera*. *J.Comp.Physiol.[A]* 164:483-494, 1989b
- Galizia C. G., McIlwrath S. L. und Menzel R.. A digital 3D atlas of the honeybee antennal lobe based on optical sections acquired using confocal microscopy. *Cell Tissue Res* 295:383-394, 1999.
- Galizia C. G., Sachse S., Rappert A. und Menzel R.. The glomerular code for odor representation is species specific in the honeybee *Apis mellifera*. *nature neuroscience* 2 (5):473-478, 1999.
- Gascuel J. und Masson C.. A quantitative ultrastructural study of the honeybee antennal lobe. *Tissue and Cell* 23:341-355, 1991.
- Goldberg F., Grünewald B., Rosenboom H. und Menzel R.. Nicotinic acetylcholine currents of cultured Kenyon cells from the mushroom bodies of the honeybee *Apis mellifera*. *J.Physiol.* 514 (3):759-768, 1999.
- Grünbaum L. und Müller U.. Induction of a specific olfactory memory leads to a long-lasting activation of protein kinase C in the antennal lobe of the honeybee. *The Journal of Neuroscience* 18 (11):4384-4392, 1998.

- Grünewald B.. Physiological properties and response modulations of mushroom body feedback neurons during olfactory learning in the honeybee *Apis mellifera*. *J.Comp.Physiol.[A]* 185:565-576, 1999.
- Hammer M. und Menzel R.. Multiple sites of associative odor learning as revealed by local brain microinjections of octopamine in honeybees. *Learning & Memory* 5:146-156, 1998.
- Hammer M., Braun G. und Mauerlshagen J.. Food-induced arousal and nonassociative learning in honeybees: dependence of sensitization on the application site and duration of food stimulation. *Behavioral and Neural Biology* 62:210-223, 1994.
- Hammer M.. An identified neuron mediates the unconditioned stimulus in associative olfactory learning in honeybees. *Nature* 366 (Nov. 4, 1993):59-63, 1993.
- Hebb D.. *The organization of behaviour*, New York:Wiley, 1949.
- Hildebrand J. G. und Shepherd G. M.. Mechanisms of olfactory discrimination: converging evidence for common principles across phyla. *Annu.Rev.Neurosci.* 20:595-631, 1997.
- Hildebrandt H. und Müller U.. Octopamine mediates rapid stimulation of protein kinase A in the antennal lobe of honeybees. *Journal of Neurobiology* 27 (1):44-50, 1995.
- Hildebrandt H. und Müller U.. PKA activity in the antennal lobe of honeybees is regulated by chemosensory stimulation in vivo. *Brain Res* 679:281-288, 1995.
- Kandel E. R., Schwartz J. H. und Jessell T. M.. *Principles of Neural Science*, New York Amsterdam London Tokyo:Elsevier. Seite 1012, 1991.

Kreissl S. und Bicker G.. Histochemistry of acetylcholinesterase and immunocytochemistry of an acetylcholine receptor-like antigen in the brain of the honey bee. *Journal of Comparative Neurology* 286:71-84, 1989.

Kreissl S., Eichmüller S., Bicker G., Rapus J. und Eckert M.. Octopamine-like immunoreactivity in the brain and suboesophageal ganglion of the honeybee. *Journal of Comparative Neurology* 348:583-595, 1994.

Kuwabara M.. Bildung des bedingten Reflexes von Pavlovs Typus bei der Honigbiene, *Apis mellifica*. *J.Fac.Sci.Hokkaido Univ.Ser.VI Zool.* 13:458-464, 1957.

Laurent G.. Olfactory network dynamics and the coding of multidimensional signals. *nature neuroscience* 3:884-895, 2002.

Lisman J.. A mechanism for the Hebb and the anti-Hebb processes underlying learning and memory. *Proc.Natl.Acad.Sci.USA* 86:9574-9578, 1989.

Mauelshagen J.. Neural correlates of olfactory learning in an identified neuron in the honey bee brain. *J.Neurophysiol.* 69:609-625, 1993.

Menzel R., Erber J. und Masuhr T.. Learning and memory in the honeybee. In: *Experimental analysis of insect behaviour*, edited by L. Barton-Browne, Berlin:Springer, 1974, p. 195-217.

Menzel R., Gaio U. C., Gerberding M., Nemrava E. A. und Wittstock S.. Formation of long-term olfactory memory in honeybees does not require protein synthesis. *Naturwiss.* 80:380-382, 1993.

Menzel R.. Learning, memory, and "cognition" in honey bees. In: *Neurobiology of comparative cognition*, edited by R. P. Kesner and D. S. Olton, Hillsdale, N.J.:Erlbaum Inc., 1990, p. 237-292.

- Menzel R.. Lernen und Gedächtnis. *Praxis der Naturwissenschaften* 38:7-37, 1989.
- Menzel R.. Memory dynamics in the honeybee. *J Comp Physiol A* 185:323-340, 1999.
- Mobbs P. G.. Brain structure. In: *Comprehensive insect physiology biochemistry and pharmacology*, edited by G. A. Kerkut and L. I. Gilbert, Oxford:Pergamon Press, 1985, p. 299-370.
- Mobbs P. G.. The brain of the honeybee *Apis mellifera* I.The connections and spatial organization of the mushroom bodies. *Phil.Trans.R.Soc.Lond.B* 298:309-354, 1982.
- Müller D.. Die olfaktorische Kodierung: Plastizität und Stabilität - Eine elektrophysiologische Analyse der Ausgangsneurone des Antennallobus der Honigbiene. *PhD-Thesis* : 1 – 132, 1999.
- Müller U.. Prolonged activation of cAMP-dependent proteinkinase during conditioning induces long-term memory in honeybees. *Neuron* 27:159-168, 2000.
- Pareto A.. Die zentrale Verteilung der Fühlerafferenz bei Arbeiterinnen der Honigbiene, *Apis mellifera* L. *Z.Zellforsch.* 131:109-140, 1972.
- Sachse S. und Galizia C. G.. Role of inhibition for temporal and spatial odor representation in olfactory output neurons: a calcium imaging study. *J Neurophysiol* 87:1106-1117, 2002.
- Sachse S., Rappert A. und Galizia C. G.. The spatial representation of chemical structures in the antennal lobe of honeybees: steps towards the olfactory code. *Europ.J.Neurosci.* 11:3970-3982, 1999.
- Sachse S.. Odor processing in the honeybee antennal lobe. *PhD-Thesis*, Chapter II: 41-52, 2002.

- Sachse S.. Odor processing in the honeybee antennal lobe. *PhD-Thesis*, Chapter IV. 69-94, 2002.
- Schäfer S. und Bicker G.. Distribution of GABA-like immunoreactivity in the brain of the honeybee. *Journal of Comparative Neurology* 246:287-300, 1986.
- Schäfer S., Bicker G., Ottersen O. P. und Storm-Mathisen J.. Taurine-Like Immunoreactivity in the Brain of the Honeybee. *Journal of Comparative Neurology* 268:60-70, 1988.
- Scheidler A., Kaulen P., Brüning G. und Erber J.. Quantitative autoradiographic localization of [<sup>125</sup>I]-bungarotoxin binding sites in the honeybee brain. *Brain Res.* 534:332-335, 1990.
- Schild D.. Principles of odor coding and a neural network for odor discrimination. *Biophys. J.* 54: 1001-1011, 1988.
- Smith B. H.. The olfactory memory of the honeybee *Apis mellifera*. I. Odorant modulation of short- and intermediate-term memory after single-trial conditioning. *Journal of Experimental Biology* 161:367-382, 1991.
- Sun X.-J., Fonta C. und Masson C.. Odour quality processing by bee antennal lobe interneurons. *Chem.Senses* 18:355-377, 1993.
- Supér H., Spekreijse H. und Lamme V.A.F.. A neural correlate of working memory in the monkey primary visual cortex. *Science* 293:120-124, 2001.
- Suzuki H.. Convergence of olfactory inputs from both antennae in the brain of the honeybee. *Journal of Experimental Biology* 62:11-26, 1975.

Vareschi E.. Duftunterscheidung bei der Honigbiene - Einzelzell-Ableitungen und Verhaltensreaktionen. *Z.vergl.Physiol.* 75:143-173, 1971.

Wilson M. A. und McNaughton B. L.. Reactivation of hippocampal ensemble memories during sleep. *Science* 265:676-679, 1994.

Witthöft W.. Absolute Anzahl und Verteilung der Zellen im Hirn der Honigbiene. *Z.Morph.Tiere* 61:160-184, 1967.

Wittstock S. und Menzel R.. Color learning and memory in honey bees are not affected by protein synthesis inhibition. *Behav.Neur.Biol.* 62:224-229, 1994.

Wittstock S., Kaatz H.-H. und Menzel R.. Inhibition of brain protein synthesis by cycloheximide does not affect formation of long-term memory in honeybees after olfactory conditioning. *J.Neurosci.* 13 (4):1379-1386, 1993.

Wüstenberg D., Gerber B. und Menzel R.. Long- but not medium-term retention of olfactory memories in honeybees is impaired by Actinomycin D and Anisomycin. *Europ.J.Neurosci.* 10 (8):2742-2745, 1998.