

9. Literatur

- Adams, J. M. and Cory, S. (2007). The Bcl-2 apoptotic switch in cancer development and therapy. *Oncogene*. 26(9):1324-37.
- Adolfsson, J., Mansson, R., Buza-Vidas, N., Hultquist, A., Liuba, K., Jensen, C. T., Bryder, D., Yang, L., Borge, O. J., Thoren, L. A., Anderson, K., Sitnicka, E., Sasaki, Y., Sigvardsson, M. and Jacobsen, S. E. (2005). Identification of Flt3+ lympho-myeloid stem cells lacking erythro-megakaryocytic potential a revised road map for adult blood lineage commitment. *Cell*. 121(2):295-306.
- Ashkenazi, A. (2002). Targeting death and decoy receptors of the tumour-necrosis factor superfamily. *Nat Rev Cancer*. 2(6):420-30.
- Bader, J. L. and Miller, R. W. (1978). Neurofibromatosis and childhood leukemia. *J Pediatr*. 92(6):925-9.
- Bakhshi, A., Jensen, J. P., Goldman, P., Wright, J. J., McBride, O. W., Epstein, A. L. and Korsmeyer, S. J. (1985). Cloning the chromosomal breakpoint of t(14;18) human lymphomas: clustering around JH on chromosome 14 and near a transcriptional unit on 18. *Cell*. 41(3):899-906.
- Ballester, R., Marchuk, D., Boguski, M., Saulino, A., Letcher, R., Wigler, M. and Collins, F. (1990). The NF1 locus encodes a protein functionally related to mammalian GAP and yeast IRA proteins. *Cell*. 63(4):851-9.
- Basu, T. N., Gutmann, D. H., Fletcher, J. A., Glover, T. W., Collins, F. S. and Downward, J. (1992). Aberrant regulation of ras proteins in malignant tumour cells from type 1 neurofibromatosis patients. *Nature*. 356(6371):713-5.
- Baum, C., Düllmann, J., Li, Z., Fehse, B., Meyer, J., Williams, D. A. and von Kalle, C. (2003). Side effects of retroviral gene transfer into hematopoietic stem cells. *Blood*. 101(6):2099-114.
- Birnbaum, R. A., O'Mearaigh, A., Wardak, Z., Zhang, Y. Y., Dranoff, G., Jacks, T., Clapp, D. W. and Shannon, K. M. (2000). Nf1 and Gmcsf interact in myeloid leukemogenesis. *Mol Cell*. 5(1):189-95.
- Boguski, M. S. and McCormick, F. (1993). Proteins regulating Ras and its relatives. *Nature*. 366(6456):643-54.
- Bollag, G., Clapp, D. W., Shih, S., Adler, F., Zhang, Y. Y., Thompson, P., Lange, B. J., Freedman, M. H., McCormick, F., Jacks, T. and Shannon, K. (1996). Loss of NF1 results in activation of the Ras signaling pathway and leads to aberrant growth in haematopoietic cells. *Nat Genet*. 12(2):144-8.
- Brannan, C. I., Perkins, A. S., Vogel, K. S., Ratner, N., Nordlund, M. L., Reid, S. W., Buchberg, A. M., Jenkins, N. A., Parada, L. F. and Copeland, N. G. (1994). Targeted disruption of the neurofibromatosis type-1 gene leads to developmental abnormalities in heart and various neural crest-derived tissues. *Genes Dev*. 8(9):1019-29.

- Bronte, V., Apolloni, E., Cabrelle, A., Ronca, R., Serafini, P., Zamboni, P., Restifo, N. P. and Zanovello, P. (2000). Identification of a CD11b(+)/Gr-1(+)/CD31(+) myeloid progenitor capable of activating or suppressing CD8(+) T cells. *Blood*. 96(12):3838-46.
- Burchert, A., Cai, D., Hofbauer, L., Samuelsson, M., Slater, E., Duyster, J., Ritter, M., Hochhaus, A., Muller, R., Eilers, M., Schmidt, M. and Neubauer, A. (2004a). Interferon consensus sequence binding protein (ICSBP; IRF-8) antagonizes BCR/ABL and down-regulates bcl-2. *Blood*. 103(9):3480-9.
- Burchert, A., Cai, D., Hofbauer, L. C., Samuelsson, M. K., Slater, E. P., Duyster, J., Ritter, M., Hochhaus, A., Muller, R., Eilers, M., Schmidt, M. and Neubauer, A. (2004b). Interferon consensus sequence binding protein (ICSBP; IRF-8) antagonizes BCR/ABL and down-regulates bcl-2. *Blood*. 103(9):3480-9.
- Calabretta, B. and Perrotti, D. (2004). The biology of CML blast crisis. *Blood*. 103(11):4010-22.
- Cambier, N., Chopra, R., Strasser, A., Metcalf, D. and Elefanty, A. G. (1998). BCR-ABL activates pathways mediating cytokine independence and protection against apoptosis in murine hematopoietic cells in a dose-dependent manner. *Oncogene*. 16(3):335-48.
- Cauley, L. S., Miller, E. E., Yen, M. and Swain, S. L. (2000). Superantigen-induced CD4 T cell tolerance mediated by myeloid cells and IFN-gamma. *J Immunol*. 165(11):6056-66.
- Cawthon, R. M., Weiss, R., Xu, G. F., Viskochil, D., Culver, M., Stevens, J., Robertson, M., Dunn, D., Gesteland, R. and O'Connell, P. e. a. (1990). A major segment of the neurofibromatosis type 1 gene: cDNA sequence, genomic structure, and point mutations. *Cell*. 62(1):193-201.
- Cichowski, K. and Jacks, T. (2001). NF1 tumor suppressor gene function: narrowing the GAP. *Cell*. 104(4):593-604.
- Cichowski, K., Santiago, S., Jardim, M., Johnson, B. W. and Jacks, T. (2003). Dynamic regulation of the Ras pathway via proteolysis of the NF1 tumor suppressor. *Genes Dev*. 17(4):449-54.
- Cleary, M. L., Smith, S. D. and Sklar, J. (1986). Cloning and structural analysis of cDNAs for bcl-2 and a hybrid bcl-2/immunoglobulin transcript resulting from the t(14;18) translocation. *Cell*. 47(1):19-28.
- Colman, S. D., Williams, C. A. and Wallace, M. R. (1995). Benign neurofibromas in type 1 neurofibromatosis (NF1) show somatic deletions of the NF1 gene. *Nat Genet*. 11(1):90-2.
- Cory, S. and Adams, J. M. (2002). The Bcl2 family: regulators of the cellular life-or-death switch. *Nat Rev Cancer*. 2(9):647-56.
- Cory, S., Huang, D. C. and Adams, J. M. (2003). The Bcl-2 family: roles in cell survival and oncogenesis. *Oncogene*. 22(53):8590-607.

- Daley, G. Q. and Baltimore, D. (1988). Transformation of an interleukin 3-dependent hematopoietic cell line by the chronic myelogenous leukemia-specific P210bcr/abl protein. *Proc Natl Sci USA.* 85(23):9312-6.
- Dameshek, W. (1951). Some speculations on the myeloproliferative syndromes. *Blood.* 6(4):372-5.
- Dasgupta, B. and Gutmann, D. H. (2003). Neurofibromatosis 1: closing the GAP between mice and men. *Curr Opin Genet Dev.* 13(1):20-7.
- DeClue, J. E., Papageorge, A. G., Fletcher, J. A., Diehl, S. R., Ratner, N., Vass, W. C. and Lowy, D. R. (1992). Abnormal regulation of mammalian p21ras contributes to malignant tumor growth in von Recklinghausen (type 1) neurofibromatosis. *Cell.* 69(2):265-73.
- Deng, M. and Daley, G. Q. (2001). Expression of interferon consensus sequence binding protein induces potent immunity against BCR/ABL-induced leukemia. *Blood.* 97(11):3491-7.
- Donnell Mc, T. J. and Korsmeyer, S. J. (1991). Progression from lymphoid hyperplasia to high-grade malignant lymphoma in mice transgenic for the t(14; 18). *Nature.* 349(6306):254-6.
- Driggers, P. H., Ennist, D. L., Gleason, S. L., Mak, W. H., Marks, M. S., Levi, B. Z., Flanagan, J. R., Appella, E. and Ozato, K. (1990). An interferon gamma-regulated protein that binds the interferon-inducible enhancer element of major histocompatibility complex class I genes. *Proc Natl Sci USA.* 87(10):3743-7.
- Emanuel, P. D., Bates, L. J., Castleberry, R. P., Gaultier, R. J. and Zuckerman, K. S. (1991). Selective hypersensitivity to granulocyte-macrophage colony-stimulating factor by juvenile chronic myeloid leukemia hematopoietic progenitors. *Blood.* 77(5):925-9.
- Fehr, T., Schoedon, G., Odermatt, B., Holtschke, T., Schneemann, M., Bachmann, M. F., Mak, T. W., Horak, I. and Zinkernagel, R. M. (1997). Crucial role of interferon consensus sequence binding protein, but neither of interferon regulatory factor 1 nor of nitric oxide synthesis for protection against murine listeriosis. *J Exp Med.* 185(5):921-31.
- Frank, O., Rudolph, C., Heberlein, C., von Neuhoff, N., Schrock, E., Schambach, A., Schlegelberger, B., Fehse, B., Ostertag, W., Stocking, C. and Baum, C. (2004). Tumor cells escape suicide gene therapy by genetic and epigenetic instability. *Blood.* 104(12):3543-9.
- Gabriele, L., Phung, J., Fukumoto, J., Segal, D., Wang, I. M., Giannakakou, P., Giese, N. A., Ozato, K. and Morse, H. C. r. (1999). Regulation of apoptosis in myeloid cells by interferon consensus sequence-binding protein. *J Exp Med.* 190(3):411-21.
- Gibrilovich, D. I., Velders, M. P., Sotomayor, E. M. and Kast, W. M. (2001). Mechanism of immune dysfunction in cancer mediated by immature Gr-1+ myeloid cells. *J Immunol.* 16(9):5398-406.
- Gaiger, A., Henn, T., Horth, E., Geissler, K., Mitterbauer, G., Maier-Dobersberger, T., Greinix, H., Mannhalter, C., Haas, O. A., Lechner, K. and Lion, T. (1995). Increase of bcr-abl chimeric mRNA expression in tumor cells of patients with chronic myeloid leukemia precedes disease progression. *Blood.* 86(6):2371-8.

- Gishizky, M. L. and Witte, O. N. (1992). Initiation of deregulated growth of multipotent progenitor cells by bcr-abl in vitro. *Science*. 256(5058):836-9.
- Guha, A., Lau, N., Huvar, I., Gutmann, D., Provias, J., Pawson, T. and Boss, G. (1996). Ras-GTP levels are elevated in human NF1 peripheral nerve tumors. *Oncogene*. 12(3):507-13.
- Gutmann, D. H. and Collins, F. S. (1993). The neurofibromatosis type 1 gene and its protein product, neurofibromin. *Neuron*. 10(3):335-43.
- Gutmann, D. H., Winkeler, E., Kabbarah, O., Hedrick, N., Dudley, S., Goodfellow, P. J. and Liskay, R. M. (2003). Mlh1 deficiency accelerates myeloid leukemogenesis in neurofibromatosis 1 (Nf1) heterozygous mice. *Oncogene*. 22(29):4581-5.
- Hanahan, D. and Weinberg, R. A. (2000). The hallmarks of cancer. *Cell*. 100(1):57-70.
- Hanenberg, H., Xiao, X. L., Diloo, D., Hashino, K., Kato, I. and Williams, D. A. (1996). Colocalization of retrovirus and target cells on specific fibronectin fragments increases genetic transduction of mammalian cells. *Nat Med*. 2(8):876-82.
- Hao, S. X. and Ren, R. (2000). Expression of interferon consensus sequence binding protein (ICSBP) is downregulated in Bcr-Abl-induced murine chronic myelogenous leukemia-like disease, and forced coexpression of ICSBP inhibits Bcr-Abl-induced myeloproliferative disorder. *Mol Cell Biol*. 20(4):1149-61.
- Hasle, H., Niemeyer, C. M., Chessells, J. M., Baumann, I., Bennett, J. M., Kerndrup, G. and Head, D. R. (2003). A pediatric approach to the WHO classification of myelodysplastic and myeloproliferative diseases. *Leukemia*. 17(2):277-82.
- Heisterkamp, N., Stam, K., Groffen, J., de Klein, A. and Grosveld, G. (1985). Structural organization of the bcr gene and its role in the Ph' translocation. *Nature*. 315(6022):758-61.
- Hockenberry, D., Nunez, G., Milliman, C., Schreiber, R. D. and Korsmeyer, S. J. (1990). Bcl-2 is an inner mitochondrial membrane protein that blocks programmed cell death. *Nature*. 348(6299):334-6.
- Holtschke, T., Lohler, J., Kanno, Y., Fehr, T., Giese, N., Rosenbauer, F., Lou, J., Knobeloch, K. P., Gabriele, L., Waring, J. F., Bachmann, M. F., Zinkernagel, R. M., Morse, H. C., Ozato, K. and Horak, I. (1996). Immunodeficiency and chronic myelogenous leukemia-like syndrome in mice with a targeted mutation of the ICSBP gene. *Cell*. 87(2):307-17.
- Honda, H., Ushijima, T., Wakazono, K., Oda, H., Tanaka, Y., Aizawa, S., Ishikawa, T., Yazaki, Y. and Hirai, H. (2000). Acquired loss of p53 induces blastic transformation in p210(bcr/abl)-expressing hematopoietic cells: a transgenic study for blast crisis of human CML. *Blood*. 95(4):1144-50.
- Hsu, Y. T. and Youle, R. J. (1998). Bax in murine thymus is a soluble monomeric protein that displays differential detergent-induced conformations. *J Biol Chem*. 273(17):10777-83.
- Huang, W., Horvath, E. and Eklund, E. A. (2007). PU.1, interferon regulatory factor 2 (IRF2) and the interferon consensus sequence binding protein (ICSBP/IRF8) cooperate to activate NF1-transcription in differentiating myeloid cells. *J Biol Chem*. [Epub ahead of print]:

- Huang, W., Saberwal, G., Horvath, E., Zhu, C., Lindsey, S. and Eklund, E. A. (2006). Leukemia-associated, constitutively active mutants of SHP2 protein tyrosine phosphatase inhibit NF1 transcriptional activation by the interferon consensus sequence binding protein. *Mol Cell Biol.* 26(17):6311-32.
- Jacks, T., Shih, T. S., Schmitt, E. M., Bronson, R. T., Bernards, A. and Weinberg, R. A. (1994). Tumour predisposition in mice heterozygous for a targeted mutation in Nf1. *Nat Genet.* 7(3):353-61.
- Jaiswal, S., Traver, D., Miyamoto, T., Akashi, K., Lagasse, E. and Weissman, I. L. (2003). Expression of BCR/ABL and BCL-2 in myeloid progenitors leads to myeloid leukemias. *Proc Natl Sci USA.* 100(17):10002-7.
- Janiak, F., Leber, B. and Andrews, D. W. (1994). Assembly of Bcl-2 into microsomal and outer mitochondrial membranes. *J Biol Chem.* 269(13):9842-9.
- Kalra, R., Paderanga, D. C., Olson, K. and Shannon, K. M. (1994). Genetic analysis is consistent with the hypothesis that NF1 limits myeloid cell growth through p21ras. *Blood.* 84(10):3435-9.
- Kantarjian, H. and Talpaz, M. (1988). Definition of the accelerated phase of chronic myelogenous leukemia. *J Clin Oncol.* 6(1):180-2.
- Kantarjian, H. M., Keating, M. J., Talpaz, M., Walters, R. S., Smith, T. L., Cork, A., McCredie, K. B. and Freireich, E. J. (1987). Chronic myelogenous leukemia in blast crisis. Analysis of 242 patients. *Am J Med.* 83(3):445-54.
- Kim, A., Morgan, K., Hasz, D. E., Wiesner, S. M., Lauchle, J. O., Geurts, J. L., Diers, M. D., Le, D. T., Kogan, S. C., Parada, L. F., Shannon, K. and Largaespada, D. A. (2007). Beta common receptor inactivation attenuates myeloproliferative disease in Nf1 mutant mice. *Blood.* 109(4):1687-91.
- Kim, H. A., Rosenbaum, T., Marchionni, M. A., Ratner, N. and DeClue, J. E. (1995). Schwann cells from neurofibromin deficient mice exhibit activation of p21ras, inhibition of cell proliferation and morphological changes. *Oncogene.* 11(2):325-35.
- Knudson, A. G. J. (1971). Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Sci USA.* 68(4):820-3.
- Kogan, S. C., Brown, D. E., Shultz, D. B., Truong, B. T., Lallemand-Breitenbach, V., Guillemin, M. C., Lagasse, E., Weissman, I. L. and Bishop, J. M. (2001). BCL-2 cooperates with promyelocytic leukemia retinoic acid receptor alpha chimeric protein (PMLRARalpha) to block neutrophil differentiation and initiate acute leukemia. *J Exp Med.* 193(4):531-43.
- Kondo, M., Wagers, A. J., Manz, M. G., Prohaska, S. S., Scherer, D. C., Beilhack, G. F., Shizuru, J. A. and Weissman, I. L. (2003). Biology of hematopoietic stem cells and progenitors: implications for clinical application. *Annu Rev Immunol.* 21:759-806.
- Kramer, A., Loffler, H., Bergmann, J., Hochhaus, A. and Hehlmann, R. (2001). Proliferating status of peripheral blood progenitor cells from patients with BCR/ABL-positive chronic myelogenous leukemia. *Leukemia.* 15(1):62-8.

- Largaespada, D. A., Brannan, C. I., Jenkins, N. A. and Copeland, N. G. (1996). Nf1 deficiency causes Ras-mediated granulocyte/macrophage colony stimulating factor hypersensitivity and chronic myeloid leukaemia. *Nat Genet.* 12(2):137-43.
- Lau, N., Feldkamp, M. M., Roncari, L., Loehr, A. H., Shannon, P., Gutmann, D. H. and Guha, A. (2000). Loss of neurofibromin is associated with activation of RAS/MAPK and PI3-K/AKT signaling in a neurofibromatosis 1 astrocytoma. *J Neuropathol Exp Neurol.* 59(9):759-67.
- Le, D. T., Kong, N., Zhu, Y., Lauchle, J. O., Aiyigari, A., Braun, B. S., Wang, E., Kogan, S. C., Le Beau, M. M., Parada, L. and Shannon, K. M. (2004). Somatic inactivation of Nf1 in hematopoietic cells results in a progressive myeloproliferative disorder. *Blood.* 103(11):4243-50.
- Lee, M. J. and Stephenson, D. A. (2007). Recent developments in neurofibromatosis type 1. *Curr Opin Neurol.* 20(2):135-41.
- Levi, B. Z., Hashmueli, S., Gleit-Kielmanowicz, M., Azriel, A. and Meraro, D. (2002). ICSBP/IRF-8 transactivation: a tale of protein-protein interaction. *J Interferon Cytokine Res.* 22(1):153-60.
- Li, H., Zhu, H., Xu, C. J. and Yuan, J. (1998). Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis. *Cell.* 94(4):491-501.
- Luo, X., Budihardjo, I., Zou, H., Slaughter, C. and Wang, X. (1998). Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors. *Cell.* 94(4):481-90.
- Mahgoub, N., Taylor, B. R., Le Beau, M. M., Gratiot, M., Carlson, K. M., Atwater, S. K., Jacks, T. and Shannon, K. M. (1999). Myeloid malignancies induced by alkylating agents in Nf1 mice. *Blood.* 93(11):3617-23.
- Martin, G. A., Viskochil, D., Bollag, G., McCabe, P. C., Crosier, W. J., Haubruck, H., Conroy, L., Clark, R., O'Connell, P., R.M., C. and al., e. (1990). The GAP-related domain of the neurofibromatosis type 1 gene product interacts with ras p21. *Cell.* 63(4):843-9.
- McLaughlin, J., Chianese, E. and Witte, O. N. (1987). In vitro transformation of immature hematopoietic cells by the P210 BCR/ABL oncogene product of the Philadelphia chromosome. *Proc Natl Sci USA.* 84(18):6558-62.
- McWhirter, J. R., Galasso, D. L. and Wang, J. Y. (1993). A coiled-coil oligomerization domain of Bcr is essential for the transforming function of Bcr-Abl oncoproteins. *Mol Cell Biol.* 13(12):7587-95.
- Melani, C., Chiodoni, C., Forni, G. and Colombo, M. P. (2003). Myeloid cell expansion elicited by the progression of spontaneous mammary carcinomas in c-erbB-2 transgenic BALB/c mice suppresses immune reactivity. *Blood.* 102(6):2138-45.
- Middleton, M. K., Zukas, A. M., Rubinstein, T., Jacob, M., Zhu, P., Zhao, L., Blair, I. and Pure, E. (2006). Identification of 12/15-lipoxygenase as a suppressor of myeloproliferative disease. *J Exp Med.* 203(11):2529-40.

- Miles, D. K., Freedman, M. H., Stephens, K., Pallavicini, M., Sievers, E. L., Weaver, M., Grunberger, T., Thompson, P. and Shannon, K. M. (1996). Patterns of hematopoietic lineage involvement in children with neurofibromatosis type 1 and malignant myeloid disorders. *Blood*. 88(11):4314-20.
- Moritz, T., Patel, V. P. and Williams, D. A. (1994). Bone marrow extracellular matrix molecules improve gene transfer into human hematopoietic cells via retroviral vectors. *J Clin Invest*. 93(4):1451-7.
- Muller, A. J., Young, J. C., Pendergast, A. M., Pondel, M., Landau, N. R., Littman, D. R. and Witte, O. N. (1991). BCR first exon sequences specifically activate the BCR/ABL tyrosine kinase oncogene of Philadelphia chromosome-positive human leukemias. *Mol Cell Biol*. 11(4):1785-92.
- Nakano, K. and Vousden, K. H. (2001). PUMA, a novel proapoptotic gene, is induced by p53. *Mol Cell*. 7(3):683-94.
- Nechushtan, A., Smith, C. L., Lamensdorf, I., Yoon, S. H. and Youle, R. J. (2001). Bax and Bak coalesce into novel mitochondria-associated clusters during apoptosis. *J Cell Biol*. 153(6):1265-76.
- Nelson, N., Kanno, Y., Hong, C., Contursi, C., Fujita, T., Fowlkes, B. J., O'Connell, E., Hu-Li, J., Paul, W. E., Jankovic, D., Sher, A. F., Coligan, J. E., Thornton, A., Appella, E., Yang, Y. and Ozato, K. (1996). Expression of IFN regulatory factor family proteins in lymphocytes. Induction of Stat-1 and IFN consensus sequence binding protein expression by T cell activation. *J Immunol*. 156(10):3711-20.
- Nguyen, H., Hiscott, J. and Pitha, P. M. (1997). The growing family of interferon regulatory factors. *Cytokine Growth Factor Rev*. 8(4):293-312.
- Nowell, P. C. and Hungerford, D. A. (1960). A minute chromosome in human chronic granulocytic leukemia. *Science*. 132:1497-99.
- Oda, E., Ohki, R., Murasawa, H., Nemoto, J., Shibue, T., Yamashita, T., Tokino, T., Taniguchi, T. and Tanaka, N. (2000). Noxa, a BH3-only member of the Bcl-2 family and candidate mediator of p53-induced apoptosis. *Science*. 288(5468):1053-8.
- Pavlov, E. V., Priault, M., Pietkiewicz, D., Cheng, E. H., Antonsson, B., Manon, S., Korsmeyer, S. J., Mannella, C. A. and Kinnally, K. W. (2001). A novel, high conductance channel of mitochondria linked to apoptosis in mammalian cells and Bax expression in yeast. *J Cell Biochem*. 155(5):725-31.
- Pelaez, B., Campillo, J. A., Lopez-Asenjo, J. A. and Subiza, J. L. (2001). Cyclophosphamide induces the development of early myeloid cells suppressing tumor cell growth by a nitric oxide-dependent mechanism. *J Immunol*. 166(11):6608-15.
- Puthalakath, H., Huang, D. C., O'Reilly, L. A., King, S. M. and Strasser, A. (1999). The proapoptotic activity of the Bcl-2 family member Bim is regulated by interaction with the dynein motor complex. *Mol Cell*. 3(3):287-96.

- Puthalakath, H., Villunger, A., O'Reilly, L. A., Beaumont, J. G., Coulter, L., Cheney, R. E., Huang, D. C. and Strasser, A. (2001). Bmf: a proapoptotic BH3-only protein regulated by interaction with the myosin V actin motor complex, activated by anoikis. *Science*. 293(5536):1829-32.
- Qian, Z., Fernald, A. A., Godley, L. A., Larson, R. A. and Le Beau, M. M. (2002). Expression profiling of CD34+ hematopoietic stem/ progenitor cells reveals distinct subtypes of therapy-related acute myeloid leukemia. *Proc Natl Sci USA*. 99(23):14925-30.
- Ren, R. (2002). The molecular mechanism of chronic myelogenous leukemia and its therapeutic implications: studies in a murine model. *Oncogene*. 21(56):8629-42.
- Rowley, J. D. (1973). Letter: A new consistent chromosomal abnormality in chronic myelogenous leukaemia identified by quinacrine fluorescence and Giemsa staining. *Nature*. 243(5405):290-3.
- Rubin, J. B. and Gutmann, D. H. (2005). Neurofibromatosis type 1 - a model for nervous system tumour formation? *Nat Rev Cancer*. 5(7):557-64.
- Sanchez-Garcia, I. and Grutz, G. (1995). Tumorigenic activity of the BCR-ABL oncogenes is mediated by BCL2. *Proc Natl Sci USA*. 92(12):5287-91.
- Sanchez-Garcia, I. and Martin-Zanca, D. (1997). Regulation of Bcl-2 gene expression by BCR-ABL is mediated by Ras. *J Mol Biol*. 267(2):225-8.
- Sander, S., Bullinger, L., Karlsson, A., Giuriato, S., Hernandez-Boussard, T., Felsher, D. W. and Pollack, J. R. (2005). Comparative genomic hybridization on mouse cDNA microarrays and its application to a murine lymphoma model. *Oncogene*. 24(40):6101-7.
- Sattler, M. and Salgia, R. (1997). Activation of hematopoietic growth factor signal transduction pathways by the human oncogene BCR/ABL. *Cytokine Growth Factor Rev*. 8(1):63-79.
- Savage, D. G., Szydlo, R. M. and Goldman, J. M. (1997). Clinical features at diagnosis in 430 patients with chronic myeloid leukaemia seen at a referral centre over a 16-year period. *Br J Haematol*. 96(1):111-6.
- Sawada, S., Florell, S., Purandare, S. M., Ota, M., Stephens, K. and Viskochil, D. (1996). Identification of NF1 mutations in both alleles of a dermal neurofibroma. *Nat Genet*. 14(1):110-2.
- Scheller, M., Foerster, J., Heyworth, C. M., Waring, J. F., Lohler, J., Gilmore, G. L., Shadduck, R. K., Dexter, T. M. and I. H. (1999). Altered development and cytokine responses of myeloid progenitors in the absence of transcription factor, interferon consensus sequence binding protein. *Blood*. 94(11):3764-71.
- Schmidt, M., Hochhaus, A., Nitsche, A., Hehlmann, R. and Neubauer, A. (2001). Expression of nuclear transcription factor interferon consensus sequence binding protein in chronic myeloid leukemia correlates with pretreatment risk features and cytogenetic response to interferon-alpha. *Blood*. 97(11):3648-50.

- Schmidt, M., Nagel, S., Proba, J., Thiede, C., Ritter, M., Waring, J. F., Rosenbauer, F., Huhn, D., Wittig, B., Horak, I. and Neubauer, A. (1998). Lack of interferon consensus sequence binding protein (ICSBP) transcripts in human myeloid leukemias. *Blood*. 91(1):22-9.
- Serafini, P., Borrello, I. and Bronte, V. (2006). Myeloid suppressor cells in cancer: recruitment, phenotype, properties, and mechanisms of immune suppression. *Semin Cancer Biol.* 16(1):53-65.
- Serra, E., Puig, S., Otero, D., Gaona, A., Kruyer, H., Ars, E., Estivill, X. and Lazaro, C. (1997). Confirmation of a double-hit model for the NF1 gene in benign neurofibromas. *Am J Hum Genet.* 61(3):512-9.
- Shannon, K. M., O'Connell, P., Martin, G. A., Paderanga, D., Olson, K., Dinndorf, P. and McCormick, F. (1994). Loss of the normal NF1 allele from the bone marrow of children with type 1 neurofibromatosis and malignant myeloid disorders. *N Engl J Med.* 330(9):597-601.
- Sherman, L. S., Atit, R., Rosenbaum, T., Cox, A. D. and Ratner, N. (2000). Single cell Ras-GTP analysis reveals altered Ras activity in a subpopulation of neurofibroma Schwann cells but not fibroblasts. *J Biol Chem.* 275(39):30740-5.
- Shet, A. S., Jahagirdar, B. N. and Verfaillie, C. M. (2002). Chronic myelogenous leukemia: mechanisms underlying disease progression. *Leukemia*. 16(8):1402-11.
- Side, L., Taylor, B., Cayouette, M., Conner, E., Thompson, P., Luce, M. and Shannon, K. (1997). Homozygous inactivation of the NF1 gene in bone marrow cells from children with neurofibromatosis type 1 and malignant myeloid disorders. *N Engl J Med.* 336(24):1713-20.
- Side, L. E., Emanuel, P. D., Taylor, B., Franklin, J., Thompson, P., Castleberry, R. P. and Shannon, K. M. (1998). Mutations of the NF1 gene in children with juvenile myelomonocytic leukemia without clinical evidence of neurofibromatosis, type 1. *Blood*. 92(1):267-72.
- Sill, H., Goldman, J. M. and Cross, N. C. (1995). Homozygous deletions of the p16 tumor-suppressor gene are associated with lymphoid transformation of chronic myeloid leukemia. *Blood*. 85(8):2013-6.
- Skorski, T., Nieborowska-Skorska, M., Wlodarski, P., Perrotti, D., Martinez, R., Wasik, M. A. and Calabretta, B. (1996). Blastic transformation of p53-deficient bone marrow cells by p210bcr/abl tyrosine kinase. *Proc Natl Sci USA*. 93(23):13137-42.
- Spiers, A. S. (1977). The clinical features of chronic granulocytic leukaemia. *Clin Haematol.* 6(1):77-95.
- Stiller, C. A., Chessells, J. M. and Fitchett, M. (1994). Neurofibromatosis and childhood leukaemia/lymphoma: a population-based UKCCSG study. *Br J Cancer*. 70(5):969-72.
- Strasser, A., Harris, A., Bath, M. and Cory, S. (1990). Novel primitive lymphoid tumours induced in transgenic mice by cooperation between myc and bcl-2. *Nature*. 348(6299):331-3.
- Strasser, A. and Harris, A. W. (1993). E mu-bcl-2 transgene facilitates spontaneous transformation of early pre-B and immunoglobulin-secreting cells but not T cells. *Oncogene*. 8(1):1-9.

- Tamura, T., Kong, H. J., Tunyaplin, C., Tsujimura, H., Calame, K. and Ozato, K. (2003). ICSBP/IRF-8 inhibits mitogenic activity of p210 Bcr/Abl in differentiating myeloid progenitor cells. *Blood*. 102(13):4547-54.
- Tamura, T., Nagamura-Inoue, T., Shmeltzer, Z., Kuwata, T. and Ozato, K. (2000). ICSBP directs bipotential myeloid progenitor cells to differentiate into mature macrophages. *Immunity*. 13(2):155-65.
- Tefferi, A. and Gilliland, D. G. (2007). Oncogenes in myeloproliferative disorders. *Cell Cycle*. 6(5):550-66.
- Terrazas, L. I., Walsh, K. L., Piskorska, D., McGuire, E. and Harn, D. A. J. (2001). The schistosome oligosaccharide lacto-N-neotetraose expands Gr1(+) cells that secrete anti-inflammatory cytokines and inhibit proliferation of naive CD4(+) cells: a potential mechanism for immune polarization in helminth infections. *J Immunol*. 167(9):5294-303.
- Terszowski, G., Waskow, C., Conradt, P., Lenze, D., Koenigsmann, J., Carstanjen, D., Horak, I. and Rodewald, H. R. (2005). Prospective isolation and global gene expression analysis of the erythrocyte colony-forming unit (CFU-E). *Blood*. 105(5):1937-45.
- The, I., Hannigan, G. E., Cowley, G. S., Reginald, S., Zhong, Y., Gusella, J. F., Hariharan, I. K. and Bernards, A. (1997). Rescue of a Drosophila NF1 mutant phenotype by protein kinase A. *Science*. 276(5313):791-4.
- Tong, J., Hannan, F., Zhu, Y., Bernards, A. and Zhong, Y. (2002). Neurofibromin regulates G protein-stimulated adenylyl cyclase activity. *Nat Neurosci*. 5(2):95-6.
- Traver, D., Akashi, K., Weissman, I. L. and Lagasse, E. (1998). Mice defective in two apoptosis pathways in the myeloid lineage develop acute myeloblastic leukemia. *Immunity*. 9(1):47-57.
- Trovo-Marqui, A. B. and Tajara, E. H. (2006). Neurofibromin: a general outlook. *Clin Genet*. 70(1):1-13.
- Tsujimoto, Y., Finger, L. R., Yunis, J., Nowell, P. C. and Croce, C. M. (1984). Cloning of the chromosome breakpoint of neoplastic B cells with the t(14;18) chromosome translocation. *Science*. 226(4678):1097-9.
- Tsujimura, H., Nagamura-Inoue, T., Tamura, T. and Ozato, K. (2002). IFN consensus sequence binding protein/IFN regulatory factor-8 guides bone marrow progenitor cells toward the macrophage lineage. *J Immunol*. 169(3):1261-9.
- Vaux, D. L., Cory, S. and Adams, J. M. (1988). Bcl-2 gene promotes haemopoietic cell survival and cooperates with c-myc to immortalize pre-B cells. *Nature*. 335(6189):440-2.
- Veis, D. J., Sorenson, C. M., Shutter, J. R. and Korsmeyer, S. J. (1993). Bcl-2-deficient mice demonstrate fulminant lymphoid apoptosis, polycystic kidneys, and hypopigmented hair. *Cell*. 75(2):229-40.

- Verfaillie, C. M., McCarthy, J. B. and McGlave, P. B. (1992). Mechanisms underlying abnormal trafficking of malignant progenitors in chronic myelogenous leukemia. Decreased adhesion to stroma and fibronectin but increased adhesion to the basement membrane components laminin and collagen type IV. *J Clin Invest.* 90(4):1232-41.
- Virchow, R. (1845). Weisses Blut (Leukämie). *Archiv Pathol Anat.* 1:563.
- Wallace, M. R., Marchuk, D. A., Andersen, L. B., Letcher, R., Odeh, H. M., Saulino, A. M., Fountain, J. W., Brereton, A., Nicholson, J. and Mitchell, A. L. e. a. (1990). Type 1 neurofibromatosis gene: identification of a large transcript disrupted in three NF1 patients. *Science.* 249(4965):181-6.
- Xu, G., O'Connell, P., Viskochil, D., Cawthon, R., Robertson, M., Culver, M., Dunn, D., Stevens, J., Gesteland, R. and White, R. e. a. (1990b). The neurofibromatosis type 1 gene encodes a protein related to GAP. *Cell.* 62(3):599-608.
- Xu, G. F., Lin, B., Tanaka, K., Dunn, D., Wood, D., Gesteland, R., White, R., Weiss, R. and Tamanoi, F. (1990). The catalytic domain of the neurofibromatosis type 1 gene product stimulates ras GTPase and complements ira mutants of *S. cerevisiae*. *Cell.* 63(4):835-41.
- Xu, W., Mulligan, L. M., Ponder, M. A., Liu, L., Smith, B. A., Mathew, C. G. and Ponder, B. A. (1992). Loss of NF1 alleles in phaeochromocytomas from patients with type I neurofibromatosis. *Genes Chromosomes Cancer.* 4(4):337-42.
- Yang, G., Khalaf, W., van de Locht, L., Jansen, J. H., Gao, M., Thompson, M. A., van der Reijden, B. A., Gutmann, D. H., Delwel, R., Clapp, D. W. and Hiebert, S. W. (2005). Transcriptional repression of the Neurofibromatosis-1 tumor suppressor by the t(8;21) fusion protein. *Mol Cell Biol.* 25(14):5869-79.
- Yu, J., Zhang, L., Hwang, P. M., Kinzler, K. W. and Vogelstein, B. (2001). PUMA induces the rapid apoptosis of colorectal cancer cells. *Mol Cell.* 7(3):673-82.
- Zamzami, N. and Kroemer, G. (2001). The mitochondrion in apoptosis: how Pandora's box opens. *Nat Rev Mol Cell Biol.* 2(1):67-71.
- Zha, J., Harada, H., Yang, E., Jockel, J. and Korsmeyer, S. J. (1996). Serine phosphorylation of death agonist BAD in response to survival factor results in binding to 14-3-3 not BCL-X(L). *Cell.* 87(4):619-28.
- Zhang, Y., Taylor, B. R., Shannon, K. and Clapp, D. W. (2001). Quantitative effects of Nf1 inactivation on in vivo hematopoiesis. *J Clin Invest.* 108(5):709-15.
- Zhang, Y. Y., Vik, T. A., Ryder, J. W., Srour, E. F., Jacks, T., Shannon, K. and Clapp, D. W. (1998). Nf1 regulates hematopoietic progenitor cell growth and ras signaling in response to multiple cytokines. *J Exp Med.* 187(11):1893-902.
- Zhu, C., Saberwal, G., Lu, Y., Platanias, L. C. and Eklund, E. A. (2004). The interferon consensus sequence-binding protein activates transcription of the gene encoding neurofibromin 1. *J Biol Chem.* 279(49):50874-85.

Zinkel, S., Gross, A. and Yang, E. (2006). BCL2 family in DNA damage and cell cycle control. *Cell Death Differ.* 13(8):1351-9.

Zion, M., Ben-Yehuda, D., Avraham, A., Cohen, O., Wetzler, M., Melloul, D. and Ben-Neriah, Y. (1994). Progressive de novo DNA methylation at the bcr-abl locus in the course of chronic myelogenous leukemia. *Proc Natl Sci USA.* 91(22):10722-6.