

4. Literaturverzeichnis

- Acuto, O., and Michel, F. (2003). CD28-mediated co-stimulation: a quantitative support for TCR signalling. *Nat Rev Immunol* 3, 939-951.
- Alderson, M. R., Smith, C. A., Tough, T. W., Davis-Smith, T., Armitage, R. J., Falk, B., Roux, E., Baker, E., Sutherland, G. R., and Din, W. S. (1994). Molecular and biological characterization of human 4-1BB and its ligand. *Eur J Immunol* 24, 2219-2227.
- Alegre, M. L., Frauwirth, K. A., and Thompson, C. B. (2001). T-cell regulation by CD28 and CTLA-4. *Nat Rev Immunol* 1, 220-228.
- Allegretta, M., Nicklas, J. A., Sriram, S., and Albertini, R. J. (1990). T cells responsive to myelin basic protein in patients with multiple sclerosis. *Science* 247, 718-721.
- Arch, R. H., and Thompson, C. B. (1998). 4-1BB and Ox40 are members of a tumor necrosis factor (TNF)-nerve growth factor receptor subfamily that bind TNF receptor-associated factors and activate nuclear factor kappaB. *Mol Cell Biol* 18, 558-565.
- Baecher-Allan, C., Brown, J. A., Freeman, G. J., and Hafler, D. A. (2001). CD4+CD25high regulatory cells in human peripheral blood. *J Immunol* 167, 1245-1253.
- Baecher-Allan, C., and Hafler, D. A. (2004). Suppressor T cells in human diseases. *J Exp Med* 200, 273-276.
- Bertin, J., Armstrong, R. C., Ottlie, S., Martin, D. A., Wang, Y., Banks, S., Wang, G. H., Senkevich, T. G., Alnemri, E. S., Moss, B., *et al.* (1997). Death effector domain-containing herpesvirus and poxvirus proteins inhibit both Fas- and TNFR1-induced apoptosis. *Proc Natl Acad Sci U S A* 94, 1172-1176.
- Bertram, E. M., Dawicki, W., Sedgmen, B., Bramson, J. L., Lynch, D. H., and Watts, T. H. (2004). A switch in costimulation from CD28 to 4-1BB during primary versus secondary CD8 T cell response to influenza in vivo. *J Immunol* 172, 981-988.
- Bieganowska, K. D., Ausubel, L. J., Modabber, Y., Slovik, E., Messersmith, W., and Hafler, D. A. (1997). Direct ex vivo analysis of activated, Fas-sensitive autoreactive T cells in human autoimmune disease. *J Exp Med* 185, 1585-1594.
- Bielekova, B., Goodwin, B., Richert, N., Cortese, I., Kondo, T., Afshar, G., Gran, B., Eaton, J., Antel, J., Frank, J. A., *et al.* (2000). Encephalitogenic potential of the myelin basic protein peptide (amino acids 83-99) in multiple sclerosis: results of a phase II clinical trial with an altered peptide ligand. *Nat Med* 6, 1167-1175.

Blankenstein, T., Cayeux, S., and Qin, Z. (1996). Genetic approaches to cancer immunotherapy. *Rev Physiol Biochem Pharmacol* *129*, 1-49.

Blattman, J. N., and Greenberg, P. D. (2004). Cancer immunotherapy: a treatment for the masses. *Science* *305*, 200-205.

Boatright, K. M., Deis, C., Denault, J. B., Sutherlin, D. P., and Salvesen, G. S. (2004). Activation of caspases-8 and -10 by FLIP(L). *Biochem J* *382*, 651-657.

Boise, L. H., Minn, A. J., Noel, P. J., June, C. H., Accavitti, M. A., Lindsten, T., and Thompson, C. B. (1995). CD28 costimulation can promote T cell survival by enhancing the expression of Bcl-XL. *Immunity* *3*, 87-98.

Borgland, S. L., Bowen, G. P., Wong, N. C., Libermann, T. A., and Muruve, D. A. (2000). Adenovirus vector-induced expression of the C-X-C chemokine IP-10 is mediated through capsid-dependent activation of NF-kappaB. *J Virol* *74*, 3941-3947.

Boussiotis, V. A., Freeman, G. J., Griffin, J. D., Gray, G. S., Gribben, J. G., and Nadler, L. M. (1994). CD2 is involved in maintenance and reversal of human alloantigen-specific clonal anergy. *J Exp Med* *180*, 1665-1673.

Brand, K., Klocke, R., Possling, A., Paul, D., and Strauss, M. (1999). Induction of apoptosis and G2/M arrest by infection with replication-deficient adenovirus at high multiplicity of infection. *Gene Ther* *6*, 1054-1063.

Bretscher, P., and Cohn, M. (1970). A theory of self-nonself discrimination. *Science* *169*, 1042-1049.

Bretscher, P. A. (1999). A two-step, two-signal model for the primary activation of precursor helper T cells. *Proc Natl Acad Sci U S A* *96*, 185-190.

Cannons, J. L., Hoeflich, K. P., Woodgett, J. R., and Watts, T. H. (1999). Role of the stress kinase pathway in signaling via the T cell costimulatory receptor 4-1BB. *J Immunol* *163*, 2990-2998.

Cannons, J. L., Lau, P., Ghuman, B., DeBenedette, M. A., Yagita, H., Okumura, K., and Watts, T. H. (2001). 4-1BB ligand induces cell division, sustains survival, and enhances effector function of CD4 and CD8 T cells with similar efficacy. *J Immunol* *167*, 1313-1324.

Cayeux, S., Beck, C., Aicher, A., Dörken, B., and Blankenstein, T. (1995). Tumor cells cotransfected with interleukin-7 and B7.1 genes induce CD25 and CD28 on tumor-infiltrating T lymphocytes and are strong vaccines. *Eur J Immunol* *25*, 2325-2331.

Cayeux, S., Beck, C., Dörken, B., and Blankenstein, T. (1996). Coexpression of interleukin-4 and B7.1 in murine tumor cells leads to improved tumor rejection and vaccine effect compared to single gene transfectants and a classical adjuvant. *Hum Gene Ther* *7*, 525-529.

- Chang, D. W., Xing, Z., Pan, Y., Algeciras-Schimnich, A., Barnhart, B. C., Yaish-Ohad, S., Peter, M. E., and Yang, X. (2002). c-FLIP(L) is a dual function regulator for caspase-8 activation and CD95-mediated apoptosis. *Embo J* 21, 3704-3714.
- Chen, C., Edelstein, L. C., and Gelinas, C. (2000). The Rel/NF-kappaB family directly activates expression of the apoptosis inhibitor Bcl-x(L). *Mol Cell Biol* 20, 2687-2695.
- Chen, Y., Kuchroo, V. K., Inobe, J., Hafler, D. A., and Weiner, H. L. (1994). Regulatory T cell clones induced by oral tolerance: suppression of autoimmune encephalomyelitis. *Science* 265, 1237-1240.
- Chllichlia, K., Moldenhauer, G., Daniel, P. T., Busslinger, M., Gazzolo, L., Schirrmacher, V., and Khazaie, K. (1995). Immediate effects of reversible HTLV-1 tax function: T-cell activation and apoptosis. *Oncogene* 10, 269-277.
- Chun, H. J., Zheng, L., Ahmad, M., Wang, J., Speirs, C. K., Siegel, R. M., Dale, J. K., Puck, J., Davis, J., Hall, C. G., et al. (2002). Pleiotropic defects in lymphocyte activation caused by caspase-8 mutations lead to human immunodeficiency. *Nature* 419, 395-399.
- Croft, M. (1994). Activation of naive, memory and effector T cells. *Curr Opin Immunol* 6, 431-437.
- Croft, M. (2003). Co-stimulatory members of the TNFR family: keys to effective T-cell immunity? *Nat Rev Immunol* 3, 609-620.
- Croft, M., Bradley, L. M., and Swain, S. L. (1994). Naive versus memory CD4 T cell response to antigen. Memory cells are less dependent on accessory cell costimulation and can respond to many antigen-presenting cell types including resting B cells. *J Immunol* 152, 2675-2685.
- Danial, N. N., and Korsmeyer, S. J. (2004). Cell death: critical control points. *Cell* 116, 205-219.
- Daniel, P. T., Kroidl, A., Cayeux, S., Bargou, R., Blankenstein, T., and Dörken, B. (1997). Costimulatory signals through B7.1/CD28 prevent T cell apoptosis during target cell lysis. *J Immunol* 159, 3808-3815.
- Daniel, P. T., Kroidl, A., Kopp, J., Sturm, I., Moldenhauer, G., Dörken, B., and Pezzutto, A. (1998). Immunotherapy of B-cell lymphoma with CD3x19 bispecific antibodies: costimulation via CD28 prevents "veto" apoptosis of antibody-targeted cytotoxic T cells. *Blood* 92, 4750-4757.
- Daniel, P. T., Scholz, C., Essmann, F., Westermann, J., Pezzutto, A., and Dörken, B. (1999). CD95/Fas-triggered apoptosis of activated T lymphocytes is prevented by dendritic cells through a CD58-dependent mechanism. *Exp Hematol* 27, 1402-1408.
- Danke, N. A., Koelle, D. M., Yee, C., Beheray, S., and Kwok, W. W. (2004). Autoreactive T cells in healthy individuals. *J Immunol* 172, 5967-5972.

- Davis, M. M., Boniface, J. J., Reich, Z., Lyons, D., Hampl, J., Arden, B., and Chien, Y. (1998). Ligand recognition by alpha beta T cell receptors. *Annu Rev Immunol* 16, 523-544.
- Davis, M. M., Krogsgaard, M., Huppa, J. B., Sumen, C., Purbhoo, M. A., Irvine, D. J., Wu, L. C., and Ehrlich, L. (2003). Dynamics of cell surface molecules during T cell recognition. *Annu Rev Biochem* 72, 717-742.
- Dawicki, W., and Watts, T. H. (2004). Expression and function of 4-1BB during CD4 versus CD8 T cell responses in vivo. *Eur J Immunol* 34, 743-751.
- Debatin, K. M., and Krammer, P. H. (2004). Death receptors in chemotherapy and cancer. *Oncogene* 23, 2950-2966.
- Dieckmann, D., Plottner, H., Berchtold, S., Berger, T., and Schuler, G. (2001). Ex vivo isolation and characterization of CD4(+)CD25(+) T cells with regulatory properties from human blood. *J Exp Med* 193, 1303-1310.
- Dranoff, G., Jaffee, E., Lazenby, A., Golumbek, P., Levitsky, H., Brose, K., Jackson, V., Hamada, H., Pardoll, D., and Mulligan, R. C. (1993). Vaccination with irradiated tumor cells engineered to secrete murine granulocyte-macrophage colony-stimulating factor stimulates potent, specific, and long-lasting anti-tumor immunity. *Proc Natl Acad Sci U S A* 90, 3539-3543.
- Dubey, C., Croft, M., and Swain, S. L. (1996). Naive and effector CD4 T cells differ in their requirements for T cell receptor versus costimulatory signals. *J Immunol* 157, 3280-3289.
- Dunn, G. P., Old, L. J., and Schreiber, R. D. (2004). The three Es of cancer immunoediting. *Annu Rev Immunol* 22, 329-360.
- Ehrenstein, M. R., Evans, J. G., Singh, A., Moore, S., Warnes, G., Isenberg, D. A., and Mauri, C. (2004). Compromised function of regulatory T cells in rheumatoid arthritis and reversal by anti-TNFalpha therapy. *J Exp Med* 200, 277-285.
- Eisen, H. N. (2001). Specificity and degeneracy in antigen recognition: yin and yang in the immune system. *Annu Rev Immunol* 19, 1-21.
- Elovaara, I., Koenig, S., Brewah, A. Y., Woods, R. M., Lehky, T., and Jacobson, S. (1993). High human T cell lymphotropic virus type 1 (HTLV-1)-specific precursor cytotoxic T lymphocyte frequencies in patients with HTLV-1-associated neurological disease. *J Exp Med* 177, 1567-1573.
- Freeman, G. J., Boussiotis, V. A., Anumanthan, A., Bernstein, G. M., Ke, X. Y., Rennert, P. D., Gray, G. S., Gribben, J. G., and Nadler, L. M. (1995). B7-1 and B7-2 do not deliver identical costimulatory signals, since B7-2 but not B7-1 preferentially costimulates the initial production of IL-4. *Immunity* 2, 523-532.
- Freitas, A. A., and Rocha, B. (1999). Peripheral T cell survival. *Curr Opin Immunol* 11, 152-156.

Fuchs, E. J., and Matzinger, P. (1992). B cells turn off virgin but not memory T cells. *Science* *258*, 1156-1159.

Fujinami, R. S., and Oldstone, M. B. (1985). Amino acid homology between the encephalitogenic site of myelin basic protein and virus: mechanism for autoimmunity. *Science* *230*, 1043-1045.

Gaur, A., Boehme, S. A., Chalmers, D., Crowe, P. D., Pahuja, A., Ling, N., Brocke, S., Steinman, L., and Conlon, P. J. (1997). Amelioration of relapsing experimental autoimmune encephalomyelitis with altered myelin basic protein peptides involves different cellular mechanisms. *J Neuroimmunol* *74*, 149-158.

Genain, C. P., and Zamvil, S. S. (2000). Specific immunotherapy: one size does not fit all. *Nat Med* *6*, 1098-1100.

Gilly, F. N., Beaujard, A., Bienvenu, J., Trillet Lenoir, V., Glehen, O., Thouvenot, D., Malcus, C., Favrot, M., Dumontet, C., Lombard-Bohas, C., *et al.* (1999). Gene therapy with Adv-IL-2 in unresectable digestive cancer: phase I-II study, intermediate report. *Hepatogastroenterology* *46 Suppl 1*, 1268-1273.

Gimmi, C. D., Freeman, G. J., Gribben, J. G., Gray, G., and Nadler, L. M. (1993). Human T-cell clonal anergy is induced by antigen presentation in the absence of B7 costimulation. *Proc Natl Acad Sci U S A* *90*, 6586-6590.

Graham, F. L. (2000). Adenovirus vectors for high-efficiency gene transfer into mammalian cells. *Immunol Today* *21*, 426-428.

Green, D. R., Droin, N., and Pinkoski, M. (2003). Activation-induced cell death in T cells. *Immunol Rev* *193*, 70-81.

Greenfield, E. A., Howard, E., Paradis, T., Nguyen, K., Benazzo, F., McLean, P., Höllsberg, P., Davis, G., Hafler, D. A., Sharpe, A. H., *et al.* (1997). B7.2 expressed by T cells does not induce CD28-mediated costimulatory activity but retains CTLA4 binding: implications for induction of antitumor immunity to T cell tumors. *J Immunol* *158*, 2025-2034.

Greenwald, R. J., Freeman, G. J., and Sharpe, A. H. (2005). The B7 Family Revisited. *Annu Rev Immunol* *23*, 515-548.

Han, D. K., Chaudhary, P. M., Wright, M. E., Friedman, C., Trask, B. J., Riedel, R. T., Baskin, D. G., Schwartz, S. M., and Hood, L. (1997). MRIT, a novel death-effector domain-containing protein, interacts with caspases and BclXL and initiates cell death. *Proc Natl Acad Sci U S A* *94*, 11333-11338.

- Harding, F. A., McArthur, J. G., Gross, J. A., Raulet, D. H., and Allison, J. P. (1992). CD28-mediated signalling co-stimulates murine T cells and prevents induction of anergy in T-cell clones. *Nature* *356*, 607-609.
- Hathcock, K. S., Laszlo, G., Pucillo, C., Linsley, P., and Hodes, R. J. (1994). Comparative analysis of B7-1 and B7-2 costimulatory ligands: expression and function. *J Exp Med* *180*, 631-640.
- Hock, H., Dorsch, M., Kunzendorf, U., Uberla, K., Qin, Z., Diamantstein, T., and Blankenstein, T. (1993). Vaccinations with tumor cells genetically engineered to produce different cytokines: effectiveness not superior to a classical adjuvant. *Cancer Res* *53*, 714-716.
- Hohlfeld, R., and Wekerle, H. (2004). Autoimmune concepts of multiple sclerosis as a basis for selective immunotherapy: from pipe dreams to (therapeutic) pipelines. *Proc Natl Acad Sci U S A* *101 Suppl 2*, 14599-14606.
- Höllsberg, P., and Hafler, D. A. (1993). Seminars in medicine of the Beth Israel Hospital, Boston. Pathogenesis of diseases induced by human lymphotropic virus type I infection. *N Engl J Med* *328*, 1173-1182.
- Höllsberg, P., Scholz, C., Anderson, D. E., Greenfield, E. A., Kuchroo, V. K., Freeman, G. J., and Hafler, D. A. (1997). Expression of a hypoglycosylated form of CD86 (B7-2) on human T cells with altered binding properties to CD28 and CTLA-4. *J Immunol* *159*, 4799-4805.
- Hou, S., Hyland, L., Ryan, K. W., Portner, A., and Doherty, P. C. (1994). Virus-specific CD8+ T-cell memory determined by clonal burst size. *Nature* *369*, 652-654.
- Hu, S., Vincenz, C., Buller, M., and Dixit, V. M. (1997). A novel family of viral death effector domain-containing molecules that inhibit both CD-95- and tumor necrosis factor receptor-1-induced apoptosis. *J Biol Chem* *272*, 9621-9624.
- Huang, A. Y., Bruce, A. T., Pardoll, D. M., and Levitsky, H. I. (1996). Does B7-1 expression confer antigen-presenting cell capacity to tumors in vivo? *J Exp Med* *183*, 769-776.
- Hunt, D. F., Henderson, R. A., Shabanowitz, J., Sakaguchi, K., Michel, H., Sevilir, N., Cox, A. L., Appella, E., and Engelhard, V. H. (1992). Characterization of peptides bound to the class I MHC molecule HLA-A2.1 by mass spectrometry. *Science* *255*, 1261-1263.
- Hurtado, J. C., Kim, Y. J., and Kwon, B. S. (1997). Signals through 4-1BB are costimulatory to previously activated splenic T cells and inhibit activation-induced cell death. *J Immunol* *158*, 2600-2609.
- Ignatowicz, L., Kappler, J., and Marrack, P. (1996). The repertoire of T cells shaped by a single MHC/peptide ligand. *Cell* *84*, 521-529.

Ignatowicz, L., Rees, W., Pacholczyk, R., Ignatowicz, H., Kushnir, E., Kappler, J., and Marrack, P. (1997). T cells can be activated by peptides that are unrelated in sequence to their selecting peptide. *Immunity* 7, 179-186.

Inohara, N., Koseki, T., Hu, Y., Chen, S., and Nunez, G. (1997). CLARP, a death effector domain-containing protein interacts with caspase-8 and regulates apoptosis. *Proc Natl Acad Sci U S A* 94, 10717-10722.

Irmler, M., Thome, M., Hahne, M., Schneider, P., Hofmann, K., Steiner, V., Bodmer, J. L., Schroter, M., Burns, K., Mattmann, C., *et al.* (1997). Inhibition of death receptor signals by cellular FLIP. *Nature* 388, 190-195.

Jenkins, M. K., and Schwartz, R. H. (1987). Antigen presentation by chemically modified splenocytes induces antigen-specific T cell unresponsiveness in vitro and in vivo. *J Exp Med* 165, 302-319.

Jones, R. G., Parsons, M., Bonnard, M., Chan, V. S., Yeh, W. C., Woodgett, J. R., and Ohashi, P. S. (2000). Protein kinase B regulates T lymphocyte survival, nuclear factor kappaB activation, and Bcl-X(L) levels in vivo. *J Exp Med* 191, 1721-1734.

Jonuleit, H., Schmitt, E., Stassen, M., Tuettenberg, A., Knop, J., and Enk, A. H. (2001). Identification and functional characterization of human CD4(+)CD25(+) T cells with regulatory properties isolated from peripheral blood. *J Exp Med* 193, 1285-1294.

June, C. H., Bluestone, J. A., Nadler, L. M., and Thompson, C. B. (1994). The B7 and CD28 receptor families. *Immunol Today* 15, 321-331.

Kane, L. P., Andres, P. G., Howland, K. C., Abbas, A. K., and Weiss, A. (2001). Akt provides the CD28 costimulatory signal for up-regulation of IL-2 and IFN-gamma but not TH2 cytokines. *Nat Immunol* 2, 37-44.

Kane, L. P., Shapiro, V. S., Stokoe, D., and Weiss, A. (1999). Induction of NF-kappaB by the Akt/PKB kinase. *Curr Biol* 9, 601-604.

Kappler, J. W., Roehm, N., and Marrack, P. (1987). T cell tolerance by clonal elimination in the thymus. *Cell* 49, 273-280.

Kappos, L., Comi, G., Panitch, H., Oger, J., Antel, J., Conlon, P., and Steinman, L. (2000). Induction of a non-encephalitogenic type 2 T helper-cell autoimmune response in multiple sclerosis after administration of an altered peptide ligand in a placebo-controlled, randomized phase II trial. The Altered Peptide Ligand in Relapsing MS Study Group. *Nat Med* 6, 1176-1182.

Kerschensteiner, M., Stadelmann, C., Dechant, G., Wekerle, H., and Hohlfeld, R. (2003). Neurotrophic cross-talk between the nervous and immune systems: implications for neurological diseases. *Ann Neurol* 53, 292-304.

- Kimachi, K., Croft, M., and Grey, H. M. (1997). The minimal number of antigen-major histocompatibility complex class II complexes required for activation of naive and primed T cells. *Eur J Immunol* 27, 3310-3317.
- Kimachi, K., Sugie, K., and Grey, H. M. (2003). Effector T cells have a lower ligand affinity threshold for activation than naive T cells. *Int Immunol* 15, 885-892.
- Kimata, J. T., Palker, T. J., and Ratner, L. (1993). The mitogenic activity of human T-cell leukemia virus type I is T-cell associated and requires the CD2/LFA-3 activation pathway. *J Virol* 67, 3134-3141.
- Kirchhoff, S., Muller, W. W., Krueger, A., Schmitz, I., and Krammer, P. H. (2000a). TCR-mediated up-regulation of c-FLIPshort correlates with resistance toward CD95-mediated apoptosis by blocking death-inducing signaling complex activity. *J Immunol* 165, 6293-6300.
- Kirchhoff, S., Muller, W. W., Li-Weber, M., and Krammer, P. H. (2000b). Up-regulation of c-FLIPshort and reduction of activation-induced cell death in CD28-costimulated human T cells. *Eur J Immunol* 30, 2765-2774.
- Kisielow, P., Teh, H. S., Bluthmann, H., and von Boehmer, H. (1988). Positive selection of antigen-specific T cells in thymus by restricting MHC molecules. *Nature* 335, 730-733.
- Korsmeyer, S. J., Wei, M. C., Saito, M., Weiler, S., Oh, K. J., and Schlesinger, P. H. (2000). Pro-apoptotic cascade activates BID, which oligomerizes BAK or BAX into pores that result in the release of cytochrome c. *Cell Death Differ* 7, 1166-1173.
- Kotera, Y., Shimizu, K., and Mule, J. J. (2001). Comparative analysis of necrotic and apoptotic tumor cells as a source of antigen(s) in dendritic cell-based immunization. *Cancer Res* 61, 8105-8109.
- Krammer, P. H. (2000). CD95's deadly mission in the immune system. *Nature* 407, 789-795.
- Kriegel, M. A., Lohmann, T., Gabler, C., Blank, N., Kalden, J. R., and Lorenz, H. M. (2004). Defective suppressor function of human CD4+ CD25+ regulatory T cells in autoimmune polyglandular syndrome type II. *J Exp Med* 199, 1285-1291.
- Krogsgaard, M., Wucherpfennig, K. W., Cannella, B., Hansen, B. E., Svejgaard, A., Pyrdol, J., Ditzel, H., Raine, C., Engberg, J., and Fugger, L. (2000). Visualization of myelin basic protein (MBP) T cell epitopes in multiple sclerosis lesions using a monoclonal antibody specific for the human histocompatibility leukocyte antigen (HLA)-DR2-MBP 85-99 complex. *J Exp Med* 191, 1395-1412.
- Krueger, A., Fas, S. C., Baumann, S., and Krammer, P. H. (2003). The role of CD95 in the regulation of peripheral T-cell apoptosis. *Immunol Rev* 193, 58-69.
- Krummel, M. F., and Allison, J. P. (1995). CD28 and CTLA-4 have opposing effects on the response of T cells to stimulation. *J Exp Med* 182, 459-465.

Krummel, M. F., and Allison, J. P. (1996). CTLA-4 engagement inhibits IL-2 accumulation and cell cycle progression upon activation of resting T cells. *J Exp Med* 183, 2533-2540.

Kuball, J., Schuler, M., Antunes Ferreira, E., Herr, W., Neumann, M., Obenauer-Kutner, L., Westreich, L., Huber, C., Wolfel, T., and Theobald, M. (2002). Generating p53-specific cytotoxic T lymphocytes by recombinant adenoviral vector-based vaccination in mice, but not man. *Gene Ther* 9, 833-843.

Kuchroo, V. K., Das, M. P., Brown, J. A., Ranger, A. M., Zamvil, S. S., Sobel, R. A., Weiner, H. L., Nabavi, N., and Glimcher, L. H. (1995). B7-1 and B7-2 costimulatory molecules activate differentially the Th1/Th2 developmental pathways: application to autoimmune disease therapy. *Cell* 80, 707-718.

Kundig, T. M., Bachmann, M. F., DiPaolo, C., Simard, J. J., Battegay, M., Lother, H., Gessner, A., Kuhlcke, K., Ohashi, P. S., Hengartner, H., and et al. (1995). Fibroblasts as efficient antigen-presenting cells in lymphoid organs. *Science* 268, 1343-1347.

Lanier, L. L., O'Fallon, S., Somoza, C., Phillips, J. H., Linsley, P. S., Okumura, K., Ito, D., and Azuma, M. (1995). CD80 (B7) and CD86 (B70) provide similar costimulatory signals for T cell proliferation, cytokine production, and generation of CTL. *J Immunol* 154, 97-105.

Lanzavecchia, A., Roosnek, E., Gregory, T., Berman, P., and Abrignani, S. (1988). T cells can present antigens such as HIV gp120 targeted to their own surface molecules. *Nature* 334, 530-532.

Lau, L. L., Jamieson, B. D., Somasundaram, T., and Ahmed, R. (1994). Cytotoxic T-cell memory without antigen. *Nature* 369, 648-652.

Lee, H. W., Nam, K. O., Park, S. J., and Kwon, B. S. (2003). 4-1BB enhances CD8+ T cell expansion by regulating cell cycle progression through changes in expression of cyclins D and E and cyclin-dependent kinase inhibitor p27kip1. *Eur J Immunol* 33, 2133-2141.

Lee, H. W., Park, S. J., Choi, B. K., Kim, H. H., Nam, K. O., and Kwon, B. S. (2002). 4-1BB promotes the survival of CD8+ T lymphocytes by increasing expression of Bcl-xL and Bfl-1. *J Immunol* 169, 4882-4888.

Lenschow, D. J., Ho, S. C., Sattar, H., Rhee, L., Gray, G., Nabavi, N., Herold, K. C., and Bluestone, J. A. (1995). Differential effects of anti-B7-1 and anti-B7-2 monoclonal antibody treatment on the development of diabetes in the nonobese diabetic mouse. *J Exp Med* 181, 1145-1155.

Lenschow, D. J., Su, G. H., Zuckerman, L. A., Nabavi, N., Jellis, C. L., Gray, G. S., Miller, J., and Bluestone, J. A. (1993). Expression and functional significance of an additional ligand for CTLA-4. *Proc Natl Acad Sci U S A* 90, 11054-11058.

Lenschow, D. J., Walunas, T. L., and Bluestone, J. A. (1996). CD28/B7 system of T cell costimulation. *Annu Rev Immunol* 14, 233-258.

- Li, C. J., Friedman, D. J., Wang, C., Metelev, V., and Pardee, A. B. (1995). Induction of apoptosis in uninfected lymphocytes by HIV-1 Tat protein. *Science* *268*, 429-431.
- Li, D., Gal, I., Vermes, C., Alegre, M. L., Chong, A. S., Chen, L., Shao, Q., Adarichev, V., Xu, X., Koreny, T., *et al.* (2004). Cutting edge: Cbl-b: one of the key molecules tuning CD28- and CTLA-4-mediated T cell costimulation. *J Immunol* *173*, 7135-7139.
- Liu, Y., and Janeway, C. A., Jr. (1992). Cells that present both specific ligand and costimulatory activity are the most efficient inducers of clonal expansion of normal CD4 T cells. *Proc Natl Acad Sci U S A* *89*, 3845-3849.
- London, C. A., Lodge, M. P., and Abbas, A. K. (2000). Functional responses and costimulator dependence of memory CD4+ T cells. *J Immunol* *164*, 265-272.
- Lovett-Racke, A. E., Trotter, J. L., Lauber, J., Perrin, P. J., June, C. H., and Racke, M. K. (1998). Decreased dependence of myelin basic protein-reactive T cells on CD28-mediated costimulation in multiple sclerosis patients. A marker of activated/memory T cells. *J Clin Invest* *101*, 725-730.
- Luo, X., Budihardjo, I., Zou, H., Slaughter, C., and Wang, X. (1998). Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors. *Cell* *94*, 481-490.
- Luqman, M., and Bottomly, K. (1992). Activation requirements for CD4+ T cells differing in CD45R expression. *J Immunol* *149*, 2300-2306.
- Madsen, L. S., Andersson, E. C., Jansson, L., krogsgaard, M., Andersen, C. B., Engberg, J., Strominger, J. L., Svejgaard, A., Hjorth, J. P., Holmdahl, R., *et al.* (1999). A humanized model for multiple sclerosis using HLA-DR2 and a human T-cell receptor. *Nat Genet* *23*, 343-347.
- Markovic-Plese, S., Cortese, I., Wandinger, K. P., McFarland, H. F., and Martin, R. (2001). CD4+CD28- costimulation-independent T cells in multiple sclerosis. *J Clin Invest* *108*, 1185-1194.
- Martin, R., Jaraquemada, D., Flerlage, M., Richert, J., Whitaker, J., Long, E. O., McFarlin, D. E., and McFarland, H. F. (1990). Fine specificity and HLA restriction of myelin basic protein-specific cytotoxic T cell lines from multiple sclerosis patients and healthy individuals. *J Immunol* *145*, 540-548.
- Martin, R., McFarland, H. F., and McFarlin, D. E. (1992). Immunological aspects of demyelinating diseases. *Annu Rev Immunol* *10*, 153-187.
- McKnight, A. J., Perez, V. L., Shea, C. M., Gray, G. S., and Abbas, A. K. (1994). Costimulator dependence of lymphokine secretion by naive and activated CD4+ T lymphocytes from TCR transgenic mice. *J Immunol* *152*, 5220-5225.

- Meuer, S. C., Hussey, R. E., Fabbi, M., Fox, D., Acuto, O., Fitzgerald, K. A., Hodgdon, J. C., Protentis, J. P., Schlossman, S. F., and Reinherz, E. L. (1984). An alternative pathway of T-cell activation: a functional role for the 50 kd T11 sheep erythrocyte receptor protein. *Cell* *36*, 897-906.
- Micheau, O., Thome, M., Schneider, P., Holler, N., Tschopp, J., Nicholson, D. W., Briand, C., and Grutter, M. G. (2002). The long form of FLIP is an activator of caspase-8 at the Fas death-inducing signaling complex. *J Biol Chem* *277*, 45162-45171.
- Miller, A., Lider, O., Roberts, A. B., Sporn, M. B., and Weiner, H. L. (1992). Suppressor T cells generated by oral tolerization to myelin basic protein suppress both in vitro and in vivo immune responses by the release of transforming growth factor beta after antigen-specific triggering. *Proc Natl Acad Sci U S A* *89*, 421-425.
- Moalem, G., Leibowitz-Amit, R., Yoles, E., Mor, F., Cohen, I. R., and Schwartz, M. (1999). Autoimmune T cells protect neurons from secondary degeneration after central nervous system axotomy. *Nat Med* *5*, 49-55.
- Morelli, A. E., Larregina, A. T., Ganster, R. W., Zahorchak, A. F., Plowey, J. M., Takayama, T., Logar, A. J., Robbins, P. D., Falo, L. D., and Thomson, A. W. (2000). Recombinant adenovirus induces maturation of dendritic cells via an NF-kappaB-dependent pathway. *J Virol* *74*, 9617-9628.
- Nam, K. O., Kang, H., Shin, S. M., Cho, K. H., Kwon, B., Kwon, B. S., Kim, S. J., and Lee, H. W. (2005). Cross-linking of 4-1BB activates TCR-signaling pathways in CD8+ T lymphocytes. *J Immunol* *174*, 1898-1905.
- Natesan, M., Razi-Wolf, Z., and Reiser, H. (1996). Costimulation of IL-4 production by murine B7-1 and B7-2 molecules. *J Immunol* *156*, 2783-2791.
- Nicholson, L. B., Murtaza, A., Hafler, B. P., Sette, A., and Kuchroo, V. K. (1997). A T cell receptor antagonist peptide induces T cells that mediate bystander suppression and prevent autoimmune encephalomyelitis induced with multiple myelin antigens. *Proc Natl Acad Sci U S A* *94*, 9279-9284.
- Ohashi, P. S., Oehen, S., Buerki, K., Pircher, H., Ohashi, C. T., Odermatt, B., Malissen, B., Zinkernagel, R. M., and Hengartner, H. (1991). Ablation of "tolerance" and induction of diabetes by virus infection in viral antigen transgenic mice. *Cell* *65*, 305-317.
- Ohno, H., Ushiyama, C., Taniguchi, M., Germain, R. N., and Saito, T. (1991). CD2 can mediate TCR/CD3-independent T cell activation. *J Immunol* *146*, 3742-3746.
- Ota, K., Matsui, M., Milford, E. L., Mackin, G. A., Weiner, H. L., and Hafler, D. A. (1990). T-cell recognition of an immunodominant myelin basic protein epitope in multiple sclerosis. *Nature* *346*, 183-187.

- Ozes, O. N., Mayo, L. D., Gustin, J. A., Pfeffer, S. R., Pfeffer, L. M., and Donner, D. B. (1999). NF-kappaB activation by tumour necrosis factor requires the Akt serine-threonine kinase. *Nature* *401*, 82-85.
- Parry, R. V., Reif, K., Smith, G., Sansom, D. M., Hemmings, B. A., and Ward, S. G. (1997). Ligation of the T cell co-stimulatory receptor CD28 activates the serine-threonine protein kinase B. *Eur J Immunol* *27*, 2495-2501.
- Peter, M. E. (2004). The flip side of FLIP. *Biochem J* *382*, e1-3.
- Peter, M. E., Kischkel, F. C., Scheuerpflug, C. G., Medema, J. P., Debatin, K. M., and Krammer, P. H. (1997). Resistance of cultured peripheral T cells towards activation-induced cell death involves a lack of recruitment of FLICE (MACH/caspase 8) to the CD95 death-inducing signaling complex. *Eur J Immunol* *27*, 1207-1212.
- Plas, D. R., Rathmell, J. C., and Thompson, C. B. (2002). Homeostatic control of lymphocyte survival: potential origins and implications. *Nat Immunol* *3*, 515-521.
- Prasad, K. V., Cai, Y. C., Raab, M., Duckworth, B., Cantley, L., Shoelson, S. E., and Rudd, C. E. (1994). T-cell antigen CD28 interacts with the lipid kinase phosphatidylinositol 3-kinase by a cytoplasmic Tyr(P)-Met-Xaa-Met motif. *Proc Natl Acad Sci U S A* *91*, 2834-2838.
- Qin, Z., and Blankenstein, T. (2004). A cancer immunosurveillance controversy. *Nat Immunol* *5*, 3-4; author reply 4-5.
- Racke, M. K., Dhib-Jalbut, S., Cannella, B., Albert, P. S., Raine, C. S., and McFarlin, D. E. (1991). Prevention and treatment of chronic relapsing experimental allergic encephalomyelitis by transforming growth factor-beta 1. *J Immunol* *146*, 3012-3017.
- Refaeli, Y., Van Parijs, L., London, C. A., Tschopp, J., and Abbas, A. K. (1998). Biochemical mechanisms of IL-2-regulated Fas-mediated T cell apoptosis. *Immunity* *8*, 615-623.
- Reif, K., and Cantrell, D. A. (1998). Networking Rho family GTPases in lymphocytes. *Immunity* *8*, 395-401.
- Richardson, J. H., Edwards, A. J., Cruickshank, J. K., Rudge, P., and Dalgleish, A. G. (1990). In vivo cellular tropism of human T-cell leukemia virus type 1. *J Virol* *64*, 5682-5687.
- Riley, J. L., and June, C. H. (2005). The CD28 family: a T-cell rheostat for therapeutic control of T-cell activation. *Blood* *105*, 13-21.
- Rittner, H. L., Zettl, A., Jendro, M. C., Bartz-Bazzanella, P., Goronzy, J. J., and Weyand, C. M. (1997). Multiple mechanisms support oligoclonal T cell expansion in rheumatoid synovitis. *Mol Med* *3*, 452-465.

- Rogers, P. R., Dubey, C., and Swain, S. L. (2000). Qualitative changes accompany memory T cell generation: faster, more effective responses at lower doses of antigen. *J Immunol* *164*, 2338-2346.
- Romashkova, J. A., and Makarov, S. S. (1999). NF-kappaB is a target of AKT in anti-apoptotic PDGF signalling. *Nature* *401*, 86-90.
- Ronchese, F., and Hausmann, B. (1993). B lymphocytes in vivo fail to prime naive T cells but can stimulate antigen-experienced T lymphocytes. *J Exp Med* *177*, 679-690.
- Sagerstrom, C. G., Kerr, E. M., Allison, J. P., and Davis, M. M. (1993). Activation and differentiation requirements of primary T cells in vitro. *Proc Natl Acad Sci U S A* *90*, 8987-8991.
- Sakaguchi, S. (2000). Regulatory T cells: key controllers of immunologic self-tolerance. *Cell* *101*, 455-458.
- Sakaguchi, S., Sakaguchi, N., Asano, M., Itoh, M., and Toda, M. (1995). Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *J Immunol* *155*, 1151-1164.
- Salomon, B., and Bluestone, J. A. (2001). Complexities of CD28/B7: CTLA-4 costimulatory pathways in autoimmunity and transplantation. *Annu Rev Immunol* *19*, 225-252.
- Sanchez-Madrid, F., Krensky, A. M., Ware, C. F., Robbins, E., Strominger, J. L., Burakoff, S. J., and Springer, T. A. (1982). Three distinct antigens associated with human T-lymphocyte-mediated cytolysis: LFA-1, LFA-2, and LFA-3. *Proc Natl Acad Sci U S A* *79*, 7489-7493.
- Saoulli, K., Lee, S. Y., Cannons, J. L., Yeh, W. C., Santana, A., Goldstein, M. D., Bangia, N., DeBenedette, M. A., Mak, T. W., Choi, Y., and Watts, T. H. (1998). CD28-independent, TRAF2-dependent costimulation of resting T cells by 4-1BB ligand. *J Exp Med* *187*, 1849-1862.
- Scaffidi, C., Fulda, S., Srinivasan, A., Friesen, C., Li, F., Tomaselli, K. J., Debatin, K. M., Krammer, P. H., and Peter, M. E. (1998). Two CD95 (APO-1/Fas) signaling pathways. *Embo J* *17*, 1675-1687.
- Scaffidi, C., Schmitz, I., Krammer, P. H., and Peter, M. E. (1999). The role of c-FLIP in modulation of CD95-induced apoptosis. *J Biol Chem* *274*, 1541-1548.
- Schirmer, M., Vallejo, A. N., Weyand, C. M., and Goronzy, J. J. (1998). Resistance to apoptosis and elevated expression of Bcl-2 in clonally expanded CD4+CD28- T cells from rheumatoid arthritis patients. *J Immunol* *161*, 1018-1025.
- Schmidt, D., Goronzy, J. J., and Weyand, C. M. (1996). CD4+ CD7- CD28- T cells are expanded in rheumatoid arthritis and are characterized by autoreactivity. *J Clin Invest* *97*, 2027-2037.

- Schmitz, I., Krueger, A., Baumann, S., Schulze-Bergkamen, H., Krammer, P. H., and Kirchhoff, S. (2003). An IL-2-dependent switch between CD95 signaling pathways sensitizes primary human T cells toward CD95-mediated activation-induced cell death. *J Immunol* *171*, 2930-2936.
- Scholz, C., Freeman, G. J., Greenfield, E. A., Hafler, D. A., and Höllsberg, P. (1996a). Activation of human T cell lymphotropic virus type I-infected T cells is independent of B7 costimulation. *J Immunol* *157*, 2932-2938.
- Scholz, C., Hafler, D. A., and Höllsberg, P. (1996b). Downregulation of IL-10 secretion and enhanced antigen-presenting abilities following HTLV-I infection of T cells. *J Neurosci Res* *45*, 786-794.
- Scholz, C., Patton, K. T., Anderson, D. E., Freeman, G. J., and Hafler, D. A. (1998). Expansion of autoreactive T cells in multiple sclerosis is independent of exogenous B7 costimulation. *J Immunol* *160*, 1532-1538.
- Scholz, C., Stärck, L., Willimsky, G., Blankenstein, T., Dörken, B., and Daniel, P. T. (2002). Adenoviral transduction of tumor cells induces apoptosis in co-cultured T lymphocytes. *Gene Ther* *9*, 1438-1446.
- Schulze-Bergkamen, H., and Krammer, P. H. (2004). Apoptosis in cancer--implications for therapy. *Semin Oncol* *31*, 90-119.
- Schwartz, R. H. (1990). A cell culture model for T lymphocyte clonal anergy. *Science* *248*, 1349-1356.
- Schwartz, R. H. (2003). T cell anergy. *Annu Rev Immunol* *21*, 305-334.
- Sebzda, E., Wallace, V. A., Mayer, J., Yeung, R. S., Mak, T. W., and Ohashi, P. S. (1994). Positive and negative thymocyte selection induced by different concentrations of a single peptide. *Science* *263*, 1615-1618.
- Sharpe, A. H., and Freeman, G. J. (2002). The B7-CD28 superfamily. *Nat Rev Immunol* *2*, 116-126.
- Shih, F. F., and Allen, P. M. (2004). T cells are not as degenerate as you think, once you get to know them. *Mol Immunol* *40*, 1041-1046.
- Shu, H. B., Halpin, D. R., and Goeddel, D. V. (1997). Casper is a FADD- and caspase-related inducer of apoptosis. *Immunity* *6*, 751-763.
- Skurk, C., Maatz, H., Kim, H. S., Yang, J., Abid, M. R., Aird, W. C., and Walsh, K. (2004). The Akt-regulated forkhead transcription factor FOXO3a controls endothelial cell viability through modulation of the caspase-8 inhibitor FLIP. *J Biol Chem* *279*, 1513-1525.
- Sloan-Lancaster, J., and Allen, P. M. (1996). Altered peptide ligand-induced partial T cell activation: molecular mechanisms and role in T cell biology. *Annu Rev Immunol* *14*, 1-27.

Sospedra, M., and Martin, R. (2005). Immunology of multiple sclerosis *. *Annu Rev Immunol* 23, 683-747.

Sprent, J., and Surh, C. D. (2002). T cell memory. *Annu Rev Immunol* 20, 551-579.

Srinivasula, S. M., Ahmad, M., Otilie, S., Bullrich, F., Banks, S., Wang, Y., Fernandes-Alnemri, T., Croce, C. M., Litwack, G., Tomaselli, K. J., *et al.* (1997). FLAME-1, a novel FADD-like anti-apoptotic molecule that regulates Fas/TNFR1-induced apoptosis. *J Biol Chem* 272, 18542-18545.

Stärck, L., Scholz, C., Blankenstein, T., Dörken, B., and Daniel, P. T. (2005a). Necrotic death but not irradiation abolishes costimulation of T-cell effector functions and survival by CD80-expressing tumor cells. *Int J Cancer* 116, 78-86.

Stärck, L., Scholz, C., Dörken, B., and Daniel, P. T. (2005b). Costimulation by CD137/4-1BB inhibits T cell apoptosis and induces Bcl-x(L) and c-FLIP(short) via phosphatidylinositol 3-kinase and AKT/protein kinase B. *Eur J Immunol* 35, 1257-1266.

Stewart, A. K., Lassam, N. J., Quirt, I. C., Bailey, D. J., Rotstein, L. E., Krajden, M., Dessureault, S., Gallinger, S., Cappe, D., Wan, Y., *et al.* (1999). Adenovector-mediated gene delivery of interleukin-2 in metastatic breast cancer and melanoma: results of a phase 1 clinical trial. *Gene Ther* 6, 350-363.

Tejada-Simon, M. V., Zang, Y. C., Hong, J., Rivera, V. M., and Zhang, J. Z. (2003). Cross-reactivity with myelin basic protein and human herpesvirus-6 in multiple sclerosis. *Ann Neurol* 53, 189-197.

Thome, M., Schneider, P., Hofmann, K., Fickenscher, H., Meinl, E., Neipel, F., Mattmann, C., Burns, K., Bodmer, J. L., Schroter, M., *et al.* (1997). Viral FLICE-inhibitory proteins (FLIPs) prevent apoptosis induced by death receptors. *Nature* 386, 517-521.

Thome, M., and Tschoopp, J. (2001). Regulation of lymphocyte proliferation and death by FLIP. *Nat Rev Immunol* 1, 50-58.

Thornton, A. M., and Shevach, E. M. (1998). CD4+CD25+ immunoregulatory T cells suppress polyclonal T cell activation in vitro by inhibiting interleukin 2 production. *J Exp Med* 188, 287-296.

Townsend, A. R., Rothbard, J., Gotch, F. M., Bahadur, G., Wraith, D., and McMichael, A. J. (1986). The epitopes of influenza nucleoprotein recognized by cytotoxic T lymphocytes can be defined with short synthetic peptides. *Cell* 44, 959-968.

Vallejo, A. N., Schirmer, M., Weyand, C. M., and Goronzy, J. J. (2000). Clonality and longevity of CD4+CD28null T cells are associated with defects in apoptotic pathways. *J Immunol* 165, 6301-6307.

Vereecke, R., Buffenoir, G., Preudhomme, C., Hetuin, D., Bauters, F., Fenaux, P., and Quesnel, B. (2000). Gene transfer of GM-CSF, CD80 and CD154 cDNA enhances survival in a murine model of acute leukemia with persistence of a minimal residual disease. *Gene Ther* 7, 1312-1316.

- Viglietta, V., Baecher-Allan, C., Weiner, H. L., and Hafler, D. A. (2004). Loss of functional suppression by CD4+CD25+ regulatory T cells in patients with multiple sclerosis. *J Exp Med* *199*, 971-979.
- Walunas, T. L., Lenschow, D. J., Bakker, C. Y., Linsley, P. S., Freeman, G. J., Green, J. M., Thompson, C. B., and Bluestone, J. A. (1994). CTLA-4 can function as a negative regulator of T cell activation. *Immunity* *1*, 405-413.
- Wang, C. Y., Mayo, M. W., Korneluk, R. G., Goeddel, D. V., and Baldwin, A. S., Jr. (1998). NF-kappaB antiapoptosis: induction of TRAF1 and TRAF2 and c-IAP1 and c-IAP2 to suppress caspase-8 activation. *Science* *281*, 1680-1683.
- Wen, T., Bukczynski, J., and Watts, T. H. (2002). 4-1BB ligand-mediated costimulation of human T cells induces CD4 and CD8 T cell expansion, cytokine production, and the development of cytolytic effector function. *J Immunol* *168*, 4897-4906.
- Wesselborg, S., Janssen, O., and Kabelitz, D. (1993). Induction of activation-driven death (apoptosis) in activated but not resting peripheral blood T cells. *J Immunol* *150*, 4338-4345.
- Willimsky, G., and Blankenstein, T. (2000). Interleukin-7/B7.1-encoding adenoviruses induce rejection of transplanted but not nontransplanted tumors. *Cancer Res* *60*, 685-692.
- Wilson, D. B., Wilson, D. H., Schroder, K., Pinilla, C., Blondelle, S., Houghten, R. A., and Garcia, K. C. (2004). Specificity and degeneracy of T cells. *Mol Immunol* *40*, 1047-1055.
- Windhagen, A., Scholz, C., Höllsberg, P., Fukaura, H., Sette, A., and Hafler, D. A. (1995). Modulation of cytokine patterns of human autoreactive T cell clones by a single amino acid substitution of their peptide ligand. *Immunity* *2*, 373-380.
- Wu, G., Chai, J., Suber, T. L., Wu, J. W., Du, C., Wang, X., and Shi, Y. (2000). Structural basis of IAP recognition by Smac/DIABLO. *Nature* *408*, 1008-1012.
- Wucherpfennig, K. W., Höllsberg, P., Richardson, J. H., Benjamin, D., and Hafler, D. A. (1992). T-cell activation by autologous human T-cell leukemia virus type I-infected T-cell clones. *Proc Natl Acad Sci U S A* *89*, 2110-2114.
- Wucherpfennig, K. W., and Strominger, J. L. (1995). Molecular mimicry in T cell-mediated autoimmunity: viral peptides activate human T cell clones specific for myelin basic protein. *Cell* *80*, 695-705.
- Yee, C., Riddell, S. R., and Greenberg, P. D. (2001). In vivo tracking of tumor-specific T cells. *Curr Opin Immunol* *13*, 141-146.
- Yu, X. Z., Martin, P. J., and Anasetti, C. (2003). CD28 signal enhances apoptosis of CD8 T cells after strong TCR ligation. *J Immunol* *170*, 3002-3006.

Zhang, J., Markovic-Plese, S., Lacet, B., Raus, J., Weiner, H. L., and Hafler, D. A. (1994). Increased frequency of interleukin 2-responsive T cells specific for myelin basic protein and proteolipid protein in peripheral blood and cerebrospinal fluid of patients with multiple sclerosis. *J Exp Med* 179, 973-984.

Zinkernagel, R. M. (2000). What is missing in immunology to understand immunity? *Nat Immunol* 1, 181-185.