



Quantitative Heat-Kernel Estimates for Diffusions with Distributional Drift

Nicolas Perkowski¹ · Willem van Zuijlen²

Received: 22 March 2021 / Accepted: 29 December 2021 / Published online: 27 January 2022
© The Author(s) 2022

Abstract

We consider the stochastic differential equation on \mathbb{R}^d given by

$$dX_t = b(t, X_t)dt + dB_t,$$

where B is a Brownian motion and b is considered to be a distribution of regularity $> -\frac{1}{2}$. We show that the martingale solution of the SDE has a transition kernel Γ_t and prove upper and lower heat-kernel estimates for Γ_t with explicit dependence on t and the norm of b .

Keywords Heat-kernel estimate · Singular diffusion · Parametrix method

Mathematics Subject Classification (2010) Primary 60H10 · Secondary 35A08

1 Introduction and Main Results

In this paper we consider the stochastic differential equation on \mathbb{R}^d given by

$$dX_t = b(t, X_t) dt + dB_t, \tag{1}$$

where B is a Brownian motion and b is a distribution of regularity $> -\frac{1}{2}$. Such *singular diffusions* (diffusions with distributional drift) appear as models for stochastic processes in random media (then b would also be random, but independent of B), for example in [4–6]. They also appear as “stochastic characteristics” in Feynman-Kac type representations of singular SPDEs, for example in [5, 13, 17]. In non-singular SPDEs, the stochastic characteristics would be formulated in terms of the Brownian motion, and they may be useful tools to infer information about the long-time behavior of the SPDE. For example, the asymptotic behavior of the total mass of the parabolic Anderson model is typically derived via

✉ Nicolas Perkowski
perkowski@math.fu-berlin.de

Willem van Zuijlen
vanzuijlen@wias-berlin.de

¹ Free University of Berlin, Arnimallee 7, 14195 Berlin, Germany

² Weierstrass Institute for Applied Analysis and Stochastics, Mohrenstraße 39, 10117 Berlin, Germany

the Feynman-Kac formula [16], and for that purpose it is important that we understand the Brownian motion and its transition probabilities very well. When studying singular variants of the parabolic Anderson model, where the Brownian motion in the Feynman-Kac representation is replaced by a singular diffusion, we thus need to understand the transition probabilities of this singular diffusion. Moreover, since we are interested in the long-time behavior, we need quantitative control of the transition probabilities on arbitrarily long time intervals. This motivates our present work.

We show that the solution to Eq. 1 possesses a transition kernel $\Gamma_t : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ for all $t > 0$. This means that under the measure \mathbb{P}_x such that $X_0 = x$ we have for all $\phi \in C_b(\mathbb{R}^d)$

$$\mathbb{E}_x[\phi(X_t)] = \int_{\mathbb{R}^d} \phi(y)\Gamma_t(x, y) dy.$$

The following theorem represents the main result of our paper, in which we show that the above transition kernel satisfies heat-kernel estimates.

For any Banach space \mathfrak{X} and $t > 0$ we write $\|\cdot\|_{C_t, \mathfrak{X}}$ for the norm on $C([0, t], \mathfrak{X})$, which is defined for $f \in C([0, t], \mathfrak{X})$ by

$$\|f\|_{C_t, \mathfrak{X}} = \sup_{s \in [0, t]} \|f(s)\|_{\mathfrak{X}}.$$

$\Delta_{-1}b$ denotes the first Littlewood-Paley block and $\Delta_{\geq 0}b$ the sum of the positive Littlewood-Paley blocks (see Section 1.2). $B_{p,q}^s$ denotes a Besov space, see [2].

Theorem 1.1 *Let $\alpha \in (0, \frac{1}{2})$ and $c > 1$. There exist a $C > 1$ and a $\kappa \in (0, 1)$ such that for all $b = (b_t)_{t \geq 0} \in C([0, \infty), B_{\infty,1}^{-\alpha}(\mathbb{R}^d, \mathbb{R}^d))$, $\mu \in \mathbb{N}_0^d$ with $|\mu| \leq 1$, and for all $t > 0$, $x, y \in \mathbb{R}^d$:*

$$|\partial_x^\mu \Gamma_t(x, y)| \leq C \exp\left(Ct\left[\|\Delta_{-1}b\|_{C_t L^\infty}^2 + \|\Delta_{\geq 0}b\|_{C_t B_{\infty,1}^{-\alpha}}^{\frac{2}{1-\alpha}}\right]\right) (t^{-\frac{|\mu|}{2}} \vee 1) p(ct, x - y), \tag{2}$$

$$\Gamma_t(x, y) \geq \frac{1}{C} \exp\left(-Ct\left[\|\Delta_{-1}b\|_{C_t L^\infty}^2 + \|\Delta_{\geq 0}b\|_{C_t B_{\infty,1}^{-\alpha}}^{\frac{2}{1-\alpha}}\right]\right) p(\kappa t, x - y), \tag{3}$$

where $p(t, x) = (2\pi t)^{-\frac{d}{2}} e^{-|x|^2/2t}$ is the standard Gaussian kernel.

As a corollary, we obtain the following estimate on the escape probability of the diffusion X to leave a ball.

Corollary 1.2 *Let $\alpha \in (0, \frac{1}{2})$. There exists a $C > 0$ such that for all $b \in C([0, \infty), B_{\infty,1}^{-\alpha}(\mathbb{R}^d, \mathbb{R}^d))$, $x \in \mathbb{R}^d$, $K > 0$ and $T \geq 1$, and for X solving (1) with $\mathbb{P}_x(X_0 = x) = 1$:*

$$\begin{aligned} & \mathbb{P}_x\left(\sup_{t \in [0, T]} |X_t - x| \geq K\right) \\ & \leq C \exp\left(CT\left[\|\Delta_{-1}b\|_{C_T L^\infty}^2 + \|\Delta_{\geq 0}b\|_{C_T B_{\infty,1}^{-\alpha}}^{\frac{2}{1-\alpha}}\right]\right) \exp\left(-\frac{K^2}{CT}\right) \end{aligned} \tag{4}$$

Remark 1.3 At least for constant b the heat-kernel estimates are sharp: If $\lambda \in \mathbb{R}^d$ and $b = \lambda$, then $\Gamma_t(x, y) = p(t, y - x - \lambda t)$ and a simple computation shows that $\sup_{x \in \mathbb{R}^d} \frac{p(t, x - \lambda t)}{p(ct, x)} = c^{\frac{d}{2}} e^{\frac{1}{2(c-1)} t \lambda^2}$ and $\inf_{x \in \mathbb{R}^d} \frac{p(t, x - \lambda t)}{p(\kappa t, x)} = \kappa^{\frac{d}{2}} e^{-\frac{1}{2(1-\kappa)} t \lambda^2}$. Since in that case $\Delta_{\geq 0}b = 0$, this corresponds exactly to our bounds Eqs. 2 and 3 (for $\mu = 0$).

Remark 1.4 As we consider a time inhomogeneous drift, we could have also formulated the heat-kernel estimates for $\Gamma_{s,t}$ (with $0 \leq s < t$), which is the transition kernel from time s to time t : If $\mathbb{P}_{s,x}$ is the probability measure under which $X_s = x$ and Eq. 1 holds (for $t > s$), then $\mathbb{E}_{s,x}[\varphi(X_t)] = \int_{\mathbb{R}^d} \varphi(y)\Gamma_{s,t}(x, y) dy$. However, to simplify notation we only consider the case $s = 0$ and we write Γ_t for $\Gamma_{0,t}$. The heat-kernel estimates for $\Gamma_{s,t}$ follow by applying Theorem 1.1 with $b'_t = b_{t+s}, t \geq 0$.

1.1 Literature

Diffusions with a distributional drift were first considered by Bass and Chen [3] and Flandoli, Russo and Wolf [8], both in the one-dimensional time-homogeneous setting. More recently, Delarue and Diel [6] used Hairer’s rough path approach to singular SPDEs [14, 15] to extend the results of [8] to the time-inhomogeneous case, and they applied this to construct a random directed polymer measure. Flandoli, Issoglio and Russo [7] were the first to consider multidimensional singular diffusions, but they require more regularity than in the previous works on the one-dimensional case (they consider the “Young regime”, i.e., the distributional drift has regularity better than $-1/2$). Zhang and Zhao [22] study the ergodicity and they derive heat-kernel estimates for singular diffusions in the Young regime. Cannizzaro and Chouk [5] use paracontrolled distributions to extend the approach of [6] to higher dimensions and the results of [7] to more singular drifts. They apply this to construct a random polymer measure that is closely related to the parabolic Anderson model.

In this paper we follow the approach of Cannizzaro and Chouk, although we restrict our attention to the more regular Young regime. This is crucial for our arguments.

As already mentioned, Zhang and Zhao [22] also prove heat-kernel estimates for SDEs with distributional drifts in the Young regime. More precisely, they prove that there exist $c, C \geq 1$ such that for all $t \in (0, T]$ and $x, y \in \mathbb{R}^d$

$$\frac{1}{c} p\left(\frac{t}{c}, x - y\right) \leq |\Gamma_t(x, y)| \leq Cp(ct, x - y).$$

Moreover, they give an upper bound on the gradient of the transition kernel, $\nabla \Gamma_t$. Here, the constant C implicitly depends on T and $\|b\|_{\mathcal{C}^{-\alpha}}$.

If b is the gradient of a function that does not dependent on time, then there is classical heat-kernel estimates for Γ , see for example Stroock [20, Theorem 4.3.9]. In that theorem we have $b = \nabla U$ for a smooth and bounded function U , but the estimate only depends on $\max U - \min U$, so by an approximation argument it extends to continuous and bounded U . This result is uniform in time, but also here the dependence of the constants on $\max U - \min U$ is implicit.

In another work by the authors together with W. König [17], our heat-kernel estimates are applied to derive the asymptotic behavior of the total mass of the parabolic Anderson model. In that application it is crucial to understand how the constant grows with t and the norm of b . Therefore, we need our “quantitative version” of the heat-kernel estimates.

1.2 Notation and Conventions

We write $\mathbb{N} = \{1, 2, \dots\}$, $\mathbb{N}_0 = \{0\} \cup \mathbb{N}$ and $\mathbb{N}_{-1} = \{-1\} \cup \mathbb{N}_0$. For the whole paper, d is an element of \mathbb{N} and will denote the dimension of the space. For families $(a_i)_{i \in \mathbb{I}}, (b_i)_{i \in \mathbb{I}}$ in \mathbb{R} for an index set \mathbb{I} , we write $a_i \lesssim b_i$ to denote the existence of a $C > 0$ such that $a_i \leq Cb_i$ for all $i \in \mathbb{I}$. We write C_b for the space of continuous bounded functions and C_b^∞ for the space of C^∞ functions for which all their derivatives are bounded functions. We abbreviate function spaces and Besov spaces by omitting “ (\mathbb{R}^d) ” in the notation, for

example we abbreviate $B_{p,q}^\beta(\mathbb{R}^d)$ to $B_{p,q}^\beta$. Moreover, we write \mathcal{C}^β for $B_{\infty,\infty}^\beta$ and \mathcal{C}_p^β for $B_{p,\infty}^\beta$. We write $u \otimes v$ for the paraproduct between u and v (with the low frequencies of u and the high frequencies of v), and $u \odot v$ for the resonance product; we adopt the notation from [19] and refer to [2] as background material.

In the rest of the paper $(\rho_i)_{i \in \mathbb{N}_{-1}}$ is a *dyadic partition of unity*, meaning that ρ_{-1} is supported in a ball around 0, ρ_0 is supported in an annulus, $\rho_i(x) = \rho_0(2^{-i}x)$ for $i \in \mathbb{N}_0$, $\sum_{i \in \mathbb{N}_{-1}} \rho_i = \mathbb{1}$, $\frac{1}{2} \leq \sum_{i \in \mathbb{N}_{-1}} \rho_i^2 \leq 1$ and $\text{supp } \rho_i \cap \text{supp } \rho_j = \emptyset$ if $|i - j| \geq 2$. For $i \in \mathbb{N}_{-1}$ we write Δ_i for the corresponding Littlewood-Paley blocks (\mathcal{F} denotes the Fourier transform)

$$\Delta_i f = \rho_i(\mathbf{D})f = \mathcal{F}^{-1}(\rho_i \mathcal{F}(f)) = \mathcal{F}^{-1}(\rho_i) * f.$$

Moreover, we define $\Delta_{\geq 0}f$ to be the sum of all the positive Littlewood-Paley blocks:

$$\Delta_{\geq 0}f = \sum_{i \in \mathbb{N}_0} \Delta_i f.$$

2 Diffusions with Distributional Drift and Their Heat-Kernel Estimates

Throughout this section we fix $T > 0$. Let $\alpha \in (0, \frac{1}{2})$. For $b \in C([0, T], B_{\infty,1}^{-\alpha}(\mathbb{R}^d, \mathbb{R}^d))$ we consider the stochastic differential equation

$$dX_t = b(t, X_t) dt + dB_t. \tag{5}$$

For $t > 0$ let \mathcal{L}_t be the operator

$$\mathcal{L}_t = \frac{1}{2} \Delta + b_t \cdot \nabla. \tag{6}$$

We consider the following Cauchy problem for $u: [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ with terminal condition ϕ :

$$\begin{cases} \partial_t u + \mathcal{L}_t u = 0 & \text{on } [0, T) \times \mathbb{R}^d, \\ u(T, \cdot) = \phi & \text{on } \mathbb{R}^d. \end{cases} \tag{7}$$

The solution theory for the Cauchy problem will be given in Proposition 2.4. We write u^ϕ for the solution to Eq. 7. But let us first discuss how to interpret (5) in terms of a martingale problem.

Definition 2.1 We say that a stochastic process $X = (X_t)_{t \in [0, T]}$ on a probability space (Ω, \mathbb{P}) is a *solution to the SDE (5) on $[0, T]$ with initial condition $X_0 = x$* if it satisfies the martingale problem for $((\mathcal{L}_t)_{t \in (0, T]}, \delta_x)$, i.e., if $\mathbb{P}(X_0 = x) = 1$ and for all $f \in C([0, T], L^\infty(\mathbb{R}^d))$, all $\phi \in C_c^\infty(\mathbb{R}^d)$ and for $u = u^\phi$ being the solution to the Cauchy problem (7), the process

$$\left(u(t, X_t) - \int_0^t f(s, X_s) ds \right)_{t \in [0, T]}$$

is a martingale.

The martingale problem has a unique solution:

Theorem 2.2 [5, Theorem 1.2] *Let $\alpha \in (0, \frac{1}{2})$. For all $x \in \mathbb{R}^d$ and $b \in C([0, T], \mathcal{C}^{-\alpha}(\mathbb{R}^d, \mathbb{R}^d))$ there exists a unique solution to the martingale problem for*

$((\mathcal{L}_t)_{t \in (0, T]}, \delta_x)$, in the sense that there is a unique probability measure \mathbb{P}_x on $\Omega = C([0, T], \mathbb{R}^d)$ such that the coordinate process $X_t(\omega) = \omega(t)$ satisfies the martingale problem for $((\mathcal{L}_t)_{t \in (0, T]}, \delta_x)$. Moreover, X is a strong Markov process under \mathbb{P}_x and the measure \mathbb{P}_x depends (weakly) continuously on the drift b .

Remark 2.3 The continuity of the solution \mathbb{P} in terms of the drift is not mentioned in [5, Theorem 1.2], but it can be extracted from their proof.

Observe that Theorem 2.2 also implies that there exists a unique probability measure $\mathbb{P}_{s,x}$ on $C([s, T], \mathbb{R}^d)$ such that the coordinate process satisfies the martingale problem for $((\mathcal{L}_t)_{t \in (s, T]}, \delta_x)$. This can be obtained by applying Theorem 2.2 to a shift of the drift, as is mentioned in Remark 1.4.

Next, our aim is to show that X admits a transition density $\Gamma_{s,t}$ for $0 \leq s < t \leq T$ (Proposition 2.9), which means that for $\varphi \in C_c(\mathbb{R}^d)$ and $x \in \mathbb{R}^d$ and with $\mathbb{P}_{s,x}$ as in Remark 2.3

$$\mathbb{E}_{s,x}[\varphi(X_t)] = \int_{\mathbb{R}^d} \varphi(y) \Gamma_{s,t}(x, y) dy. \tag{8}$$

We do this by showing that $\Gamma_{t,T}(x, y) = u^{\delta_y}(t, x)$ for the solution u^{δ_y} to Eq. 7 with terminal condition $u(T, \cdot) = \delta_y$.

In order to construct the solution u^{δ_y} we have to slightly extend the results of [5]. Indeed, in [5, Theorem 3.1 and 3.2] the well-posedness of the Cauchy problem is shown for $\phi \in \mathcal{C}^\beta$ with $\beta \in (1 + \alpha, 2 - \alpha)$, and δ_z is not in this space. The solution theory in [5] is formulated in terms of mild solutions: A *mild solution* of Eq. 7 is a fixed point u of Φ , i.e., $\Phi u = u$, where Φ is defined on $C([0, T], \mathcal{S}') \cap [\bigcup_{p \in [1, \infty)} C([0, T], \mathcal{C}_p^\beta(\mathbb{R}^d))]$ for $\beta > 1 + \alpha$ by

$$(\Phi u)_s = P_{T-s} \phi - \int_s^T P_{r-s} (b_r \cdot \nabla u_r) dr, \tag{9}$$

where $P_t \phi := p(t, \cdot) * \phi$ for $t > 0$ and $P_0 \phi = \phi$ (that Φ is well-defined follows by 2.6).

In order to allow δ_y as a terminal condition, we will consider a different space that “allows a blowup as $t \uparrow T$ ”. However, for notational elegance, we instead consider a space with “a blowup at 0” and mention that u is a fixed point of Φ if and only if v given by $v(t, \cdot) = u(T - t, \cdot)$ is a fixed point of Θ , given by

$$(\Theta v)_s = P_s \phi + \int_0^s P_{s-r} (b_{T-r} \cdot \nabla v_r) dr, \tag{10}$$

so that we call v a *mild solution* of

$$\begin{cases} \partial_t v - \mathcal{L}_{T-t} v = 0 & \text{on } (0, T] \times \mathbb{R}^d, \\ v(0, \cdot) = \phi & \text{on } \mathbb{R}^d. \end{cases} \tag{11}$$

We will show that Θ has a fixed point in the following space (for suitable δ, β). For $\delta \geq 0, \beta \in \mathbb{R}$ and $t > 0$ we define

$$\begin{aligned} \|u\|_{M_t^\delta \mathcal{C}_p^\beta} &= \sup_{s \in (0, t]} s^\delta \|u_s\|_{\mathcal{C}_p^\beta}, \\ M_t^\delta \mathcal{C}_p^\beta &= \{u \in C((0, t], \mathcal{C}_p^\beta) : \|u\|_{M_t^\delta \mathcal{C}_p^\beta} < \infty\}. \end{aligned}$$

The following proposition is a slight extension of [5, Theorem 3.1 and 3.2].

Proposition 2.4 *Let $\alpha \in (0, \frac{1}{2})$, $p \in [1, \infty]$ and $\gamma > \alpha - 1$. For $\phi \in \mathcal{C}_p^\gamma$, $b \in C([0, T], B_{\infty,1}^{-\alpha})$, $\beta \in (1 + \alpha, 2 - \alpha)$ and $\varepsilon > 0$ the Cauchy problem (7) has a unique mild solution $u^{\phi,b}$ in $C([0, T], \mathcal{C}_p^{(\gamma-\varepsilon)\wedge\beta}) \cap C([0, T], \mathcal{C}_p^\beta)$ such that $u^{\phi,b}(t) \in \mathcal{C}^\beta$ for all $t \in [0, T)$. Moreover, for all $t > 0$ the map $\mathcal{C}_p^\gamma \times C([0, T], B_{\infty,1}^{-\alpha}) \rightarrow \mathcal{C}^\beta$ given by $(\phi, b) \mapsto u^{\phi,b}(t, \cdot)$ is locally Lipschitz.*

Another difference with [5] is that we consider $b \in C([0, T], B_{\infty,1}^{-\alpha})$ instead of $b \in C([0, T], \mathcal{C}^{-\alpha})$. Since $B_{\infty,1}^{-\alpha} \subset \mathcal{C}^{-\alpha} \subset B_{\infty,1}^{-\alpha-\varepsilon}$ (as continuous embeddings), this does not make much of a difference. But our heat-kernel estimates depend on the $B_{\infty,1}^{-\alpha}$ -norm and for their derivation it is more convenient to work with $B_{\infty,1}^{-\alpha}$.

Before we prove Proposition 2.4 we present two auxiliary facts, Lemma 2.5 and 2.6.

We write B for the beta function (see e.g. [1, Section 1.1]), which is the function $B : (0, \infty)^2 \rightarrow (0, \infty)$ given by

$$B(\beta, \gamma) = \int_0^1 r^{\gamma-1}(1-r)^{\beta-1} dr. \tag{12}$$

Lemma 2.5 *Let $p \in [1, \infty]$, $\kappa \geq 0$, $\delta \in [0, 1)$, $\alpha, \gamma \in \mathbb{R}$ and $\beta \in [-\alpha, 2 - \alpha)$.*

There exists a $C > 0$ such that for all $t \in (0, 1]$,

$$\|s \mapsto P_s \phi\|_{M_t^{\frac{\kappa}{2}} \mathcal{C}_p^{\gamma+\kappa}} \leq C \|\phi\|_{\mathcal{C}_p^\gamma}, \quad \|P_t \phi - \phi\|_{\mathcal{C}_p^{\gamma-2\delta}} \leq C t^\delta \|\phi\|_{\mathcal{C}_p^\gamma}, \tag{13}$$

$$\left\| s \mapsto \int_0^s P_{s-r} w_r dr \right\|_{M_t^\delta \mathcal{C}_p^\beta} \leq C t^{1-\frac{\alpha+\beta}{2}} \|w\|_{M_t^\delta \mathcal{C}_p^{-\alpha}}. \tag{14}$$

Proof In [12, Lemma A.7] it is proven (for $p = \infty$, but can be carried on mutatis mutandis for general $p \in [1, \infty]$) that for all $\kappa \geq 0$ and $\gamma \in \mathbb{R}$ there exists a $C > 0$ such that for all $t \in (0, 1]$

$$\|P_t \phi\|_{\mathcal{C}_p^{\gamma+\kappa}} \leq C t^{-\frac{\kappa}{2}} \|\phi\|_{\mathcal{C}_p^\gamma}, \tag{15}$$

which implies the first bound in Eq. 13. The second bound in Eq. 14 follows by Eq. 15 as

$$\begin{aligned} \|P_t \phi - \phi\|_{\mathcal{C}_p^{\gamma-2\delta}} &= \left\| \int_0^t \partial_s P_s \phi ds \right\|_{\mathcal{C}_p^{\gamma-2\delta}} \leq \int_0^t \|P_s \Delta \phi\|_{\mathcal{C}_p^{\gamma-2\delta}} ds \\ &\lesssim \int_0^t s^{-\frac{2-2\delta}{2}} ds \|\Delta \phi\|_{\mathcal{C}_p^{\gamma-2}} \lesssim t^\delta \|\phi\|_{\mathcal{C}_p^\gamma}. \end{aligned}$$

The bound in Eq. 14 is also proven in [12, Lemma A.9], we give the proof to be self-contained. By applying Eq. 15 we obtain for $t \in (0, 1]$

$$\begin{aligned} \left\| \int_0^t P_{t-s} w_s ds \right\|_{\mathcal{C}_p^\beta} &\lesssim \int_0^t (t-s)^{-\frac{\alpha+\beta}{2}} s^{-\delta} ds \|w\|_{M_t^\delta \mathcal{C}_p^{-\alpha}} \\ &\lesssim t^{-\delta+1-\frac{\alpha+\beta}{2}} B\left(1 - \frac{\alpha+\beta}{2}, 1 - \delta\right) \|w\|_{M_t^\delta \mathcal{C}_p^{-\alpha}}. \end{aligned} \tag{16}$$

This proves Eq. 14. □

2.6 Let $\alpha > 0$ and let $\beta > 1 + \alpha$ and $\varepsilon > 0$ be such that $1 + \alpha + \varepsilon \leq \beta$. Then we have by Theorem A.1 together with Bernstein’s inequality ([2, Lemma 2.1 or 2.78]):

$$\|a \cdot \nabla w\|_{B_{p,\infty}^{-\alpha}} \lesssim \|a\|_{B_{\infty,1}^{-\alpha}} \|\nabla w\|_{B_{p,\infty}^{\alpha+\varepsilon}} \lesssim \|a\|_{B_{\infty,1}^{-\alpha}} \|w\|_{B_{p,\infty}^\beta}.$$

Proof of Proposition 2.4 If $\gamma \geq \beta$, then the statement follows directly from [5, Theorem 3.2]. Therefore, we assume that $\gamma < \beta$ and it is sufficient to show that the statement holds for “ t_0 ” instead of “ T ”, where t_0 will be chosen small, as we can extend the solution to $[t_0, T]$ by [5, Theorem 3.2].

As mentioned before, it is sufficient to consider the fixed point problem for Θ as in Eq. 10 instead of Φ . Let us write Θ_t^ϕ for Θ as in Eq. 10 but with “ T ” replaced by “ t ”. We will show that there exists a t_0 such that

- (a) $\Theta_{t_0}^\phi$ has a unique fixed point in $M_{t_0}^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta$,
- (b) $\Theta_{t_0}^\phi$ has a unique fixed point in $C([0, t_0], \mathcal{C}_p^{\gamma-\varepsilon})$,
- (c) $\Theta_{t_0}^\phi$ maps $C((0, t_0], \mathcal{C}_p^\beta)$ and thus $M_{t_0}^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta$ into $C([0, t_0], \mathcal{C}_p^{\gamma-\varepsilon})$, so that the fixed point in $C([0, t_0], \mathcal{C}_p^{\gamma-\varepsilon})$ agrees with the fixed point in $M_{t_0}^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta$,
- (d) for all $t > 0$ the map $\mathcal{C}_p^\gamma \times C([0, T], B_{\infty,1}^{-\alpha}) \rightarrow \mathcal{C}_p^\beta$ given by $(\phi, b) \mapsto u^{\phi,b}(t, \cdot)$ is locally Lipschitz,
- (e) the fixed point v satisfies $v(t) \in \mathcal{C}^\beta$ for $t \in (0, t_0]$ and the continuity in d can be shown for $p = \infty$, by showing that we can “increase the integrability parameter p to ∞ ”.

First, we assume that $\gamma > -\alpha$ and show (a)–(e). After that we show how one can treat $\gamma \in (\alpha - 1, -\alpha]$ too.

(a) By combining the observation in 2.6 with Lemma 2.5 with $\kappa = \beta - \gamma$ and $\delta = \frac{\beta-\gamma}{2}$ (observe that by assumption $\kappa > 0$ and $\delta \in (0, 1)$, because $0 < \beta - \gamma < 2 - \alpha + \alpha$); for $t \in (0, 1]$

$$\begin{aligned} \|\Theta_t^\phi v\|_{M_t^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta} &\lesssim \|\phi\|_{\mathcal{C}_p^\gamma} + \|s \mapsto \int_0^s P_{s-r}(b_{t-r} \cdot \nabla v_r) dr\|_{M_t^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta} \\ &\lesssim \|\phi\|_{\mathcal{C}_p^\gamma} + t^{1-\frac{\alpha+\beta}{2}} \|s \mapsto b_{t-s} \cdot \nabla v_s\|_{M_t^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^{-\alpha}} \\ &\lesssim \|\phi\|_{\mathcal{C}_p^\gamma} + t^{1-\frac{\alpha+\beta}{2}} \|b\|_{C_1 B_{\infty,1}^{-\alpha}} \|v\|_{M_t^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta}, \end{aligned} \tag{17}$$

and, moreover

$$\|\Theta_t^\phi v - \Theta_t^\phi \tilde{v}\|_{M_t^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta} \lesssim t^{1-\frac{\alpha+\beta}{2}} \|b\|_{C_1 B_{\infty,1}^{-\alpha}} \|v - \tilde{v}\|_{M_t^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta}. \tag{18}$$

That $\Theta_t^\phi v$ forms an element of $C((0, t], \mathcal{C}_p^\beta)$ follows by Lemma 2.5. Therefore, with Eq. 17 it follows that Θ_t^ϕ maps $M_t^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta$ to itself. By Eq. 18 then follows that for sufficiently small t_0 the map $\Theta_{t_0}^\phi$ is a contraction on the Banach space $M_{t_0}^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta$ and it has a unique fixed point in that space.

(b) When t_0 is as above, then $\Theta_{t_0}^\phi$ has a unique fixed point in $C([0, t_0], \mathcal{C}_p^{\gamma-\varepsilon})$ which follows from the following estimates which follow similarly as the above ones (use that $\gamma > \beta$ and Eq. 14 with $\beta = -\alpha$ and $\delta = 0$)

$$\begin{aligned} \|\Theta_t^\phi v\|_{C([0,t], \mathcal{C}_p^{\gamma-\varepsilon})} &\lesssim \|\phi\|_{\mathcal{C}_p^{\gamma-\varepsilon}} + t \|b\|_{C_1 B_{\infty,1}^{-\alpha}} \|v\|_{C([0,t], \mathcal{C}_p^{\gamma-\varepsilon})}, \\ \|\Theta_t^\phi v - \Theta_t^\phi \tilde{v}\|_{C([0,t], \mathcal{C}_p^{\gamma-\varepsilon})} &\lesssim t \|b\|_{C_1 B_{\infty,1}^{-\alpha}} \|v - \tilde{v}\|_{C([0,t], \mathcal{C}_p^{\gamma-\varepsilon})}. \end{aligned}$$

(c) That Θ_0^ϕ maps $C((0, t_0], \mathcal{C}_p^\beta)$ into $C([0, t_0], \mathcal{C}_p^{\gamma-\varepsilon})$, which means that $\Theta_0^\phi(v)(t)$ converges to ϕ in $\mathcal{C}_p^{\gamma-\varepsilon}$ as $t \downarrow 0$ for any $v \in C((0, t_0], \mathcal{C}_p^\beta)$, follows from the second bound in Eq. 13 and the following estimate (by 2.6, which follows similarly to Eq. 17)

$$\|\Theta_0^\phi(v)(t) - \phi\|_{\mathcal{C}_p^{\gamma-\varepsilon}} \leq \|P_t\phi - \phi\|_{\mathcal{C}_p^{\gamma-\varepsilon}} + t^{1-\frac{\alpha+\beta}{2}} \|b\|_{C_1 B_{\infty,1}^{-\alpha}} \|v\|_{C((0,t_0], \mathcal{C}_p^\beta)}.$$

(d) Let us write $v^{\phi,b}$ for the solution of Eq. 11 (with \mathcal{L}_t as in Eq. 6). To see the continuity of the solution with respect to b and ϕ , let $b_1, b_2 \in C([0, t_0], B_{\infty,1}^{-\alpha})$ and $\phi_1, \phi_2 \in \mathcal{C}_p^\gamma$. Let $v_i = v^{\phi_i, b_i}$ for $i \in \{1, 2\}$. By Lemma 2.5 and by 2.6 we have

$$\begin{aligned} \|v_1 - v_2\|_{M_t^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta} &\lesssim \|\phi_1 - \phi_2\|_{\mathcal{C}_p^\gamma} + t^{1-\frac{\alpha+\beta}{2}} \|b_1\|_{C_t B_{\infty,1}^{-\alpha}} \|v_1 - v_2\|_{M_t^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta} \\ &\quad + t^{1-\frac{\alpha+\beta}{2}} \|b_1 - b_2\|_{C_t B_{\infty,1}^{-\alpha}} \|v_2\|_{M_t^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta}. \end{aligned}$$

Hence there exists a $\delta \in (0, t_0)$ (small enough, e.g., $\delta^{1-\frac{\alpha+\beta}{2}} \|b_1\|_{C_\delta B_{\infty,1}^{-\alpha}} < \frac{1}{2}$) such that

$$\|v_1 - v_2\|_{M_\delta^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta} \lesssim \|\phi_1 - \phi_2\|_{\mathcal{C}_p^\gamma} + \|b_1 - b_2\|_{C_{t_0} B_{\infty,1}^{-\alpha}} \|v_2\|_{M_{t_0}^{\frac{\beta-\gamma}{2}} \mathcal{C}_p^\beta}. \tag{19}$$

So for $t \in (0, \delta]$ we obtain the desired continuity. By an iteration argument we can obtain the continuity for all $t \in (0, t_0]$, as for example for $t \in (\delta, 2\delta]$ we have $v_i(t) = v^{v_i(\delta), b}(t - \delta)$.

(e) It remains to show that we can increase the integrability from p to ∞ , i.e., that $v_t \in \mathcal{C}^\beta$ for all $t > 0$ and that also as an element of \mathcal{C}^β the solution v_t for fixed $t > 0$ depends continuously on b and ϕ . First we show that if $t > 0$, then $v_s \in \mathcal{C}^\beta$ for all $s > t$. To simplify notation we only consider the most extreme case $p = 1$, but the argument for general p is essentially the same. Let $n \in \mathbb{N}_0$ be such that

$$n(\beta - \gamma) < d, \quad (n + 1)(\beta - \gamma) \geq d.$$

Write $p_0 = 1$ and for $i \in \{1, \dots, n\}$

$$p_i = \frac{d}{d - i(\beta - \gamma)} \in (1, \infty).$$

Then $\beta - \frac{d}{p_n} \geq \gamma$ and $\beta - d(\frac{1}{p_{i-1}} - \frac{1}{p_i}) = \gamma$ for all $i \in \{1, \dots, n - 1\}$, hence the Besov embedding theorem [2, Proposition 2.71] gives $\mathcal{C}_{p_{i-1}}^\beta \subset \mathcal{C}_{p_i}^\gamma$ for all $i \in \{1, \dots, n - 1\}$, and $\mathcal{C}_{p_n}^\beta \subset \mathcal{C}^\gamma$. We have $v_{\frac{t}{n}} \in \mathcal{C}_1^\beta \subset \mathcal{C}_{p_1}^\gamma$. By considering the Eq. 11 with initial condition $v_{\frac{t}{n}}$ we obtain that v_s is in $\mathcal{C}_{p_1}^\beta$ for $s > \frac{t}{n}$, in particular $v_{\frac{2t}{n}} \in \mathcal{C}_{p_2}^\gamma$. Repeating the argument we obtain $v_{\frac{i}{n}t} \in \mathcal{C}_{p_i}^\gamma$ for all $i \in \{1, \dots, n - 1\}$ and $v_t \in \mathcal{C}^\beta$, so indeed $v_s \in \mathcal{C}^\beta$ for all $s > t$. As t was arbitrary, we have shown that $v_t \in \mathcal{C}^\beta$ for all $t > 0$. As all the inclusions \subset above are given by continuous embeddings, the continuity of the solution with respect to ϕ and b follows from the continuity shown in (d).

We are left to show that we can also treat $\gamma \in (\alpha - 1, -\alpha]$. Let γ be as such. We choose $\tilde{\beta} \in (1 + \alpha, 2 - \alpha)$ such that $\tilde{\beta} - \gamma < 2$. Then we have $\tilde{\beta} > \gamma$ and $\frac{\tilde{\beta}-\gamma}{2} \in (0, 1)$, so that the conditions of observation 2.6 and Lemma 2.5 are satisfied. Hence we obtain also Eq. 17 and Eq. 18 with “ $\tilde{\beta}$ ” instead of “ β ”. So then we find a $\tilde{t}_0 \in (0, t_0)$ such that Θ_0^ϕ has a fixed point \tilde{v} in $M_{\tilde{t}_0}^{\frac{\tilde{\beta}-\gamma}{2}} \mathcal{C}_p^\beta$. Let w be the fixed point of $\Phi_{t_0-\tilde{t}_0}^{\tilde{v}(\tilde{t}_0)}$ in $M_{t_0-\tilde{t}_0}^{\frac{\beta-\tilde{\beta}}{2}}$, which exists by (a) because $\tilde{\beta} > -\alpha$. Then $v(t) := \tilde{v}(t)$ for $t \in (0, \tilde{t}_0]$ and $v(t) := w(t - t_0)$ for $t \in (\tilde{t}_0, t_0]$ is a fixed

point of Φ_0^ϕ such that $(\tilde{t}_0, t_0] \mapsto \mathcal{C}_p^\beta, t \mapsto v(t)$ is continuous. As \tilde{t}_0 can be taken arbitrarily small, we conclude that $v \in C((0, t_0], \mathcal{C}_p^\beta)$. Similarly, we can obtain the continuity of the solution by using Eq. 19 with “ (β, γ) ” replaced by “ $(\beta, \tilde{\beta})$ ” and using Eq. 19 with “ (β, γ) ” replaced by “ $(\tilde{\beta}, \gamma)$ ”. \square

2.7 A direct computation using that $\Delta_i \delta_z(x) = \mathcal{F}^{-1}(\rho(2^{-i} \cdot))(x-z) = 2^{id} \mathcal{F}^{-1}(\rho)(2^i(x-z))$ for $i \geq 0$ shows that the Dirac delta δ_z is in $\mathcal{C}_p^{-d(1-\frac{1}{p})}$ for all $p \in [1, \infty]$, so in particular $\delta_z \in \mathcal{C}_1^0$. Moreover, $\mathcal{F}^{-1} \rho_i$ is a Schwartz function for fixed $i \geq -1$, and therefore $\mathbb{R}^d \ni z \mapsto \Delta_i \delta_z \in L^p$ is continuous. This easily yields that for $\epsilon > 0$ the map $\mathbb{R}^d \ni z \mapsto \delta_z \in \mathcal{C}_1^{-\epsilon}$ is continuous.

Corollary 2.8 (of Proposition 2.4) *Let $\alpha \in (0, \frac{1}{2})$ and $b \in C([0, T], B_{\infty,1}^{-\alpha}(\mathbb{R}^d, \mathbb{R}^d))$.*

For $t \in [0, T]$ and $n \in \mathbb{N}$ let $b_t^{(n)} = \sum_{i=1}^n \Delta_i b_t \in C_b^\infty(\mathbb{R}^d, \mathbb{R}^d)$ and let $\Gamma_{t,T}(x, y) = u^{\delta_y, b}(t, x)$ and $\Gamma_{t,T}^{(n)}(x, y) = u^{\delta_y, b^{(n)}}(t, x)$ (notation as in Proposition 2.4). Then $\Gamma_{t,T}$ and $\Gamma_{t,T}^{(n)}$ are continuous on $\mathbb{R}^d \times \mathbb{R}^d$ and we have for all $t \in [0, T]$ and $\mu \in \mathbb{N}_0^d$ with $|\mu| \leq 1$:

$$\sup_{x,y \in \mathbb{R}^d} |\partial_x^\mu [\Gamma_{t,T}(x, y) - \Gamma_{t,T}^{(n)}(x, y)]| \xrightarrow{n \rightarrow \infty} 0.$$

Proof The continuity follows from Proposition 2.4.

Because there exists a $C > 0$ such that $\|b_s^{(n)} - b_r^{(n)}\|_{B_{\infty,1}^{-\alpha}} \leq C \|b_s - b_r\|_{B_{\infty,1}^{-\alpha}}$ for all $n \in \mathbb{N}$, $s, r \in [0, \infty)$ and $\|b_s^{(n)} - b_s\|_{B_{\infty,1}^{-\alpha}} \rightarrow 0$ for all $s \in [0, \infty)$ we obtain by a “ 3ϵ argument” that

$$\|b^{(n)} - b\|_{C_t B_{\infty,1}^{-\alpha}} \rightarrow 0$$

As moreover $\sup_{y \in \mathbb{R}^d} \|\delta_y\|_{B_{1,\infty}^0} \lesssim 1$, Proposition 2.4 yields

$$\sup_{y \in \mathbb{R}^d} \|\Gamma_{t,T}(\cdot, y) - \Gamma_{t,T}^{(n)}(\cdot, y)\|_{\mathcal{C}^\beta} \rightarrow 0,$$

for all $\beta < 2 - \alpha$. \square

Proposition 2.9 *Let $\alpha \in (0, \frac{1}{2})$ and $b \in C([0, T], B_{\infty,1}^{-\alpha}(\mathbb{R}^d, \mathbb{R}^d))$. For $t \in [0, T]$ let $\Gamma_{t,T}: \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ be defined by $\Gamma_{t,T}(x, y) = u^{\delta_y}(t, x)$. Let $\mathbb{P}_{t,x}$ be the unique probability measure on $C([t, T], \mathbb{R}^d)$ such that the coordinate process X is a solution to the SDE Eq. 5 on $[t, T]$ with initial condition $X_t = x$. Then $\Gamma_{t,T}(x, \cdot)$ is the density of X_T under $\mathbb{P}_{t,x}$, i.e., $\mathbb{E}_{t,x}[\phi(X_T)] = \int_{\mathbb{R}^d} \phi(y) \Gamma_{t,T}(x, y) dy$ for all $\phi \in C_c(\mathbb{R}^d)$.*

Proof For b with values in C_b^∞ this is classical, see for example [10, Theorem 6.5.4]. So let $b^{(n)}$ and $\Gamma_{t,T}^{(n)}$ be as in Corollary 2.8 and for $x \in \mathbb{R}^d$ let $\mathbb{P}_{t,x}^{(n)}$ be the unique probability measure on $C([t, T], \mathbb{R}^d)$ such that the coordinate process X is a solution to the martingale problem for $((\mathcal{L}_s^{(n)})_{s \in (t, T]}, \delta_x)$, where $\mathcal{L}_s^{(n)} = \frac{1}{2} \Delta + b_{T-s}^{(n)} \cdot \nabla$. Using that $\mathbb{P}_{t,x}^{(n)}$ weakly converges to $\mathbb{P}_{t,x}$ (Theorem 2.2) and the uniform convergence in Corollary 2.8 we obtain for $\phi \in C_c(\mathbb{R}^d)$:

$$\mathbb{E}_{t,x}[\phi(X_T)] = \lim_{n \rightarrow \infty} \mathbb{E}_{t,x}^{(n)}[\phi(X_T)] = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} \phi(y) \Gamma_{t,T}^{(n)}(x, y) dy = \int_{\mathbb{R}^d} \phi(y) \Gamma_{t,T}(x, y) dy.$$

\square

3 Heat-Kernel Upper Bounds

Here we prove the upper bound (2) of the heat-kernel estimates. We follow the ‘‘parametrix’’ approach from Friedman’s book [9] to prove the heat-kernel estimates presented in Theorem 1.1. This means that we write Γ_t as a series (see Lemma 3.3) and bound each term in that series to obtain a bound for the whole series and thus for Γ_t . Usually the point of the parametrix is to deal with non-constant diffusion coefficients, but the approach is still useful for us despite the fact that we deal with constant diffusion coefficients.

Because of Corollary 2.8 we can restrict our attention to b in $C([0, T], C_b^\infty(\mathbb{R}^d, \mathbb{R}^d))$ and then extend the bounds to b in $C([0, T], B_{\infty,1}^{-\alpha}(\mathbb{R}^d, \mathbb{R}^d))$ by a limiting argument.

For the rest of this section we fix $\alpha \in (0, \frac{1}{2})$, and $c > 1$ as in Theorem 1.1 and $b \in C([0, \infty), C_b^\infty(\mathbb{R}^d, \mathbb{R}^d))$. (Instead of $[0, T]$ we consider $[0, \infty)$ for notational convenience.)

3.1 Let $g \in L^1(\mathbb{R}^d, \mathbb{R}^d)$ and $a \in C_b^\infty(\mathbb{R}^d, \mathbb{R}^d)$. Let $(\tilde{\rho}_i)_{i \in \mathbb{N}_{>1}}$ be another dyadic partition of unity, but such that $\text{supp } \tilde{\rho}_{-1} \cap \text{supp } \rho_i = \emptyset$ for $i \in \mathbb{N}_0$ so that

$$\begin{aligned} \int_{\mathbb{R}^d} (\Delta_i a)(z)(\tilde{\Delta}_{-1} g)(z) \, dz &= \int_{\mathbb{R}^d} \mathcal{F}^{-1}(\rho_i \hat{a})(z) \mathcal{F}^{-1}(\tilde{\rho}_{-1} \hat{g})(z) \, dz \\ &= \int_{\mathbb{R}^d} \hat{a}(-z) \rho_i(z) \tilde{\rho}_{-1}(z) \hat{g}(z) \, dz = 0, \end{aligned}$$

and thus

$$\int_{\mathbb{R}^d} (\Delta_{\geq 0} a)(z) g(z) \, dz = \int_{\mathbb{R}^d} (\Delta_{\geq 0} a)(z)(\tilde{\Delta}_{\geq 0} g)(z) \, dz.$$

By duality and Bernstein’s inequality, see [2, Proposition 2.76 and Lemma 2.1], we have

$$\begin{aligned} \left| \int_{\mathbb{R}^d} a(z) \cdot g(z) \, dz \right| &\leq \left| \int_{\mathbb{R}^d} \Delta_{-1} a(z) \cdot g(z) \, dz \right| + \left| \int_{\mathbb{R}^d} \Delta_{\geq 0} a(z) \cdot g(z) \, dz \right| \\ &\lesssim \|\Delta_{-1} a\|_{L^\infty} \|g\|_{L^1} + \|\Delta_{\geq 0} a\|_{B_{\infty,1}^{-\alpha}} \|\tilde{\Delta}_{\geq 0} g\|_{B_{1,\infty}^\alpha} \\ &\lesssim \|\Delta_{-1} a\|_{L^\infty} \|g\|_{L^1} + \|\Delta_{\geq 0} a\|_{B_{\infty,1}^{-\alpha}} \left(\sup_{j \geq 0} \left\{ \|\tilde{\Delta}_j g\|_{L^1}^{1-\alpha} (2^j \|\tilde{\Delta}_j g\|_{L^1})^\alpha \right\} \right) \\ &\lesssim \|\Delta_{-1} a\|_{L^\infty} \|g\|_{L^1} + \|\Delta_{\geq 0} a\|_{B_{\infty,1}^{-\alpha}} \|g\|_{L^1}^{1-\alpha} \|\nabla g\|_{L^1}^\alpha. \end{aligned} \tag{20}$$

We will apply the above bound for functions g that are Gaussian, therefore we will need estimates for derivatives of Gaussian functions. So we recall the following bound:

3.2 Let $p(t, x) = (2\pi t)^{-\frac{d}{2}} e^{-\frac{1}{2t}|x|^2}$ for $(t, x) \in (0, \infty) \times \mathbb{R}^d$ be the standard Gaussian kernel. For the space derivatives $\partial^\mu p$ we have the following estimate:

$$\forall \mu \in \mathbb{N}_0^d \exists C > 0 \forall (t, x) \in (0, \infty) \times \mathbb{R}^d : |\partial^\mu p(t, x)| \leq C t^{-\frac{|\mu|}{2}} p(ct, x), \tag{21}$$

The proof of the upper bound (2) essentially follows by iterating the previous two observations. To carry out the argument we need the following result, which allows us to write Γ as an infinite series.

Lemma 3.3 Let $t > 0$ and $y \in \mathbb{R}^d$. For $s \in [0, t)$ and $x \in \mathbb{R}^d$ we define

$$\Psi_{s,t}^{y,1}(x) = -b(t - s, x) \cdot \nabla p(s, x - y). \tag{22}$$

Then for all $k \in \mathbb{N}$ the map $s \mapsto \Psi_{s,t}^{y,k}$ is in $L^1([0, t], L^1(\mathbb{R}^d))$, where

$$\Psi_{s,t}^{y,k+1}(x) = -\int_0^s \int_{\mathbb{R}^d} b(t-s, x) \cdot \nabla p(s-r, x-z) \Psi_{r,t}^{y,k}(z) \, dz \, dr. \tag{23}$$

Moreover, (with $\Gamma_{s,t}$ as in Proposition 2.9)

$$\Gamma_{s,t}(x, y) = p(t-s, x-y) + \sum_{k=1}^{\infty} \int_0^{t-s} \int_{\mathbb{R}^d} p(t-s-r, x-z) \Psi_{r,t}^{y,k}(z) \, dz \, dr. \tag{24}$$

Proof By Eq. 21 we know that $\|\Psi_{s,t}^{y,1}\|_{L^1(\mathbb{R}^d)} \lesssim \|\nabla p(s, \cdot)\|_{L^1(\mathbb{R}^d)} \lesssim s^{-\frac{1}{2}}$ and therefore $s \mapsto \Psi_{s,t}^{y,1}$ is in $L^1([0, t], L^1(\mathbb{R}^d))$. Observe that $\Psi_{s,t}^{y,k+1}$ equals the inner product of $-b(t-s, x)$ with a convolution in space and time. Therefore, by applying the L^1 inequality for convolutions (Young’s inequality) for the space as well for the time convolution, we obtain

$$\begin{aligned} \|\Psi_{s,t}^{y,k+1}\|_{L^1(\mathbb{R}^d)} &\lesssim \int_0^s \|\nabla p(s-r, \cdot) * \Psi_{r,t}^{y,k}\|_{L^1(\mathbb{R}^d)} \, dr \\ &\lesssim \int_0^s (s-r)^{-\frac{1}{2}} \, dr \int_0^s \|\Psi_{r,t}^{y,k}\|_{L^1(\mathbb{R}^d)} \, dr \lesssim s^{\frac{1}{2}} \int_0^t \|\Psi_{r,t}^{y,k}\|_{L^1(\mathbb{R}^d)} \, dr, \end{aligned}$$

from which we conclude that $\int_0^t \|\Psi_{r,t}^{y,k}\|_{L^1(\mathbb{R}^d)} \, dr$ is finite (actually it is $\lesssim t^{\frac{1+3k}{2}}$) for all $k \in \mathbb{N}$.

It remains to show Eq. 24. As $\Gamma_{s,t}(x, y) = u^{\delta_y}(s, x)$ where u^{δ_y} being the fixed point of the map Φ as in Eq. 9 with $\phi = \delta_y$, that is, with $u = u^{\delta_y}$,

$$\begin{aligned} (\Phi u)_s &= P_{t-s} \delta_y - \int_s^t P_{q-s} (b_q \cdot \nabla u_q) \, dq \\ &= P_{t-s} \delta_y - \int_0^{t-s} P_{t-s-r} (b_{t-r} \cdot \nabla u_{t-r}) \, dr. \end{aligned}$$

From a Picard iteration it follows that Γ is the limit of the sequence $\Gamma_t^0 = 0$,

$$\begin{aligned} \Gamma_{s,t}^{k+1}(x, y) &= p(t-s, x-y) - \int_0^{t-s} \int_{\mathbb{R}^d} p(t-s-r, x-z) (b(t-r, z) \cdot \nabla_z \Gamma_{t-r,t}^k(z, y)) \, dz \, dr. \end{aligned}$$

Therefore, $\Gamma_{s,t}^1(x, y) = p(t-s, x-y)$ and we obtain recursively (see also [9, Chapter 1.4])

$$\Gamma_{s,t}^{k+1}(x, y) = p(t-s, x-y) + \sum_{\ell=1}^k \int_0^{t-s} \int_{\mathbb{R}^d} p(t-s-r, x-z) \Psi_{r,t}^{y,\ell}(z) \, dz \, dr.$$

This proves (24). □

3.4 Now let us get back to Remark 1.4. Observe that in the right-hand side in Eq. 24 the dependence on t is in the $\Psi^{y,k}$ functions, and we see that the rest is a function of $t-s$. This allows us to take the first time variable, s , equal to zero, and proof the heat-kernel bounds as in Theorem 1.1. From now on we write “ Γ_t ” for “ $\Gamma_{0,t}$ ”.

Note that the first term appearing in the right-hand side of Eq. 24 is already bounded by the right-hand side of Eq. 2. Therefore, we will recursively estimate

$$\int_0^t \int_{\mathbb{R}^d} p(t-s, x-z) \Psi_{s,t}^{y,k}(z) \, dz \, ds.$$

This will be done with the help of some auxiliary lemmas, which follow below.

3.5 Let $\mu \in \mathbb{N}_0^d$, $t > 0$, $k \in \mathbb{N}$, $x, y \in \mathbb{R}^d$ and $g \in L^1(\mathbb{R}^d)$. As we write $P_t g = p(t, \cdot) * g$ (see Eq. 9), we have $\partial^\mu P_t g = \partial^\mu p(t, \cdot) * g$.

For any given norm $\| \cdot \|$ we will write $\|\nabla f\| = \sum_{i=1}^d \|\partial_i f\|$ and $\|\nabla^2 f\| = \sum_{i,j=1}^d \|\partial_i \partial_j f\|$.

Lemma 3.6 *There exists a $C > 0$ (independent of b) such that for all $\mu \in \mathbb{N}_0^d$ with $|\mu| \leq 2$, $y \in \mathbb{R}^d$ and $t, s, r \in (0, \infty)$ with $t > s > r$ and all $f \in L^1(\mathbb{R}^d)$, with $g_{t,s,r}(z) = b(t-s, z) \cdot \int_{\mathbb{R}^d} \nabla p(s-r, z-w) f(w) dw$*

$$\begin{aligned}
 |\partial^\mu P_{t-s} g_{t,s,r}(x)| &\leq C(t-s)^{-\frac{|\mu|}{2}} p(ct, x-y) \left(\|\Delta_{-1} b_{t-s}\|_{L^\infty} \left\| \frac{\nabla P_{s-r} f}{p(cs, \cdot - y)} \right\|_{L^\infty} \right. \\
 &\quad \left. + \|\Delta_{\geq 0} b_{t-s}\|_{B_{\infty,1}^{-\alpha}} \left[(t-s)^{-\frac{\alpha}{2}} \left\| \frac{\nabla P_{s-r} f}{p(cs, \cdot - y)} \right\|_{L^\infty} + \left\| \frac{\nabla P_{s-r} f}{p(cs, \cdot - y)} \right\|_{L^\infty}^{1-\alpha} \left\| \frac{\nabla^2 P_{s-r} f}{p(cs, \cdot - y)} \right\|_{L^\infty}^\alpha \right] \right). \tag{25}
 \end{aligned}$$

Proof We abbreviate $g_{t,s,r}$ by g . Observe that $g(z) = b(t-s, z) \cdot \nabla P_{s-r} f(z)$. Then, with $h : \mathbb{R}^d \rightarrow \mathbb{R}^d$, $h(z) = \partial^\mu p(t-s, x-z) \nabla P_{s-r} f(z)$, by Eq. 20

$$\begin{aligned}
 |\partial^\mu P_{t-s} g(x)| &= \left| \int_{\mathbb{R}^d} \partial^\mu p(t-s, x-z) b(t-s, z) \cdot \nabla P_{s-r} f(z) dz \right| \\
 &\lesssim \|\Delta_{-1} b_{t-s}\|_{L^\infty} \|h\|_{L^1} + \|\Delta_{\geq 0} b_{t-s}\|_{B_{\infty,1}^{-\alpha}} \|h\|_{L^1}^{1-\alpha} \|\nabla h\|_{L^1}^\alpha.
 \end{aligned}$$

We estimate both $\|h\|_{L^1}$ and $\|\nabla h\|_{L^1}$. We use Eq. 21 and $\int_{\mathbb{R}^d} p(c(t-s), x-z) p(cs, z-y) dz = p(c(t-s), \cdot) * p(cs, \cdot)(x-y) = p(ct, x-y)$ to obtain

$$\begin{aligned}
 \|h\|_{L^1} &= \int_{\mathbb{R}^d} |\partial^\mu p(t-s, x-z) \nabla P_{s-r} f(z)| dz \\
 &\lesssim \int_{\mathbb{R}^d} (t-s)^{-\frac{|\mu|}{2}} p(c(t-s), x-z) p(cs, z-y) \left\| \frac{\nabla P_{s-r} f}{p(cs, \cdot - y)} \right\|_{L^\infty} dz \\
 &= (t-s)^{-\frac{|\mu|}{2}} p(ct, x-y) \left\| \frac{\nabla P_{s-r} f}{p(cs, \cdot - y)} \right\|_{L^\infty}.
 \end{aligned}$$

Similarly, in combination with Leibniz’s rule, we obtain

$$\begin{aligned}
 \|\nabla h\|_{L^1} &= \|\nabla(\partial^\mu p(t-s, x-\cdot) \nabla P_{s-r} f)\|_{L^1} \\
 &\leq \sum_{i=1}^d \|\partial^\mu \partial_i p(t-s, x-\cdot) \nabla P_{s-r} f\|_{L^1} + \|\partial^\mu p(t-s, x-\cdot) \nabla^2 P_{s-r} f\|_{L^1} \\
 &\lesssim (t-s)^{-\frac{|\mu|}{2}} p(ct, x-y) \left[(t-s)^{-\frac{1}{2}} \left\| \frac{\nabla P_{s-r} f}{p(cs, \cdot - y)} \right\|_{L^\infty} + \left\| \frac{\nabla^2 P_{s-r} f}{p(cs, \cdot - y)} \right\|_{L^\infty} \right].
 \end{aligned}$$

Using the above and that $(a+b)^\alpha \leq a^\alpha + b^\alpha$ for $a, b \geq 0$ we obtain Eq. 25. □

3.7 Now we apply the above lemma to our setting. But first, let us introduce some notation. For $k \in \mathbb{N}$, $t \geq 0$, $i \in \{0, 1\}$, and $\beta \in \{0, \alpha\}$ we write

$$\mathcal{J}_{i,k}^\beta(t) = \sup_{y \in \mathbb{R}^d} \int_0^t \left\| \frac{\nabla^i P_{t-s} [\Psi_{s,t}^{y,k}]}{p(ct, \cdot - y)} \right\|_{L^\infty}^{1-\beta} \left\| \frac{\nabla^{i+1} P_{t-s} [\Psi_{s,t}^{y,k}]}{p(ct, \cdot - y)} \right\|_{L^\infty}^\beta ds.$$

We are interested in the bounds for $\mathcal{J}_{i,k}^0$ only. But in order to describe a recursive relation for them, as we will see in the next lemma, we also need the $\mathcal{J}_{i,k}^\alpha$'s.

Lemma 3.8 *Let $C > 0$ be as in Lemma 3.6. For all $k \in \mathbb{N}$, $t \geq 0$, $i \in \{0, 1\}$ and $\beta \in \{0, \alpha\}$*

$$\mathcal{J}_{i,k+1}^\beta(t) \leq C \int_0^t (t-s)^{-\frac{i+\beta}{2}} \left(\|\Delta_{-1}b\|_{C_t L^\infty} \mathcal{J}_{1,k}^0(s) + \|\Delta_{\geq 0}b\|_{C_t B_{\infty,1}^{-\alpha}} [(t-s)^{-\frac{\alpha}{2}} \mathcal{J}_{1,k}^0(s) + \mathcal{J}_{1,k}^\alpha(s)] \right) ds. \tag{26}$$

Proof We claim that the following holds. For all $k \in \mathbb{N}$, $y \in \mathbb{R}^d$ and $i \in \{0, 1, 2\}$

$$\begin{aligned} \left\| \frac{\nabla^i P_{t-s}[\Psi_{s,t}^{y,k+1}]}{p(ct, \cdot - y)} \right\|_{L^\infty} &\leq C(t-s)^{-\frac{i}{2}} \left(\|\Delta_{-1}b\|_{C_t L^\infty} \int_0^s \left\| \frac{\nabla P_{s-r}[\Psi_{r,t}^{y,k}]}{p(cs, \cdot - y)} \right\|_{L^\infty} dr \right. \\ &\quad + \|\Delta_{\geq 0}b\|_{C_t B_{\infty,1}^{-\alpha}} \left[(t-s)^{-\frac{\alpha}{2}} \int_0^s \left\| \frac{\nabla P_{s-r}[\Psi_{r,t}^{y,k}]}{p(cs, \cdot - y)} \right\|_{L^\infty} dr \right. \\ &\quad \left. \left. + \int_0^s \left\| \frac{\nabla P_{s-r}[\Psi_{r,t}^{y,k}]}{p(cs, \cdot - y)} \right\|_{L^\infty}^{1-\alpha} \left\| \frac{\nabla^2 P_{s-r}[\Psi_{r,t}^{y,k}]}{p(cs, \cdot - y)} \right\|_{L^\infty}^\alpha dr \right] \right). \tag{27} \end{aligned}$$

From this Eq. 26 follows by definition of \mathcal{J}_k^β . Now let us prove Eq. 27. Let $g_{t,s,r}$ be as in Lemma 3.6 with $f = \Psi_{r,t}^{y,k}$. Observe that by definition of $\Psi_{s,t}^{y,k+1}$ Eq. 23 we can write

$$\Psi_{s,t}^{y,k+1}(z) = \int_0^s b(t-s, z) \cdot \nabla P_{s-r}[\Psi_{r,t}^{y,k}](z) dr = \int_0^s g_{t,s,r}(z) dr,$$

so that (one can verify the interchange of integrals by Fubini's theorem and using Lemma 3.3)

$$|\nabla^i P_{t-s}[\Psi_{s,t}^{y,k+1}](x)| \leq \int_0^s |\nabla^i P_{t-s} g_{t,s,r}(x)| dr.$$

With this, Eq. 27 follows from Eq. 25. □

In the proof of Lemma 3.10 we will use the following bound for the beta function (see Eq. 12).

Lemma 3.9 *Let $\delta \in (0, 1]$. Then $M_\delta := \sup\{B(\beta, \gamma)\gamma^\beta : (\beta, \gamma) \in [\delta, 1] \times [\delta, \infty)\} < \infty$. Hence, for all $(\beta, \gamma) \in [\delta, 1] \times [\delta, \infty)$,*

$$B(\beta, \gamma) = B(\gamma, \beta) \leq M_\delta \gamma^{-\beta}.$$

Proof By [1, Theorem 1.1.4 and Theorem 1.4.1] we have for $\gamma, \beta > 0$

$$B(\beta, \gamma) = \frac{\Gamma(\gamma)\Gamma(\beta)}{\Gamma(\gamma + \beta)}, \quad \text{and} \quad \lim_{\gamma \rightarrow \infty} \frac{\Gamma(\gamma)}{\sqrt{2\pi}\gamma^{\gamma-\frac{1}{2}}e^{-\gamma}} = 1.$$

From this we deduce the following. Let $\beta_n \rightarrow \beta$ for some $\beta \in [\delta, 1]$ and $\gamma_n \rightarrow \infty$. Then

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{B(\beta_n, \gamma_n) \gamma_n^{\beta_n}}{\Gamma(\beta_n)} &= \lim_{n \rightarrow \infty} \frac{\sqrt{2\pi} \gamma_n^{\gamma_n - \frac{1}{2}} e^{-\gamma_n} \gamma_n^{\beta_n}}{\sqrt{2\pi} (\gamma_n + \beta_n)^{\gamma_n + \beta_n - \frac{1}{2}} e^{-(\gamma_n + \beta_n)}} \\ &= \lim_{n \rightarrow \infty} \left(1 + \frac{\beta_n}{\gamma_n}\right)^{-(\gamma_n + \beta_n - \frac{1}{2})} e^{\beta_n} \\ &= \lim_{\gamma \rightarrow \infty} \left(1 + \frac{\beta_n}{\gamma_n}\right)^{-\gamma_n} e^{\beta_n} = e^{-\beta_n} e^{\beta_n} = 1. \end{aligned}$$

Therefore

$$\lim_{n \rightarrow \infty} B(\beta_n, \gamma_n) \gamma_n^{\beta_n} = \Gamma(\beta),$$

so that from the continuity of Γ it follows that $(\beta, \gamma) \mapsto B(\beta, \gamma) \gamma^\beta$ is a bounded function on $[\delta, 1] \times [\delta, \infty)$. □

Let us now use the recursive relation for $\mathcal{I}_{i,k}^\beta$ and the bounds on the beta function to obtain estimates for $\mathcal{I}_{i,k}^\beta$:

Lemma 3.10 *Let $C > 0$ be as in Lemma 3.6 and let $M = 8M_{\frac{1}{2}-\alpha}$ with M_δ as in Lemma 3.9. There exists a $K > 0$ (independent of b) such that for all $k \in \mathbb{N}$, $t > 0$, $\beta \in \{0, \alpha\}$ and $i \in \{0, 1\}$*

$$\mathcal{I}_{i,k}^\beta(t) \leq K \sum_{\substack{m, n \in \mathbb{N}_0 \\ m+n=k}} t^{-\frac{i+\beta}{2}} \frac{(CM \|\Delta_{-1}b\|_{C_t L^\infty} t^{\frac{1}{2}})^m}{(m!)^{\frac{1-\beta}{2}}} \frac{(CM \|\Delta_{\geq 0}b\|_{C_t B_{\infty,1}^{-\alpha}} t^{\frac{1-\alpha}{2}})^n}{(n!)^{\frac{1-\alpha-\beta}{2}}}. \tag{28}$$

Proof We give a proof by induction. Instead of “ $\|\Delta_{-1}b\|_{C_t L^\infty}$ ” and “ $\|\Delta_{\geq 0}b\|_{C_t B_{\infty,1}^{-\alpha}}$ ” we will write “ X ” and “ Y ”, respectively.

- The induction start, $k = 1$:

We have for $\mu \in \mathbb{N}_0^d$ with $|\mu| \leq 2$

$$\partial^\mu P_{t-s}[\Psi_{s,t}^{y,1}](x) = \int_{\mathbb{R}^d} \partial^\mu p(t-s, x-z) \Psi_{s,t}^{y,1}(z) dz = \int_{\mathbb{R}^d} b(z) \cdot g_\mu(z) dz$$

with $g_\mu(z) = \nabla p(s, z-y) \partial^\mu p(t-s, x-z)$. By Eq. 21 there exists a $K > 0$ such that for all $\mu, v \in \mathbb{N}_0^d$ with $|\mu| \leq 2$ and $|v| \leq 1$:

$$\begin{aligned} |g_\mu(z)| &\leq K(t-s)^{-\frac{|\mu|}{2}} s^{-\frac{1}{2}} p(cs, z-y) p(c(t-s), x-z), \\ |\partial^v g_\mu(z)| &\leq K(t-s)^{-\frac{|\mu|}{2}} s^{-\frac{1}{2}} [(t-s)^{-\frac{1}{2}} + s^{-\frac{1}{2}}] p(cs, z-y) p(c(t-s), x-z). \end{aligned}$$

Therefore, by Eq. 20, for $j \in \{0, 1, 2\}$

$$\left\| \frac{\nabla^j P_{t-s}[\Psi_{s,t}^{y,1}]}{p(ct, \cdot - y)} \right\|_{L^\infty} \leq K(t-s)^{-\frac{j}{2}} s^{-\frac{1}{2}} \left(X + Y[(t-s)^{-\frac{\alpha}{2}} + s^{-\frac{\alpha}{2}}] \right),$$

so that for $i \in \{0, 1\}$

$$\begin{aligned} \mathcal{I}_{i,1}^\beta(t) &\leq K \int_0^t (t-s)^{-\frac{i+\beta}{2}} s^{-\frac{1}{2}} \left(X + Y[(t-s)^{-\frac{\alpha}{2}} + s^{-\frac{\alpha}{2}}] \right) ds \\ &\leq t^{-\frac{i+\beta}{2}} K \left(B\left(\frac{2-i-\beta}{2}, \frac{1}{2}\right) X t^{\frac{1}{2}} + \left[B\left(\frac{2-i-\alpha-\beta}{2}, \frac{1}{2}\right) + B\left(\frac{2-i-\beta}{2}, \frac{1-\alpha}{2}\right) \right] Y t^{\frac{1-\alpha}{2}} \right). \end{aligned}$$

Hence, for $k = 1$, the inequality (28) follows by applying Lemma 3.9 for the beta functions and using that $\delta \mapsto M_\delta$ is decreasing:

$$\begin{aligned} B\left(\frac{2-i-\beta}{2}, \frac{1}{2}\right) &\leq M_{\frac{2-i-\beta}{2}} \left(\frac{1}{2}\right)^{-\frac{2-i-\beta}{2}} \leq 2M_{\frac{1}{2}-\alpha} \leq M, \\ B\left(\frac{2-i-\alpha-\beta}{2}, \frac{1}{2}\right) &\leq M_{\frac{2-i-\alpha-\beta}{2}} 2^{\frac{1-\alpha-\beta}{2}} \leq M, \\ B\left(\frac{2-i-\beta}{2}, \frac{1-\alpha}{2}\right) &\leq M_{\frac{2-i-\beta}{2}} \left(\frac{1-\alpha}{2}\right)^{-\frac{1-\beta}{2}} \leq M_{\frac{1}{2}-\alpha} 4^{\frac{1-\beta}{2}} \leq M. \end{aligned}$$

- The induction step, from k to $k + 1$:

Let $k \in \mathbb{N}$ and assume that Eq. 28 holds. Then by Lemma 3.8

$$\begin{aligned} \mathcal{I}_{i,k+1}^\beta(t) &\leq C \int_0^t (t-s)^{-\frac{i+\beta}{2}} \left(X \mathcal{I}_{1,k}^0(s) + Y[(t-s)^{-\frac{\alpha}{2}} \mathcal{I}_{1,k}^0(s) + \mathcal{I}_{1,k}^\alpha(s)] \right) ds \\ &\leq KC \sum_{\substack{m,n \in \mathbb{N}_0: \\ m+n=k}} \frac{(CMX)^m}{(m!)^{\frac{1-\beta}{2}}} \frac{(CMY)^n}{(n!)^{\frac{1-\alpha-\beta}{2}}} \\ &\quad \times \int_0^t (t-s)^{-\frac{i+\beta}{2}} s^{-\frac{1}{2} + \frac{m}{2} + n\frac{1-\alpha}{2}} \left(X + Y[(t-s)^{-\frac{\alpha}{2}} + s^{-\frac{\alpha}{2}}] \right) ds. \end{aligned}$$

We bound the latter integral, for which we have the following identity:

$$\begin{aligned} &\int_0^t (t-s)^{-\frac{i+\beta}{2}} s^{-\frac{1}{2} + \frac{m}{2} + n\frac{1-\alpha}{2}} \left(X + Y[(t-s)^{-\frac{\alpha}{2}} + s^{-\frac{\alpha}{2}}] \right) ds \\ &= t^{-\frac{i+\beta}{2}} t^{\frac{m}{2} + n\frac{1-\alpha}{2}} \left(X t^{\frac{1}{2}} B\left(\frac{1-\beta}{2}, \frac{m+1+n(1-\alpha)}{2}\right) \right. \\ &\quad \left. + Y t^{\frac{1-\alpha}{2}} \left[B\left(\frac{1-\alpha-\beta}{2}, \frac{m+1+n(1-\alpha)}{2}\right) + B\left(\frac{1-\beta}{2}, \frac{m+(n+1)(1-\alpha)}{2}\right) \right] \right). \end{aligned}$$

This shows that the power of t is the right one. We bound the beta function terms to finish the proof. By Lemma 3.9 we have

$$\begin{aligned} B\left(\frac{1-\beta}{2}, \frac{m+1+n(1-\alpha)}{2}\right) &\leq M_{\frac{1-\beta}{2}} \left(\frac{m+1+n(1-\alpha)}{2}\right)^{-\frac{1-\beta}{2}} \leq 4M_{\frac{1}{2}-\alpha} (m+1)^{-\frac{1-\beta}{2}}, \\ B\left(\frac{1-\alpha-\beta}{2}, \frac{m+1+n(1-\alpha)}{2}\right) &\leq M_{\frac{1-\alpha-\beta}{2}} \left(\frac{m+1+n(1-\alpha)}{2}\right)^{-\frac{1-\alpha-\beta}{2}} \leq 4M_{\frac{1}{2}-\alpha} (n+1)^{-\frac{1-\alpha-\beta}{2}}, \\ B\left(\frac{1-\beta}{2}, \frac{m+(n+1)(1-\alpha)}{2}\right) &\leq M_{\frac{1-\beta}{2}} \left(\frac{m+(n+1)(1-\alpha)}{2}\right)^{-\frac{1-\beta}{2}} \leq 4M_{\frac{1}{2}-\alpha} (n+1)^{-\frac{1-\alpha-\beta}{2}}. \end{aligned}$$

□

Remark 3.11 The restriction $\alpha \in (0, \frac{1}{2})$ in Lemma 3.10 is necessary since $M = 4M_{\frac{1}{2}-\alpha}$ diverges as $\alpha \uparrow \frac{1}{2}$ (see the definition of M_δ in Lemma 3.9). This is not unexpected, since for $\alpha > \frac{1}{2}$ we are no longer in the Young regime and we would need techniques like paracontrolled distributions or regularity structures to solve the equation for Γ .

Lemma 3.10 together with the following basic inequality constitutes the proof of Theorem 1.1.

Lemma 3.12 *Let $\beta \in (0, 1)$. Then there exists an $L > 0$ such that for $z \geq 0$*

$$\sum_{k=0}^{\infty} \frac{z^k}{(k!)^\beta} \leq L \exp(Lz^{\frac{1}{\beta}}).$$

Proof Let $\delta > 0$. By writing $z^k = ((1 + \delta)z)^k (1 + \delta)^{-k}$ we get with Hölder’s inequality

$$\sum_{k=0}^{\infty} \frac{z^k}{(k!)^\beta} \leq \left(\sum_{k=0}^{\infty} \left(\frac{((1+\delta)z)^k}{(k!)^\beta} \right)^{\frac{1}{\beta}} \right)^\beta \left(\sum_{k=0}^{\infty} (1 + \delta)^{-\frac{k}{1-\beta}} \right)^{1-\beta} \simeq \exp(\beta(1 + \delta)^{\frac{1}{\beta}} z^{\frac{1}{\beta}}).$$

□

Lemma 3.13 *There exists a $C > 0$ (independent of b) such that for all $\mu \in \mathbb{N}_0^d$ with $|\mu| \leq 1$, and for all $t > 0, x, y \in \mathbb{R}^d$,*

$$\begin{aligned} \partial_x^\mu \Gamma_t(x, y) &= \partial_x^\mu p(t, x - y) + \sum_{k=1}^{\infty} \int_0^t \int_{\mathbb{R}^d} \partial_x^\mu p(t - s, x - z) \Psi_{s,t}^{y,k}(z) \, dz \, ds, \quad (29) \\ |\partial_x^\mu \Gamma_t(x, y) - \partial_x^\mu p(t, x - y)| &\leq Ct^{-\frac{|\mu|}{2}} p(ct, x - y) (\|\Delta_{-1}b\|_{C_t L^\infty} t^{\frac{1}{2}} \vee \|\Delta_{\geq 0}b\|_{C_t B_{\infty,1}^{-\alpha}} t^{\frac{1-\alpha}{2}}) \\ &\quad \times \exp\left(Ct \left[\|\Delta_{-1}b\|_{C_t L^\infty}^2 + \|\Delta_{\geq 0}b\|_{C_t B_{\infty,1}^{-\alpha}}^{\frac{2}{1-\alpha}} \right]\right). \quad (30) \end{aligned}$$

Proof To show both Eqs. 29 and 30 it is sufficient to estimate the series with the modulus of each term in the series in the right-hand side of Eq. 29 by the right-hand side of Eq. 30.

Let K, C, M be as in Lemma 3.10. Again, we will write “ X ” and “ Y ” instead of “ $\|\Delta_{-1}b\|_{C_t L^\infty}$ ” and “ $\|\Delta_{\geq 0}b\|_{C_t B_{\infty,1}^{-\alpha}}$ ”. With $i = |\mu|$

$$\begin{aligned} \sum_{k=1}^{\infty} \int_0^t \left| \int_{\mathbb{R}^d} \partial_x^\mu p(t - s, x - z) \Psi_{s,t}^{y,k}(z) \, dz \right| \, ds &\leq \left(\sum_{k=1}^{\infty} \mathcal{J}_{i,k}^0(t) \right) p(ct, x - y) \\ &\leq Kt^{-\frac{i}{2}} p(ct, x - y) \sum_{\substack{m,n \in \mathbb{N}_0: \\ m+n \geq 1}} \frac{(CMXt^{\frac{1}{2}})^m}{(m!)^{\frac{1}{2}}} \frac{(CMYt^{\frac{1-\alpha}{2}})^n}{(n!)^{\frac{1-\alpha}{2}}} \\ &\leq Kt^{-\frac{i}{2}} p(ct, x - y) CM(Xt^{\frac{1}{2}} + Yt^{\frac{1-\alpha}{2}}) \\ &\quad \times \left(\sum_{m \in \mathbb{N}_0} \frac{(CMXt^{\frac{1}{2}})^m}{(m!)^{\frac{1}{2}}} \right) \left(\sum_{n \in \mathbb{N}_0} \frac{(CMYt^{\frac{1-\alpha}{2}})^n}{(n!)^{\frac{1-\alpha}{2}}} \right). \end{aligned}$$

Indeed, for $a, b > 0$

$$\begin{aligned} \sum_{\substack{m, n \in \mathbb{N}_0 \\ m+n \geq 1}} \frac{a^m}{(m!)^{\frac{1}{2}}} \frac{b^n}{(n!)^{\frac{1-\alpha}{2}}} &\leq \sum_{m, n \in \mathbb{N}_0} \frac{a^{m+1}}{((m+1)!)^{\frac{1}{2}}} \frac{b^n}{(n!)^{\frac{1-\alpha}{2}}} + \sum_{m, n \in \mathbb{N}_0} \frac{a^m}{(m!)^{\frac{1}{2}}} \frac{b^{n+1}}{((n+1)!)^{\frac{1-\alpha}{2}}} \\ &\leq (a+b) \sum_{m, n \in \mathbb{N}_0} \frac{a^m}{(m!)^{\frac{1}{2}}} \frac{b^n}{(n!)^{\frac{1-\alpha}{2}}}. \end{aligned}$$

Now by applying Lemma 3.12 we obtain the desired bound. □

Proof of the heat-kernel upper bound (2) of Theorem 1.1 This is a direct consequence of Lemma 3.13, as there exists a $K > 0$ such that for all $t \geq 0$

$$Ct(Xt^{\frac{1}{2}} \vee Yt^{\frac{1-\alpha}{2}}) \leq \exp\left(Kt[X^2 + Y^{1-\alpha}]\right).$$

□

4 Heat-Kernel Lower Bounds

The lower bound follows from Lemma 3.13 together with the next result, which is a small variation of [20, Lemma 4.3.8].

Lemma 4.1 *Let $q_t : \mathbb{R}^d \times \mathbb{R}^d \rightarrow [0, \infty)$ for all $t \in [0, \infty)$. Suppose that $(q_t)_{t \in [0, \infty)}$ satisfies the Chapman-Kolmogorov equations, i.e., $q_{t+s}(x, y) = \int_{\mathbb{R}^d} q_t(x, z)q_s(z, y) dz$. Let $a, b > 0$. Suppose that $q_t(x, y) \geq bt^{-\frac{d}{2}}$ for all $t \in (0, a]$ and $x, y \in \mathbb{R}^d$ with $|x - y| \leq \sqrt{t}$. Then there exist a $\kappa \in (0, 1)$ and an $M > 1$, which only depends on b and d , such that for all $t \in [0, \infty)$ and $x, y \in \mathbb{R}^d$*

$$q_t(x, y) \geq M^{-1-\frac{1}{a}} p(\kappa t, x - y).$$

Proof By following the first step of the proof of [20, Lemma 4.3.8] we find a $\kappa \in (0, 1)$ and a $M > 1$ which depend only on b and d such that for all $t \in (0, a]$ and $x, y \in \mathbb{R}^d$

$$q_t(x, y) \geq M^{-1} p(\kappa t, x - y).$$

Let $t > a$ and $n = \lceil \frac{t}{a} \rceil$. Then for all $x, y \in \mathbb{R}^d$

$$\begin{aligned} q_t(x, y) &= \int_{(\mathbb{R}^d)^{n-1}} q_{\frac{t}{n}}(x, z_1)q_{\frac{t}{n}}(z_1, z_2) \cdots q_{\frac{t}{n}}(z_{n-1}, y) dz \\ &\geq \int_{(\mathbb{R}^d)^{n-1}} M^{-n} p(\kappa \frac{t}{n}, x - z_1)p(\kappa \frac{t}{n}, z_1 - z_2) \cdots p(\kappa \frac{t}{n}, z_{n-1} - y) dz \\ &\geq M^{-1-\frac{1}{a}} p(\kappa t, x - y). \end{aligned}$$

□

Now we can prove the heat-kernel lower bounds:

Proof of the heat-kernel lower bound (3) of Theorem 1.1 We want to apply Lemma 4.1. Therefore we will find an a such that the condition is satisfied. Once more we will write “ X ” and “ Y ” instead of “ $\|\Delta_{-1}b\|_{C_L^\infty}$ ” and “ $\|\Delta_{\geq 0}b\|_{C_B^\infty}$ ”. Let us also take

$X = \|\Delta_{-1}b\|_{C_t L^\infty}$ and $Y = \|\Delta_{\geq 0}b\|_{C_t B_{\infty,1}^{-\alpha}}$. Let $\alpha \in (0, \frac{1}{2})$, $c > 1$ and $C > 0$ be as in Lemma 3.13. Then Eq. 30 gives for $a > 0$, $t \in (0, a]$ and $x, y \in \mathbb{R}^d$ with $|x - y| \leq \sqrt{t}$:

$$\begin{aligned} \Gamma_t(x, y) &\geq p(t, x - y) - C(Xt^{\frac{1}{2}} \vee Yt^{\frac{1-\alpha}{2}}) \exp\left(Ct[X^2 + Y^{\frac{2}{1-\alpha}}]\right) p(ct, x - y) \\ &\geq (2\pi t)^{-\frac{d}{2}} e^{-\frac{1}{2}} - C((X^2a)^{\frac{1}{2}} \vee (Y^{\frac{2}{1-\alpha}}a)^{\frac{1-\alpha}{2}}) \exp\left(Ca[X^2 + Y^{\frac{2}{1-\alpha}}]\right) c^{-\frac{d}{2}} (2\pi t)^{-\frac{d}{2}}. \end{aligned}$$

Therefore, it holds that $\Gamma_t(x, y) \geq \frac{1}{2}(2\pi t)^{-\frac{d}{2}} e^{-\frac{1}{2}}$ if

$$C((X^2a)^{\frac{1}{2}} \vee (Y^{\frac{2}{1-\alpha}}a)^{\frac{1-\alpha}{2}}) \exp\left(Ca[X^2 + Y^{\frac{2}{1-\alpha}}]\right) c^{-\frac{d}{2}} \leq \frac{e^{-\frac{1}{2}}}{2}.$$

Hence there exists a $K \in (0, 1)$ (which only depends on c, C and α) such that the choice $a = K[X^2 + Y^{\frac{2}{1-\alpha}}]^{-1}$ works. So by Lemma 4.1 there exist a $\kappa \in (0, 1)$ and a $M > 1$ such that for all $t \in [0, \infty)$ and $x, y \in \mathbb{R}^d$,

$$\Gamma_t(x, y) \geq M^{-1-\frac{1}{a}} p(\kappa t, x - y) = \frac{1}{M} \exp\left(-t \frac{\log M}{K}[X^2 + Y^{\frac{2}{1-\alpha}}]\right) p(\kappa t, x - y).$$

This proves that Eq. 3 holds for a large enough C . □

5 Proof of Corollary 1.2

As before, we consider $b \in C([0, T], B_{\infty,1}^{-\alpha})$ for some $\alpha \in (0, \frac{1}{2})$ and we let $X = (X_t)_{t \in [0, T]}$ be the solution to the martingale problem for $((\mathcal{L}_t)_{t \in (0, T]}, \delta_x)$. We prove Corollary 1.2, which means that we estimate the probability that X escapes a box of size K before time T . The estimate is a consequence of our heat-kernel estimates (Theorem 1.1), Markov’s inequality and the Garsia-Rademich-Rumsey inequality. By the latter (see [21, Theorem 2.1.3]) we have for $\kappa > 0$

$$\kappa |X_t - X_s| \leq 4 \int_0^{t-s} u^{-\frac{1}{2}} \sqrt{\log\left(1 + \frac{4(F_{T,\kappa} - T^2)}{u^2}\right)} du, \tag{31}$$

where

$$F_{T,\kappa} = \int_0^T \int_0^T \exp\left(\kappa \left(\frac{|X_{r_2} - X_{r_1}|}{|r_2 - r_1|^{\frac{1}{2}}}\right)^2\right) dr_1 dr_2. \tag{32}$$

In the proof of Corollary 5.2 we will bound the right-hand side of Eq. 31 in terms of a function ζ . In the next lemma we start by gathering some auxiliary facts about ζ .

Lemma 5.1 *Let $\zeta, \psi : (0, \infty) \rightarrow (0, \infty)$ be given by*

$$\zeta(r) := \int_0^r u^{-\frac{1}{2}} \left(\sqrt{\log(1 + u^{-2})} \vee 1\right) du, \quad \psi(r) := r^{\frac{1}{2}} \sqrt{(\log(\frac{1}{r})) \vee 1}.$$

There exist $m, M > 0$ such that $m\zeta(r) \leq \psi(r) \leq M\zeta(r)$ for all $r > 0$. Moreover, $\psi(rs) \leq \sqrt{2}\psi(r)\psi(s)$ for all $r, s > 0$ and ψ is strictly increasing.

Proof That ψ is strictly increasing on $(\frac{1}{e}, \infty)$ will be clear, whereas on $[0, \frac{1}{e})$ it follows by calculating its derivative. Since ψ and ζ are continuous and bounded away from 0 and ∞ on compact subintervals of $(0, \infty)$, the existence of such m and M follows once we show

that $\lim_{r \rightarrow 0} \frac{\zeta(r)}{\psi(r)}$ and $\lim_{r \rightarrow \infty} \frac{\zeta(r)}{\psi(r)}$ exist and are in $(0, \infty)$. By applying L'Hospital's rule we obtain

$$\lim_{r \rightarrow 0} \frac{\zeta(r)}{\psi(r)} = \lim_{r \rightarrow 0} \frac{\int_0^r u^{-\frac{1}{2}} \sqrt{\log(1+u^{-2})} du}{r^{\frac{1}{2}} \sqrt{\log(\frac{1}{r})}} \in (0, \infty).$$

And also for $r \rightarrow \infty$ we have

$$\lim_{r \rightarrow \infty} \frac{\zeta(r)}{\psi(r)} = \lim_{r \rightarrow \infty} \frac{\int_0^{\sqrt{e-1}} u^{-\frac{1}{2}} \sqrt{\log(1+u^{-2})} du + \int_{\sqrt{e-1}}^r u^{-\frac{1}{2}} du}{r^{\frac{1}{2}}} \in (0, \infty).$$

Furthermore

$$\psi(rs) = (rs)^{\frac{1}{2}} \left(\sqrt{(\log(\frac{1}{r}) + \log(\frac{1}{s})) \vee 1} \right)$$

and for all $x, y \in \mathbb{R}$ we have $(x + y) \vee 1 \leq x \vee 1 + y \vee 1 \leq 2(x \vee 1)(y \vee 1)$. Therefore,

$$\psi(rs) \leq \sqrt{2}(rs)^{\frac{1}{2}} \left(\sqrt{\log(\frac{1}{r})} \vee 1 \right) \left(\sqrt{\log(\frac{1}{s})} \vee 1 \right) = \sqrt{2}\psi(r)\psi(s).$$

□

Corollary 5.2 *Let ψ be as in Lemma 5.1 and let $C > 0$ be as in Theorem 1.1. Then there exists an $M > 0$ such that for all $T \geq 1$*

$$\begin{aligned} & \mathbb{E}_x \left[\exp \left(\frac{1}{M} \left(\sup_{\substack{s,t \in [0,T] \\ s < t}} |X_t - X_s| \right)^2 \right) \right] \\ & \leq M \exp \left(CT \left[\|\Delta_{-1} b\|_{C_T L^\infty}^2 + \|\Delta_{\geq 0} b\|_{C_T B_{\infty,1}^{-\alpha}}^{\frac{2}{1-\alpha}} \right] \right). \end{aligned} \tag{33}$$

Proof The proof is inspired by [11, Corollary A.5]. Unfortunately we cannot directly apply that result, because the constant they derive depends on the time interval $[0, T]$ (even though this is not explicitly stated).

Let us define $G_{T,\kappa} := 2\sqrt{F_{T,\kappa} \vee 4}$, where $F_{T,\kappa}$ is as in Eq. 32. Let ζ be as in Lemma 5.1. By Eq. 31 and using $4(F_{T,\kappa} - T^2) \leq G_{T,\kappa}^2$ we have by a substitution and by Lemma 5.1 (observe that $G_{T,\kappa} \geq 4 \geq e$) that for $T \geq 1, \kappa > 0, s, t \in [0, T]$ with $s < t$ and by writing $G = G_{T,\kappa}$

$$\begin{aligned} \kappa |X_t - X_s| & \leq 4\sqrt{G} \int_0^{\frac{t-s}{G}} u^{-\frac{1}{2}} \sqrt{\log\left(1 + \frac{1}{u^2}\right)} du \lesssim \sqrt{G} \zeta\left(\frac{t-s}{G}\right) \\ & \lesssim \sqrt{G} \psi\left(\frac{t-s}{G}\right) \lesssim \sqrt{G} \psi(t-s) \psi\left(\frac{1}{G}\right) \lesssim \psi(t-s) \sqrt{\log G}. \end{aligned}$$

Let $M > 0$ be such that $\kappa |X_t - X_s| \leq \sqrt{M} \psi(t-s) \sqrt{\log G_{T,\kappa}}$ for all $T \geq 1, \kappa > 0$ and $s, t \in [0, T]$ with $s < t$. Then

$$\mathbb{E}_x \left[\exp \left(\frac{\kappa^2}{M} \left(\sup_{\substack{s,t \in [0,T] \\ s < t}} |X_t - X_s| \right)^2 \right) \right] \leq \mathbb{E}_x [G_{T,\kappa}].$$

As by Jensen's inequality $\mathbb{E}_x [G_{T,\kappa}] = 2\mathbb{E}_x [\sqrt{F_{T,\kappa} \vee 4}] \leq 2\sqrt{\mathbb{E}_x [F_{T,\kappa}] + 4}$ we will obtain a bound of $\mathbb{E}_x [G_{T,\kappa}]$, by estimating $\mathbb{E}_x [F_{T,\kappa}]$. Let $c \in (0, 1)$ and $\kappa > 0$ be such that $\kappa < \frac{1}{2c}$.

Then for all $r_2, r_1 > 0$ with $r_2 \neq r_1$

$$\int_{\mathbb{R}^d} p(c|r_2 - r_1|, y) \exp(\kappa \left(\frac{|y|}{|r_2 - r_1|^{\frac{1}{2}}}\right)^2) dy = \left(\frac{1}{1-2c\kappa}\right)^{\frac{d}{2}} < \infty. \tag{34}$$

Hence, by Theorem 1.1

$$\begin{aligned} \mathbb{E}_x[F_{T,\kappa}] &= \int_0^T \int_0^T \mathbb{E}_x \left[\int_{\mathbb{R}^d} \Gamma_{|r_2-r_1|}(y, X_{r_1}) \exp(\kappa \left(\frac{|y - X_{r_1}|}{|r_2 - r_1|^{\frac{1}{2}}}\right)^2) dy \right] dr_1 dr_2 \\ &\leq C \left(\frac{1}{1-2c\kappa}\right)^{\frac{d}{2}} \int_0^T \int_0^T \exp \left(C|r_2 - r_1| \left[\|\Delta_{-1}b\|_{C_t L^\infty}^2 + \|\Delta_{\geq 0}b\|_{C_t B_{\infty,1}^{-\alpha}}^{\frac{2}{1-\alpha}} \right] \right) dr_1 dr_2. \end{aligned}$$

The proof is completed by observing that for $A \geq 1$

$$\int_0^T \int_0^T \exp(A|r_2 - r_1|) dr_1 dr_2 = 2 \int_0^T \int_0^t e^{A(t-s)} ds dt \lesssim e^{AT}.$$

□

Proof of Corollary 1.2 As $T \geq 1 \geq e^{-1}$ we have $\psi(T) = \sqrt{T}$. Therefore, by Markov’s inequality for all $M, K > 0$ and the fact that ψ is strictly increasing:

$$\begin{aligned} \mathbb{P}_x \left(\sup_{t \in [0, T]} |X_t - x| \geq K \right) &\leq \mathbb{E}_x \left[\exp \left(\frac{1}{MT} \sup_{t \in [0, T]} |X_t - x|^2 \right) \right] \exp \left(- \frac{K^2}{MT} \right) \\ &\leq \mathbb{E}_x \left[\exp \left(\frac{1}{M} \left(\sup_{\substack{s, t \in [0, T] \\ s < t}} \frac{|X_t - X_s|}{\psi(t-s)} \right)^2 \right) \right] \exp \left(- \frac{K^2}{MT} \right). \end{aligned}$$

So Eq. 4 follows from Corollary 5.2. □

A Appendix

Theorem A.1 *Suppose $\alpha < 0$ and $\beta > 0$ are such that $\alpha + \beta > 0$. Let $p, p_1, p_2, q_1, q_2 \in [1, \infty]$ be such that*

$$\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}. \tag{35}$$

For all $r \geq q_1$

$$\|u \cdot v\|_{B_{p,r}^\alpha} \lesssim \|u\|_{B_{p_1,q_1}^\alpha} \|v\|_{B_{p_2,q_2}^\beta}. \tag{36}$$

Proof For the proof see also [18, Corollary 2.1.35]. By slightly adapting [2, Theorem 2.82] and by using the Hölder inequality and [2, Theorem 2.79] (for Eq. 38), we obtain implies the following two estimates.

$$\|u \otimes v\|_{B_{p,q}^{\alpha+\beta}} \lesssim \|u\|_{B_{p_1,q_1}^\alpha} \|v\|_{B_{p_2,q_2}^\beta}, \tag{37}$$

$$\|u \otimes v\|_{B_{p,r}^\alpha} \lesssim \|v\|_{L^{p_2}} \|u\|_{B_{p_1,r}^\alpha} \lesssim \|v\|_{B_{p_2,q_2}^\beta} \|u\|_{B_{p_1,q_1}^\alpha}. \tag{38}$$

As [2, Theorem 2.52] implies $\|u \odot v\|_{B_{p,q}^{\alpha+\beta}} \lesssim \|u\|_{B_{p_1,q_1}^\alpha} \|v\|_{B_{p_2,q_2}^\beta}$, combining the above inequalities proves Eq. 36. □

Acknowledgements This work was supported by the German Science Foundation (DFG) via the Forschergruppe FOR2402 “Rough paths, stochastic partial differential equations and related topics”. WvZ was supported by the DFG through SPPI1590 “Probabilistic Structures in Evolution”. NP thanks the DFG for financial support through the Heisenberg program. The main part of the work was done while NP was employed at Humboldt-Universität zu Berlin and Max-Planck-Institute for Mathematics in the Sciences, Leipzig. The authors are also grateful to the anonymous referees for their valuable feedback, suggestions and careful reading.

Funding Open Access funding enabled and organized by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Andrews, G.E., Askey, R., Roy, R.: Special functions volume 71 of encyclopedia of mathematics and its applications. Cambridge University Press, Cambridge (1999)
2. Bahouri, H., Chemin, J.Y., Danchin, R.: Fourier analysis and nonlinear partial differential equations, volume 343 of grundlehren der mathematischen wissenschaften [fundamental principles of mathematical Sciences]. Springer, Heidelberg (2011)
3. Bass, R.F., Chen, Z.Q.: Stochastic differential equations for Dirichlet processes. *Probab. Theory Related Fields* **121**(3), 422–446 (2001)
4. Brox, T.: A one-dimensional diffusion process in a Wiener medium. *Ann. Probab.* **14**(4), 1206–1218 (1986)
5. Cannizzaro, G., Chouk, K.: Multidimensional SDEs with singular drift and universal construction of the polymer measure with white noise potential. *Ann. Probab.* **46**(3), 1710–1763 (2018)
6. Delarue, F., Diel, R.: Rough paths and 1d SDE with a time dependent distributional drift: application to polymers. *Probab. Theory Related Fields* **165**(1-2), 1–63 (2016)
7. Flandoli, F., Issoglio, E., Russo, F.: Multidimensional stochastic differential equations with distributional drift. *Trans. Amer. Math. Soc.* **369**(3), 1665–1688 (2017)
8. Flandoli, F., Russo, F., Wol, J.: Some SDEs with distributional drift. I. General calculus. *Osaka J. Math.* **40**(2), 493–542 (2003)
9. Friedman, A.: Partial differential equations of parabolic type. Prentice-Hall, Inc., Englewood Cliffs, N.J. (1964)
10. Friedman, A.: Stochastic differential equations and applications. Vol. 1. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, Probability and mathematical statistics, Vol. 28 (1975)
11. Friz, P.K., Victoir, N.B.: Multidimensional stochastic processes as rough paths, Volume 120 of Cambridge Studies in Advanced Mathematics. Cambridge, Universit Press (2010). Theory and applications
12. Gubinelli, M., Imkeller, P., Perkowski, N.: Paracontrolled distributions and singular PDEs. *Forum Math. Pi* **3**(e6), 75 (2015)
13. Gubinelli, M., Perkowski, N.: KPZ, reloaded. *Comm. Math Phys.* **349**(1), 165–269 (2017)
14. Hairer, M.: Rough stochastic PDEs. *Comm Pure Appl. Math.* **64**(11), 1547–1585 (2011)
15. Hairer, M.: Solving the KPZ equation. *Ann. Math.* **178**(2), 559–664 (2013)
16. König, W.: The Parabolic Anderson Model. Pathways in Mathematics. Birkhäuser, Springer (2016). Random walk in random potential
17. König, W., Perkowski, N., van Zuijlen, W.B.: Long-time asymptotics of the two-dimensional parabolic Anderson model with white-noise potential. [arXiv:2009.11611](https://arxiv.org/abs/2009.11611) (2020)
18. Martin, J.: Refinements of the solution theory for singular SPDEs. PhD thesis Humboldt-Universität zu Berlin (2018)
19. Martin, J., Perkowski, N.: Paracontrolled distributions on Bravais lattices and weak universality of the 2d parabolic Anderson model. *Ann. Inst. Henri Poincaré Probab. Stat.* **55**(4), 2058–2110 (2019)

20. Stroock, D.W.: Partial differential equations for probabilists, volume 112 of cambridge studies in advanced mathematics. Cambridge University Press, Cambridge (2008)
21. Stroock, D.W., Varadhan, S.R.S.: Multidimensional diffusion processes. Classics in Mathematics. Springer-Verlag, Berlin (2006). Reprint of the 1997 edition
22. Zhang, X., Zhao, G.: Heat kernel and ergodicity of SDEs with distributional drifts. arXiv:[1710.10537](https://arxiv.org/abs/1710.10537) (2017)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.