

Appendix

List of figures

<i>Figure 1. Mechanoreceptors of mouse skin innervated by the saphenous nerve</i>	7
<i>Figure 2. Schematic drawing of different molecules involved in transduction and modulation of nociceptive stimuli</i>	10
<i>Figure 3. Overview of the Affymetrix experimental procedure</i>	42
<i>Figure 4. Overview of the anti-NGF production</i>	48
<i>Figure 5. Overview of the skin-nerve preparation</i>	49
<i>Figure 6. Overview of the patch perfusion system used to record Iheat</i>	51
<i>Figure 7. Schematic drawing of the mechanical parameters studied for evaluating mechanoreceptor function in mice</i>	55
<i>Figure 8. Standardized stimulus response protocol</i>	57
<i>Figure 9. Stimulus response function of high and low threshold mechanoreceptors</i>	59
<i>Figure 10. Mechanical latency</i>	61
<i>Figure 11. Von Frey threshold of mechanically sensitive receptors</i>	63
<i>Figure 12. Standardized stimulus response protocol used to record velocity response function</i>	64
<i>Figure 13. Velocity response function of low threshold mechanoreceptors</i>	65
<i>Figure 14. Mechanical latency for characteristic displacement at 1.4 and 2.9 m/s velocity of moving probe for all types of cutaneous mechanoreceptors</i>	65
<i>Figure 15. Determination of the effective dose of anti-NGF</i>	67
<i>Figure 16. anti-NGF treatment during the early postnatal development induce permanent change in proportion of cutaneous sensory fibers</i>	69
<i>Figure 17. Mechanosensitivity of Aδ fibers after anti-NGF treatment</i>	70
<i>Figure 18. Mechanosensitivity of Aβ fibers after anti-NGF treatment</i>	70
<i>Figure 19. Mechanosensitivity of C fibers after anti-NGF treatment</i>	71
<i>Figure 20. Mechanical threshold of C-fibers after the anti-NGF treatment</i>	72
<i>Figure 21. The comaparison of the stimulus response function of CMH and CM fibers - CM were more sensitive to mechanical stimuli compared to the CMH</i>	72
<i>Figure 22. Standardized heating protocol</i>	73
<i>Figure 23. Effect of anti-NGF treatment on heat sensitivity of CMH</i>	74
<i>Figure 24. Electron microscopy of the saphenous nerve</i>	76

<i>Figure 25. anti-NGF treatment led to behavioral thermal hypoalgesia</i>	77
<i>Figure 26. Schematic diagram of changes in cell population in DRG after the anti-NGF treatment</i>	81
<i>Figure 27. Specific expression pattern of transmembrane channel genes with altered gene expression after anti-NGF treatment</i>	83
<i>Figure 28. Specific expression pattern of transmembrane receptor genes with altered gene expression after anti-NGF treatment</i>	84
<i>Figure 29. Specific expression pattern of transmembrane protein genes with altered gene expression after anti-NGF treatment</i>	85
<i>Figure 30. Specific expression pattern of transmembrane protein gene with altered gene expression after anti-NGF treatment</i>	86
<i>Figure 31. Specific expression pattern of neurotransmitter related gene with altered gene expression after anti-NGF treatment</i>	87
<i>Figure 32. Specific expression pattern of cytoskeletal binding genes with altered gene expression after anti-NGF treatment</i>	87
<i>Figure 33. Specific expression pattern of GPCR gene with altered gene expression after anti-NGF treatment</i>	88
<i>Figure 34. Detailed view of Gpr177 coding region and the position of the gene trap</i>	89
<i>Figure 35. Specific expression pattern of G protein signaling related genes with altered gene expression after anti-NGF treatment</i>	90
<i>Figure 36. Specific expression pattern of signaling protein genes with altered gene expression after anti-NGF treatment</i>	91
<i>Figure 37. Specific expression pattern of kinase receptor gene with altered gene expression after anti-NGF treatment</i>	92
<i>Figure 38. Specific expression pattern of transferase genes with altered gene expression after anti-NGF treatment</i>	93
<i>Figure 39. Specific expression pattern of transcription factor gene with altered gene expression after anti-NGF treatment</i>	94
<i>Figure 40. Gene expression change in anti-NGF treated mice relative to control</i>	95
<i>Figure 41. Mice expressing the c-Kit null mutation have an altered proportion of cutaneous sensory fibers</i>	98
<i>Figure 42. Mechanosensitivity of D-hairs in c-Kit^{-/-} mice</i>	100
<i>Figure 43. Mechanosensitivity of SAMs in c-Kit^{-/-} mice</i>	101
<i>Figure 44. Mechanosensitivity of mechanonociceptors in c-Kit^{-/-} mice</i>	102
<i>Figure 45. Von Frey threshold of mechanically sensitive receptors in c-Kit^{-/-} mice</i>	103
<i>Figure 46. Electrophysiological analysis of sensory neurons in control and c-Kit^{-/-} mice</i>	103
<i>Figure 47. Effect of c-Kit null mutation on heat sensitivity of CMH</i>	104
<i>Figure 48. c-Kit null mutation led to behavioral thermal hypoalgesia and mechanical hyperalgesia</i>	105
<i>Figure 49. Effect of c-Kit haploinsufficiency on heat sensitivity of CMH</i>	106
<i>Figure 50. Potentiation of I_{heat} in DRG neurons by c-Kit ligand SCF</i>	107

<i>Figure 51. Approximately 50% of heat sensitive cultivated DRG cells were affected by SCF application</i>	108
<i>Figure 52. SCF alters the threshold temperature of I_{heat}</i>	109
<i>Figure 53. Schematic diagram of the changes in the phenotype of DRG primary sensory neurons and alternations in gene expression levels after temporal neonatal NGF deprivation</i>	119
<i>Figure 54. Schematic diagram of the abundance of distinct DRG neuron subtypes after the temporal neonatal NGF deprivation and in c-Kit null mutant.</i>	126

List of tables

<i>Table 1. Detailed breakdown of the single fibre recordings from C57BL/6 mice at 32°C and 24°C</i>	56
<i>Table 2. Detailed breakdown of the single fiber recordings from control and anti-NGF treated mice</i>	75
<i>Table 3. List of transcripts whose expression level changed after anti-NGF treatment during the critical period of postnatal development</i>	80
<i>Table 4. Tabular overview of transcripts expressed in DRG</i>	82
<i>Table 5. Detailed breakdown of the single fiber recordings from c-Kit^{-/-} and control c-Kit^{+/+} mice</i>	99
<i>Table 6. Similarities and differences between the phenotype of the neonatally NGF deprived mice and SorL1, calca and c-Kit null mutants.</i>	124