

5 Literaturverzeichnis

- [1] Langhorst, P.; Lambertz, M.; Kluge, W. und Rittweger, J. (1992): Different modes of dampening influence from baroreceptors are determined by the functional organization of the NTS neuronal network, *J.Auton.Nerv.Syst.* (volume 41), issue 1-2, pp. 141-156. URL: PM:1491110
- [2] DiBona, G. F. (2004): The sympathetic nervous system and hypertension: recent developments, *Hypertension* (volume 43), issue 2, pp. 147-150. URL: PM:14707153
- [3] Lohmeier, T. E.; Irwin, E. D.; Rossing, M. A.; Serdar, D. J. und Kieval, R. S. (2004): Prolonged activation of the baroreflex produces sustained hypotension, *Hypertension* (volume 43), issue 2, pp. 306-311. URL: PM:14707159
- [4] Robertson, D.; Hollister, A. S.; Biaggioni, I.; Netterville, J. L.; Mosqueda-Garcia, R. und Robertson, R. M. (1993): The diagnosis and treatment of baroreflex failure, *N.Engl.J.Med.* (volume 329), issue 20, pp. 1449-1455. URL: PM:8413455
- [5] Bradbury, S. und Eggleston, C. (1925): Postural hypotension. A report of three cases, *Am.Heart J.* (volume 16), issue 1, pp. 73-86.
- [6] SHY, G. M. und DRAGER, G. A. (1960): A neurological syndrome associated with orthostatic hypotension: a clinical-pathologic study, *Arch.Neurol.* (volume 2), pp. 511-527. URL: PM:14446364
- [7] Ali, Y. S.; Daamen, N.; Jacob, G.; Jordan, J.; Shannon, J. R.; Biaggioni, I. und Robertson, D. (2000): Orthostatic intolerance: a disorder of young women, *Obstet.Gynecol.Surv.* (volume 55), issue 4, pp. 251-259. URL: PM:10758621
- [8] Furlan, R.; Jacob, G.; Snell, M.; Robertson, D.; Porta, A.; Harris, P. und Mosqueda-Garcia, R. (1998): Chronic orthostatic intolerance: a disorder with discordant cardiac and vascular sympathetic control, *Circulation* (volume 98), issue 20, pp. 2154-2159. URL: PM:9815870
- [9] Hagbarth, K. E. (1979): Exteroceptive, proprioceptive, and sympathetic activity recorded with microelectrodes from human peripheral nerves, *Mayo Clin.Proc.* (volume 54), issue 6, pp. 353-365. URL: PM:156292
- [10] Esler, M. (1993): Clinical application of noradrenaline spillover methodology: delineation of regional human sympathetic nervous responses, *Pharmacol.Toxicol.* (volume 73), issue 5, pp. 243-253. URL: PM:8115306
- [11] Esler, M.; Lambert, G.; Brunner-La Rocca, H. P.; Vaddadi, G. und Kaye, D. (2003): Sympathetic nerve activity and neurotransmitter release in humans: translation from pathophysiology into clinical practice, *Acta Physiol Scand.* (volume 177), issue 3, pp. 275-284. URL: PM:12608997

- [12] Packer, M.; Coats, A. J.; Fowler, M. B.; Katus, H. A.; Krum, H.; Mohacsi, P.; Rouleau, J. L.; Tendera, M.; Castaigne, A.; Roecker, E. B.; Schultz, M. K. und DeMets, D. L. (2001): Effect of carvedilol on survival in severe chronic heart failure, *N.Engl.J.Med.* (volume 344), issue 22, pp. 1651-1658. URL: PM:11386263
- [13] La Rovere, M. T.; Bigger, J. T., Jr.; Marcus, F. I.; Mortara, A. und Schwartz, P. J. (1998): Baroreflex sensitivity and heart-rate variability in prediction of total cardiac mortality after myocardial infarction. ATRAMI (Autonomic Tone and Reflexes After Myocardial Infarction) Investigators, *Lancet* (volume 351), issue 9101, pp. 478-484. URL: PM:9482439
- [14] Rathmann, W.; Ziegler, D.; Jahnke, M.; Haastert, B. und Gries, F. A. (1993): Mortality in diabetic patients with cardiovascular autonomic neuropathy, *Diabet.Med.* (volume 10), issue 9, pp. 820-824. URL: PM:8281726
- [15] Shannon, J. R.; Flattem, N. L.; Jordan, J.; Jacob, G.; Black, B. K.; Biaggioni, I.; Blakely, R. D. und Robertson, D. (2000): Orthostatic intolerance and tachycardia associated with norepinephrine-transporter deficiency, *N.Engl.J.Med.* (volume 342), issue 8, pp. 541-549. URL: PM:10684912
- [16] Mathias, C. J.; Deguchi, K. und Schatz, I. (2001): Observations on recurrent syncope and presyncope in 641 patients, *Lancet* (volume 357), issue 9253, pp. 348-353. URL: PM:11210997
- [17] Schroeder, C.; Tank, J.; Boschmann, M.; Diedrich, A.; Sharma, A. M.; Biaggioni, I.; Luft, F. C. und Jordan, J. (2002): Selective norepinephrine reuptake inhibition as a human model of orthostatic intolerance, *Circulation* (volume 105), issue 3, pp. 347-353. URL: PM:11804991
- [18] Mosqueda-Garcia, R.; Fernandez-Violante, R.; Tank, J.; Snell, M.; Cunningham, G. und Furlan, R. (1998): Yohimbine in neurally mediated syncope. Pathophysiological implications, *J.Clin.Invest* (volume 102), issue 10, pp. 1824-1830. URL: PM:9819368
- [19] Brierley, E. J.; Jackson, M. J.; Clark, R. S. und Kenny, R. A. (2001): Alarming asystole, *Lancet* (volume 357), issue 9274, pp. 2100. URL: PM:11445105
- [20] Somers, V. K.; Dyken, M. E.; Mark, A. L. und Abboud, F. M. (1993): Sympathetic-nerve activity during sleep in normal subjects, *N.Engl.J.Med.* (volume 328), issue 5, pp. 303-307. URL: PM:8419815
- [21] Wittstein, I. S.; Thiemann, D. R.; Lima, J. A.; Baughman, K. L.; Schulman, S. P.; Gerstenblith, G.; Wu, K. C.; Rade, J. J.; Bivalacqua, T. J. und Champion, H. C. (2005): Neurohumoral features of myocardial stunning due to sudden emotional stress, *N.Engl.J.Med.* (volume 352), issue 6, pp. 539-548. URL: PM:15703419
- [22] Yeragani, V. K.; Tancer, M. und Uhde, T. (2003): Heart rate and QT interval

- variability: abnormal alpha-2 adrenergic function in patients with panic disorder, *Psychiatry Res.* (volume 121), issue 2, pp. 185-196. URL: PM:14656453
- [23] Delius, W.; Hagbarth, K. E.; Hongell, A. und Wallin, B. G. (1972): General characteristics of sympathetic activity in human muscle nerves, *Acta Physiol Scand.* (volume 84), issue 1, pp. 65-81. URL: PM:5029385
- [24] Mark, A. L.; Correia, M. L.; Rahmouni, K. und Haynes, W. G. (2002): Selective leptin resistance: a new concept in leptin physiology with cardiovascular implications, *J.Hypertens.* (volume 20), issue 7, pp. 1245-1250. URL: PM:12131511
- [25] Weyer, C.; Pratley, R. E.; Snitker, S.; Spraul, M.; Ravussin, E. und Tataranni, P. A. (2000): Ethnic differences in insulinemia and sympathetic tone as links between obesity and blood pressure, *Hypertension* (volume 36), issue 4, pp. 531-537. URL: PM:11040231
- [26] Rayner, D. V. und Trayhurn, P. (2001): Regulation of leptin production: sympathetic nervous system interactions, *J Mol Med* (volume 79), issue 1, pp. 8-20.
- [27] Rosenbaum, M.; Leibel, R. L. und Hirsch, J. (1997): Obesity, *N Engl J Med* (volume 337), issue 6, pp. 396-407.
- [28] Brabant, G.; Nave, H.; Mayr, B.; Behrend, M.; Van-Harmelen, V und Arner, P. (2002): Secretion of free and protein-bound leptin from subcutaneous adipose tissue of lean and obese women, *J.Clin.Endocrinol.Metab* (volume 87), issue 8, pp. 3966-3970. URL: PM:12161541
- [29] Eckberg, D. L. und Sleight, P. (1992): *Human Baroreflex in Health and Disease.*, Oxford, Clarendon Press, 1992.
- [30] Smyth, H. S.; Sleight, P. und Pickering, G. W. (1969): Reflex regulation of arterial pressure during sleep in man. A quantitative method of assessing baroreflex sensitivity, *Circ.Res.* (volume 24), issue 1, pp. 109-121. URL: PM:4303309
- [31] Penaz, J. (1973): Photoelectric measurement of blood pressure, volume and flow in the finger. Dresden, Conference Proceedings, pp. 162-164.
- [32] Mayer, S. (1874): Studien zur Physiologie des Herzens und der Blutgefäße. 5. Abhandlung. Über spontane Blutdruckschwankungen., *Sber.Akad.Wiss.Wien* (volume 74), pp. 281-307.
- [33] Hering, E. (1869): Über den Einfluß der Atmung auf den Kreislauf. 1.Mitteilung. Über Atembewegungen des Gefäßsystems., *Sber.Akad.Wiss.Wien* (volume 60), pp. 829-855.
- [34] Cyon, E. (1874): Zur Physiologie des Gefäßnervencentrums, *Pflugers Arch.* (volume 9), pp. 499-513.
- [35] Traube, L. (1865): Ueber periodische Thätigkeits-Aeusserungen des

vasomotorischen und Hemmungs-Nervencentrums, Cbl.Med.Wiss. (volume 56), pp. 881-885.

- [36] Penaz, J. (1978): Mayer Waves: History and Methodology, *Automedica* (volume 2), pp. 135-141.
- [37] Preiss, G. und Polosa, C. (1974): Patterns of sympathetic neuron activity associated with Mayer waves, *Am J.Physiol* (volume 226), issue 3, pp. 724-730. URL: PM:4817426
- [38] Polosa, C. (1984): Rhythms in the Activity of the Autonomic Nervous System: Their Role in the Generation of Systemic Arterial Pressure Waves., Miyakawa, K., *Mechanisms of Blood Pressure Waves*, pp. 27-41, Japan Sci.Soc.Press, Springer-Verlag, Tokyo, Berlin.
- [39] Tank, J.; Baevski, R. M.; Fender, A.; Baevski, A. R.; Graves, K. F.; Ploewka, K. und Weck, M. (2000): Reference values of indices of spontaneous baroreceptor reflex sensitivity, *Am J Hypertens* (volume 13), issue 3, pp. 268-275.
- [40] Tank, J.; Neuke, A.; Molle, A.; Jordan, J. und Weck, M. (2001): Spontaneous baroreflex sensitivity and heart rate variability are not superior to classic autonomic testing in older patients with type 2 diabetes, *Am J Med Sci* (volume 322), issue 1, pp. 24-30.
- [41] Ludbrook, J. (1984): Concern about gain: is this the best measure of performance of cardiovascular reflexes?, *Clin.Exp.Pharmacol.Physiol* (volume 11), issue 4, pp. 385-389. URL: PM:6518667
- [42] Grassi, G. (2004): Sympathetic and baroreflex function in hypertension: implications for current and new drugs, *Curr.Pharm.Des* (volume 10), issue 29, pp. 3579-3589. URL: PM:15579055
- [43] Christou, D. D.; Jones, P. P.; Jordan, J.; Diedrich, A.; Robertson, D. und Seals, D. R. (2005): Women have lower tonic autonomic support of arterial blood pressure and less effective baroreflex buffering than men, *Circulation* (volume 111), issue 4, pp. 494-498. URL: PM:15687139
- [44] Barnett, S. R.; Morin, R. J.; Kiely, D. K.; Gagnon, M.; Azhar, G.; Knight, E. L.; Nelson, J. C. und Lipsitz, L. A. (1999): Effects of age and gender on autonomic control of blood pressure dynamics, *Hypertension* (volume 33), issue 5, pp. 1195-1200. URL: PM:10334811
- [45] Narkiewicz, K.; Phillips, B. G.; Kato, M.; Hering, D.; Bieniaszewski, L. und Somers, V. (2005): A Gender Selective Interaction Between Aging, Blood Pressure and Sympathetic Nerve Activity, *Hypertension* (volume 45), issue 4, pp. 522-525.
- [46] Ng, A. V.; Callister, R.; Johnson, D. G. und Seals, D. R. (1993): Age and gender

influence muscle sympathetic nerve activity at rest in healthy humans, *Hypertension* (volume 21), issue 4, pp. 498-503. URL: PM:8458648

- [47] Shoemaker, J. K.; Hogeman, C. S.; Khan, M.; Kimmerly, D. S. und Sinoway, L. I. (2001): Gender affects sympathetic and hemodynamic response to postural stress, *Am.J.Physiol Heart Circ.Physiol* (volume 281), issue 5, pp. H2028-H2035. URL: PM:11668064
- [48] Brenner, R.; Perez, G. J.; Bonev, A. D.; Eckman, D. M.; Kosek, J. C.; Wiler, S. W.; Patterson, A. J.; Nelson, M. T. und Aldrich, R. W. (2000): Vasoregulation by the beta1 subunit of the calcium-activated potassium channel, *Nature* (volume 407), issue 6806, pp. 870-876. URL: PM:11057658
- [49] Lambert, G. W.; Kaye, D. M.; Thompson, J. M.; Turner, A. G.; Cox, H. S.; Vaz, M.; Jennings, G. L.; Wallin, B. G. und Esler, M. D. (1998): Internal jugular venous spillover of noradrenaline and metabolites and their association with sympathetic nervous activity, *Acta Physiol Scand.* (volume 163), issue 2, pp. 155-163. URL: PM:9648634
- [50] Tangri, K. K.; Gupta, S. K.; Vrat, S. und Husain, M. (1993): A study of effects of putative neurotransmitters injected into the lateral cerebral ventricle of man, *Prog.Neuropsychopharmacol.Biol.Psychiatry* (volume 17), issue 5, pp. 731-745. URL: PM:7902993
- [51] Jordan, J.; Tank, J.; Shannon, J. R.; Diedrich, A.; Lipp, A.; Schroder, C.; Arnold, G.; Sharma, A. M.; Biaggioni, I.; Robertson, D. und Luft, F. C. (2002): Baroreflex buffering and susceptibility to vasoactive drugs, *Circulation* (volume 105), issue 12, pp. 1459-1464. URL: PM:11914255
- [52] Grassi, G.; Turri, C.; Seravalle, G.; Bertinieri, G.; Pierini, A. und Mancia, G. (2001): Effects of chronic clonidine administration on sympathetic nerve traffic and baroreflex function in heart failure, *Hypertension* (volume 38), issue 2, pp. 286-291. URL: PM:11509491
- [53] Mancia, G.; Ferrari, A.; Gregorini, L. und Zanchetti, A. (1979): Clonidine and carotid baroreflex in essential hypertension, *Hypertension* (volume 1), issue 4, pp. 362-370. URL: PM:541039
- [54] Luft, F. C.; Veelken, R.; Becker, H.; Ganten, D.; Lang, R. E. und Unger, T. (1986): Effect of urapidil, clonidine, and prazosin on sympathetic tone in conscious rats, *Hypertension* (volume 8), issue 4, pp. 303-311. URL: PM:3957416
- [55] Head, G. A. (1991): Cardiovascular functions of central noradrenergic neurons in rabbits, *Clin.Exp.Pharmacol.Physiol* (volume 18), issue 1, pp. 51-54. URL: PM:1674453
- [56] Robertson, H. A. und Leslie, R. A. (1985): Noradrenergic alpha 2 binding sites in vagal dorsal motor nucleus and nucleus tractus solitarius: autoradiographic localization, *Can.J.Physiol Pharmacol.* (volume 63), issue 9, pp. 1190-1194. URL:

PM:2996733

- [57] Parlow, J.; Viale, J. P.; Annat, G.; Hughson, R. und Quintin, L. (1995): Spontaneous cardiac baroreflex in humans. Comparison with drug-induced responses, *Hypertension* (volume 25), issue 5, pp. 1058-1068. URL: PM:7737717
- [58] Gross, V.; Plehm, R.; Tank, J.; Jordan, J.; Diedrich, A.; Obst, M. und Luft, F. C. (2002): Heart rate variability and baroreflex function in AT2 receptor-disrupted mice, *Hypertension* (volume 40), issue 2, pp. 207-213. URL: PM:12154115
- [59] Janssen, B. J.; Leenders, P. J. und Smits, J. F. (2000): Short-term and long-term blood pressure and heart rate variability in the mouse, *Am.J.Physiol Regul.Integr.Comp Physiol* (volume 278), issue 1, pp. R215-R225. URL: PM:10644642
- [60] Gehrman, J.; Hammer, P. E.; Maguire, C. T.; Wakimoto, H.; Tiedman, J. K. und Berul, C. I. (2000): Phenotypic screening for heart rate variability in the mouse, *Am.J.Physiol Heart Circ.Physiol* (volume 279), issue 2, pp. H733-H740. URL: PM:10924073
- [61] Brabant, G.; Horn, R.; von-zur-Muhlen, A.; Mayr, B.; Wurster, U.; Heidenreich, F.; Schnabel, D.; Gruters, Kieslich A.; Zimmermann, Belsing T. und Feldt, Rasmussen U. (2000): Free and protein bound leptin are distinct and independently controlled factors in energy regulation, *Diabetologia* (volume 43), issue 4, pp. 438-442.
- [62] Gross, V.; Tank, J.; Obst, M.; Plehm, R.; Blumer, K. J.; Diedrich, A.; Jordan, J. und Luft, F. C. (2005): Autonomic Nervous System and Blood Pressure Regulation in RGS2-deficient Mice, *Am J.Physiol Regul.Integr.Comp Physiol* (volume 288), issue 5, pp. R1134-R1142. URL: PM:15661972
- [63] Lowell, B. B. und Shulman, G. I. (2005): Mitochondrial dysfunction and type 2 diabetes, *Science* (volume 307), issue 5708, pp. 384-387. URL: PM:15662004
- [64] Bierhaus, A.; Haslbeck, K. M.; Humpert, P. M.; Liliensiek, B.; Dehmer, T.; Morcos, M.; Sayed, A. A.; Andrassy, M.; Schiekofer, S.; Schneider, J. G.; Schulz, J. B.; Heuss, D.; Neundorfer, B.; Dierl, S.; Huber, J.; Tritschler, H.; Schmidt, A. M.; Schwaninger, M.; Haering, H. U.; Schleicher, E.; Kasper, M.; Stern, D. M.; Arnold, B. und Nawroth, P. P. (2004): Loss of pain perception in diabetes is dependent on a receptor of the immunoglobulin superfamily, *J.Clin.Invest* (volume 114), issue 12, pp. 1741-1751. URL: PM:15599399
- [65] Rahmouni, K.; Correia, M. L.; Haynes, W. G. und Mark, A. L. (2005): Obesity-associated hypertension: new insights into mechanisms, *Hypertension* (volume 45), issue 1, pp. 9-14. URL: PM:15583075
- [66] Gollasch, M. und Dubrovskaja, G. (2004): Paracrine role for periaortic adipose tissue in the regulation of arterial tone, *Trends Pharmacol.Sci.* (volume 25), issue 12,

pp. 647-653. URL: PM:15530643