

## Literaturverzeichnis

- [1] C. J. Cummings, H. Y. Zoghbi, Trinucleotide repeats: mechanisms and pathophysiology., *Annu Rev Genomics Hum Genet* 1 (2000) 281–328.
- [2] R. L. Margolis, The spinocerebellar ataxias: order emerges from chaos., *Curr Neurol Neurosci Rep* 2 (5) (2002) 447–456.
- [3] A. Michalik, C. V. Broeckhoven, Pathogenesis of polyglutamine disorders: aggregation revisited., *Hum Mol Genet* 12 Spec No 2 (2003) R173–R186.
- [4] P. S. Reddy, D. E. Housman, The complex pathology of trinucleotide repeats., *Curr Opin Cell Biol* 9 (3) (1997) 364–372.
- [5] L. Schöls, P. Bauer, T. Schmidt, T. Schulte, O. Riess, Autosomal dominant cerebellar ataxias: clinical features, genetics, and pathogenesis., *Lancet Neurol* 3 (5) (2004) 291–304.
- [6] G. Stevanin, A. Dürr, A. Brice, Spinocerebellar ataxias caused by polyglutamine expansions., *Adv Exp Med Biol* 516 (2002) 47–77.
- [7] R. Butler, G. P. Bates, Histone deacetylase inhibitors as therapeutics for polyglutamine disorders, *Nature Reviews* 7 (2006) 784–796.
- [8] Y. Trottier, V. Biancalana, J. L. Mandel, Instability of cag repeats in huntington's disease: relation to parental transmission and age of onset., *J Med Genet* 31 (5) (1994) 377–382.

## Literaturverzeichnis

---

- [9] G. Cancel, N. Abbas, G. Stevanin, A. Dürr, H. Chneiweiss, C. Néri, C. Duyckaerts, C. Penet, H. M. Cann, Y. Agid, Marked phenotypic heterogeneity associated with expansion of a cag repeat sequence at the spinocerebellar ataxia 3/machado-joseph disease locus., *Am J Hum Genet* 57 (4) (1995) 809–816.
- [10] S. S. Chong, A. E. McCall, J. Cota, S. H. Subramony, H. T. Orr, M. R. Hughes, H. Y. Zoghbi, Gametic and somatic tissue-specific heterogeneity of the expanded sca1 cag repeat in spinocerebellar ataxia type 1., *Nat Genet* 10 (3) (1995) 344–350.
- [11] R. Truant, L. A. Raymond, J. Xia, D. Pinchev, A. Burtnik, R. S. Atwal, Canadian association of neurosciences review: polyglutamine expansion neurodegenerative diseases., *Can J Neurol Sci* 33 (3) (2006) 278–291.
- [12] J. S. Steffan, A. Kazantsev, O. Spasic-Boskovic, M. Greenwald, Y. Z. Zhu, H. Gohler, E. E. Wanker, G. P. Bates, D. E. Housman, L. M. Thompson, The huntington's disease protein interacts with p53 and creb-binding protein and represses transcription., *Proc Natl Acad Sci U S A* 97 (12) (2000) 6763–6768.
- [13] A. P. Bemelmans, P. Horellou, L. Pradier, I. Brunet, P. Colin, J. Mallet, Brain-derived neurotrophic factor-mediated protection of striatal neurons in an excitotoxic rat model of huntington's disease, as demonstrated by adenoviral gene transfer., *Hum Gene Ther* 10 (18) (1999) 2987–2997.
- [14] E. Cattaneo, D. Rigamonti, C. Zuccato, The enigma of huntington's disease., *Sci Am* 287 (6) (2002) 92–97.
- [15] E. Pérez-Navarro, A. M. Canudas, P. Akerund, J. Alberch, E. Arenas, Brain-derived neurotrophic factor, neurotrophin-3, and neurotrophin-4/5 prevent the death of striatal projection neurons in a rodent model of huntington's disease., *J Neurochem* 75 (5) (2000) 2190–2199.
- [16] C. Zuccato, A. Ciampola, D. Rigamonti, B. R. Leavitt, D. Goffredo, L. Conti, M. E.

- MacDonald, R. M. Friedlander, V. Silani, M. R. Hayden, T. Timmusk, S. Sipione, E. Cattaneo, Loss of huntingtin-mediated bdnf gene transcription in huntington's disease., *Science* 293 (5529) (2001) 493–498.
- [17] A. O. Brinkmann, J. Trapman, Androgen receptor mutants that affect normal growth and development., *Cancer Surv* 14 (1992) 95–111.
- [18] S. Irwin, M. Vandelft, D. Pinchev, J. L. Howell, J. Graczyk, H. T. Orr, R. Truant, Rna association and nucleocytoplasmic shuttling by ataxin-1., *J Cell Sci* 118 (Pt 1) (2005) 233–242.
- [19] S. Yue, H. G. Serra, H. Y. Zoghbi, H. T. Orr, The spinocerebellar ataxia type 1 protein, ataxin-1, has rna-binding activity that is inversely affected by the length of its polyglutamine tract., *Hum Mol Genet* 10 (1) (2001) 25–30.
- [20] D. Helmlinger, S. Hardy, S. Sasorith, F. Klein, F. Robert, C. Weber, L. Miguet, N. Pottier, A. Van-Dorsellaer, J.-M. Wurtz, J.-L. Mandel, L. Tora, D. Devys, Ataxin-7 is a subunit of gcn5 histone acetyltransferase-containing complexes., *Hum Mol Genet* 13 (12) (2004) 1257–1265.
- [21] S. Chen, G.-H. Peng, X. Wang, A. C. Smith, S. K. Grote, B. L. Sopher, A. R. L. Spada, Interference of crx-dependent transcription by ataxin-7 involves interaction between the glutamine regions and requires the ataxin-7 carboxy-terminal region for nuclear localization., *Hum Mol Genet* 13 (1) (2004) 53–67.
- [22] I. A. Klement, P. J. Skinner, M. D. Kaytor, H. Yi, S. M. Hersch, H. B. Clark, H. Y. Zoghbi, H. T. Orr, Ataxin-1 nuclear localization and aggregation: role in polyglutamine-induced disease in sca1 transgenic mice., *Cell* 95 (1) (1998) 41–53.
- [23] V. R. Lingappa, G. Blobel, Early events in the biosynthesis of secretory and membrane proteins: the signal hypothesis., *Recent Prog Horm Res* 36 (1980) 451–475.

- [24] G. Schilling, J. D. Wood, K. Duan, H. H. Slunt, V. Gonzales, M. Yamada, J. K. Cooper, R. L. Margolis, N. A. Jenkins, N. G. Copeland, H. Takahashi, S. Tsuji, D. L. Price, D. R. Borchelt, C. A. Ross, Nuclear accumulation of truncated atrophin-1 fragments in a transgenic mouse model of drpla., *Neuron* 24 (1) (1999) 275–286.
- [25] J. Taylor, S. K. Grote, J. Xia, M. Vandelft, J. Graczyk, L. M. Ellerby, A. R. L. Spada, R. Truant, Ataxin-7 can export from the nucleus via a conserved exportin-dependent signal., *J Biol Chem* 281 (5) (2006) 2730–2739.
- [26] J. Xia, D. H. Lee, J. Taylor, M. Vandelft, R. Truant, Huntingtin contains a highly conserved nuclear export signal., *Hum Mol Genet* 12 (12) (2003) 1393–1403.
- [27] P. Günther, A. Storch, J. Schwarz, O. Sabri, P. Steinbach, A. Wagner, S. Hesse, Basal ganglia involvement of a patient with sca 17—a new form of autosomal dominant spinocerebellar ataxia., *J Neurol* 251 (7) (2004) 896–897.
- [28] J. M. Hagenah, C. Zühlke, Y. Hellenbroich, W. Heide, C. Klein, Focal dystonia as a presenting sign of spinocerebellar ataxia 17., *Mov Disord* 19 (2) (2004) 217–220.
- [29] A. C. Bruni, J. Takahashi-Fujigasaki, F. Maltecca, J. F. Foncin, A. Servadio, G. Casari, P. D'Adamo, R. Maletta, S. A. M. Curcio, G. D. Michele, A. Filli, K. H. E. Hachimi, C. Duyckaerts, Behavioral disorder, dementia, ataxia, and rigidity in a large family with tata box-binding protein mutation., *Arch Neurol* 61 (8) (2004) 1314–1320.
- [30] T. R. Hebbes, A. W. Thorne, C. Crane-Robinson, A direct link between core histone acetylation and transcriptionally active chromatin., *EMBO J* 7 (5) (1988) 1395–1402.
- [31] D. Y. Lee, J. J. Hayes, D. Pruss, A. P. Wolffe, A positive role for histone acetylation in transcription factor access to nucleosomal dna., *Cell* 72 (1) (1993) 73–84.

- [32] V. G. Norton, K. W. Marvin, P. Yau, E. M. Bradbury, Nucleosome linking number change controlled by acetylation of histones h3 and h4., *J Biol Chem* 265 (32) (1990) 19848–19852.
- [33] B. G. Burnett, R. N. Pittman, The polyglutamine neurodegenerative protein ataxin 3 regulates aggresome formation., *Proc Natl Acad Sci U S A* 102 (12) (2005) 4330–4335.
- [34] H. T. McMahon, I. G. Mills, Cop and clathrin-coated vesicle budding: different pathways, common approaches., *Curr Opin Cell Biol* 16 (4) (2004) 379–391.
- [35] J. Rohrbough, K. Broadie, Lipid regulation of the synaptic vesicle cycle., *Nat Rev Neurosci* 6 (2) (2005) 139–150.
- [36] L.-G. Wu, Kinetic regulation of vesicle endocytosis at synapses., *Trends Neurosci* 27 (9) (2004) 548–554.
- [37] M. DiFiglia, E. Sapp, K. Chase, C. Schwarz, A. Meloni, C. Young, E. Martin, J. P. Vonsattel, R. Carraway, S. A. Reeves, Huntingtin is a cytoplasmic protein associated with vesicles in human and rat brain neurons., *Neuron* 14 (5) (1995) 1075–1081.
- [38] K. E. D. Rooij, J. C. Dorsman, M. A. Smoor, J. T. D. Dunnen, G. J. V. Ommen, Subcellular localization of the huntington's disease gene product in cell lines by immunofluorescence and biochemical subcellular fractionation., *Hum Mol Genet* 5 (8) (1996) 1093–1099.
- [39] A. Pal, F. Severin, B. Lommer, A. Shevchenko, M. Zerial, Huntington-hap40 complex is a novel rab5 effector that regulates early endosome motility and is up-regulated in huntington's disease., *J Cell Biol* 172 (4) (2006) 605–618.
- [40] I. Bezprozvanny, M. R. Hayden, Deranged neuronal calcium signaling and huntington disease., *Biochem Biophys Res Commun* 322 (4) (2004) 1310–1317.

## Literaturverzeichnis

---

- [41] A. V. Panov, S. Lund, J. T. Greenamyre, Ca<sup>2+</sup>-induced permeability transition in human lymphoblastoid cell mitochondria from normal and huntington's disease individuals., *Mol Cell Biochem* 269 (1-2) (2005) 143–152.
- [42] T.-S. Tang, E. Slow, V. Lupu, I. G. Stavrovskaya, M. Sugimori, R. Llinás, B. S. Kristal, M. R. Hayden, I. Bezprozvanny, Disturbed ca<sup>2+</sup> signaling and apoptosis of medium spiny neurons in huntington's disease., *Proc Natl Acad Sci U S A* 102 (7) (2005) 2602–2607.
- [43] O. Zhuchenko, J. Bailey, P. Bonnen, T. Ashizawa, D. W. Stockton, C. Amos, W. B. Dobyns, S. H. Subramony, H. Y. Zoghbi, C. C. Lee, Autosomal dominant cerebellar ataxia (sca6) associated with small polyglutamine expansions in the alpha 1a-voltage-dependent calcium channel., *Nat Genet* 15 (1) (1997) 62–69.
- [44] D. P. Huynh, M. R. D. Bigio, D. H. Ho, S. M. Pulst, Expression of ataxin-2 in brains from normal individuals and patients with alzheimer's disease and spinocerebellar ataxia 2., *Ann Neurol* 45 (2) (1999) 232–241.
- [45] S. Koyano, T. Uchihara, H. Fujigasaki, A. Nakamura, S. Yagishita, K. Iwabuchi, Neuronal intranuclear inclusions in spinocerebellar ataxia type 2: triple-labeling immunofluorescent study., *Neurosci Lett* 273 (2) (1999) 117–120.
- [46] M. L. Moseley, K. A. Benzow, L. J. Schut, T. D. Bird, C. M. Gomez, P. E. Barkhaus, K. A. Blindauer, M. Labuda, M. Pandolfo, M. D. Koob, L. P. Ranum, Incidence of dominant spinocerebellar and friedreich triplet repeats among 361 ataxia families., *Neurology* 51 (6) (1998) 1666–1671.
- [47] A. Fillia, C. Mariotti, G. Caruso, G. Coppola, S. Cocozza, I. Castaldo, O. Calabrese, E. Salvatore, G. D. Michele, M. C. Riggio, D. Pareyson, C. Gellera, S. D. Donato, Relative frequencies of cag expansions in spinocerebellar ataxia and dentatorubro-pallidoluysian atrophy in 116 italian families., *Eur Neurol* 44 (1) (2000) 31–36.

- [48] A. Fillà, G. D. Michele, L. Santoro, O. Calabrese, I. Castaldo, S. Giuffrida, D. Restivo, L. Serlenga, D. F. Condorelli, U. Bonuccelli, R. Scala, G. Coppola, G. Caruso, S. Cocozza, Spinocerebellar ataxia type 2 in southern Italy: a clinical and molecular study of 30 families., *J Neurol* 246 (6) (1999) 467–471.
- [49] P. Giunti, G. Sabbadini, M. G. Sweeney, M. B. Davis, L. Veneziano, E. Mantuano, A. Federico, R. Plasmati, M. Frontali, N. W. Wood, The role of the *sca2* trinucleotide repeat expansion in 89 autosomal dominant cerebellar ataxia families. frequency, clinical and genetic correlates., *Brain* 121 (Pt 3) (1998) 459–467.
- [50] O. Riess, F. A. Laccone, S. Gispert, L. Schöls, C. Zühlke, A. M. Vieira-Saecker, S. Herlt, K. Wessel, J. T. Epplen, B. H. Weber, F. Kreuz, S. Chahrokh-Zadeh, A. Meindl, A. Lunkes, J. Aguiar, M. Macek, A. Krebsová, M. Macek, K. Bürk, S. Tinschert, I. Schreyer, S. M. Pulst, G. Auburger, Sca2 trinucleotide expansion in german *sca* patients., *Neurogenetics* 1 (1) (1997) 59–64.
- [51] A. H. Koeppen, The hereditary ataxias., *J Neuropathol Exp Neurol* 57 (6) (1998) 531–543.
- [52] A. H. Koeppen, The pathogenesis of spinocerebellar ataxia., *Cerebellum* 4 (1) (2005) 62–73.
- [53] A. H. Koeppen, The purkinje cell and its afferents in human hereditary ataxia., *J Neuropathol Exp Neurol* 50 (4) (1991) 505–514.
- [54] K. Araki, H. Meguro, E. Kushiya, C. Takayama, Y. Inoue, M. Mishina, Selective expression of the glutamate receptor channel delta 2 subunit in cerebellar purkinje cells., *Biochem Biophys Res Commun* 197 (3) (1993) 1267–1276.
- [55] N. Kashiwabuchi, K. Ikeda, K. Araki, T. Hirano, K. Shibuki, C. Takayama, Y. Inoue, T. Kutsuwada, T. Yagi, Y. Kang, Impairment of motor coordination, purkinje cell syn-

- apse formation, and cerebellar long-term depression in glur delta 2 mutant mice., Cell 81 (2) (1995) 245–252.
- [56] H. Mori, M. Mishina, Roles of diverse glutamate receptors in brain functions elucidated by subunit-specific and region-specific gene targeting., Life Sci 74 (2-3) (2003) 329–336.
- [57] T. Hirano, Cerebellar regulation mechanisms learned from studies on glurdelta2., Mol Neurobiol 33 (1) (2006) 1–16.
- [58] G. Imbert, F. Saudou, G. Yvert, D. Devys, Y. Trottier, J. M. Garnier, C. Weber, J. L. Mandel, G. Cancel, N. Abbas, A. Dürr, O. Didierjean, G. Stevanin, Y. Agid, A. Brice, Cloning of the gene for spinocerebellar ataxia 2 reveals a locus with high sensitivity to expanded cag/glutamine repeats., Nat Genet 14 (3) (1996) 285–291.
- [59] S. M. Pulst, A. Nechiporuk, T. Nechiporuk, S. Gispert, X. N. Chen, I. Lopes-Cendes, S. Pearlman, S. Starkman, G. Orozco-Diaz, A. Lunkes, P. DeJong, G. A. Rouleau, G. Auburger, J. R. Korenberg, C. Figueroa, S. Sahba, Moderate expansion of a normally biallelic trinucleotide repeat in spinocerebellar ataxia type 2., Nat Genet 14 (3) (1996) 269–276.
- [60] K. Sanpei, H. Takano, S. Igarashi, T. Sato, M. Oyake, H. Sasaki, A. Wakisaka, K. Tashiro, Y. Ishida, T. Ikeuchi, R. Koide, M. Saito, A. Sato, T. Tanaka, S. Hanyu, Y. Takiyama, M. Nishizawa, N. Shimizu, Y. Nomura, M. Segawa, K. Iwabuchi, I. Eguuchi, H. Tanaka, H. Takahashi, S. Tsuji, Identification of the spinocerebellar ataxia type 2 gene using a direct identification of repeat expansion and cloning technique, direct., Nat Genet 14 (3) (1996) 277–284.
- [61] S. Sahba, A. Nechiporuk, K. P. Figueroa, T. Nechiporuk, S. M. Pulst, Genomic structure of the human gene for spinocerebellar ataxia type 2 (sca2) on chromosome 12q24.1., Genomics 47 (3) (1998) 359–364.

- [62] S. Choudhry, M. Mukerji, A. K. Srivastava, S. Jain, S. K. Brahmachari, Cag repeat instability at sca2 locus: anchoring caa interruptions and linked single nucleotide polymorphisms., *Hum Mol Genet* 10 (21) (2001) 2437–2446.
- [63] R. Wiedemeyer, F. Westermann, I. Wittke, J. Nowock, M. Schwab, Ataxin-2 promotes apoptosis of human neuroblastoma cells., *Oncogene* 22 (3) (2003) 401–411.
- [64] D. P. Huynh, H.-T. Yang, H. Vakharia, D. Nguyen, S. M. Pulst, Expansion of the polyq repeat in ataxin-2 alters its golgi localization, disrupts the golgi complex and causes cell death., *Hum Mol Genet* 12 (13) (2003) 1485–1496.
- [65] D. P. Huynh, K. Figueroa, N. Hoang, S. M. Pulst, Nuclear localization or inclusion body formation of ataxin-2 are not necessary for sca2 pathogenesis in mouse or human., *Nat Genet* 26 (1) (2000) 44–50.
- [66] J. Aguiar, J. Fernández, A. Aguilar, Y. Mendoza, M. Vázquez, J. Suárez, J. Berlanga, S. Cruz, G. Guillén, L. Herrera, L. Velázquez, N. Santos, N. Merino, Ubiquitous expression of human sca2 gene under the regulation of the sca2 self promoter cause specific purkinje cell degeneration in transgenic mice., *Neurosci Lett* 392 (3) (2006) 202–206.
- [67] T.-R. Kiehl, A. Nechiporuk, K. P. Figueroa, M. T. Keating, D. P. Huynh, S.-M. Pulst, Generation and characterization of sca2 (ataxin-2) knockout mice., *Biochem Biophys Res Commun* 339 (1) (2006) 17–24.
- [68] M. Waldherr, A. Ragnini, B. Jank, R. Teply, G. Wiesenberger, R. J. Schweyen, A multitude of suppressors of group ii intron-splicing defects in yeast., *Curr Genet* 24 (4) (1993) 301–306.
- [69] M. Ralser, M. Albrecht, U. Nonhoff, T. Lengauer, H. Lehrach, S. Krobitsch, An integrative approach to gain insights into the cellular function of human ataxin-2., *J Mol Biol* 346 (1) (2005) 203–214.

## Literaturverzeichnis

---

- [70] D. A. Mangus, N. Amrani, A. Jacobson, Pbp1p, a factor interacting with *saccharomyces cerevisiae* poly(a)-binding protein, regulates polyadenylation., *Mol Cell Biol* 18 (12) (1998) 7383–7396.
- [71] D. A. Mangus, M. C. Evans, A. Jacobson, Poly(a)-binding proteins: multifunctional scaffolds for the post-transcriptional control of gene expression., *Genome Biol* 4 (7) (2003) 223.
- [72] D. A. Mangus, M. M. Smith, J. M. McSweeney, A. Jacobson, Identification of factors regulating poly(a) tail synthesis and maturation., *Mol Cell Biol* 24 (10) (2004) 4196–4206.
- [73] D. A. Mangus, M. C. Evans, N. S. Agrin, M. Smith, P. Gongidi, A. Jacobson, Positive and negative regulation of poly(a) nuclease., *Mol Cell Biol* 24 (12) (2004) 5521–5533.
- [74] J. M. Coller, M. Tucker, U. Sheth, M. A. Valencia-Sanchez, R. Parker, The dead box helicase, dhh1p, functions in mrna decapping and interacts with both the decapping and deadenylase complexes., *RNA* 7 (12) (2001) 1717–1727.
- [75] N. Fischer, K. Weis, The dead box protein dhh1 stimulates the decapping enzyme dcp1., *EMBO J* 21 (11) (2002) 2788–2797.
- [76] S. S.-I. Tseng-Rogenski, J.-L. Chong, C. B. Thomas, S. Enomoto, J. Berman, T.-H. Chang, Functional conservation of dhh1p, a cytoplasmic dexd/h-box protein present in large complexes., *Nucleic Acids Res* 31 (17) (2003) 4995–5002.
- [77] U. Sheth, R. Parker, Targeting of aberrant mrnas to cytoplasmic processing bodies., *Cell* 125 (6) (2006) 1095–1109.
- [78] C. Mazzoni, P. Mancini, L. Verdone, F. Madeo, A. Serafini, E. Herker, C. Falcone, A truncated form of kllsm4p and the absence of factors involved in mrna decapping trigger apoptosis in yeast., *Mol Biol Cell* 14 (2) (2003) 721–729.

- [79] W. He, R. Parker, Functions of lsm proteins in mrna degradation and splicing., *Curr Opin Cell Biol* 12 (3) (2000) 346–350.
- [80] J. D. Beggs, Lsm proteins and rna processing., *Biochem Soc Trans* 33 (Pt 3) (2005) 433–438.
- [81] T. F. Satterfield, S. M. Jackson, L. J. Pallanck, A drosophila homolog of the polyglutamine disease gene sca2 is a dosage-sensitive regulator of actin filament formation., *Genetics* 162 (4) (2002) 1687–1702.
- [82] T. F. Satterfield, L. J. Pallanck, Ataxin-2 and its drosophila homolog, atx2, physically assemble with polyribosomes., *Hum Mol Genet* 15 (16) (2006) 2523–2532.
- [83] R. Ciosk, M. DePalma, J. R. Priess, Atx-2, the *c. elegans* ortholog of ataxin 2, functions in translational regulation in the germline., *Development* 131 (19) (2004) 4831–4841.
- [84] E. M. Maine, D. Hansen, D. Springer, V. E. Vought, *Caenorhabditis elegans* atx-2 promotes germline proliferation and the oocyte fate., *Genetics* 168 (2) (2004) 817–830.
- [85] K. Sanpei, [the function of spinocerebellar ataxia type 2 (sca2) gene product, ataxin-2 and the mechanism of pathogenesis for sca2], *Nippon Rinsho* 57 (4) (1999) 822–824.
- [86] M. Albrecht, T. Lengauer, Novel sm-like proteins with long c-terminal tails and associated methyltransferases., *FEBS Lett* 569 (1-3) (2004) 18–26.
- [87] G. Kozlov, G. D. Crescenzo, N. S. Lim, N. Siddiqui, D. Fantus, A. Kahvejian, J.-F. Trempe, D. Elias, I. Ekiel, N. Sonenberg, M. O'Connor-McCourt, K. Gehring, Structural basis of ligand recognition by pabc, a highly specific peptide-binding domain found in poly(a)-binding protein and a hect ubiquitin ligase., *EMBO J* 23 (2) (2004) 272–281.

## Literaturverzeichnis

---

- [88] G. Kozlov, J. F. Trempe, K. Khaleghpour, A. Kahvejian, I. Ekiel, K. Gehring, Structure and function of the c-terminal pabc domain of human poly(a)-binding protein., Proc Natl Acad Sci U S A 98 (8) (2001) 4409–4413.
- [89] H. Shibata, D. P. Huynh, S. M. Pulst, A novel protein with rna-binding motifs interacts with ataxin-2., Hum Mol Genet 9 (9) (2000) 1303–1313.
- [90] C. A. Beelman, R. Parker, Degradation of mrna in eukaryotes., Cell 81 (2) (1995) 179–183.
- [91] P. Bernstein, J. Ross, Poly(a), poly(a) binding protein and the regulation of mrna stability., Trends Biochem Sci 14 (9) (1989) 373–377.
- [92] C. J. Decker, R. Parker, Mechanisms of mrna degradation in eukaryotes., Trends Biochem Sci 19 (8) (1994) 336–340.
- [93] S. W. Peltz, G. Brewer, P. Bernstein, P. A. Hart, J. Ross, Regulation of mrna turnover in eukaryotic cells., Crit Rev Eukaryot Gene Expr 1 (2) (1991) 99–126.
- [94] J. S. Butler, The yin and yang of the exosome., Trends Cell Biol 12 (2) (2002) 90–96.
- [95] E. van Dijk, N. Cougot, S. Meyer, S. Babajko, E. Wahle, B. Séraphin, Human dcp2: a catalytically active mrna decapping enzyme located in specific cytoplasmic structures., EMBO J 21 (24) (2002) 6915–6924.
- [96] N. Cougot, S. Babajko, B. Séraphin, Cytoplasmic foci are sites of mrna decay in human cells., J Cell Biol 165 (1) (2004) 31–40.
- [97] E. Bouveret, G. Rigaut, A. Shevchenko, M. Wilm, B. Séraphin, A sm-like protein complex that participates in mrna degradation., EMBO J 19 (7) (2000) 1661–1671.
- [98] T. Eystathioy, E. K. L. Chan, S. A. Tenenbaum, J. D. Keene, K. Griffith, M. J. Fritzler, A phosphorylated cytoplasmic autoantigen, gw182, associates with a unique

- population of human mrnas within novel cytoplasmic speckles., Mol Biol Cell 13 (4) (2002) 1338–1351.
- [99] D. Ingelfinger, D. J. Arndt-Jovin, R. Lührmann, T. Achsel, The human lsm1-7 proteins colocalize with the mRNA-degrading enzymes dcp1/2 and xrn1 in distinct cytoplasmic foci., RNA 8 (12) (2002) 1489–1501.
- [100] J. Lykke-Andersen, Identification of a human decapping complex associated with hupf proteins in nonsense-mediated decay., Mol Cell Biol 22 (23) (2002) 8114–8121.
- [101] J. H. Yu, W.-H. Yang, T. Gulick, K. D. Bloch, D. B. Bloch, Ge-1 is a central component of the mammalian cytoplasmic mRNA processing body., RNA 11 (12) (2005) 1795–1802.
- [102] M. Fenger-Grøn, C. Fillman, B. Norrild, J. Lykke-Andersen, Multiple processing body factors and the ARE binding protein ttp activate mRNA decapping., Mol Cell 20 (6) (2005) 905–915.
- [103] J. Liu, F. V. Rivas, J. Wohlschlegel, J. R. Yates, R. Parker, G. J. Hannon, A role for the P-body component gw182 in microRNA function., Nat Cell Biol 7 (12) (2005) 1261–1266.
- [104] G. L. Sen, H. M. Blau, Argonaute 2/risc resides in sites of mammalian mRNA decay known as cytoplasmic bodies., Nat Cell Biol 7 (6) (2005) 633–636.
- [105] W.-H. Yang, J. H. Yu, T. Gulick, K. D. Bloch, D. B. Bloch, RNA-associated protein 55 (rap55) localizes to mRNA processing bodies and stress granules., RNA 12 (4) (2006) 547–554.
- [106] D. Teixeira, U. Sheth, M. A. Valencia-Sánchez, M. Brengues, R. Parker, Processing bodies require RNA for assembly and contain nontranslating mRNAs., RNA 11 (4) (2005) 371–382.

## Literaturverzeichnis

---

- [107] Z. Yang, A. Jakymiw, M. R. Wood, T. Eystathioy, R. L. Rubin, M. J. Fritzler, E. K. L. Chan, Gw182 is critical for the stability of gw bodies expressed during the cell cycle and cell proliferation., *J Cell Sci* 117 (Pt 23) (2004) 5567–5578.
- [108] A. Eulalio, I. Behm-Ansmant, E. Izaurralde, P bodies: at the crossroads of post-transcriptional pathways., *Nat Rev Mol Cell Biol* 8 (1) (2007) 9–22.
- [109] R. Parker, U. Sheth, P bodies and the control of mrna translation and degradation., *Mol Cell* 25 (5) (2007) 635–646.
- [110] M. Brengues, D. Teixeira, R. Parker, Movement of eukaryotic mrnas between poly-somes and cytoplasmic processing bodies., *Science* 310 (5747) (2005) 486–489.
- [111] P. Anderson, N. Kedersha, Rna granules., *J Cell Biol* 172 (6) (2006) 803–808.
- [112] N. Kedersha, M. R. Cho, W. Li, P. W. Yacono, S. Chen, N. Gilks, D. E. Golan, P. Anderson, Dynamic shuttling of tia-1 accompanies the recruitment of mrna to mammalian stress granules., *J Cell Biol* 151 (6) (2000) 1257–1268.
- [113] L. Nover, K. D. Scharf, D. Neumann, Cytoplasmic heat shock granules are formed from precursor particles and are associated with a specific set of mrnas., *Mol Cell Biol* 9 (3) (1989) 1298–1308.
- [114] L. Nover, K. D. Scharf, D. Neumann, Formation of cytoplasmic heat shock granules in tomato cell cultures and leaves., *Mol Cell Biol* 3 (9) (1983) 1648–1655.
- [115] N. C. Collier, M. J. Schlesinger, The dynamic state of heat shock proteins in chicken embryo fibroblasts., *J Cell Biol* 103 (4) (1986) 1495–1507.
- [116] N. C. Collier, J. Heuser, M. A. Levy, M. J. Schlesinger, Ultrastructural and biochemical analysis of the stress granule in chicken embryo fibroblasts., *J Cell Biol* 106 (4) (1988) 1131–1139.

- [117] A. P. Arrigo, J. P. Suhan, W. J. Welch, Dynamic changes in the structure and intracellular locale of the mammalian low-molecular-weight heat shock protein., *Mol Cell Biol* 8 (12) (1988) 5059–5071.
- [118] N. L. Kedersha, M. Gupta, W. Li, I. Miller, P. Anderson, Rna-binding proteins tia-1 and tiar link the phosphorylation of eif-2 alpha to the assembly of mammalian stress granules., *J Cell Biol* 147 (7) (1999) 1431–1442.
- [119] D. A. Mangiardi, K. McLaughlin-Williamson, K. E. May, E. P. Messana, D. C. Mountain, D. A. Cotanche, Progression of hair cell ejection and molecular markers of apoptosis in the avian cochlea following gentamicin treatment., *J Comp Neurol* 475 (1) (2004) 1–18.
- [120] B. J. Moeller, Y. Cao, C. Y. Li, M. W. Dewhirst, Radiation activates hif-1 to regulate vascular radiosensitivity in tumors: role of reoxygenation, free radicals, and stress granules., *Cancer Cell* 5 (5) (2004) 429–441.
- [121] R. I. Morimoto, Regulation of the heat shock transcriptional response: cross talk between a family of heat shock factors, molecular chaperones, and negative regulators., *Genes Dev* 12 (24) (1998) 3788–3796.
- [122] R. I. Morimoto, K. D. Sarge, K. Abravaya, Transcriptional regulation of heat shock genes. a paradigm for inducible genomic responses., *J Biol Chem* 267 (31) (1992) 21987–21990.
- [123] N. Kedersha, P. Anderson, Stress granules: sites of mrna triage that regulate mrna stability and translatability., *Biochem Soc Trans* 30 (Pt 6) (2002) 963–969.
- [124] K. Asano, J. Clayton, A. Shalev, A. G. Hinnebusch, A multifactor complex of eukaryotic initiation factors, eif1, eif2, eif3, eif5, and initiator trna(met) is an important translation initiation intermediate in vivo., *Genes Dev* 14 (19) (2000) 2534–2546.

## Literaturverzeichnis

---

- [125] T. E. Dever, Translation initiation: adept at adapting., *Trends Biochem Sci* 24 (10) (1999) 398–403.
- [126] T. V. Pestova, C. U. Hellen, Ribosome recruitment and scanning: what's new?, *Trends Biochem Sci* 24 (3) (1999) 85–87.
- [127] L. Phan, L. W. Schoenfeld, L. Valásek, K. H. Nielsen, A. G. Hinnebusch, A subcomplex of three eif3 subunits binds eif1 and eif5 and stimulates ribosome binding of mrna and trna(i)met., *EMBO J* 20 (11) (2001) 2954–2965.
- [128] P. Anderson, N. Kedersha, Stressful initiations., *J Cell Sci* 115 (Pt 16) (2002) 3227–3234.
- [129] S. R. Kimball, M. J. Clemens, V. J. Tilleray, R. C. Wek, R. L. Horetsky, L. S. Jefferson, The double-stranded rna-activated protein kinase pkr is dispensable for regulation of translation initiation in response to either calcium mobilization from the endoplasmic reticulum or essential amino acid starvation., *Biochem Biophys Res Commun* 280 (1) (2001) 293–300.
- [130] B. R. Williams, Signal integration via pkr., *Sci STKE* 2001 (89) (2001) RE2.
- [131] P. Anderson, N. Kedersha, Visibly stressed: the role of eif2, tia-1, and stress granules in protein translation., *Cell Stress Chaperones* 7 (2) (2002) 213–221.
- [132] Y. Dang, N. Kedersha, W.-K. Low, D. Romo, M. Gorospe, R. Kaufman, P. Anderson, J. O. Liu, Eukaryotic initiation factor 2alpha-independent pathway of stress granule induction by the natural product pateamine a., *J Biol Chem* 281 (43) (2006) 32870–32878.
- [133] R. Mazroui, R. Sukarieh, M.-E. Bordeleau, R. J. Kaufman, P. Northcote, J. Tanaka, I. Gallouzi, J. Pelletier, Inhibition of ribosome recruitment induces stress granule formation independently of eukaryotic initiation factor 2alpha phosphorylation., *Mol Biol Cell* 17 (10) (2006) 4212–4219.

- [134] N. Kedersha, G. Stoecklin, M. Ayodele, P. Yacono, J. Lykke-Andersen, M. J. Fritzler, D. Scheuner, R. J. Kaufman, D. E. Golan, P. Anderson, Stress granules and processing bodies are dynamically linked sites of mrnp remodeling., *J Cell Biol* 169 (6) (2005) 871–884.
- [135] A. Wilczynska, C. Aigueperse, M. Kress, F. Dautry, D. Weil, The translational regulator cpeb1 provides a link between dcp1 bodies and stress granules., *J Cell Sci* 118 (Pt 5) (2005) 981–992.
- [136] J. Coller, R. Parker, General translational repression by activators of mrna decapping., *Cell* 122 (6) (2005) 875–886.
- [137] M. Ralser, U. Nonhoff, M. Albrecht, T. Lengauer, E. E. Wanker, H. Lehrach, S. Krobitsch, Ataxin-2 and huntingtin interact with endophilin-a complexes to function in plastin-associated pathways., *Hum Mol Genet* 14 (19) (2005) 2893–2909.
- [138] U. Nonhoff, M. Ralser, F. Welzel, I. Piccini, D. Balzereit, M.-L. Yaspo, H. Lehrach, S. Krobitsch, Ataxin-2 interacts with the dead/h-box rna helicase ddx6 and interferes with p-bodies and stress granules., *Mol Biol Cell* 18 (4) (2007) 1385–1396.
- [139] M. M. Bradford, A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding., *Anal Biochem* 72 (1976) 248–254.
- [140] U. K. Laemmli, Cleavage of structural proteins during the assembly of the head of bacteriophage t4., *Nature* 227 (5259) (1970) 680–685.
- [141] H. Towbin, T. Staehelin, J. Gordon, Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications., *Proc Natl Acad Sci U S A* 76 (9) (1979) 4350–4354.
- [142] A. Henschel, F. Buchholz, B. Habermann, Deqr: a web-based tool for the design

- and quality control of sirnas., Nucleic Acids Res 32 (Web Server issue) (2004) W113–W120.
- [143] N. Gilks, N. Kedersha, M. Ayodele, L. Shen, G. Stoecklin, L. M. Dember, P. Anderson, Stress granule assembly is mediated by prion-like aggregation of tia-1., Mol Biol Cell 15 (12) (2004) 5383–5398.
- [144] M. G. Thomas, L. J. M. Tosar, M. Loschi, J. M. Pasquini, J. Correale, S. Kindler, G. L. Boccaccio, Staufen recruitment into stress granules does not affect early mrna transport in oligodendrocytes., Mol Biol Cell 16 (1) (2005) 405–420.
- [145] A. Ménoret, D. Chaillot, M. Callahan, C. Jacquin, Hsp70, an immunological actor playing with the intracellular self under oxidative stress., Int J Hyperthermia 18 (6) (2002) 490–505.
- [146] G. J. Hannon, Rna interference., Nature 418 (6894) (2002) 244–251.
- [147] A. M. Denli, G. J. Hannon, Rnai: an ever-growing puzzle., Trends Biochem Sci 28 (4) (2003) 196–201.
- [148] V. Delanote, J. Vandekerckhove, J. Gettemans, Plastins: versatile modulators of actin organization in (patho)physiological cellular processes., Acta Pharmacol Sin 26 (7) (2005) 769–779.
- [149] V. Delanote, K. V. Impe, V. D. Corte, E. Bruyneel, G. Vetter, C. Boucherie, M. Maereel, J. Vandekerckhove, E. Friederich, J. Gettemans, Molecular basis for dissimilar nuclear trafficking of the actin-bundling protein isoforms t- and l-plastin., Traffic 6 (4) (2005) 335–345.
- [150] B. Blagoev, I. Kratchmarova, S.-E. Ong, M. Nielsen, L. J. Foster, M. Mann, A proteomics strategy to elucidate functional protein-protein interactions applied to egf signaling., Nat Biotechnol 21 (3) (2003) 315–318.

- [151] Y. Ho, A. Gruhler, A. Heilbut, G. D. Bader, L. Moore, S.-L. Adams, A. Millar, P. Taylor, K. Bennett, K. Boutilier, L. Yang, C. Wolting, I. Donaldson, S. Schandorff, J. Shewnarane, M. Vo, J. Taggart, M. Goudreault, B. Muskat, C. Alfarano, D. Dewar, Z. Lin, K. Michalickova, A. R. Willem, H. Sassi, P. A. Nielsen, K. J. Rasmussen, J. R. Andersen, L. E. Johansen, L. H. Hansen, H. Jespersen, A. Podtelejnikov, E. Nielsen, J. Crawford, V. Poulsen, B. D. Sørensen, J. Matthiesen, R. C. Hendrickson, F. Gleeson, T. Pawson, M. F. Moran, D. Durocher, M. Mann, C. W. V. Hogue, D. Figeys, M. Tyers, Systematic identification of protein complexes in *saccharomyces cerevisiae* by mass spectrometry., *Nature* 415 (6868) (2002) 180–183.
- [152] T. Tadauchi, T. Inada, K. Matsumoto, K. Irie, Posttranscriptional regulation of ho expression by the mkt1-pbp1 complex., *Mol Cell Biol* 24 (9) (2004) 3670–3681.
- [153] C. G. Burd, G. Dreyfuss, Conserved structures and diversity of functions of rna-binding proteins., *Science* 265 (5172) (1994) 615–621.
- [154] R. C. Deo, J. B. Bonanno, N. Sonenberg, S. K. Burley, Recognition of polyadenylate rna by the poly(a)-binding protein., *Cell* 98 (6) (1999) 835–845.
- [155] U. Kühn, T. Pieler, Xenopus poly(a) binding protein: functional domains in rna binding and protein-protein interaction., *J Mol Biol* 256 (1) (1996) 20–30.
- [156] R. C. Deo, N. Sonenberg, S. K. Burley, X-ray structure of the human hyperplastic discs protein: an ortholog of the c-terminal domain of poly(a)-binding protein., *Proc Natl Acad Sci U S A* 98 (8) (2001) 4414–4419.
- [157] J. M. Huibregtse, M. Scheffner, S. Beaudenon, P. M. Howley, A family of proteins structurally and functionally related to the e6-ap ubiquitin-protein ligase., *Proc Natl Acad Sci U S A* 92 (7) (1995) 2563–2567.
- [158] M. J. Callaghan, A. J. Russell, E. Woollatt, G. R. Sutherland, R. L. Sutherland, C. K.

## Literaturverzeichnis

---

- Watts, Identification of a human hect family protein with homology to the drosophila tumor suppressor gene hyperplastic discs., *Oncogene* 17 (26) (1998) 3479–3491.
- [159] R. Oughtred, N. Bedard, O. A. J. Adegoke, C. R. Morales, J. Trasler, V. Rajapurohitam, S. S. Wing, Characterization of rat100, a 300-kilodalton ubiquitin-protein ligase induced in germ cells of the rat testis and similar to the drosophila hyperplastic discs gene., *Endocrinology* 143 (10) (2002) 3740–3747.
- [160] A. B. Sachs, R. W. Davis, The poly(a) binding protein is required for poly(a) shortening and 60s ribosomal subunit-dependent translation initiation., *Cell* 58 (5) (1989) 857–867.
- [161] N. Amrani, M. Minet, M. L. Gouar, F. Lacroute, F. Wyers, Yeast pab1 interacts with rna15 and participates in the control of the poly(a) tail length in vitro., *Mol Cell Biol* 17 (7) (1997) 3694–3701.
- [162] S. Z. Tarun, A. B. Sachs, Association of the yeast poly(a) tail binding protein with translation initiation factor eif-4g., *EMBO J* 15 (24) (1996) 7168–7177.
- [163] S. Z. Tarun, A. B. Sachs, Binding of eukaryotic translation initiation factor 4e (eif4e) to eif4g represses translation of uncapped mrna., *Mol Cell Biol* 17 (12) (1997) 6876–6886.
- [164] A. Kahvejian, Y. V. Svitkin, R. Sukarieh, M.-N. M'Boutchou, N. Sonenberg, Mammalian poly(a)-binding protein is a eukaryotic translation initiation factor, which acts via multiple mechanisms., *Genes Dev* 19 (1) (2005) 104–113.
- [165] M. Ptushkina, T. von der Haar, S. Vasilescu, R. Frank, R. Birkenhäger, J. E. McCarthy, Cooperative modulation by eif4g of eif4e-binding to the mrna 5' cap in yeast involves a site partially shared by p20., *EMBO J* 17 (16) (1998) 4798–4808.
- [166] A. Haghigat, N. Sonenberg, eif4g dramatically enhances the binding of eif4e to the mrna 5'-cap structure., *J Biol Chem* 272 (35) (1997) 21677–21680.

- [167] O. Cordin, J. Banroques, N. K. Tanner, P. Linder, The dead-box protein family of rna helicases., *Gene* 367 (2006) 17–37.
- [168] N. K. Tanner, O. Cordin, J. Banroques, M. Doère, P. Linder, The q motif: a newly identified motif in dead box helicases may regulate atp binding and hydrolysis., *Mol Cell* 11 (1) (2003) 127–138.
- [169] N. K. Tanner, P. Linder, Dexd/h box rna helicases: from generic motors to specific dissociation functions., *Mol Cell* 8 (2) (2001) 251–262.
- [170] J. M. Caruthers, D. B. McKay, Helicase structure and mechanism., *Curr Opin Struct Biol* 12 (1) (2002) 123–133.
- [171] H. Hata, H. Mitsui, H. Liu, Y. Bai, C. L. Denis, Y. Shimizu, A. Sakai, Dhh1p, a putative rna helicase, associates with the general transcription factors pop2p and ccr4p from *saccharomyces cerevisiae*., *Genetics* 148 (2) (1998) 571–579.
- [172] R. E. Navarro, T. K. Blackwell, Requirement for p granules and meiosis for accumulation of the germline rna helicase cgh-1., *Genesis* 42 (3) (2005) 172–180.
- [173] D. A. Smillie, J. Sommerville, Rna helicase p54 (ddx6) is a shuttling protein involved in nuclear assembly of stored mrnp particles., *J Cell Sci* 115 (Pt 2) (2002) 395–407.
- [174] M. Ladomery, E. Wade, J. Sommerville, Xp54, the *xenopus* homologue of human rna helicase p54, is an integral component of stored mrnp particles in oocytes., *Nucleic Acids Res* 25 (5) (1997) 965–973.
- [175] A. Nakamura, R. Amikura, K. Hanyu, S. Kobayashi, Me31b silences translation of oocyte-localizing rnas through the formation of cytoplasmic rnp complex during *drosophila* oogenesis., *Development* 128 (17) (2001) 3233–3242.
- [176] N. Minshall, G. Thom, N. Standart, A conserved role of a dead box helicase in mrna masking., *RNA* 7 (12) (2001) 1728–1742.

## Literaturverzeichnis

---

- [177] J. J. Cotto, R. I. Morimoto, Stress-induced activation of the heat-shock response: cell and molecular biology of heat-shock factors., *Biochem Soc Symp* 64 (1999) 105–118.
- [178] L. Sistonen, K. D. Sarge, R. I. Morimoto, Human heat shock factors 1 and 2 are differentially activated and can synergistically induce hsp70 gene transcription., *Mol Cell Biol* 14 (3) (1994) 2087–2099.
- [179] G. Carrard, A.-L. Bulteau, I. Petropoulos, B. Friguet, Impairment of proteasome structure and function in aging., *Int J Biochem Cell Biol* 34 (11) (2002) 1461–1474.
- [180] J. Emerit, M. Edeas, F. Bricaire, Neurodegenerative diseases and oxidative stress., *Biomed Pharmacother* 58 (1) (2004) 39–46.
- [181] I. L. de Silanes, S. Galbán, J. L. Martindale, X. Yang, K. Mazan-Mamczarz, F. E. Indig, G. Falco, M. Zhan, M. Gorospe, Identification and functional outcome of mrnas associated with rna-binding protein tia-1., *Mol Cell Biol* 25 (21) (2005) 9520–9531.
- [182] K. Phillips, N. Kedersha, L. Shen, P. J. Blackshear, P. Anderson, Arthritis suppressor genes tia-1 and ttp dampen the expression of tumor necrosis factor alpha, cyclooxygenase 2, and inflammatory arthritis., *Proc Natl Acad Sci U S A* 101 (7) (2004) 2011–2016.
- [183] P. Anderson, K. Phillips, G. Stoecklin, N. Kedersha, Post-transcriptional regulation of proinflammatory proteins., *J Leukoc Biol* 76 (1) (2004) 42–47.
- [184] S. Ma, T. Musa, J. Bag, Reduced stability of mitogen-activated protein kinase kinase-2 mrna and phosphorylation of poly(a)-binding protein (pabp) in cells over-expressing pabp., *J Biol Chem* 281 (6) (2006) 3145–3156.
- [185] B. Cosson, A. Couturier, S. Chabelskaya, D. Kiktev, S. Inge-Vechtomov, M. Philippe, G. Zhouravleva, Poly(a)-binding protein acts in translation termination via eukaryo-

- tic release factor 3 interaction and does not influence [psi(+)] propagation., Mol Cell Biol 22 (10) (2002) 3301–3315.
- [186] V. A. Tallada, R. R. Daga, C. Palomeque, A. Garzón, J. Jimenez, Genome-wide search of schizosaccharomyces pombe genes causing overexpression-mediated cell cycle defects., Yeast 19 (13) (2002) 1139–1151.
- [187] M. Wormington, A. M. Searfoss, C. A. Hurney, Overexpression of poly(a) binding protein prevents maturation-specific deadenylation and translational inactivation in xenopus oocytes., EMBO J 15 (4) (1996) 900–909.
- [188] M. Verlaet, V. Deregowki, G. Denis, C. Humblet, M. T. Stalmans, V. Bours, V. Castronovo, J. Boniver, M. P. Defresne, Genetic imbalances in preleukemic thymuses., Biochem Biophys Res Commun 283 (1) (2001) 12–18.
- [189] L. Zhang, W. Zhou, V. E. Velculescu, S. E. Kern, R. H. Hruban, S. R. Hamilton, B. Vogelstein, K. W. Kinzler, Gene expression profiles in normal and cancer cells., Science 276 (5316) (1997) 1268–1272.
- [190] M. A. Andrei, D. Ingelfinger, R. Heintzmann, T. Achsel, R. Rivera-Pomar, R. Lührmann, A role for eif4e and eif4e-transporter in targeting mrnps to mammalian processing bodies., RNA 11 (5) (2005) 717–727.
- [191] C. ying Chu, T. M. Rana, Translation repression in human cells by microrna-induced gene silencing requires rck/p54., PLoS Biol 4 (7) (2006) e210.
- [192] K. M. Pauley, T. Eystathioy, A. Jakymiw, J. C. Hamel, M. J. Fritzler, E. K. L. Chan, Formation of gw bodies is a consequence of microrna genesis., EMBO Rep 7 (9) (2006) 904–910.
- [193] T. Eystathioy, E. K. L. Chan, K. Takeuchi, M. Mahler, L. M. Luft, D. W. Zochodne, M. J. Fritzler, Clinical and serological associations of autoantibodies to gw bodies and a novel cytoplasmic autoantigen gw182., J Mol Med 81 (12) (2003) 811–818.

- [194] M. Yoshida, K. Yoshida, G. Kozlov, N. S. Lim, G. D. Crescenzo, Z. Pang, J. J. Berlanga, A. Kahvejian, K. Gehring, S. S. Wing, N. Sonenberg, Poly(a) binding protein (pabp) homeostasis is mediated by the stability of its inhibitor, paip2., *EMBO J* 25 (9) (2006) 1934–1944.
- [195] M. Albrecht, T. Lengauer, Survey on the pabc recognition motif pam2., *Biochem Biophys Res Commun* 316 (1) (2004) 129–138.
- [196] M. M. Karim, Y. V. Svitkin, A. Kahvejian, G. D. Crescenzo, M. Costa-Mattioli, N. Sonenberg, A mechanism of translational repression by competition of paip2 with eif4g for poly(a) binding protein (pabp) binding., *Proc Natl Acad Sci U S A* 103 (25) (2006) 9494–9499.
- [197] J. L. Clancy, M. J. Henderson, A. J. Russell, D. W. Anderson, R. J. Bova, I. G. Campbell, D. Y. H. Choong, G. A. Macdonald, G. J. Mann, T. Nolan, G. Brady, O. I. Olopade, E. Woollatt, M. J. Davies, D. Segara, N. F. Hacker, S. M. Henshall, R. L. Sutherland, C. K. W. Watts, Edd, the human orthologue of the hyperplastic discs tumour suppressor gene, is amplified and overexpressed in cancer., *Oncogene* 22 (32) (2003) 5070–5081.
- [198] T. J. Fuja, F. Lin, K. E. Osann, P. J. Bryant, Somatic mutations and altered expression of the candidate tumor suppressors csnk1 epsilon, dlg1, and edd/hhyd in mammary ductal carcinoma., *Cancer Res* 64 (3) (2004) 942–951.
- [199] N. S. Lim, G. Kozlov, T.-C. Chang, O. Groover, N. Siddiqui, L. Volpon, G. D. Crescenzo, A.-B. Shyu, K. Gehring, Comparative peptide binding studies of the pabc domains from the ubiquitin-protein isopeptide ligase hyd and poly(a)-binding protein. implications for hyd function., *J Biol Chem* 281 (20) (2006) 14376–14382.
- [200] C. S. Lin, R. H. Aebersold, S. B. Kent, M. Varma, J. Leavitt, Molecular cloning and characterization of plastin, a human leukocyte protein expressed in transformed human fibroblasts., *Mol Cell Biol* 8 (11) (1988) 4659–4668.

- [201] M. V. de Arruda, S. Watson, C. S. Lin, J. Leavitt, P. Matsudaira, Fimbrin is a homologue of the cytoplasmic phosphoprotein plastin and has domains homologous with calmodulin and actin gelation proteins., *J Cell Biol* 111 (3) (1990) 1069–1079.
- [202] E. Korenbaum, F. Rivero, Calponin homology domains at a glance., *J Cell Sci* 115 (Pt 18) (2002) 3543–3545.
- [203] C. S. Lin, Z. P. Chen, T. Park, K. Ghosh, J. Leavitt, Characterization of the human I-plastin gene promoter in normal and neoplastic cells., *J Biol Chem* 268 (4) (1993) 2793–2801.
- [204] M. Arpin, E. Friederich, M. Algrain, F. Vernel, D. Louvard, Functional differences between I- and t-plastin isoforms., *J Cell Biol* 127 (6 Pt 2) (1994) 1995–2008.
- [205] A. Giganti, J. Plastino, B. Janji, M. V. Troys, D. Lentz, C. Ampe, C. Sykes, E. Friederich, Actin-filament cross-linking protein t-plastin increases arp2/3-mediated actin-based movement., *J Cell Sci* 118 (Pt 6) (2005) 1255–1265.
- [206] T. M. Sandrock, S. M. Brower, K. A. Toenjes, A. E. Adams, Suppressor analysis of fimbrin (sac6p) overexpression in yeast., *Genetics* 151 (4) (1999) 1287–1297.
- [207] K. Nakano, K. Satoh, A. Morimatsu, M. Ohnuma, I. Mabuchi, Interactions among a fimbrin, a capping protein, and an actin-depolymerizing factor in organization of the fission yeast actin cytoskeleton., *Mol Biol Cell* 12 (11) (2001) 3515–3526.
- [208] I. Kondo, H. Hamaguchi, Evidence for the close linkage between lymphocyte cytosol polypeptide with molecular weight of 64,000 (lcp1) and esterase d., *Am J Hum Genet* 37 (6) (1985) 1106–1111.
- [209] M. wan Su, I. Dorocicz, W. H. Dragowska, V. Ho, G. Li, N. Voss, R. Gascoyne, Y. Zhou, Aberrant expression of t-plastin in sezary cells., *Cancer Res* 63 (21) (2003) 7122–7127.

## Literaturverzeichnis

---

- [210] L. Kari, A. Loboda, M. Nebozhyn, A. H. Rook, E. C. Vonderheid, C. Nichols, D. Virok, C. Chang, W.-H. Horng, J. Johnston, M. Wysocka, M. K. Showe, L. C. Showe, Classification and prediction of survival in patients with the leukemic phase of cutaneous t cell lymphoma., *J Exp Med* 197 (11) (2003) 1477–1488.
- [211] Y. Sasaki, F. Itoh, T. Kobayashi, T. Kikuchi, H. Suzuki, M. Toyota, K. Imai, Increased expression of t-fimbrin gene after dna damage in cho cells and inactivation of t-fimbrin by cpg methylation in human colorectal cancer cells., *Int J Cancer* 97 (2) (2002) 211–216.
- [212] R. M. Rao, S. Rama, A. J. Rao, Changes in t-plastin expression with human trophoblast differentiation., *Reprod Biomed Online* 7 (2) (2003) 235–242.
- [213] H. Tourrière, K. Chebli, L. Zekri, B. Courseaud, J. M. Blanchard, E. Bertrand, J. Tazi, The rasgap-associated endoribonuclease g3bp assembles stress granules., *J Cell Biol* 160 (6) (2003) 823–831.
- [214] T. J. Sweet, B. Boyer, W. Hu, K. E. Baker, J. Coller, Microtubule disruption stimulates p-body formation., *RNA* 13 (4) (2007) 493–502.
- [215] P. Fernandez-Funez, M. L. Nino-Rosales, B. de Gouyon, W. C. She, J. M. Luchak, P. Martinez, E. Turiegano, J. Benito, M. Capovilla, P. J. Skinner, A. McCall, I. Canal, H. T. Orr, H. Y. Zoghbi, J. Botas, Identification of genes that modify ataxin-1-induced neurodegeneration., *Nature* 408 (6808) (2000) 101–106.
- [216] R. Mazroui, M.-E. Huot, S. Tremblay, C. Filion, Y. Labelle, E. W. Khandjian, Trapping of messenger rna by fragile x mental retardation protein into cytoplasmic granules induces translation repression., *Hum Mol Genet* 11 (24) (2002) 3007–3017.
- [217] Y. Hua, J. Zhou, Survival motor neuron protein facilitates assembly of stress granules., *FEBS Lett* 572 (1-3) (2004) 69–74.