

VII. REFERENCES

- Adams**, S., van der Laan, L.J., Vernon-Wilson, E., Renardel de Lavalette, C., Dopp, E.A., Dijkstra, C.D., Simmons, D.L. and van den Berg, T.K. (1998) Signal-regulatory protein is selectively expressed by myeloid and neuronal cells. *J Immunol* **161**, 1853-9.
- Aderem**, A. and Ulevitch, R.J. (2000) Toll-like receptors in the induction of the innate immune response. *Nature* **406**, 782-7.
- Akbar**, A.N., Terry, L., Timms, A., Beverley, P.C. and Janossy, G. (1988) Loss of CD45R and gain of UCHL1 reactivity is a feature of primed T cells. *J Immunol* **140**, 2171-8.
- Alexander**, D.R. (2000) The CD45 tyrosine phosphatase: a positive and negative regulator of immune cell function. *Semin Immunol* **12**, 349-59.
- Alvarez**, B., Sanchez, C., Bullido, R., Marina, A., Lunney, J., Alonso, F., Ezquerra, A. and Dominguez, J. (2000) A porcine cell surface receptor identified by monoclonal antibodies to SWC3 is a member of the signal regulatory protein family and associates with protein-tyrosine phosphatase SHP-1. *Tissue Antigens* **55**, 342-51.
- Andrews**, R.G., Singer, J.W. and Bernstein, I.D. (1986) Monoclonal antibody 12-8 recognizes a 115-kd molecule present on both unipotent and multipotent hematopoietic colony-forming cells and their precursors. *Blood* **67**, 842-5.
- Antczak**, D.F. (1989) Biology of the major histocompatibility complex. In: *Veterinary clinical Immunology*. (Halliwell, R.E.W., Gorman, N.T., eds.) W.B. Saunders Company, Philadelphia, PA, USA. p. 473-492.
- Antczak**, D.F. (1992) The major histocompatibility complex of the horse. In: *Equine Infectious diseases VI: Proceedings of the sixth international Conference, Cambridge, 1991*. (W. Plowright, P.D. Ross-dale and J.F. Wade, eds.) R & W Publications, Newmarket, p. 99-112.
- Aristoteli**, L.P., Moller, H.J., Bailey, B., Moestrup, S.K. and Kritharides, L. (2006) The monocytic lineage specific soluble CD163 is a plasma marker of coronary atherosclerosis. *Atherosclerosis* **184**, 342-7.
- Aruffo**, A. and Seed, B. (1987) Molecular cloning of a CD28 cDNA by a high-efficiency COS cell expression system. *Proc Natl Acad Sci U S A* **84**, 8573-7.
- Azuma**, M., Ito, D., Yagita, H., Okumura, K., Phillips, J.H., Lanier, L.L. and Somoza, C. (1993) B70 antigen is a second ligand for CTLA-4 and CD28. *Nature* **366**, 76-9.
- Backe**, E., Schwarting, R., Gerdes, J., Ernst, M. and Stein, H. (1991) Ber-MAC3: new monoclonal antibody that defines human monocyte/macrophage differentiation antigen. *J Clin Pathol* **44**, 936-45.
- Bacon**, K., Gauchat, J.F., Aubry, J.P., Pochon, S., Gruber, P., Henchoz, S. and Bonnefoy, J.Y. (1993) CD21 expressed on basophilic cells is involved in histamine release triggered by CD23 and anti-CD21 antibodies. *Eur J Immunol* **23**, 2721-4.

- Baeten, D., Moller, H.J., Delanghe, J., Veys, E.M., Moestrup, S.K. and De Keyser, F.** (2004) Association of CD163+ macrophages and local production of soluble CD163 with decreased lymphocyte activation in spondylarthropathy synovitis. *Arthritis Rheum* **50**, 1611-23.
- Barbis, D.P., Bainbridge, D., Crump, A.L., Zhang, C.H. and Antczak, D.F.** (1994b) Variation in expression of MHC class II antigens on horse lymphocytes determined by MHC haplotype. *Vet Immunol Immunopathol* **42**, 103-14.
- Bernoco, M., Liu, I.K., Wuest-Ehlert, C.J., Miller, M.E. and Bowers, J.** (1987a) Chemotactic and phagocytic function of peripheral blood polymorphonuclear leucocytes in newborn foals. *J Reprod Fertil Suppl* **35**, 599-605.
- Beyersdorf, N., Hanke, T., Kerkau, T. and Hünig, T.** (2005) Superagonistic anti-CD28 antibodies : potent activators of regulatory T cells for the therapy of autoimmune diseases. *Ann Rheum Dis* **64** Suppl 4, iv91-5.
- Blanchard-Channell, M., Moore, P.F. and Stott, J.L.** (1994) Characterization of monoclonal antibodies specific for equine homologues of CD3 and CD5. *Immunology* **82**, 548-54.
- Bosshart, H. and Jarrett, R.F.** (1998) Deficient major histocompatibility complex class II antigen presentation in a subset of Hodgkin's disease tumor cells. *Blood* **92**, 2252-9.
- Braun, M., Melchers, I., Peter, H.H. and Illges, H.** (1998) Human B and T lymphocytes have similar amounts of CD21 mRNA, but differ in surface expression of the CD21 glycoprotein. *Int Immunol* **10**, 1197-202.
- Brodersen, R., Bijlsma, F., Gori, K., Jensen, K.T., Chen, W., Dominguez, J., Haverson, K., Moore, P.F., Saalmüller, A., Sachs, D., Slierendrecht, W.J., Stokes, C., Vainio, O., Zuckermann, F. and Aasted, B.** (1998) Analysis of the immunological cross reactivities of 213 well characterized monoclonal antibodies with specificities against various leucocyte surface antigens of human and 11 animal species. *Vet Immunol Immunopathol* **64**, 1-13.
- Burt, R.K.** (1999) Clinical utility in maximizing CD34+ cell count in stem cell grafts. *Oncologist* **4**, 265-8.
- Byrne, K.M., Davis, W.C., Holmes, M.A., Brassfield, A.L. and McGuire, T.C.** (1997) Cytokine RNA expression in an equine CD4+ subset differentiated by expression of a novel 46-kDa surface protein. *Vet Immunol Immunopathol* **56**, 191-204.
- Civin, C.I., Trischmann, T., Kadan, N.S., Davis, J., Noga, S., Cohen, K., Duffy, B., Groenewegen, I., Wiley, J., Law, P., Hardwick, A., Oldham, F. and Gee, A.** (1996) Highly purified CD34-positive cells reconstitute hematopoiesis. *J Clin Oncol* **14**, 2224-33.
- Clement, L.T., Yamashita, N. and Martin, A.M.** (1988) The functionally distinct subpopulations of human CD4+ helper/inducer T lymphocytes defined by anti-CD45R antibodies derive sequentially from a differentiation pathway that is regulated by activation-dependent post-thymic differentiation. *J Immunol* **141**, 1464-70.

- Cobbold**, S.P. and Metcalfe, S. (1994) Monoclonal antibodies that define canine homologues of human CD antigens: Summary of the First International Canine Leukocyte Antigen Workshop (CLAW). *Tissue Antigens* **43**:137-154.
- Crawford**, K., Gabuzda, D., Pantazopoulos, V., Xu, J., Clement, C., Reinherz, E. and Alper, C.A. (1999) Circulating CD2+ monocytes are dendritic cells. *J Immunol* **163**, 5920-8.
- Crepaldi**, T., Crump, A., Newman, M., Ferrone, S. and Antczak, D.F. (1986) Equine T lymphocytes express MHC class II antigens. *J Immunogenet* **13**, 349-60.
- Crump**, A.L., Davis, W. and Antczak, D.F. (1988) A monoclonal antibody identifying a T-cell marker in the horse. *Anim Genet* **19**, 349-57.
- Davis**, W.C., Marusic, S., Lewin, H.A., Splitter, G.A., Perryman, L.E., McGuire, T.C. and Gorham, J.R. (1987) The development and analysis of species specific and cross reactive monoclonal antibodies to leukocyte differentiation antigens and antigens of the major histocompatibility complex for use in the study of the immune system in cattle and other species. *Vet Immunol Immunopathol* **15**, 337-76.
- Deans**, J.P., Boyd, A.W. and Pilarski, L.M. (1989) Transitions from high to low molecular weight isoforms of CD45 (T200) involve rapid activation of alternate mRNA splicing and slow turnover of surface CD45R. *J Immunol* **143**, 1233-8.
- DeFranco**, A.L. (1991) Between B cells and T cells. *Nature* **351**, 603-4.
- de La Selle**, V., Gluckman, E. and Bruley-Rosset, M. (1996) Newborn blood can engraft adult mice without inducing graft-versus-host disease across non H-2 antigens. *Blood* **87**, 3977-83.
- Dieckmann**, D., Plottner, H., Berchtold, S., Berger, T. and Schuler, G. (2001) Ex vivo isolation and characterization of CD4(+)CD25(+) T cells with regulatory properties from human blood. *J Exp Med* **193**, 1303-10.
- Donaldson**, W.L., Crump, A.L., Zhang, C.H., Kornbluth, J., Kamoun, M., Davis, W. and Antczak, D.F. (1988) At least two loci encode polymorphic class I MHC antigens in the horse. *Anim Genet* **19**, 379-90.
- Drbal**, K., Angelisova, P., Hilgert, I., Cerny, J., Novak, P. and Horejsi, V. (2001) A proteolytically truncated form of free CD18, the common chain of leukocyte integrins, as a novel marker of activated myeloid cells. *Blood* **98**, 1561-6.
- Drillenburg**, P. and Pals, S.T. (2000) Cell adhesion receptors in lymphoma dissemination. *Blood* **95**, 1900-10.
- Ezekowitz**, R.A., Sastry, K., Bailly, P. and Warner, A. (1990) Molecular characterization of the human macrophage mannose receptor: demonstration of multiple carbohydrate recognition-like domains and phagocytosis of yeasts in Cos-1 cells. *J Exp Med* **172**, 1785-94.
- Fackler**, M.J., Civin, C.I. and May, W.S. (1992) Up-regulation of surface CD34 is associated with protein kinase C-mediated hyperphosphorylation of CD34. *J Biol Chem* **267**, 17540-6.

- Fischer**, E., Delibrias, C. and Kazatchkine, M.D. (1991) Expression of CR2 (the C3dg/EBV receptor, CD21) on normal human peripheral blood T lymphocytes. *J Immunol* **146**, 865-9.
- Fujioka**, Y., Matozaki, T., Noguchi, T., Iwamatsu, A., Yamao, T., Takahashi, N., Tsuda, M., Takada, T. and Kasuga, M. (1996) A novel membrane glycoprotein, SHPS-1, that binds the SH2-domain-containing protein tyrosine phosphatase SHP-2 in response to mitogens and cell adhesion. *Mol Cell Biol* **16**, 6887-99.
- Fukuda**, M. (1991) Lysosomal membrane glycoproteins. Structure, biosynthesis, and intracellular trafficking. *J Biol Chem* **266**, 21327-30.
- Geissler**, K., Geissler, W., Hinterberger, W., Lechner, K. and Wurnig, P. (1986) Circulating committed and pluripotent haemopoietic progenitor cells, in infants. *Acta Haematol* **75**, 18-22.
- George**, M., Jr. and Ryder, O.A. (1986) Mitochondrial DNA evolution in the genus Equus. *Mol Biol Evol* **3**, 535-46.
- Goyert**, S.M. (2002b) Cluster report: CD68. In: *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uggioni, E. van der Schoot, E. Vivier, and H. Zola, eds.) Oxford University Press, Oxford, p. 821.
- Goyert**, S.M. (2002a) Cluster report: CD14. In: *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uggioni, E. van der Schoot, E. Vivier, and H. Zola, eds.) Oxford University Press, Oxford, p. 761-763.
- Greaves**, M., Titley, I., Colman, S., Buhrig, H-J, Campos, L., castoldi, G., Garrido, F., Gaudnack, G., GirardJ-P, Ingles-Esteve, J., Invernizzi, R., Knapp, W., Landsorp, P., lanza, F., Merle-Beral, H., Parravicini, C., Razak, K., Ruiz-Cabello, F., Springer, T., Van der schoot, C., Sutherland, D. (1995) M10 CD34 cluster workshop section. In: *Leukocyte Typing V: White Cell Differentiation Antigen.*, Vol. 1. (S. F. Schlossman, L. Boumsell, W. Gilks, J. M. Harlan, T. Kashimoto, C. Morimoto, J. Ritz, S. Shaw, R. Silverstein, T. Springer, T. F. Tedder, R. F. Todd, eds.) Oxford University Press, New York, p. 840.
- Greenlee**, P.G., Calvano, S.E., Quimby, F.W. and Hurvitz, A.I. (1987) Investigation of cross-reactivity between commercially available antibodies directed against human, mouse, and rat lymphocyte surface antigens and surface markers on canine cells. *Vet Immunol Immunopathol* **15**, 285-96.
- Groves**, C.P. (2002) Taxonomy of living Equidae. In: *Equids: Zebras, Assess and Horses: Status Survey and Conservation Action Plan*. IUCN/SSC Equid Specialist Group, IUCN (The World ConservationUnion), Gland, Switzerland and Cambridge. (<http://www.iucn.org/themes/ssc/actionplans/equids/equidactionplanlowres.pdf>) P. 94-107.

- Hathcock, K.S., Laszlo, G., Dickler, H.B., Sharrow, S.O., Johnson, P., Trowbridge, I.S. and Hodes, R.J.** (1992) Expression of variable exon A-, B-, and C-specific CD45 determinants on peripheral and thymic T cell populations. *J Immunol* **148**, 19-28.
- Hammond, S.A., Horohov, D. and Montelaro, R.C.** (1999) Functional characterization of equine dendritic cells propagated ex vivo using recombinant human GM-CSF and recombinant equine IL-4. *Vet Immunol Immunopathol* **71**, 197-214.
- Hanawa, H., Ma, Y., Mikolajczak, S.A., Charles, M.L., Yoshida, T., Yoshida, R., Strathdee, C.A., Litchfield, D.W. and Ochi, A.** (2002) A novel costimulatory signaling in human T lymphocytes by a splice variant of CD28. *Blood* **99**, 2138-45.
- Handgretinger, R., Schumm, M., Lang, P., Greil, J., Reiter, A., Bader, P., Niethammer, D. and Klingebiel, T.** (1999) Transplantation of megadoses of purified haploididentical stem cells. *Ann N Y Acad Sci* **872**, 351-62
- Haziot, A., Tsuberi, B.Z. and Goyert, S.M.** (1993) Neutrophil CD14: biochemical properties and role in the secretion of tumor necrosis factor-alpha in response to lipopolysaccharide. *J Immunol* **150**, 5556-65.
- Hermiston, M.L., Xu, Z. and Weiss, A.** (2003) CD45: a critical regulator of signaling thresholds in immune cells. *Annu Rev Immunol* **21**, 107-37.
- Herz, J., Kowal, R.C., Goldstein, J.L. and Brown, M.S.** (1990) Proteolytic processing of the 600 kd low density lipoprotein receptor-related protein (LRP) occurs in a trans-Golgi compartment. *Embo J* **9**, 1769-76.
- Hildreth, J.E. and August, J.T.** (1985) The human lymphocyte function-associated (HLFA) antigen and a related macrophage differentiation antigen (HMac-1): functional effects of subunit-specific monoclonal antibodies. *J Immunol* **134**, 3272-80.
- Hogarth, P.M. and McKenzie, I.F.** (1992) Ly-15.2 and Ly-21.2 are distinct polymorphisms of the LFA-1 heavy chain. *Immunogenetics* **36**, 396-9.
- Hogg, N.** (2002a) Cluster report: CD11b. In: *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uggioni, E. van der Schoot, E. Vivier, and H. Zola, eds.) Oxford University Press, Oxford, p. 758-759.
- Hogg, N.** (2002b) Cluster report: CD11c. In: *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uggioni, E. van der Schoot, E. Vivier, and H. Zola, eds.) Oxford University Press, Oxford, p. 759-760.
- Hogger, P., Dreier, J., Droste, A., Buck, F. and Sorg, C.** (1998) Identification of the integral membrane protein RM3/1 on human monocytes as a glucocorticoid-inducible member of the scavenger receptor cysteine-rich family (CD163). *J Immunol* **161**, 1883-90.
- Holmes, M.A. and Lunn, D.P.** (1994) Variation of MHC II expression on canine lymphocytes with age. *Tissue Antigens* **43**, 179-83.

- Holness**, C.L. and Simmons, D.L. (1993) Molecular cloning of CD68, a human macrophage marker related to lysosomal glycoproteins. *Blood* **81**, 1607-13.
- Horejsi**, V. (1991) Surface antigens of human leukocytes. *Adv Immunol* **49**, 75-147.
- Hormanski**, C.E., Truax, R., Pourciau, S.S., Folsom, R.W. and Horohov, D.W. (1992) Induction of lymphokine-activated killer cells of equine origin: specificity for equine target cells. *Vet Immunol Immunopathol* **32**, 25-36.
- Hume**, D.A. and Gordon, S. (1983) Mononuclear phagocyte system of the mouse defined by immunohistochemical localization of antigen F4/80. Identification of resident macrophages in renal medullary and cortical interstitium and the juxtaglomerular complex. *J Exp Med* **157**, 1704-9.
- Jacobsen**, C.N., Aasted, B., Broe, M.K. and Petersen, J.L. (1993) Reactivities of 20 anti-human monoclonal antibodies with leucocytes from ten different animal species. *Vet Immunol Immunopathol* **39**, 461-6.
- Janeway**, C.A., Jr. (1992) The T cell receptor as a multicomponent signalling machine: CD4/CD8 coreceptors and CD45 in T cell activation. *Annu Rev Immunol* **10**, 645-74.
- Jonuleit**, H., Schmitt, E., Stassen, M., Tuettenberg, A., Knop, J. and Enk, A.H. (2001) Identification and functional characterization of human CD4(+)CD25(+) T cells with regulatory properties isolated from peripheral blood. *J Exp Med* **193**, 1285-94.
- June**, C.H., Bluestone, J.A., Nadler, L.M. and Thompson, C.B. (1994) The B7 and CD28 receptor families. *Immunol Today* **15**, 321-31.
- Katz**, F.E., Tindle, R., Sutherland, D.R. and Greaves, M.F. (1985) Identification of a membrane glycoprotein associated with haemopoietic progenitor cells. *Leuk Res* **9**, 191-8.
- Keogh**, M.C., Elliot, J., Norton, T. and Lake, R.A. (1997) A function for CD2 on murine B cells? *Immunol Cell Biol* **75**, 333-9.
- Kharitonov**, A., Chen, Z., Sures, I., Wang, H., Schilling, J. and Ullrich, A. (1997) A family of proteins that inhibit signalling through tyrosine kinase receptors. *Nature* **386**, 181-6.
- Kipps**, T.J. (1989) The CD5 B cell. *Adv Immunol* **47**, 117-85.
- Kishimoto**, T.K., O'Connor, K., Lee, A., Roberts, T.M. and Springer, T.A. (1987) Cloning of the beta subunit of the leukocyte adhesion proteins: homology to an extracellular matrix receptor defines a novel supergene family. *Cell* **48**, 681-90.
- Köhler**, G. and Milstein, C. (1975) Continuous cultures of fused cells secreting antibody of predefined specificity. *Nature* **256**, 495-7.
- Kozlow**, E.J., Wilson, G.L., Fox, C.H. and Kehrl, J.H. (1993) Subtractive cDNA cloning of a novel member of the Ig gene superfamily expressed at high levels in activated B lymphocytes. *Blood* **81**, 454-61.

- Krause**, D.S., Fackler, M.J., Civin, C.I. and May, W.S. (1996) CD34: structure, biology, and clinical utility. *Blood* **87**, 1-13.
- Krause**, D.S., Fackler, M.J., Smith, O.M., May W. (1993) Two forms of CD34 protein are expressed in human KMT2 and KG1a cells. *Blood* **82**: 427a (abstr, suppl 1).
- Krause**, D.S., Ito, T., Fackler, M.J., Smith, O.M., Collector, M.I., Sharkis, S.J. and May, W.S. (1994) Characterization of murine CD34, a marker for hematopoietic progenitor and stem cells. *Blood* **84**, 691-701.
- Kürzinger**, K., Reynolds, T., Germain, R.N., Davignon, D., Martz, E. and Springer, T.A. (1981) A novel lymphocyte function-associated antigen (LFA-1): cellular distribution, quantitative expression, and structure. *J Immunol* **127**, 596-602.
- Kydd**, J., Antczak, D.F., Allen, W.R., Barbis, D., Butcher, G., Davis, W., Duffus, W.P., Edington, N., Grunig, G., Holmes, M.A. and et al. (1994) Report of the First International Workshop on Equine Leucocyte Antigens, Cambridge, UK, July 1991. *Vet Immunol Immunopathol* **42**, 3-60.
- Kydd**, J.H., Butcher, G.W., Antczak, D.F. and Allen, W.R. (1991) Expression of major histocompatibility complex (MHC) class 1 molecules on early trophoblast. *J Reprod Fertil Suppl* **44**, 463-77.
- Lafferty**, K.J. and Cunningham, A.J. (1975) A new analysis of allogeneic interactions. *Aust J Exp Biol Med Sci* **53**, 27-42.
- Lauener**, R.P., Goyert, S.M., Geha, R.S. and Vercelli, D. (1990) Interleukin 4 down-regulates the expression of CD14 in normal human monocytes. *Eur J Immunol* **20**, 2375-81.
- Lazary**, S., Antczak, D.F., Bailey, E., Bell, T.K., Bernoco, D., Byrns, G. and McClure, J.J. (1988) Joint Report of the Fifth International Workshop on Lymphocyte Alloantigens of the Horse, Baton Rouge, Louisiana, 31 October-1 November 1987. *Anim Genet* **19**, 447-56.
- Ledbetter**, J.A., Evans, R.L., Lipinski, M., Cunningham-Rundles, C., Good, R.A. and Herzenberg, L.A. (1981) Evolutionary conservation of surface molecules that distinguish T lymphocyte helper/inducer and cytotoxic/suppressor subpopulations in mouse and man. *J Exp Med* **153**, 310-23.
- Lee**, K.P., Taylor, C., Petryniak, B., Turka, L.A., June, C.H. and Thompson, C.B. (1990) The genomic organization of the CD28 gene. Implications for the regulation of CD28 mRNA expression and heterogeneity. *J Immunol* **145**, 344-52.
- Leung**, H.T., Bradshaw, J., Cleaveland, J.S. and Linsley, P.S. (1995) Cytotoxic T lymphocyte-associated molecule-4, a high-avidity receptor for CD80 and CD86, contains an intracellular localization motif in its cytoplasmic tail. *J Biol Chem* **270**, 25107-14.
- Li**, K., Li, C.K., Fok, T.F., Liu, J. and Yuen, P.M. (1998) Neonatal blood: a source of haematopoietic stem cells for transplantation? *Lancet* **351**, 647-8.

- Lin**, G.-X., Yang X., Hollemweguer E., Yu J.-F., Li L., Wu X.-W., Ward T., Chen Z.. (2002) Cross-reactivity of CD antibodies in eight animal species. In: *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uguccioni, E. van der Schoot, E. Vivier, and H. Zola, eds.) Oxford University Press, Oxford, p. 519-523.
- Linsley**, P.S. and Ledbetter, J.A. (1993) The role of the CD28 receptor during T cell responses to antigen. *Annu Rev Immunol* **11**, 191-212.
- Lu**, P., Wang, Y.L. and Linsley, P.S. (1997) Regulation of self-tolerance by CD80/CD86 interactions. *Curr Opin Immunol* **9**, 858-62.
- Lühder**, F., Huang, Y., Dennehy, K.M., Guntermann, C., Muller, I., Winkler, E., Kerkau, T., Ikemizu, S., Davis, S.J., Hanke, T. and Hünig, T. (2003) Topological requirements and signaling properties of T cell-activating, anti-CD28 antibody superagonists. *J Exp Med* **197**, 955-66.
- Lunn**, D.P. (1993) A comparative review of human and equine leucocyte differentiation antigens. *Br Vet J* **149**, 31-49.
- Lunn**, D.P., Holmes, M.A., Antczak, D.F., Agerwal, N., Baker, J., Bendali-Ahcene, S., Blanchard-Channell, M., Byrne, K.M., Cannizzo, K., Davis, W., Hamilton, M.J., Hannant, D., Kondo, T., Kydd, J.H., Monier, M.C., Moore, P.F., O'Neil, T., Schram, B.R., Sheoran, A., Stott, J.L., Sugiura, T. and Vagnoni, K.E. (1998) Report of the Second Equine Leucocyte Antigen Workshop, Squaw valley, California, July 1995. *Vet Immunol Immunopathol* **62**, 101-43.
- Lunney**, J.K., Walker, K., Goldman, T., Aasted, B., Bianchi, A., Binns, R., Licence, S., Bischof, R., Brandon, M., Blecha, F. and et al. (1994) Overview of the First International Workshop to Define Swine Leukocyte Cluster of Differentiation (CD) Antigens. *Vet Immunol Immunopathol* **43**, 193-206.
- Lydyard**, P.M. and Mackenzie, L. (1989) Lymphocyte activation by CD5 mAb. In: *Leucocyte Typing IV: White Cell Differentiation Antigens*. (Knapp W., Dorken B., Rieber E.P., Stein H., Gilks W.R., Schmidt R.E. and Von dem Borne A.E.G.Kr., eds.) Oxford University Press, Oxford, p. 338.
- Lynch**, K.W. (2004) Consequences of regulated pre-mRNA splicing in the immune system. *Nat Rev Immunol* **4**, 931-40.
- Mao**, M., Fu, G., Wu, J.S., Zhang, Q.H., Zhou, J., Kan, L.X., Huang, Q.H., He, K.L., Gu, B.W., Han, Z.G., Shen, Y., Gu, J., Yu, Y.P., Xu, S.H., Wang, Y.X., Chen, S.J. and Chen, Z. (1998) Identification of genes expressed in human CD34 (+) hematopoietic stem/progenitor cells by expressed sequence tags and efficient full-length cDNA cloning. *Proc Natl Acad Sci U S A* **95**, 8175-80.
- Marchant**, A., Duchow, J., Delville, J.P. and Goldman, M. (1992) Lipopolysaccharide induces up-regulation of CD14 molecule on monocytes in human whole blood. *Eur J Immunol* **22**, 1663-5.

- Margulies, D.H.** (2003) CD28, costimulator or agonist receptor? *J Exp Med* **197**, 949-53.
- Marvel, J.** and **Mayer, A.** (1988) CD45R gives immunofluorescence and transduces signals on mouse T cells. *Eur J Immunol* **18**, 825-8.
- Mason, D.**, Andre, P., Bensussan, A., Buckley, C., Civin, C., Clark, E., de Haas, M., Goyert, S., Hadam, M., Hart, D., Horejsi, V., Meuer, S., Morrissey, J., Schwartz-Albiez, R., Shaw, S., Simmons, D., Uguccioni, M., van der Schoot, E., Vivier, E. and Zola, H. (2002) Reference: CD Antigens 2002. *J Immunol* **168**, 2083-6.
- Mauel, S.**, Steinbach, F. and Ludwig, H. (2006) Monocyte-derived dendritic cells from horses differ from dendritic cells of humans and mice. *Immunology* **117**, 463-73.
- May G.**, Healy, L., Gale, K., Greaves, M (1994) Functional analysis of stem cell antigen CD34 in transgenic mice. *Blood* **84**: 871.
- Mayall, S.**, Siedek, E. and Hamblin, A.S. (2001) The anti-human CD21 antibody, BU33, identifies equine B cells. *J Comp Pathol* **124**, 83-7.
- Merant, C.**, Bonnefont, C., Desbos, A., Greenland, T., Cadore, J.L. and Monier, J.C. (2003) Cross-species reactivity of seven monoclonal antibodies with equine lymphocytes by flow cytometry. *Vet Res* **34**, 791-801.
- Meuer, S.C.**, Hussey, R.E., Fabbri, M., Fox, D., Acuto, O., Fitzgerald, K.A., Hodgdon, J.C., Protentis, J.P., Schlossman, S.F. and Reinherz, E.L. (1984) An alternative pathway of T-cell activation: a functional role for the 50 kd T11 sheep erythrocyte receptor protein. *Cell* **36**, 897-906.
- Miyake, K.**, Underhill, C.B., Lesley, J. and Kincade, P.W. (1990) Hyaluronate can function as a cell adhesion molecule and CD44 participates in hyaluronate recognition. *J Exp Med* **172**, 69-75.
- Moehlman, P.D.** (ed.) (2002) *Equids: Zebras, Assess and Horses: Status Survey and Conservation Action Plan*. IUCN/SSC Equid Specialist Group, IUCN (The World ConservationUnion), Gland, Switzerland and Cambridge. (<http://www.iucn.org/themes/ssc/actionplans/equids/equidactionplanlowres.pdf>).
- Moingeon, P.**, Chang, H.C., Sayre, P.H., Clayton, L.K., Alcover, A., Gardner, P. and Reinherz, E.L. (1989) The structural biology of CD2. *Immunol Rev* **111**, 111-44.
- Moestrup, S.** (2002) Cluster report: CD91. In: *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uguccioni, E. van der Schoot, E. Vivier, and H. Zola, eds.) Oxford University Press, Oxford, p. 836-837.
- Moller, H.J.**, Aerts, H., Gronbaek, H., Peterslund, N.A., Hyltoft Petersen, P., Hornung, N., Rejnmark, L., Jabbarpour, E. and Moestrup, S.K. (2002) Soluble CD163: a marker molecule for monocyte/macrophage activity in disease. *Scand J Clin Lab Invest Suppl* **237**, 29-33.

- Moller**, H.J., de Fost, M., Aerts, H., Hollak, C. and Moestrup, S.K. (2004) Plasma level of the macrophage-derived soluble CD163 is increased and positively correlates with severity in Gaucher's disease. *Eur J Haematol* **72**, 135-9.
- Moller**, H.J., Petersen, P.H., Rejnmark, L. and Moestrup, S.K. (2003) Biological variation of soluble CD163. *Scand J Clin Lab Invest* **63**, 15-21.
- Monos**, D.S., Wolf, B., Radka, S.F., Rifat, S., Donawick, W.J., Soma, L.R., Zmijewski, C.M. and Kamoun, M. (1989) Equine class II MHC antigens: identification of two sets of epitopes using anti-human monoclonal antibodies. *Tissue Antigens* **34**, 111-20.
- Muto**, S., Vetvicka, V. and Ross, G.D. (1993) CR3 (CD11b/CD18) expressed by cytotoxic T cells and natural killer cells is upregulated in a manner similar to neutrophil CR3 following stimulation with various activating agents. *J Clin Immunol* **13**, 175-84.
- Nakamura**, S., Nagahama, M., Kagami, Y., Yatabe, Y., Takeuchi, T., Kojima, M., Motoori, T., Suzuki, R., Taji, H., Ogura, M., Mizoguchi, Y., Okamoto, M., Suzuki, H., Oyama, A., Seto, M., Morishima, Y., Koshikawa, T., Takahashi, T., Kurita, S. and Suchi, T. (1999) Hodgkin's disease expressing follicular dendritic cell marker CD21 without any other B-cell marker: a clinicopathologic study of nine cases. *Am J Surg Pathol* **23**, 363-76.
- Nakazawa**, T., Agematsu, K. and Yabuhara, A. (1997) Later development of Fas ligand-mediated cytotoxicity as compared with granule-mediated cytotoxicity during the maturation of natural killer cells. *Immunology* **92**, 180-7.
- Nauseef**, W.M., Root, R.K. and Malech, H.L. (1983) Biochemical and immunologic analysis of hereditary myeloperoxidase deficiency. *J Clin Invest* **71**, 1297-307.
- Nishimura**, Y., Bierer, B.E. and Burakoff, S.J. (1988a) Expression of CD5 regulates responsiveness to IL-1. *J Immunol* **141**, 3438-44.
- Nishimura**, Y., Bierer, B.E., Jones, W.K., Jones, N.H., Strominger, J.L. and Burakoff, S.J. (1988b) Expression and function of a CD5 cDNA in human and murine T cells. *Eur J Immunol* **18**, 747-53.
- Oakenfull**, E.A. and Clegg, J.B. (1998) Phylogenetic relationships within the genus *Equus* and the evolution of alpha and theta globin genes. *J Mol Evol* **47**, 772-83.
- Oakenfull**, E.A., Lim, H.N. and Ryder, O.A. (2000) A survey of equid mitochondrial DNA: Implications for the evolution, genetic diversity and conservation of *Equus*. *Conservation Genetics* **1**, 341-355.
- Oldenborg**, P.A., Gresham, H.D. and Lindberg, F.P. (2001) CD47-signal regulatory protein alpha (SIRPalpha) regulates Fc γ and complement receptor-mediated phagocytosis. *J Exp Med* **193**, 855-62.
- Parsons**, K.R., Young, J.R., Collins, B.A. and Howard, C.J. (1996) Cattle CTLA-4, CD28 and chicken CD28 bind CD86: MYPPPY is not conserved in cattle CD28. *Immunogenetics* **43**, 388-91.

- Peach, R.J., Bajorath, J., Brady, W., Leytze, G., Greene, J., Naemura, J. and Linsley, P.S.** (1994) Complementarity determining region 1 (CDR1)- and CDR3-analogous regions in CTLA-4 and CD28 determine the binding to B7-1. *J Exp Med* **180**, 2049-58.
- Pedersen, L.G., Castelruiz, Y., Jacobsen, S. and Aasted, B.** (2002) Identification of monoclonal antibodies that cross-react with cytokines from different animal species. *Vet Immunol Immunopathol* **88**, 111-22
- Penninger, J.M., Irie-Sasaki, J., Sasaki, T. and Oliveira-dos-Santos, A.J.** (2001) CD45: new jobs for an old acquaintance. *Nat Immunol* **2**, 389-96.
- Perussia, B., Fanning, V. and Trinchieri, G.** (1983) A human NK and K cell subset shares with cytotoxic T cells expression of the antigen recognized by antibody OKT8. *J Immunol* **131**, 223-31.
- Philippidis, P., Mason, J.C., Evans, B.J., Nadra, I., Taylor, K.M., Haskard, D.O. and Landis, R.C.** (2004) Hemoglobin scavenger receptor CD163 mediates interleukin-10 release and heme oxygenase-1 synthesis: antiinflammatory monocyte-macrophage responses in vitro, in resolving skin blisters in vivo, and after cardiopulmonary bypass surgery. *Circ Res* **94**, 119-26.
- Pintado, C.O., Friend, M. and Llanes, D.** (1995) Characterisation of a membrane receptor on ruminants and equine platelets and peripheral blood leukocytes similar to the human integrin receptor glycoprotein IIb/IIIa (CD41/61). *Vet Immunol Immunopathol* **44**, 359-68.
- Powrie, F., Carlino, J., Leach, M.W., Mauze, S. and Coffman, R.L.** (1996) A critical role for transforming growth factor-beta but not interleukin 4 in the suppression of T helper type 1-mediated colitis by CD45RB(low) CD4+ T cells. *J Exp Med* **183**, 2669-74.
- Pugin, J., Schurer-Maly, C.C., Leturcq, D., Moriarty, A., Ulevitch, R.J. and Tobias, P.S.** (1993) Lipopolysaccharide activation of human endothelial and epithelial cells is mediated by lipopolysaccharide-binding protein and soluble CD14. *Proc Natl Acad Sci U S A* **90**, 2744-8.
- Ramkumar, P., Martinez-Pomares, L., Gordon, S., Fraser P., Ezekowitz, A., Stahl, D.** (2002) cluster report: CD206 (MMR). In: *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uggioni, E. van der Schoot, E. Vivier, and H. Zola, eds.) Oxford University Press, Oxford, p. 917-918.
- Raulet, D.H. and Eisen, H.N.** (1990) Cellular basis for immune responses. In: *general Immunology*, 2nd edn (Eisen, H. N. ed.), J.B. Lippincott Company, Philadelphia, USA. p. 81-124.
- Reimann, K.A., Waite, B.C., Lee-Parritz, D.E., Lin, W., Uchanska-Ziegler, B., O'Connell, M.J. and Letvin, N.L.** (1994) Use of human leukocyte-specific monoclonal antibodies for clinically immunophenotyping lymphocytes of rhesus monkeys. *Cytometry* **17**, 102-8.

- Reinherz**, E.L., Nadler, L.M., Rosenthal, D.S., Moloney, W.C. and Schlossman, S.F. (1979) T-cell-subset characterization of human T-CLL. *Blood* **53**, 1066-75.
- Ritter**, M., Buechler, C., Langmann, T. and Schmitz, G. (1999) Genomic organization and chromosomal localization of the human CD163 (M130) gene: a member of the scavenger receptor cysteine-rich superfamily. *Biochem Biophys Res Commun* **260**, 466-74.
- Ritz**, J., Schmidt, R.E., Michon, J., Hercend, T. and Schlossman, S.F. (1988) Characterization of functional surface structures on human natural killer cells. *Adv Immunol* **42**, 181-211.
- Rodeberg**, D.A., Morris, R.E. and Babcock, G.F. (1997) Azurophilic granules of human neutrophils contain CD14. *Infect Immun* **65**, 4747-53.
- Rodig**, S.J., Shahsafaei, A., Li, B. and Dorfman, D.M. (2005) The CD45 isoform B220 identifies select subsets of human B cells and B-cell lymphoproliferative disorders. *Hum Pathol* **36**, 51-7.
- Saalmüller**, A., Lunney, J.K., Daubенberger, C., Davis, W., Fischer, U., Gobel, T.W., Griebel, P., Hollemweguer, E., Lasco, T., Meister, R., Schuberth, H.J., Sestak, K., Sopp, P., Steinbach, F., Xiao-Wei, W. and Aasted, B. (2005) Summary of the animal homologue section of HLDA8. *Cell Immunol* **236**, 51-8.
- Saginario**, C., Sterling, H., Beckers, C., Kobayashi, R., Solimena, M., Ullu, E. and Vignery, A. (1998) MFR, a putative receptor mediating the fusion of macrophages. *Mol Cell Biol* **18**, 6213-23.
- Sakaguchi**, S., Sakaguchi, N., Asano, M., Itoh, M. and Toda, M. (1995) Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *J Immunol* **155**, 1151-64.
- Saito**, N., Pulford, K.A., Breton-Gorius, J., Masse, J.M., Mason, D.Y. and Cramer, E.M. (1991) Ultrastructural localization of the CD68 macrophage-associated antigen in human blood neutrophils and monocytes. *Am J Pathol* **139**, 1053-9.
- Sallusto**, F. and Lanzavecchia, A. (1994) Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin 4 and downregulated by tumor necrosis factor alpha. *J Exp Med* **179**, 1109-18.
- Salmon**, M., Kitas, G.D. and Bacon, P.A. (1989) Production of lymphokine mRNA by CD45R+ and CD45R- helper T cells from human peripheral blood and by human CD4+ T cell clones. *J Immunol* **143**, 907-12.
- Salmon**, M., Pilling, D., Borthwick, N.J., Viner, N., Janossy, G., Bacon, P.A. and Akbar, A.N. (1994) The progressive differentiation of primed T cells is associated with an increasing susceptibility to apoptosis. *Eur J Immunol* **24**, 892-9.

- Sambrook**, J.E., Fritsch F., and Maniatis T. (1989) *Molecular cloning: a laboratory manual*. Second edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, USA. E. 5.
- Sanchez**, C., Domenech, N., Vazquez, J., Alonso, F., Ezquerra, A. and Dominguez, J. (1999) The porcine 2A10 antigen is homologous to human CD163 and related to macrophage differentiation. *J Immunol* **162**, 5230-7.
- Sanchez-Madrid**, F., Nagy, J.A., Robbins, E., Simon, P. and Springer, T.A. (1983) A human leukocyte differentiation antigen family with distinct alpha-subunits and a common beta-subunit: the lymphocyte function-associated antigen (LFA-1), the C3bi complement receptor (OKM1/Mac-1), and the p150,95 molecule. *J Exp Med* **158**, 1785-1803.
- Sano**, S., Ito, S., Nakamura, M. and Nakagawa, H. (1990) Oligosaccharide-related epitope specific for a brain-specific glycoprotein, 1D4 antigen. *J Neurochem* **55**, 1252-7.
- Sano**, S., Matsuda, Y. and Nakagawa, H. (1989) A novel brain-specific antigen: a glycoprotein electrophoretically similar to but immunochemically different from type B nucleoside diphosphatase. *J Biochem (Tokyo)* **105**, 457-60.
- Sano**, S., Ohnishi, H., Omori, A., Hasegawa, J. and Kubota, M. (1997) BIT, an immune antigen receptor-like molecule in the brain. *FEBS Lett* **411**, 327-34.
- Sato**, S. and Tedder, T.F. (2002) Cluster report: CD19. In: *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uguccioni, E. van der Schoot, E. Vivier, and H. Zola, eds.) Oxford University Press, Oxford, p. 764-765.
- Schütt**, C., Witt, S., Grunwald, U., Stelter, F., Schilling, T., Fan, X., Marquart, B., Bassarab, P., Krüger, C. (1995) Epitope mapping of CD14 glycoprotein. in *Leukocyte typing V* (S. F. Schlossman, L. Boumsell, W. Gilks, J. M. Harlan, T. Kashimoto, C. Morimoto, J. Ritz, S. Shaw, R. Silverstein, T. Springer, T. F. Tedder, R. F. Todd, eds.) Oxford University Press, Oxford, 785-788
- Seiffert**, M., Brossart, P., Cant, C., Cella, M., Colonna, M., Brugger, W., Kanz, L., Ullrich, A. and Buhring, H.J. (2001) Signal-regulatory protein alpha (SIRPalpha) but not SIRPbeta is involved in T-cell activation, binds to CD47 with high affinity, and is expressed on immature CD34 (+) CD38 (-) hematopoietic cells. *Blood* **97**, 2741-9.
- Seiffert**, M., Cant, C., Chen, Z., Rappold, I., Brugger, W., Kanz, L., Brown, E.J., Ullrich, A. and Buhring, H.J. (1999) Human signal-regulatory protein is expressed on normal, but not on subsets of leukemic myeloid cells and mediates cellular adhesion involving its counterreceptor CD47. *Blood* **94**, 3633-43.
- Sen**, G.C. and Sarkar, S.N. (2005) Transcriptional signaling by double-stranded RNA: role of TLR3. *Cytokine Growth Factor Rev* **16**, 1-14.
- Serra**, H.M., Krowka, J.F., Ledbetter, J.A. and Pilarski, L.M. (1988) Loss of CD45R (Lp220) represents a post-thymic T cell differentiation event. *J Immunol* **140**, 1435-41.

- Sharon**, M., Klausner, R.D., Cullen, B.R., Chizzonite, R. and Leonard, W.J. (1986) Novel interleukin-2 receptor subunit detected by cross-linking under high-affinity conditions. *Science* **234**, 859-63.
- Shaw**, S. (1995) BP1 concepts in cross-lineage (blind panel) analysis of expression of differentiation antigen. In: *Leukocyte Typing V: White Cell Differentiation Antigens Vol. 1*. (S. F. Schlossman, L. Boumsell, W. Gilks, J. M. Harlan, T. Kashimoto, C. Morimoto, J. Ritz, S. Shaw, R. Silverstein, T. Springer, T. F. Tedder, R. F. Todd, eds.) Oxford University Press, New York, P. 51.
- Shepherd**, V.L., Campbell, E.J., Senior, R.M. and Stahl, P.D. (1982) Characterization of the mannose/fucose receptor on human mononuclear phagocytes. *J Reticuloendothel Soc* **32**, 423-31.
- Siedek**, E.M., Honnah-Symns, N., Fincham, S.C., Mayall, S. and Hamblin, A.S. (2000) Equine macrophage identification with an antibody (Ki-M6) to human CD68 and a new monoclonal antibody (JB10). *J Comp Pathol* **122**, 145-54.
- Simmons**, D.L. and Vernon-Wilson, E. (2002) Structure and function of the signal regulator proteins (SIRPs). In *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uggioni, E. van der Schoot, E. Vivier, and H. Zola, eds.) Oxford University Press, Oxford, p. 35-38.
- Songyang**, Z., Shoelson, S.E., Chaudhuri, M., Gish, G., Pawson, T., Haser, W.G., King, F., Roberts, T., Ratnofsky, S., Lechleider, R.J. and et al. (1993) SH2 domains recognize specific phosphopeptide sequences. *Cell* **72**, 767-78.
- Sopp**, P. and Howard, C.J. (1997) Cross-reactivity of monoclonal antibodies to defined human leucocyte differentiation antigens with bovine cells. *Vet Immunol Immunopathol* **56**, 11-25.
- Springer**, T.A. (1994) Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm. *Cell* **76**, 301-14.
- Springer**, T.A., Teplow, D.B. and Dreyer, W.J. (1985) Sequence homology of the LFA-1 and Mac-1 leukocyte adhesion glycoproteins and unexpected relation to leukocyte interferon. *Nature* **314**, 540-2.
- Stacker**, S.A. and Springer, T.A. (1991) Leukocyte integrin P150,95 (CD11c/CD18) functions as an adhesion molecule binding to a counter-receptor on stimulated endothelium. *J Immunol* **146**, 648-55.
- Stahl**, P. and Gordon, S. (1982) Expression of a mannosyl-fucosyl receptor for endocytosis on cultured primary macrophages and their hybrids. *J Cell Biol* **93**, 49-56.
- Stein**, P.H., Fraser, J.D. and Weiss, A. (1994) The cytoplasmic domain of CD28 is both necessary and sufficient for costimulation of interleukin-2 secretion and association with phosphatidylinositol 3'-kinase. *Mol Cell Biol* **14**, 3392-402.

- Steinbach**, F., Deeg, C., Mael, S. and Wagner, B. (2002) Equine immunology: offspring of the serum horse. *Trends Immunol* **23**, 223-5.
- Steinbach**, F., Stark, R., Ibrahim, S., Gawad, E.A., Ludwig, H., Walter, J., Commandeur, U. and Mael, S. (2005) Molecular cloning and characterization of markers and cytokines for equid myeloid cells. *Vet Immunol Immunopathol* **108**, 227-36.
- Steinbach**, F. and Thiele, B. (1994) Phenotypic investigation of mononuclear phagocytes by flow cytometry. *J Immunol Methods* **174**, 109-22.
- Steinman**, R.M. (1991) The dendritic cell system and its role in immunogenicity. *Annu Rev Immunol* **9**, 271-96.
- Stelter**, F., Bernheiden, M., Menzel, R., Jack, R.S., Witt, S., Fan, X., Pfister, M. and Schutt, C. (1997) Mutation of amino acids 39-44 of human CD14 abrogates binding of lipopolysaccharide and Escherichia coli. *Eur J Biochem* **243**, 100-9.
- Stockinger**, H. (1989) Cluster report: CD68. In: *Leucocyte Typing IV: White Cell Differentiation Antigens*. (W. Knapp, B. Dorken, E.P. Rieber, H. Stein, W.R. Gilks, R.E. Schmidt and A.E.G.Kr. Von dem Borne, eds.) Oxford University Press, Oxford, pp. 841-843.
- Stoll**, M., Dalchau, R., Schmidt, R.E. (1989) Report on the CD44 cluster workshop. In: *Leucocyte Typing IV: white cell differentiation antigens*. (W. Knapp, B. Dorken, E.P. Rieber, H. Stein, W.R. Gilks, R.E. Schmidt and A.E.G.Kr. Von dem Borne, eds.) Oxford University Press, Oxford, p. 619-662.
- Sugamura**, K. (2002) Cluster report: CD25. In: *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uggioni, E. van der Schoot, E. Vivier, and H. Zola, eds.) Oxford University Press, Oxford, p. 770-771.
- Sulahian**, T.H., Hogger, P., Wahner, A.E., Wardwell, K., Goulding, N.J., Sorg, C., Droste, A., Stehling, M., Wallace, P.K., Morganelli, P.M. and Guyre, P.M. (2000) Human monocytes express CD163, which is upregulated by IL-10 and identical to p155. *Cytokine* **12**, 1312-21.
- Szalai**, G., Antczak, D.F., Gerber, H. and Lazary, S. (1994a) Molecular cloning and characterization of horse DQA cDNA. *Immunogenetics* **40**, 457.
- Szalai**, G., Antczak, D.F., Gerber, H. and Lazary, S. (1994b) Molecular cloning and characterization of horse DQB cDNA. *Immunogenetics* **40**, 458.
- Tacke**, M., Hanke, G., Hanke, T. and Hünig, T. (1997) CD28-mediated induction of proliferation in resting T cells in vitro and in vivo without engagement of the T cell receptor: evidence for functionally distinct forms of CD28. *Eur J Immunol* **27**, 239-47.

- Tanaka**, Y. (2002) Cluster report: CD49d. In: *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uguccioni, E. van der Schoot, E. Vivier, and H. Zola, eds). Oxford University Press, Oxford, p. 798-799.
- Tavernor**, A.S., Deverson, E.V., Coadwell, W.J., Lunn, D.P., Zhang, C., Davis, W. and Butcher, G.W. (1993) Molecular cloning of equine CD44 cDNA by a COS cell expression system. *Immunogenetics* **37**, 474-7.
- Tavernor**, A.S., Kydd, J.H., Bodian, D.L., Jones, E.Y., Stuart, D.I., Davis, S.J. and Butcher, G.W. (1994) Expression cloning of an equine T-lymphocyte glycoprotein CD2 cDNA. Structure-based analysis of conserved sequence elements. *Eur J Biochem* **219**, 969-76.
- Tedder**, T.F., Inaoki, M. and Sato, S. (1997) The CD19-CD21 complex regulates signal transduction thresholds governing humoral immunity and autoimmunity. *Immunity* **6**, 107-18.
- Tedder**, T.F., Zhou, L.J. and Engel, P. (1994) The CD19/CD21 signal transduction complex of B lymphocytes. *Immunol Today* **15**, 437-42.
- Thompson**, C.B. (1995) Distinct roles for the costimulatory ligands B7-1 and B7-2 in T helper cell differentiation? *Cell* **81**, 979-82.
- Timens**, W. (2002) Cluster report: CD21. In: *Leucocyte typing VII*. (D. Mason, P. André, A. Bensussan, C. Buckley, C. Civin, E. Clark, M. de Haas, S. Goyert, M. Hadam, D. Hart, V. Horejsi, S. Meurer, J. Morrissey, R. Schwartz-Albiez, S. Shaw, D. Simmons, L. Turni, M. Uguccioni, E. van der Schoot, E. Vivier, and H. Zola, eds.) Oxford University Press, Oxford, p. 766-767.
- Timms**, J.F., Carlberg, K., Gu, H., Chen, H., Kamatkar, S., Nadler, M.J., Rohrschneider, L.R. and Neel, B.G. (1998) Identification of major binding proteins and substrates for the SH2-containing protein tyrosine phosphatase SHP-1 in macrophages. *Mol Cell Biol* **18**, 3838-50.
- Trowbridge**, I.S. and Thomas, M.L. (1994) CD45: an emerging role as a protein tyrosine phosphatase required for lymphocyte activation and development. *Annu Rev Immunol* **12**, 85-116.
- Tumas**, D.B., Brassfield, A.L., Travenor, A.S., Hines, M.T., Davis, W.C. and McGuire, T.C. (1994) Monoclonal antibodies to the equine CD2 T lymphocyte marker, to a pan-granulocyte/monocyte marker and to a unique pan-B lymphocyte marker. *Immunobiology* **192**, 48-64.
- Tuveson**, D.A., Ahearn, J.M., Matsumoto, A.K. and Fearon, D.T. (1991) Molecular interactions of complement receptors on B lymphocytes: a CR1/CR2 complex distinct from the CR2/CD19 complex. *J Exp Med* **173**, 1083-9.
- Van de Velde**, H., von Hoegen, I., Luo, W., Parnes, J.R. and Thielemans, K. (1991) The B-cell surface protein CD72/Lyb-2 is the ligand for CD5. *Nature* **351**, 662-5.

- van den Berg**, T.K., Yoder, J.A. and Litman, G.W. (2004) On the origins of adaptive immunity: innate immune receptors join the tale. *Trends Immunol* **25**, 11-6.
- van den Nieuwenhof**, I.M., Renardel de Lavalette, C., Diaz, N., van Die, I. and van den Berg, T.K. (2001) Differential galactosylation of neuronal and haematopoietic signal regulatory protein-alpha determines its cellular binding-specificity. *J Cell Sci* **114**, 1321-9.
- van Spriel**, A.B., Leusen, J.H., van Egmond, M., Dijkman, H.B., Assmann, K.J., Mayadas, T.N. and van de Winkel, J.G. (2001) Mac-1 (CD11b/CD18) is essential for Fc receptor-mediated neutrophil cytotoxicity and immunologic synapse formation. *Blood* **97**, 2478-86.
- Veillette**, A., Thibaudeau, E. and Latour, S. (1998) High expression of inhibitory receptor SHPS-1 and its association with protein-tyrosine phosphatase SHP-1 in macrophages. *J Biol Chem* **273**, 22719-28.
- Waldmann**, H. (1989) Manipulation of T-cell responses with monoclonal antibodies. *Annu Rev Immunol* **7**, 407-44.
- Waldmann**, T.A. (1986) The structure, function, and expression of interleukin-2 receptors on normal and malignant lymphocytes. *Science* **232**, 727-32.
- Waldmann**, T.A. (1991) The interleukin-2 receptor. *J Biol Chem* **266**, 2681-4.
- Wang**, S., Zhu, G., Tamada, K., Chen, L. and Bajorath, J. (2002) Ligand binding sites of inducible costimulator and high avidity mutants with improved function. *J Exp Med* **195**, 1033-41.
- Werfel**, T., Witter, W. and Gotze, O. (1991) CD11b and CD11c antigens are rapidly increased on human natural killer cells upon activation. *J Immunol* **147**, 2423-7.
- Werners**, A.H., Bull, S., Fink-Gremmels, J. and Bryant, C.E. (2004) Generation and characterization of an equine macrophage cell line (e-CAS cells) derived from equine bone marrow cells. *Vet Immunol Immunopathol* **97**, 65-76.
- Wesch**, D., Beetz, S., Oberg, H.H., Marget, M., Krengel, K. and Kabelitz, D. (2006) Direct costimulatory effect of TLR3 ligand poly(I:C) on human gamma delta T lymphocytes. *J Immunol* **176**, 1348-54.
- Wright**, S.D., Ramos, R.A., Tobias, P.S., Ulevitch, R.J. and Mathison, J.C. (1990) CD14, a receptor for complexes of lipopolysaccharide (LPS) and LPS binding protein. *Science* **249**, 1431-3.
- Yamamoto**, M., Sato, S., Mori, K., Hoshino, K., Takeuchi, O., Takeda, K. and Akira, S. (2002) Cutting edge: a novel Toll/IL-1 receptor domain-containing adapter that preferentially activates the IFN-beta promoter in the Toll-like receptor signaling. *J Immunol* **169**, 6668-72.
- Zarling**, J.M., Clouse, K.A., Biddison, W.E. and Kung, P.C. (1981) Phenotypes of human natural killer cell populations detected with monoclonal antibodies. *J Immunol* **127**, 2575-80.

- Zhang**, C.H., Davis, W.C., Grunig, G. and Antczak, D.F. (1998) The equine homologue of LFA-1 (CD11a/CD18): cellular distribution and differential determinants. *Vet Immunol Immunopathol* **62**, 167-83.
- Zhou**, L.J., Schwarting, R., Smith, H.M. and Tedder, T.F. (1992) A novel cell-surface molecule expressed by human interdigitating reticulum cells, Langerhans cells, and activated lymphocytes is a new member of the Ig superfamily. *J Immunol* **149**, 735-42.
- Zhou**, L.J. and Tedder, T.F. (1995) A distinct pattern of cytokine gene expression by human CD83+ blood dendritic cells. *Blood* **86**, 3295-301.
- Ziegler-Heitbrock**, H.W., Thiel, E., Futterer, A., Herzog, V., Wirtz, A. and Riethmuller, G. (1988) Establishment of a human cell line (Mono Mac 6) with characteristics of mature monocytes. *Int J Cancer* **41**, 456-61.
- Ziegler-Heitbrock**, H.W. and Ulevitch, R.J. (1993) CD14: cell surface receptor and differentiation marker. *Immunol Today* **14**, 121-5.
- Zola**, H., Swart, B., Nicholson, I., Aasted, B., Bensussan, A., Boumsell, L., Buckley, C., Clark, G., Drbal, K., Engel, P., Hart, D., Horejsi, V., Isacke, C., Macardle, P., Malavasi, F., Mason, D., Olive, D., Saalmueller, A., Schlossman, S.F., Schwartz-Albiez, R., Simmons, P., Tedder, T.F., Uguccioni, M. and Warren, H. (2005) CD molecules 2005: human cell differentiation molecules. *Blood* **106**, 3123-6.