

## 8. QUELLENNACHWEIS

- (1) **Ballard DW.** & Bothwell A. (1986) Mutational analysis of the immunoglobulin heavy chain promoter region. *Proc Natl Acad Sci USA* 83: 9626-9630
- (2) **Bargou RC.**, Leng C., Krappmann D. et al. (1996) High-level nuclear NF-kappaB and Oct-2 is a common feature of cultured Hodgkin/Reed-Sternberg cells. *Blood* 87: 4340-4347
- (3) **Bergman Y.**, Rice D., Grosschedl R. et al. (1984) Two regulatory elements for immunoglobulin κ light chain gene expression. *Proc Natl Acad Sci USA* 81: 7041-7045
- (4) **Botfield MC.**, Jansco A., Weiss MA. (1994) An invariant asparagine in the POU-specific homeodomain regulates the specificity of the Oct-2 POU motif. *Biochemistry* 33: 8113-8121
- (5) **Bräuninger A.**, Hansmann ML., Strickler JG. et al. (1999) Identification of common Germinal Centre B-Cell Precursors in two Patients with both Hodgkin's Disease and Non Hodgkin Lymphoma. *N Engl J Med* 340: 1239-1247
- (6) **Canellos GP.**, Lister TA. and Sclar J. (1998) The Lymphomas. Philadelphia: WB Saunders: 287
- (7) **Cepek KL.**, Chasman PI., Sharp PA. (1996) Sequence-specific DNA binding of the B-cell specific co-activator OCA-B. *Genes & Dev* 10: 2079-2088
- (8) **Cordell JL.**, Falini B., Erber WN. et al. (1984) Immunoenzymatic labelling of monoclonal antibodies using immune complexes of alkaline phosphatase and monoclonal anti-alkaline phosphatase (APAAP complexes). *J Histochem Cytochem* 32: 219-229
- (9) **Dallenbach FE** & Stein H (1989) Expression of T-cell-receptor β chain in Reed-Sternberg cells. *Lancet* 2: 828-830
- (10) **Delabie J.**, Tierens A., Gavriil T. et al. (1996) Phenotype, genotype and clonality of Reed-Sternberg cells in nodular sclerosis Hodgkin's disease: Results of a single cell study. *Br J Haematol* 94: 198-205

- (11) **Diehl V.**, Kirchner HH., Schaadt M. et al. (1981) Hodgkin's disease: establishment and characterisation of four in vitro cell lines. *J Cancer Res Clin Oncol* 101: 111-124
- (12) **Dreyfus M.**, Doyen N., Rougeon F. (1987) The conserved decanucleotide from the immunoglobulin heavy chain promoter induces a very high transcriptional activity in B-cells when introduced into heterologous promoter. *EMBO J* 6: 1685-1690
- (13) **Eaton S.** & Calame K. (1987) Multiple DNA sequence elements are necessary for the function of an immunoglobulin heavy chain promoter. *Proc Natl Acad Sci USA* 84: 7634-7638
- (14) **Falini B.**, Stein H., Pileri S. et al. (1987) Expression of lymphoid-associated antigens on Hodgkin's and Reed-Sternberg cells of Hodgkin's disease. An immunocytochemical study on lymph node cytopsins using monoclonal antibodies. *Histopathology* 11: 1229-1242
- (15) **Feldhaus AL.**, Klug CA., Arvin K. et al. (1993) Targeted disruption of the Oct-2 locus in a B-cell provides genetic evidence for two distinct cell-type specific pathways of Octamer-element mediated gene activation. *EMBO J* 12: 2763-2772
- (16) **Freedman AS. & Nadler LM.** (1998) Harrison's Principles of Internal Medicine. McGraw-Hill Companies Inc., New York 14<sup>th</sup> Edition, Editors: Fauci AS., Braunwald E., Isselbacher KJ. et al.: 707-712
- (17) **Gstaiger M.**, Knoepfel L., Georgiev O. et al. (1995) A B cell coactivator of Octamer-binding transcription factors. *Nature* 373: 360-362
- (18) **Haino M.**, Hayashida H., Miyata T. et al. (1994) Comparison and evolution of human immunoglobulin VH segments located in the 3' 0.8-megabase region: evidence for unidirectional transfer of segmental gene sequences. *J Biol Chem* 269: 2619-2626
- (19) **Harris NL.**, Jaffe ES., Stein H. et al. (1994) A revised European-American classification of lymphoid neoplasm's: a proposal from the International Lymphoma Study Group. *Blood* 84: 1361-1392
- (20) **Hodgkin T.** (1832) On same morbid appearances of the absorbent glands and spleens. *Med Chir Trans* 17: 68-114

- (21) **Hummel M.**, Anagnostopoulos I., Dallenbach F. et al. (1992) EBV Infection patterns in Hodgkins disease and normal lymphoid tissue: expression and cellular localisation of EBV gene products. *Br J Haematol* 82: 689-694
- (22) **Hummel M.**, Ziemann K., Lammert H. et al. (1995) Hodgkin's disease with monoclonal and polyclonal populations of Reed-Sternberg cells. *N Engl J Med* 333: 901-906
- (23) **Hummel M.**; Marafioti T., Stein H. (1999) Clonality of Reed-Sternberg cells in Hodgkin's disease. *N Engl J Med* 340: 394-395
- (24) **Hummel M.** & Stein H. (2000) Clinical relevance of immunoglobulin mutation analysis. *Curr Opin Oncol* 12: 395-402
- (25) **Janeway CA.** & Travers PA. (1995) Immunologie. *Spectrum Akademischer Verlag Heidelberg • Berlin • Oxford*
- (26) **Jenuwein T.** & Grosschedl R. (1991) Complex Pattern of Immunoglobulin  $\mu$  gene Expression in normal and transgenic mice: nonoverlapping regulatory sequences govern distinct tissue specificities. *Genes & Dev* 5: 932-943
- (27) **Jox A.**, Zander T., Küppers R. et al. (1999) Somatic mutations within the untranslated regions of rearranged Ig genes in a case of classical Hodgkin's disease as a potential cause for the absence of Ig in lymphoma cells. *Blood* 93: 1-10
- (28) **Kanzler H.**, Küppers R., Hansmann ML. et al. (1996) Hodgkin and Reed-Sternberg cells in Hodgkin's disease represent the outgrowth of a dominant tumor clone derived from (crippled) germinal centre B cells. *J Exp Med* 184: 1495-1505
- (29) **Kim U.**, Qin XF., Gong S. et al. (1996) The B-cell-specific transcription co-activator OCA-B/OBF-1/BOB-1 is essential for normal production of immunoglobulin isotypes. *Nature* 383: 542-547
- (30) **Klein U.**, Goossens T., Fischer M. et al. (1998) Somatic hypermutation in normal and transformed human B cells. *Immunol Rev* 162: 261-280
- (31) **Küppers R.**, Rajewsky K., Zhao M. et al. (1994) Hodgkin's disease: Hodgkin and Reed-Sternberg cells picked from histological sections show clonal immunoglobulin

- gene rearrangements and appear to be derived from B cells at various stages of development. *Proc Natl Acad Sci USA* 91: 10962-10966
- (32) **Küppers R.** & Rajewsky K. (1998) The origin of Hodgkin and Reed/Sternberg cells in Hodgkin's disease. *Annu Rev Immunol* 16: 471-493
  - (33) **Küppers R.**, Klein U., Hansmann ML. et al. (1999) Cellular origin of human B-cell lymphomas. *N Engl J Med* 341: 1520-1529
  - (34) **Lefranc MP.** & Lefranc G. (2001) "The Immunoglobulin Factsbook", *Academic Press, Hartcourt Publishers Limited, London, UK*: 1-458
  - (35) **Lukes RJ.** & Butler JJ. (1966) The pathology and nomenclature of Hodgkin's disease. *Cancer Res* 26: 1063-1083
  - (36) **Luo Y.** & Roeder RG. (1995) Cloning, functional characterisation, and mechanism of action of the B-cell-specific transcriptional co-activator OCA-B. *Mol Cell Biol* 15: 4115-4124
  - (37) **Marafioti T.**, Hummel M., Anagnostopoulos I. et al. (1997) Origin of nodular lymphocyte predominant Hodgkin's disease from a clonal expansion of highly mutated germinal-centre B cells. *N Engl J Med* 337: 453-458
  - (38) **Marafioti T.**, Hummel M., Anagnostopoulos I. et al. (1999) Classical Hodgkin's disease and follicular lymphoma originating from the same germinal centre B-cell. *J Clin Oncol* 17: 3804-3809
  - (39) **Marafioti T.**, Hummel M., Foss HD. et al. (2000) Hodgkin and Reed-Sternberg cells represent an expansion of a single clone originating from a germinal centre B-cell with functional immunoglobulin gene rearrangements but defective immunoglobulin transcription. *Blood* 95: 1443-1450
  - (40) **Matsuda F.**, Ishii K., Bourvagnet P. et al. (1998) The complete nucleotide sequence of the human immunoglobulin heavy chain variable region locus. *J Exp Med* 188: 2151-2162
  - (41) **Matthias P.** (1998) Lymphoid-specific transcription mediated by the conserved Octamer site: Who is doing what? *Semin Immunol* 10: 155-163

- (42) **Papadimitriou CS.**, Stein H., Lennert K. (1978) The complexity of immunohistochemical staining pattern of Hodgkin and Reed-Sternberg cells – Demonstration of immunoglobulin, albumin, alpha-1 antichymotrypsin and lysozyme. *Int J Cancer* 21: 531-541
- (43) **Parslow TG.**, Blair DL., Murphy WJ. et al. (1984) Structure of the 5' ends of the immunoglobulin genes: a novel conserved sequence. *Proc Natl Acad Sci USA* 81: 2650-2654
- (44) **Pinkus GS.** & Said JW. (1988) Hodgkin's disease, lymphocyte predominant type, nodular – further evidence for a B cell derivation. L & H variants of Reed-Sternberg cells express L26, a pan B cell marker. *Am J Pathol* 133: 211-217
- (45) **Pinkus GS.**, Pinkus Jl., Langhoff E. et al. (1997) Fascin, a sensitive new marker for Reed-Sternberg cells of Hodgkins disease. Evidence for a dendritic or B cell derivation? *Am J Pathol* 150: 543-562
- (46) **Poppema S.** (1980) The diversity of immunohistological staining pattern of Sternberg-Reed cells. *J Histochem Cytochem* 28: 788-791
- (47) **Rajewsky K.** (1996) Clonal selection and learning in the antibody system. *Nature* 351: 751-758
- (48) **Re D.**, Müschen M., Ahmadi T. et al. (2001) (a) Oct-2 and Bob-1 Deficiency in Hodgkin and Reed Sternberg cells. *Cancer Res* 61:2080-2084
- (49) **Re D.**, Staratschek-Jox A., Holtick U. et al. (2001) (b) Deregulation of immunoglobulin gene transcription in the Hodgkin-Reed Sternberg cell line L1236. *Br J Haematol* 115: 326-328
- (50) **Reed D.** (1902) On the pathological changes in Hodgkin's disease, with special reference to its relationship to tuberculosis. *J Hopkins Hosp Rep* 10: 133-196
- (51) **Regula DP Jr.**, Hoppe RT., Weiss LM. (1988) Nodular and diffuse types of lymphocyte predominance Hodgkins disease. *N Engl J Med* 318: 214-219
- (52) **Roth J.**, Daus H., Trümper L. et al. (1994) Detection of immunoglobulin heavy-chain gene rearrangement at the single-cell level in malignant lymphomas: No rearrangement is found in Hodgkin and Reed-Sternberg cells. *Int J Cancer* 57: 799-804

- (53) **Ryan AK.** & Rosenfeld MG. (1997) POU domain family values: flexibility, partnerships, and developmental codes. *Genes & Dev* 11: 1207-1225
- (54) **Saiki RK.**, Gelfand DH., Stoffel S. et al. (1988) Primer directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239: 487-491
- (55) **Sauter P.** & Matthias P. (1998) Coactivator OBF-1 makes selective contacts with both the POU-specific domain and the POU-homeodomain and acts as a molecular clamp on DNA. *Mol Cell Biol* 18: 7397-7409
- (56) **Scheidereit C.**, Heguy A., Roeder RG. (1987) Identification and purification of a human lymphoid-specific Octamer binding protein (OTF-2) that activates transcription of an immunoglobulin promoter in vitro. *Cell* 51: 783-793
- (57) **Schubart K.**, Massa S., Schubart D. et al. (2001) B-cell development and immunoglobulin transcription in the absence of Oct-2 and OBF-1. *Nat Immunol* 2: 69-74
- (58) **Schwab U.**, Stein H., Gerdes J. et al. (1982) Production of a monoclonal antibody specific for Sternberg-Reed cells of Hodgkin's disease and a subset of normal lymphoid cells. *Nature* 299: 65-67
- (59) **Seitz V.**, Hummel M., Marafioti T. et al. (2000) Detection of clonal T-cell receptor gamma-chain gene rearrangements in Reed-Sternberg cells of classic Hodgkin disease. *Blood* 95: 3020-3024
- (60) **Shah PC.**, Bertolino E., Singh H. (1997) Using altered specificity Oct-1 and Oct-2 mutants to analyse the regulation of immunoglobulin gene transcription. *EMBO J* 16: 7105-7117
- (61) **Staudt LM** (1991) Immunoglobulin gene transcription. *Annu Rev Immunol* 9: 373-398
- (62) **Stein H.**, Gerdes J., Kirchner H. et al. (1981) Immunohistological analysis of Hodgkin's and Sternberg-Reed cells: Detection of a new antigen and evidence for selective IgG uptake in the absence of B cell, T cell and histiocytic markers. *J Cancer Res Clin Oncol* 101: 125-134

- (63) **Stein H.**, Gerdes J., Schwab U. et al. (1982) Identification of Hodgkin and Sternberg-Reed cells as a unique cell type derived from a newly-detected small-cell population. *Int J Cancer* 30: 445-459
- (64) **Stein H.**, Hansmann ML., Lennert K. et al. (1986) Reed-Sternberg and Hodgkin cells in lymphocyte-predominant Hodgkin's disease of nodular subtype contain J chain. *Am J Clin Pathol* 86: 292-297
- (65) **Stein H.**, Hummel M., Marafioti T. et al. (1997) Hodgkin's disease: Its mystery is being revealed. *Verh Dtsch Ges Path* 81: 327-338
- (66) **Stein H.** & Hummel M. (1999) Cellular origin and clonality of classic Hodgkin's lymphoma: immunophenotypic and molecular studies. *Semin Haematol* 36: 233-241
- (67) **Stein H.**, Marafioti T., Foss HD. et al. (2001) Down-regulation of BOB.1/OBF.1 and Oct-2 in classical Hodgkin disease but not in lymphocyte predominant Hodgkin disease correlates with immunoglobulin transcription. *Blood* 97: 496-501
- (68) **Stein H.**, Delsol G., Pileri S et al. (2001a) WHO classification of tumours – Pathology and genetics of tumours of haematopoietic and lymphoid tissues. Editors: Jaffe ES., Harris NL., Stein H. and Vardiman JW. *IARC Press Lyon*: 239-253
- (69) **Sternberg C.** (1898) Über eine eigenartige unter dem Bilde der Pseudoleukämie verlaufende Tuberkulose des lymphatischen Apparates. *Z Heilkunde* 19: 21-90
- (70) **Strubin M.**, Newell JW., Matthias P. (1995) OBF-1, a novel b-cell-specific coactivator that stimulates immunoglobulin promoter activity through association with Octamer-binding proteins. *Cell* 80: 497-506
- (71) **Sturm RA.**, Das G., Herr W. (1988) The ubiquitous Octamer-binding protein Oct-1 contains a POU domain with a homeobox subdomain. *Genes & Dev* 2: 1582-1599
- (72) **Tamaru J.**, Hummel M., Zemlin M. et al. (1994) Hodgkin's disease with a B cell phenotype often shows a VDJ rearrangement and somatic mutations in the VH genes. *Blood* 84: 708-715
- (73) **Tamaru J.**, Hummel M., Marafioti T. et al. (1995) Burkitt's lymphomas express VH genes with a moderate number of antigen-selected somatic mutations. *Am J Pathol* 147: 1398-1407

- (74) **Tang H.** & Sharp PA. (1999) Transcriptional regulation of the murine 3' IgH enhancer by Oct-2. *Immunity* 11: 517-526
- (75) **Theil J.**, Laumen H., Marafioti T. et al. (2001) Defective Octamer-dependent transcription is responsible for silenced immunoglobulin transcription in Reed-Sternberg cells. *Blood* 97: 3191-3196
- (76) **Tomlinson IM.**, Walter G., Marks JD. et al. (1992) The repertoire of human germline VH sequences reveals about fifty groups of VH segments with different hypervariable loops. *J Mol Biol* 227: 776-798
- (77) **Trümpfer LH.**, Brady G., Bagg A. et al. (1993) Single-cell analysis of Hodgkin and Reed-Sternberg cells: molecular heterogeneity of gene expression and p53 mutations. *Blood* 81: 3097-3115
- (78) **Verrijzer CP.**, Alkema MJ., van Weperen WW. et al. (1992) The DNA binding specificity of the biparte POU domain and its subdomains. *EMBO J* 11: 4993-5003
- (79) Gendatenbank des DKFZ (Deutsches Krebsforschungszentrum) Heidelberg und National Center of Biotechnology Information der USA mit Gene Bank, Ref Seq und PDB Datenbank ([www.ncbi.nlm.nih.gov.BLAST](http://www.ncbi.nlm.nih.gov.BLAST))