

6. Bibliographie

1. UNAIDS/WHO. AIDS epidemic update: 2006. Geneva: Joint United Nations programme on HIV/AIDS (UNAIDS) and World Health Organization (WHO), 2006.
2. Barre-Sinoussi F, Chermann JC, Rey F, Nugeyre MT, Chamaret S, Gruest J, et al. Isolation of a T-lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS). *Science* 1983;220(4599):868-71.
3. Gallo RC, Sarin PS, Gelmann EP, Robert-Guroff M, Richardson E, Kalyanaraman VS, et al. Isolation of human T-cell leukemia virus in acquired immune deficiency syndrome (AIDS). *Science* 1983;220(4599):865-7.
4. UNAIDS/WHO. AIDS epidemic update: 2003. Geneva: Joint United Nations programme on HIV/AIDS (UNAIDS) and World Health Organization (WHO), 2003.
5. AIDS Vaccine Trials - Getting the Global House in Order: AVAC, 2004.
6. Cohen J. AIDS vaccine trial produces disappointment and confusion. *Science* 2003(299):1290-91.
7. Singh M. No vaccine against HIV yet--are we not perfectly equipped? *Virology* 2006;3:60.
8. Das erworbene Immunschwächesyndrom. In: Janeway CA, Travers, P., Walport, M., Shlomchik, M., editor. *Immunologie*. 5. ed. Heidelberg-Berlin: Spektrum Akademischer Verlag, 2002:484-499.
9. Rubbert A. Pathophysiologie der HIV-Infektion. In: Kamps BS, editor. *HIV.NET2005*. Wuppertal: Steinhäuser Verlag, 2005:58-79.

10. Dalglish AG, Beverley PC, Clapham PR, Crawford DH, Greaves MF, Weiss RA. The CD4 (T4) antigen is an essential component of the receptor for the AIDS retrovirus. *Nature* 1984;312(5996):763-7.
11. Klatzmann D, Barre-Sinoussi F, Nugeyre MT, Danquet C, Vilmer E, Griscelli C, et al. Selective tropism of lymphadenopathy associated virus (LAV) for helper-inducer T lymphocytes. *Science* 1984;225(4657):59-63.
12. Janeway CA, Travers, P., Walport, M., Shlomchik, M. Die angeborene Immunität. In: Janeway CA, Travers, P., Walport, M., Shlomchik, M., editor. *Immunologie*. 5. Auflage ed. Heidelberg-Berlin: Spektrum Akademischer Verlag, 2002:75-78.
13. Maddon PJ, Dalglish AG, McDougal JS, Clapham PR, Weiss RA, Axel R. The T4 gene encodes the AIDS virus receptor and is expressed in the immune system and the brain. *Cell* 1986;47(3):333-48.
14. Chesebro B, Buller R, Portis J, Wehrly K. Failure of human immunodeficiency virus entry and infection in CD4-positive human brain and skin cells. *J Virol* 1990;64(1):215-21.
15. Ashorn PA, Berger EA, Moss B. Human immunodeficiency virus envelope glycoprotein/CD4-mediated fusion of nonprimate cells with human cells. *J Virol* 1990;64(5):2149-56.
16. Tersmette M, de Goede RE, Al BJ, Winkel IN, Gruters RA, Cuypers HT, et al. Differential syncytium-inducing capacity of human immunodeficiency virus isolates: frequent detection of syncytium-inducing isolates in patients with acquired immunodeficiency syndrome (AIDS) and AIDS-related complex. *J Virol* 1988;62(6):2026-32.
17. Schuitemaker H, Kootstra NA, de Goede RE, de Wolf F, Miedema F, Tersmette M. Monocytotropic human immunodeficiency virus type 1 (HIV-1) variants detectable in all stages of HIV-1 infection lack T-cell line tropism and syncytium-inducing ability in primary T-cell culture. *J Virol* 1991;65(1):356-63.

18. Schuitemaker H, Koot M, Kootstra NA, Dercksen MW, de Goede RE, van Steenwijk RP, et al. Biological phenotype of human immunodeficiency virus type 1 clones at different stages of infection: progression of disease is associated with a shift from monocyctotropic to T-cell-tropic virus population. *J Virol* 1992;66(3):1354-60.
19. Feng Y, Broder CC, Kennedy PE, Berger EA. HIV-1 entry cofactor: functional cDNA cloning of a seven-transmembrane, G protein-coupled receptor. *Science* 1996;272(5263):872-7.
20. Loetscher M, Geiser T, O'Reilly T, Zwahlen R, Baggiolini M, Moser B. Cloning of a human seven-transmembrane domain receptor, LESTR, that is highly expressed in leukocytes. *J Biol Chem* 1994;269(1):232-7.
21. Bleul CC, Farzan M, Choe H, Parolin C, Clark-Lewis I, Sodroski J, et al. The lymphocyte chemoattractant SDF-1 is a ligand for LESTR/fusin and blocks HIV-1 entry. *Nature* 1996;382(6594):829-33.
22. Oberlin E, Amara A, Bachelier F, Bessia C, Virelizier JL, Arenzana-Seisdedos F, et al. The CXC chemokine SDF-1 is the ligand for LESTR/fusin and prevents infection by T-cell-line-adapted HIV-1. *Nature* 1996;382(6594):833-5.
23. Cocchi F, DeVico AL, Garzino-Demo A, Arya SK, Gallo RC, Lusso P. Identification of RANTES, MIP-1 alpha, and MIP-1 beta as the major HIV-suppressive factors produced by CD8+ T cells. *Science* 1995;270(5243):1811-5.
24. Alkhatib G, Combadiere C, Broder CC, Feng Y, Kennedy PE, Murphy PM, et al. CC CKR5: a RANTES, MIP-1alpha, MIP-1beta receptor as a fusion cofactor for macrophage-tropic HIV-1. *Science* 1996;272(5270):1955-8.
25. Choe H, Farzan M, Sun Y, Sullivan N, Rollins B, Ponath PD, et al. The beta-chemokine receptors CCR3 and CCR5 facilitate infection by primary HIV-1 isolates. *Cell* 1996;85(7):1135-48.
26. Deng H, Liu R, Ellmeier W, Choe S, Unutmaz D, Burkhart M, et al. Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 1996;381(6584):661-6.

27. Dragic T, Litwin V, Allaway GP, Martin SR, Huang Y, Nagashima KA, et al. HIV-1 entry into CD4+ cells is mediated by the chemokine receptor CC-CKR-5. *Nature* 1996;381(6584):667-73.
28. Doranz BJ, Rucker J, Yi Y, Smyth RJ, Samson M, Peiper SC, et al. A dual-tropic primary HIV-1 isolate that uses fusin and the beta-chemokine receptors CKR-5, CKR-3, and CKR-2b as fusion cofactors. *Cell* 1996;85(7):1149-58.
29. Samson M, Labbe O, Mollereau C, Vassart G, Parmentier M. Molecular cloning and functional expression of a new human CC-chemokine receptor gene. *Biochemistry* 1996;35(11):3362-7.
30. Dean M, Carrington M, Winkler C, Huttley GA, Smith MW, Allikmets R, et al. Genetic restriction of HIV-1 infection and progression to AIDS by a deletion allele of the CKR5 structural gene. Hemophilia Growth and Development Study, Multicenter AIDS Cohort Study, Multicenter Hemophilia Cohort Study, San Francisco City Cohort, ALIVE Study. *Science* 1996;273(5283):1856-62.
31. Liu R, Paxton WA, Choe S, Ceradini D, Martin SR, Horuk R, et al. Homozygous defect in HIV-1 coreceptor accounts for resistance of some multiply-exposed individuals to HIV-1 infection. *Cell* 1996;86(3):367-77.
32. Huang Y, Paxton WA, Wolinsky SM, Neumann AU, Zhang L, He T, et al. The role of a mutant CCR5 allele in HIV-1 transmission and disease progression. *Nat Med* 1996;2(11):1240-3.
33. Connor RI, Paxton WA, Sheridan KE, Koup RA. Macrophages and CD4+ T lymphocytes from two multiply exposed, uninfected individuals resist infection with primary non-syncytium-inducing isolates of human immunodeficiency virus type 1. *J Virol* 1996;70(12):8758-64.
34. Murphy PM. The molecular biology of leukocyte chemoattractant receptors. *Annu Rev Immunol* 1994;12:593-633.

35. Menten P, Wuyts A, Van Damme J. Macrophage inflammatory protein-1. *Cytokine Growth Factor Rev* 2002;13(6):455-81.
36. D'Souza MP, Harden VA. Chemokines and HIV-1 second receptors. Confluence of two fields generates optimism in AIDS research. *Nat Med* 1996;2(12):1293-300.
37. Premack BA, Schall TJ. Chemokine receptors: gateways to inflammation and infection. *Nat Med* 1996;2(11):1174-8.
38. Ward SG, Westwick J. Chemokines: understanding their role in T-lymphocyte biology. *Biochem J* 1998;333 (Pt 3):457-70.
39. Mason D, Andre P, Bensussan A, Buckley C, Civin C, Clark E, et al. CD antigens 2001. *J Leukoc Biol* 2001;70(5):685-90.
40. Lu Z, Berson JF, Chen Y, Turner JD, Zhang T, Sharron M, et al. Evolution of HIV-1 coreceptor usage through interactions with distinct CCR5 and CXCR4 domains. *Proc Natl Acad Sci U S A* 1997;94(12):6426-31.
41. Bleul CC, Wu L, Hoxie JA, Springer TA, Mackay CR. The HIV coreceptors CXCR4 and CCR5 are differentially expressed and regulated on human T lymphocytes. *Proc Natl Acad Sci U S A* 1997;94(5):1925-30.
42. Bleul CC, Fuhlbrigge RC, Casasnovas JM, Aiuti A, Springer TA. A highly efficacious lymphocyte chemoattractant, stromal cell-derived factor 1 (SDF-1). *J Exp Med* 1996;184(3):1101-9.
43. Nagasawa T, Hirota S, Tachibana K, Takakura N, Nishikawa S, Kitamura Y, et al. Defects of B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the CXC chemokine PBSF/SDF-1. *Nature* 1996;382(6592):635-8.
44. Lee B, Sharron M, Blanpain C, Doranz BJ, Vakili J, Setoh P, et al. Epitope mapping of CCR5 reveals multiple conformational states and distinct but overlapping structures involved in chemokine and coreceptor function. *J Biol Chem* 1999;274(14):9617-26.

45. Horuk R. Chemokine receptors and HIV-1: the fusion of two major research fields. *Immunol Today* 1999;20(2):89-94.
46. Loetscher P, Uguccioni M, Bordoli L, Baggiolini M, Moser B, Chizzolini C, et al. CCR5 is characteristic of Th1 lymphocytes. *Nature* 1998;391(6665):344-5.
47. Chemokine/chemokine receptor nomenclature. *Cytokine* 2003;21(1):48-9.
48. Blanpain C, Migeotte I, Lee B, Vakili J, Doranz BJ, Govaerts C, et al. CCR5 binds multiple CC-chemokines: MCP-3 acts as a natural antagonist. *Blood* 1999;94(6):1899-905.
49. Zhu T, Mo H, Wang N, Nam DS, Cao Y, Koup RA, et al. Genotypic and phenotypic characterization of HIV-1 patients with primary infection. *Science* 1993;261(5125):1179-81.
50. van't Wout AB, Kootstra NA, Mulder-Kampinga GA, Albrecht-van Lent N, Scherpbier HJ, Veenstra J, et al. Macrophage-tropic variants initiate human immunodeficiency virus type 1 infection after sexual, parenteral, and vertical transmission. *J Clin Invest* 1994;94(5):2060-7.
51. Connor RI, Sheridan KE, Ceradini D, Choe S, Landau NR. Change in coreceptor use correlates with disease progression in HIV-1--infected individuals. *J Exp Med* 1997;185(4):621-8.
52. de Roda Husman AM, van Rij RP, Blaak H, Broersen S, Schuitemaker H. Adaptation to promiscuous usage of chemokine receptors is not a prerequisite for human immunodeficiency virus type 1 disease progression. *J Infect Dis* 1999;180(4):1106-15.
53. Lee B, Montaner LJ. Chemokine immunobiology in HIV-1 pathogenesis. *J Leukoc Biol* 1999;65(5):552-65.
54. Mowat AM, Viney JL. The anatomical basis of intestinal immunity. *Immunol Rev* 1997;156:145-66.

55. Ullrich R, Schmidt W, Zippel T, Schneider T, Zeitz M, Riecken EO. Mucosal HIV infection. *Pathobiology* 1998;66(3-4):145-50.
56. Pabst R. Das darmassoziierte lymphatische Gewebe. In: Zenker W, editor. *Benninghoff Anatomie*. 15. ed. München; Wien; Baltimore: Urban&Schwarzenberg, 1994:751-53.
57. Veazey RS, Lackner A. The mucosal immune system and HIV-1 infection. *AIDS Rev* 2003;5:245-252.
58. Zeitz M, Ullrich R, Schneider T, Schieferdecker HL, Riecken EO. Cell differentiation and proliferation in the gastrointestinal tract with respect to the local immune system. *Ann N Y Acad Sci* 1994;733:75-86.
59. Brandtzaeg P, Farstad IN, Helgeland L. Phenotypes of T cells in the gut. *Chem Immunol* 1998;71:1-26.
60. Ullrich R, Schieferdecker HL, Jahn HU, Zeitz M. Gamma-delta T cells in the human intestine. *Immunol Res* 1991;10(3-4):306-9.
61. Jarry A, Cerf-Bensussan N, Brousse N, Selz F, Guy-Grand D. Subsets of CD3+ (T cell receptor alpha/beta or gamma/delta) and CD3- lymphocytes isolated from normal human gut epithelium display phenotypical features different from their counterparts in peripheral blood. *Eur J Immunol* 1990;20(5):1097-103.
62. Zeitz M, Schieferdecker HL, Ullrich R, Jahn HU, James SP, Riecken EO. Phenotype and function of lamina propria T lymphocytes. *Immunol Res* 1991;10(3-4):199-206.
63. Zeitz M, Schieferdecker HL, James SP, Riecken EO. Special functional features of T-lymphocyte subpopulations in the effector compartment of the intestinal mucosa and their relation to mucosal transformation. *Digestion* 1990;46 Suppl 2:280-9.

64. Schieferdecker HL, Ullrich R, Hirseland H, Zeitz M. T cell differentiation antigens on lymphocytes in the human intestinal lamina propria. *J Immunol* 1992;149(8):2816-22.
65. Cerf-Bensussan N, Jarry A, Brousse N, Lisowska-Grospierre B, Guy-Grand D, Griscelli C. A monoclonal antibody (HML-1) defining a novel membrane molecule present on human intestinal lymphocytes. *Eur J Immunol* 1987;17(9):1279-85.
66. Larson RS, Springer TA. Structure and function of leukocyte integrins. *Immunol Rev* 1990;114:181-217.
67. Lundqvist C, Baranov V, Hammarstrom S, Athlin L, Hammarstrom ML. Intra-epithelial lymphocytes. Evidence for regional specialization and extrathymic T cell maturation in the human gut epithelium. *Int Immunol* 1995;7(9):1473-87.
68. Brandtzaeg P, Halstensen TS, Kett K, Krajci P, Kvale D, Rognum TO, et al. Immunobiology and immunopathology of human gut mucosa: humoral immunity and intraepithelial lymphocytes. *Gastroenterology* 1989;97(6):1562-84.
69. Sarnacki S, Begue B, Jarry A, Cerf-Bensussan N. Human intestinal intraepithelial lymphocytes, a distinct population of activated T cells. *Immunol Res* 1991;10(3-4):302-5.
70. Ullrich R, Schieferdecker HL, Ziegler K, Riecken EO, Zeitz M. gamma delta T cells in the human intestine express surface markers of activation and are preferentially located in the epithelium. *Cell Immunol* 1990;128(2):619-27.
71. Ebert EC, Roberts AI. Costimulation of the CD3 pathway by CD28 ligation in human intestinal lymphocytes. *Cell Immunol* 1996;171(2):211-6.
72. Cepek KL, Shaw SK, Parker CM, Russell GJ, Morrow JS, Rimm DL, et al. Adhesion between epithelial cells and T lymphocytes mediated by E-cadherin and the alpha E beta 7 integrin. *Nature* 1994;372(6502):190-3.

73. Cerf-Bensussan N, Guy-Grand D, Griscelli C. Intraepithelial lymphocytes of human gut: isolation, characterisation and study of natural killer activity. *Gut* 1985;26(1):81-8.
74. Selby WS, Janossy G, Bofill M, Jewell DP. Lymphocyte subpopulations in the human small intestine. The findings in normal mucosa and in the mucosa of patients with adult coeliac disease. *Clin Exp Immunol* 1983;52(1):219-28.
75. Cerf-Bensussan N, Jarry A, Gneragbe T, Brousse N, Lisowska-Grospierre B, Griscelli C, et al. Monoclonal antibodies specific for intestinal lymphocytes. *Monogr Allergy* 1988;24:167-72.
76. Ebert EC. Proliferative responses of human intraepithelial lymphocytes to various T-cell stimuli. *Gastroenterology* 1989;97(6):1372-81.
77. Ebert EC. Activation of human intraepithelial lymphocytes reduces CD3 expression. *Clin Exp Immunol* 2003;132(3):424-9.
78. Ebert EC, Roberts AI, Brolin RE, Raska K. Examination of the low proliferative capacity of human jejunal intraepithelial lymphocytes. *Clin Exp Immunol* 1986;65(1):148-57.
79. Lefrancois L, Puddington, L. Basic aspects of intraepithelial lymphocyte immunobiology. In: Ogra PL, Mestecky J., Lamm M.E., Strober W., Bienstock J., McGhee, J.R., editor. *Mucosal Immunology*. 2. ed: San Diego Academic Press, 1999:977-899.
80. Hayday A, Theodoridis E, Ramsburg E, Shires J. Intraepithelial lymphocytes: exploring the Third Way in immunology. *Nat Immunol* 2001;2(11):997-1003.
81. Bucy RP, Chen CL, Cooper MD. Tissue localization and CD8 accessory molecule expression of T gamma delta cells in humans. *J Immunol* 1989;142(9):3045-9.
82. Brandtzaeg P, Bosnes V, Halstensen TS, Scott H, Sollid LM, Valnes KN. T lymphocytes in human gut epithelium preferentially express the alpha/beta

- antigen receptor and are often CD45/UCHL1-positive. *Scand J Immunol* 1989;30(1):123-8.
83. Cerf-Bensussan N, Guy-Grand D. Intestinal intraepithelial lymphocytes. *Gastroenterol Clin North Am* 1991;20(3):549-76.
84. Groh V, Porcelli S, Fabbi M, Lanier LL, Picker LJ, Anderson T, et al. Human lymphocytes bearing T cell receptor gamma/delta are phenotypically diverse and evenly distributed throughout the lymphoid system. *J Exp Med* 1989;169(4):1277-94.
85. Sillett HK, Southgate J, Howdle PD, Trejdosiewicz LK. Expression of activation and costimulatory elements by human intestinal intraepithelial lymphocytes. *Scand J Immunol* 1999;50(1):52-60.
86. Schon MP, Arya A, Murphy EA, Adams CM, Strauch UG, Agace WW, et al. Mucosal T lymphocyte numbers are selectively reduced in integrin alpha E (CD103)-deficient mice. *J Immunol* 1999;162(11):6641-9.
87. Lefrancois L, Parker CM, Olson S, Muller W, Wagner N, Schon MP, et al. The role of beta7 integrins in CD8 T cell trafficking during an antiviral immune response. *J Exp Med* 1999;189(10):1631-8.
88. Ericsson A, Svensson M, Arya A, Agace WW. CCL25/CCR9 promotes the induction and function of CD103 on intestinal intraepithelial lymphocytes. *Eur J Immunol* 2004;34(10):2720-9.
89. Hadley G. Role of integrin CD103 in promoting destruction of renal allografts by CD8 T cells. *Am J Transplant* 2004;4(7):1026-32.
90. Lundqvist C, Melgar S, Yeung MM, Hammarstrom S, Hammarstrom ML. Intraepithelial lymphocytes in human gut have lytic potential and a cytokine profile that suggest T helper 1 and cytotoxic functions. *J Immunol* 1996;157(5):1926-34.

91. Chang F, Mahadeva U, Deere H. Pathological and clinical significance of increased intraepithelial lymphocytes (IELs) in small bowel mucosa. *Apmis* 2005;113(6):385-99.
92. Halstensen TS, Scott H, Brandtzaeg P. Intraepithelial T cells of the TcR gamma/delta+ CD8- and V delta 1/J delta 1+ phenotypes are increased in coeliac disease. *Scand J Immunol* 1989;30(6):665-72.
93. Scott KG, Yu LC, Buret AG. Role of CD8+ and CD4+ T lymphocytes in jejunal mucosal injury during murine giardiasis. *Infect Immun* 2004;72(6):3536-42.
94. Madi K, Trajman A, da Silva CF, Barroso PF, Belo MT, Cunha JM, et al. Jejunal biopsy in HIV-infected patients. *J Acquir Immune Defic Syndr* 1991;4(10):930-7.
95. Anton PA, Mitsuyasu RT, Deeks SG, Scadden DT, Wagner B, Huang C, et al. Multiple measures of HIV burden in blood and tissue are correlated with each other but not with clinical parameters in aviremic subjects. *Aids* 2003;17(1):53-63.
96. Schneider T, Jahn HU, Schmidt W, Riecken EO, Zeitz M, Ullrich R. Loss of CD4 T lymphocytes in patients infected with human immunodeficiency virus type 1 is more pronounced in the duodenal mucosa than in the peripheral blood. Berlin Diarrhea/Wasting Syndrome Study Group. *Gut* 1995;37(4):524-9.
97. Meng G, Wei X, Wu X, Sellers MT, Decker JM, Moldoveanu Z, et al. Primary intestinal epithelial cells selectively transfer R5 HIV-1 to CCR5+ cells. *Nat Med* 2002;8(2):150-6.
98. Alfsen A, Iniguez P, Bouguyon E, Bomsel M. Secretory IgA specific for a conserved epitope on gp41 envelope glycoprotein inhibits epithelial transcytosis of HIV-1. *J Immunol* 2001;166(10):6257-65.
99. Zeitz M, Ullrich R, Schneider T, Kewenig S, Riecken E. Mucosal immunodeficiency in HIV/SIV infection. *Pathobiology* 1998;66(3-4):151-7.

100. Schneider T, Ullrich R, Bergs C, Schmidt W, Riecken EO, Zeitz M. Abnormalities in subset distribution, activation, and differentiation of T cells isolated from large intestine biopsies in HIV infection. The Berlin Diarrhoea/Wasting Syndrome Study Group. *Clin Exp Immunol* 1994;95(3):430-5.
101. Bull DM, Bookman MA. Isolation and functional characterization of human intestinal mucosal lymphoid cells. *J Clin Invest* 1977;59(5):966-74.
102. Raffael A, Nebe CT, G. V. Grundlagen der Durchflußzytometrie. In: Rothe G, editor. *Durchflußzytometrie in der Klinischen Diagnostik*. Stuttgart: Schattauer, 1994:3-48.
103. Ebert EC, Roberts AI. Lamina propria lymphocytes produce interferon-gamma and develop suppressor activity in response to lactoglobulin. *Dig Dis Sci* 2001;46(3):661-7.
104. Ostrowski MA, Justement SJ, Catanzaro A, Hallahan CA, Ehler LA, Mizell SB, et al. Expression of chemokine receptors CXCR4 and CCR5 in HIV-1-infected and uninfected individuals. *J Immunol* 1998;161(6):3195-201.
105. Lapenta C, Boirivant M, Marini M, Santini SM, Logozzi M, Viora M, et al. Human intestinal lamina propria lymphocytes are naturally permissive to HIV-1 infection. *Eur J Immunol* 1999;29(4):1202-8.
106. Meng G, Sellers MT, Mosteller-Barnum M, Rogers TS, Shaw GM, Smith PD. Lamina propria lymphocytes, not macrophages, express CCR5 and CXCR4 and are the likely target cell for human immunodeficiency virus type 1 in the intestinal mucosa. *J Infect Dis* 2000;182(3):785-91.
107. Lee B, Sharron M, Montaner LJ, Weissman D, Doms RW. Quantification of CD4, CCR5, and CXCR4 levels on lymphocyte subsets, dendritic cells, and differentially conditioned monocyte-derived macrophages. *Proc Natl Acad Sci U S A* 1999;96(9):5215-20.

108. Bermejo M, Martin-Serrano J, Oberlin E, Pedraza MA, Serrano A, Santiago B, et al. Activation of blood T lymphocytes down-regulates CXCR4 expression and interferes with propagation of X4 HIV strains. *Eur J Immunol* 1998;28(10):3192-204.
109. Peters B, Whittall T, Babaahmady K, Gray K, Vaughan R, Lehner T. Effect of heterosexual intercourse on mucosal alloimmunisation and resistance to HIV-1 infection. *Lancet* 2004;363(9408):518-24.
110. Zack JA, Arrigo SJ, Weitsman SR, Go AS, Haislip A, Chen IS. HIV-1 entry into quiescent primary lymphocytes: molecular analysis reveals a labile, latent viral structure. *Cell* 1990;61(2):213-22.
111. Lapham CK, Zaitseva MB, Lee S, Romanstseva T, Golding H. Fusion of monocytes and macrophages with HIV-1 correlates with biochemical properties of CXCR4 and CCR5. *Nat Med* 1999;5(3):303-8.
112. Mengozzi M, Malipatlolla M, De Rosa SC, Herzenberg LA, Roederer M. Naive CD4 T cells inhibit CD28-costimulated R5 HIV replication in memory CD4 T cells. *Proc Natl Acad Sci U S A* 2001;98(20):11644-9.
113. Zaitseva M, Blauvelt A, Lee S, Lapham CK, Klaus-Kovtun V, Mostowski H, et al. Expression and function of CCR5 and CXCR4 on human Langerhans cells and macrophages: implications for HIV primary infection. *Nat Med* 1997;3(12):1369-75.
114. Wilkin T, Su Z, Kuritzkes D, et al. Co-receptor tropism in patients screening for ACTG 5211, a phase 2 study of vicriviroc, a CCR5 inhibitor. *CROI 2006*. Denver, 2006.
115. Tachibana K, Hirota S, Iizasa H, Yoshida H, Kawabata K, Kataoka Y, et al. The chemokine receptor CXCR4 is essential for vascularization of the gastrointestinal tract. *Nature* 1998;393(6685):591-4.

116. Zou YR, Kottmann AH, Kuroda M, Taniuchi I, Littman DR. Function of the chemokine receptor CXCR4 in haematopoiesis and in cerebellar development. *Nature* 1998;393(6685):595-9.
117. Chen JD, Bai X, Yang AG, Cong Y, Chen SY. Inactivation of HIV-1 chemokine co-receptor CXCR-4 by a novel intrakine strategy. *Nat Med* 1997;3(10):1110-6.
118. Anderson J, Akkina R. HIV-1 resistance conferred by siRNA cosuppression of CXCR4 and CCR5 coreceptors by a bispecific lentiviral vector. *AIDS Res Ther* 2005;2(1):1.
119. Levine BL, Mosca JD, Riley JL, Carroll RG, Vahey MT, Jagodzinski LL, et al. Antiviral effect and ex vivo CD4+ T cell proliferation in HIV-positive patients as a result of CD28 costimulation. *Science* 1996;272(5270):1939-43.
120. Roederer M, Raju PA, Mitra DK, Herzenberg LA. HIV does not replicate in naive CD4 T cells stimulated with CD3/CD28. *J Clin Invest* 1997;99(7):1555-64.
121. Levine BL, Bernstein WB, Aronson NE, Schlienger K, Cotte J, Perfetto S, et al. Adoptive transfer of costimulated CD4+ T cells induces expansion of peripheral T cells and decreased CCR5 expression in HIV infection. *Nat Med* 2002;8(1):47-53.